



Divergent Gemycircularvirus in HIV-Positive Blood, France.

Rathviro Uch, Pierre-Edouard Fournier, Catherine Robert, Caroline Blanc-Tailleur, Vital Galicher, Romain Barre, François Jordier, Philippe de Micco, Didier Raoult, Philippe Biagini

► To cite this version:

Rathviro Uch, Pierre-Edouard Fournier, Catherine Robert, Caroline Blanc-Tailleur, Vital Galicher, et al.. Divergent Gemycircularvirus in HIV-Positive Blood, France.. *Emerging Infectious Diseases*, 2015, 21 (11), pp.2096-8. 10.3201/eid2111.150486 . hal-01235994

HAL Id: hal-01235994

<https://hal.science/hal-01235994>

Submitted on 31 May 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

9. Adenis AA, Aznar C, Couppié P. Histoplasmosis in HIV-infected patients: a review of new developments and remaining gaps. *Curr Trop Med Rep*. 2014;1:119–28.
10. Adenis A, Nacher M, Hanf M, Basurko C, Dufour J, Huber F, et al. Tuberculosis and histoplasmosis among human immunodeficiency virus–infected patients: a comparative study. *Am J Trop Med Hyg*. 2014;90:216–23. <http://dx.doi.org/10.4269/ajtmh.13-0084>

Address for correspondence: Christine E. Mandengue, Department of Internal Medicine (Dermatology), Université des Montagnes, Cliniques Universitaires des Montagnes, PO Box 208, Bangangté, Cameroon; email: cmmandengue@udesmontagnes.org

Divergent Gemycircularvirus in HIV-Positive Blood, France

Rathviro Uch, Pierre-Edouard Fournier, Catherine Robert, Caroline Blanc-Tailleur, Vital Galicher, Romain Barre, François Jordier, Philippe de Micco, Didier Raoult, Philippe Biagini

Author affiliations: Centre National de la Recherche Scientifique, Marseilles (R. Uch, P.-E. Fournier, C. Robert, C. Blanc-Tailleur, V. Galicher, R. Barre, F. Jordier, P. de Micco, D. Raoult, P. Biagini); Institut Hospitalo-Universitaire Méditerranée-Infection, Marseille, France (R. Uch, P.-E. Fournier, C. Robert, C. Blanc-Tailleur, R. Barre, F. Jordier, P. de Micco, D. Raoult, P. Biagini); Aix-Marseille Université, Marseille (R. Uch, P.-E. Fournier, C. Robert, C. Blanc-Tailleur, V. Galicher, R. Barre, F. Jordier, P. de Micco, D. Raoult, P. Biagini); Unité Mixte de Recherche 7268 ADES, Marseilles (R. Uch, V. Galicher, R. Barre, F. Jordier, P. de Micco, P. Biagini); Etablissement Français du Sang, Marseilles (R. Uch, V. Galicher, R. Barre, F. Jordier, P. de Micco, P. Biagini); Unité de Recherche sur les Maladies Infectieuses et Tropicales Emergentes, Marseilles (P.-E. Fournier, C. Robert, C. Blanc-Tailleur, D. Raoult)

DOI: <http://dx.doi.org/10.3201/eid2111.150486>

To the Editor: Gemycircularviruses are a group of recently discovered single-stranded DNA viruses, found initially in fungi in 2010 (*1*). These “myco-like” viruses have a genome ranging from 2.1 to 2.3 kb, containing 2 opposite open reading frames that probably code for a capsid protein (CP) and a spliced replication-associated protein (Rep). Related viruses have been subsequently identified in animal blood and fecal matter, raw and treated sewage, and insects and plant material, suggesting that gemycircularviruses may represent a large group of viruses exhibiting considerable genetic diversity (2–8). The presence of these viruses was recently extended to humans after gemycircularvirus sequences were identified in human blood and brain tissue


(multiple sclerosis patient), cerebrospinal fluid, and fecal matter (8,9).

While investigating the virome content of an HIV-positive blood donation, we identified several gemycircularvirus-related sequences. The initial metagenomic approach involved an HIV-1–positive plasma sample (B genotype, ≈530 copies/mL) obtained from the French blood agency national plasma bank in Tours, France. A 4-mL aliquot was prepared for metagenomic analysis after filtration, concentration, and nucleases treatment. Next, particle-protected nucleic acids were recovered and used for the preparation of a next-generation sequencing library and its subsequent analysis (online Technical Appendix, <http://wwwnc.cdc.gov/EID/article/21/11/15-0486-Techapp1.pdf>). Gemycircularvirus sequences identified among reads (1,680 vs. 82,560 reads total; ≈2%) were assembled into a resulting full-length sequence (GemyC1c) by using CodonCode Aligner version 5.1 (CodonCode Corporation, Centerville, MA, USA). This sequence was verified by using back-to-back specific primers, and the amplicon was cloned and sequenced according to the Sanger method.

The analysis of the GemyC1c sequence (2,109 nt, GenBank accession no. KP987887) revealed a genome divergent from those already available in databases, despite a similar genomic organization (Figure, panel A) and assignment to gemycircularviruses after BLAST (<http://blast.ncbi.nlm.nih.gov/Blast.cgi>) analysis of putative CP and Rep proteins. This divergence was demonstrated by the phylogenetic analysis of the deduced CP (Figure, panel B), which exhibited ≈72% and ≈44% aa pairwise identity with the 2 closest gemycircularvirus CP sequences available in GenBank (gemycircularvirus c from mongoose feces [Conceicao-Neto N., unpub. data] and HCBI8.215 from cattle blood, respectively). Moreover, GemyC1c CP exhibited ≈30% pairwise identity with viral sequences identified previously in humans (BZ1 from feces, SL1 from cerebrospinal fluid, MSS12.225 from blood). The deduced spliced Rep (major Rep1 and minor Rep2), seen in such viruses, contained putative rolling circle motifs I (LFTYS), II (HLHAFVD), and III (YATKD) retrieved from gemycircularviruses (4).

We subsequently investigated the presence of GemyC1c DNA in 128 HIV-positive plasma samples (French blood agency national plasma bank) along with 256 HIV-negative plasma samples (healthy blood donors, southeastern France; mean donor age 38 years; 136 men; 1 man:1.13 women). Plasma samples were prepared as described previously (*10*), and extracted nucleic acids were tested for GemyC1c DNA by using a specific PCR that included negative, positive, and extraction controls (online Technical Appendix).

Application of the above GemyC1c DNA detection system did not generate any positive signal in the 384 plasma samples in this study, suggesting that the presence



Gemycircularviruses are potentially very stable in the environment. Because an unknown part of this group is able to infect fungi, possible contamination from the laboratory environment or nucleic acid extraction methods must be considered. The fact that the same genome was never identified in other libraries generated in our laboratory supports the absence of local contamination; of note, we were also able to detect GemyC1c DNA with PCR by using another plasma aliquot extracted by an alternative method (NuLi-SENS magnetic extraction; bioMérieux, Marcy l'Étoile, France) In addition, the systematic elimination of the first

Our discovery of the GemyC1c by a sequence-independent molecular approach was informative for several reasons: 1) this viral sequence would have been undetectable by PCR according to the high genetic divergence existing between GemyC1c and other gemycircularviruses

identified; 2) this finding adds clues to the identification of potential new co-infections occurring in HIV-infected persons; and 3) this finding underlines the need to investigate the virome content of blood samples in a research context of new microbes as potential threats for transfusion. Further studies aimed at exploring genetic diversity and natural history of gemycircularviruses in human hosts are needed.

This work was supported by grant APR 2013.10 from the Établissement Français du Sang (Paris, France) and by the Méditerranée-Infection foundation.

References

1. Yu X, Li B, Fu Y, Jiang D, Ghabrial SA, Li G, et al. A geminivirus-related DNA mycovirus that confers hypovirulence to a plant pathogenic fungus. *Proc Natl Acad Sci U S A*. 2010;107:8387–92. <http://dx.doi.org/10.1073/pnas.0913535107>
2. Rosario K, Dayaram A, Marinov M, Ware J, Kraberger S, Staiton D, et al. Diverse circular ssDNA viruses discovered in dragonflies (Odonata: Epiprocta). *J Gen Virol*. 2012;93:2668–81. <http://dx.doi.org/10.1099/vir.0.045948-0>
3. Dayaram A, Opong A, Jaschke A, Hadfield J, Baschiera M, Dobson RC, et al. Molecular characterization of a novel cassava associated circular ssDNA virus. *Virus Res*. 2012;166:130–5. <http://dx.doi.org/10.1016/j.virusres.2012.03.009>
4. Sikorski A, Massaro M, Kraberger S, Young LM, Smalley D, Martin DP, et al. Novel myco-like DNA viruses discovered in the faecal matter of various animals. *Virus Res*. 2013;177:209–16. <http://dx.doi.org/10.1016/j.virusres.2013.08.008>
5. Ng TF, Chen LF, Zhou Y, Shapiro B, Stiller M, Heintzman PD, et al. Preservation of viral genomes in 700-y-old caribou feces from a subarctic ice patch. *Proc Natl Acad Sci U S A*. 2014;111:16842–7. <http://dx.doi.org/10.1073/pnas.1410429111>
6. Kraberger S, Argüello-Astorga GR, Greenfield LG, Galilee C, Law D, Martin DP, et al. Characterisation of a diverse range of circular replication-associated protein encoding DNA viruses recovered from a sewage treatment oxidation pond. *Infect Genet Evol*. 2015;31:73–86. <http://dx.doi.org/10.1016/j.meegid.2015.01.001>
7. Kraberger S, Farkas K, Bernardo P, Booker C, Argüello-Astorga GR, Mesléard F, et al. Identification of novel *Bromus*- and *Trifolium*-associated circular DNA viruses. *Arch Virol*. 2015;160:1303–11. <http://dx.doi.org/10.1007/s00705-015-2358-6>
8. Lamberto I, Gunst K, Müller H, zur Hauzen H, de Villiers EM. Mycovirus-like DNA virus sequences from cattle serum and human brain samples from multiple sclerosis patients. *Genome Announc*. 2014;2: pii: e00848–14. <http://dx.doi.org/10.1128/genomeA.00848-14>
9. Phan TG, Mori D, Deng X, Rajindrajith S, Ranawaka U, Fan Ng TF, et al. Small circular single stranded DNA viral genomes in unexplained cases of human encephalitis, diarrhea, and in untreated sewage. *Virology*. 2015;482:98–104. <http://dx.doi.org/10.1016/j.virol.2015.03.011>
10. Biagini P, Touinssi M, Galicher V, de Micco P. KIs virus and blood donors, France. *Emerg Infect Dis*. 2012;18:1374–5. <http://dx.doi.org/10.3201/eid1808.120442>

Address for correspondence: Philippe Biagini, Equipe “Emergence et co-évolution virale,” UMR 7268 ADES, Aix-Marseille Université, EFS, CNRS, 27 Blvd Jean Moulin, 13005 Marseille, France; email: philippe.biagini@efs.sante.fr

Co-infection with Drug-Susceptible and Reactivated Latent Multidrug-Resistant *Mycobacterium tuberculosis*

Laura Pérez-Lago, Miguel Martínez Lirola, Yurena Navarro, Marta Herranz, María Jesús Ruiz-Serrano, Emilio Bouza, Darío García-de-Viedma

Author affiliations: Gregorio Marañón University General Hospital, Madrid, Spain (L. Pérez-Lago, Y. Navarro, M. Herranz, M.J. Ruiz-Serrano, E. Bouza, D. García-de-Viedma); Gregorio Marañón Health Research Institute, Madrid (L. Pérez-Lago, Y. Navarro, M. Herranz, M.J. Ruiz-Serrano, E. Bouza, D. García-de-Viedma); Centro de Investigación Biomédica en Red en Enfermedades Respiratorias, Madrid (L. Pérez-Lago, Y. Navarro, M. Herranz, M.J. Ruiz-Serrano, E. Bouza, D. García-de-Viedma); Torrecárdenas Hospital, Almería, Spain (M. Martínez Lirola); Complutense University of Madrid, Spain (E. Bouza); Moncloa International Campus of Excellence, Complutense University-Politécnica University, Madrid (Y. Navarro, D. García-de-Viedma)

DOI: <http://dx.doi.org/10.3201/eid2111.150683>

To the Editor: Genotyping of *Mycobacterium tuberculosis* (MTB) has identified mixed infections involving >1 MTB strain (1–4), which are clinically relevant when different susceptibility patterns are involved (4–7). We describe a tuberculosis (TB) case-patient with mixed infection in an area of moderate incidence. In a low-resistance setting (monoresistance 4.6%; multiresistance 1.7%), 1 of the strains was drug susceptible and the other was multidrug-resistant (MDR). Molecular fingerprinting and epidemiologic research revealed that the infection corresponded to a recent infection by a susceptible strain and reactivation of an MDR TB strain. The patient was an HIV-negative woman, 47 years of age, who had immigrated to Spain from Romania and had been living in Almería for >3 years. TB was diagnosed in May 2014; she had experienced symptoms for 2 months. Her diagnosis was confirmed 3 years after being studied in Almería as a close contact of her husband, also from Romania, who had tested positive for MDR TB (resistant to rifampin and isoniazid). When she was observed in the contact trace, she tested positive for purified protein derivative, had been vaccinated against the *M. bovis* bacillus Calmette-Guérin strain, and had no radiologic findings or clinical symptoms. Based on the susceptibility profile of her husband, prophylaxis was not prescribed. Her husband adhered to anti-TB treatment for 20 months; all microbiological control test results had been negative since 2 months after starting therapy.