



HAL
open science

Separator Algebra for State Estimation

Luc Jaulin

► **To cite this version:**

Luc Jaulin. Separator Algebra for State Estimation. SMART 2015, Université de Manchester, Sep 2015, Manchester, United Kingdom. hal-01236498

HAL Id: hal-01236498

<https://hal.science/hal-01236498>

Submitted on 1 Dec 2015

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Separator Algebra for State Estimation

Luc Jaulin, ENSTA-Bretagne, LabSTICC.

1 Introduction

Consider the following state estimation problem [Jau15]

$$\begin{aligned} \text{(i)} \quad & \dot{\mathbf{x}}(t) = \mathbf{f}(\mathbf{x}(t)), t \in \mathbb{R} \\ \text{(ii)} \quad & \mathbf{g}(t_k) \in \mathbb{Y}(k), k \in \mathbb{N} \end{aligned} \quad (1)$$

Our objective is to find an inner and an outer approximation of the set $\mathbb{X}(t)$ of all state vectors that are consistent with (1) at time t . If we define by flow map φ_{t_1, t_2} as follows:

$$(\mathbf{x}(t_1) = \mathbf{x}_1 \text{ and } \dot{\mathbf{x}}(t) = \mathbf{f}(\mathbf{x}(t)) \Rightarrow \mathbf{x}_2 = \varphi_{t_1, t_2}(\mathbf{x}_1)). \quad (2)$$

The set of all *causal feasible states* at time t is defined by

$$\mathbb{X}(t) = \bigcap_{t_k \leq t} \varphi_{t_k, t} \circ \mathbf{g}^{-1}(\mathbb{Y}(k)). \quad (3)$$

In this paper, we show how it is possible to find both an inner and an outer approximations for $\mathbb{X}(t)$. Some existing methods are able to find an outer approximation [KJWM99] [GRMA13], but, to my knowledge, none of them is able to get an inner approximation. The main idea is to copy a classical contractor approach [CJ09] for state estimation, but to use separators [JD14] instead of contractors.

2 Separators

In this section, we present separators and show how they can be used by a paver in order to bracket the solution sets. An *interval* of \mathbb{R} is a closed connected set of \mathbb{R} . A box $[\mathbf{x}]$ of \mathbb{R}^n is the Cartesian product of n intervals. The set of all boxes of \mathbb{R}^n is denoted by $\mathbb{I}\mathbb{R}^n$. A *contractor* \mathcal{C} is an operator $\mathbb{I}\mathbb{R}^n \mapsto \mathbb{I}\mathbb{R}^n$ such that $\mathcal{C}([\mathbf{x}]) \subset [\mathbf{x}]$ and $[\mathbf{x}] \subset [\mathbf{y}] \Rightarrow \mathcal{C}([\mathbf{x}]) \subset \mathcal{C}([\mathbf{y}])$. A set \mathbb{S} is *consistent* with the contractor \mathcal{C} (we will write $\mathbb{S} \sim \mathcal{C}$) if for all $[\mathbf{x}]$, we have $\mathcal{C}([\mathbf{x}]) \cap \mathbb{S} = [\mathbf{x}] \cap \mathbb{S}$. A *separator* \mathcal{S} is pair of contractors $\{\mathcal{S}^{\text{in}}, \mathcal{S}^{\text{out}}\}$ such that, for all $[\mathbf{x}] \in \mathbb{I}\mathbb{R}^n$, we have $\mathcal{S}^{\text{in}}([\mathbf{x}]) \cup \mathcal{S}^{\text{out}}([\mathbf{x}]) = [\mathbf{x}]$. A set \mathbb{S} is *consistent* with the separator \mathcal{S} (we write $\mathbb{S} \sim \mathcal{S}$), if $\mathbb{S} \sim \mathcal{S}^{\text{out}}$ and $\overline{\mathbb{S}} \sim \mathcal{S}^{\text{in}}$, where $\overline{\mathbb{S}} = \{\mathbf{x} \mid \mathbf{x} \notin \mathbb{S}\}$. Using a separator inside a *paver* we can easily to classify part of the search space that are inside or outside a solution set \mathbb{S} associated with \mathcal{S} .

The algebra for separators is a direct extension of contractor algebra [CJ09]. If $\mathcal{S}_i = \{\mathcal{S}_i^{\text{in}}, \mathcal{S}_i^{\text{out}}\}$, $i \in \{1, 2\}$ are separators, we define

$$\begin{aligned} \mathcal{S}_1 \cap \mathcal{S}_2 &= \{\mathcal{S}_1^{\text{in}} \cup \mathcal{S}_2^{\text{in}}, \mathcal{S}_1^{\text{out}} \cap \mathcal{S}_2^{\text{out}}\} && \text{(intersection)} \\ \mathcal{S}_1 \cup \mathcal{S}_2 &= \{\mathcal{S}_1^{\text{in}} \cap \mathcal{S}_2^{\text{in}}, \mathcal{S}_1^{\text{out}} \cup \mathcal{S}_2^{\text{out}}\} && \text{(union)} \\ \mathbf{f}^{-1}(\mathcal{S}_1) &= \{\mathbf{f}^{-1}(\mathcal{S}_1^{\text{in}}), \mathbf{f}^{-1}(\mathcal{S}_1^{\text{out}})\} && \text{(inverse)} \end{aligned} \quad (4)$$

If \mathcal{S}_i are sets of \mathbb{R}^n , we have [JSD14]

$$\begin{aligned} \text{(i)} \quad & \mathcal{S}_1 \cap \mathcal{S}_2 \sim \mathcal{S}_1 \cap \mathcal{S}_2 \\ \text{(ii)} \quad & \mathcal{S}_1 \cup \mathcal{S}_2 \sim \mathcal{S}_1 \cup \mathcal{S}_2 \\ \text{(iii)} \quad & \mathbf{f}^{-1}(\mathcal{S}_1) \sim \mathbf{f}^{-1}(\mathcal{S}_1). \end{aligned} \tag{5}$$

Interval analysis [Moo66] [KK96] combined with contractors [CJ09] has been shown to be able to give an outer approximation of set. For the inner subpaving, the *De Morgan* rules make it possible to express the complementary set $\overline{\mathbb{X}}$ of \mathbb{X} . Then, basic contractor techniques can be used to get an inner characterization \mathbb{X}^- . Now, the task is not so easy and the role of *separators* is to make it automatic.

3 Transformation of separators

A transformation is an invertible function \mathbf{f} such as an analytical expression if known for both \mathbf{f} and \mathbf{f}^{-1} . The set of transformation from \mathbb{R}^n to \mathbb{R}^n is a group with respect to the composition \circ . Symmetries, translations, homotheties, rotations, ... are linear transformations.

Theorem. Consider a set \mathbb{X} and a transformation \mathbf{f} . Denote by $[\mathbf{f}]$ and $[\mathbf{f}^{-1}]$ two inclusion functions for \mathbf{f} and \mathbf{f}^{-1} . If $\mathcal{S}_{\mathbb{X}}$ is a separator for \mathbb{X} then a separator $\mathcal{S}_{\mathbb{Y}}$ for $\mathbb{Y} = \mathbf{f}(\mathbb{X})$ is

$$[\mathbf{y}] \rightarrow \{([\mathbf{f}] \circ \mathcal{S}_{\mathbb{X}}^{\text{in}} \circ [\mathbf{f}^{-1}])([\mathbf{y}] \cap [\mathbf{y}]), ([\mathbf{f}] \circ \mathcal{S}_{\mathbb{X}}^{\text{out}} \circ [\mathbf{f}^{-1}])([\mathbf{y}] \cap [\mathbf{y}])\} \tag{6}$$

or equivalently

$$\mathbf{f}(\mathbb{X}) \sim \{[\mathbf{f}] \circ \mathcal{S}_{\mathbb{X}}^{\text{in}} \circ [\mathbf{f}^{-1}] \cap \text{Id}, [\mathbf{f}] \circ \mathcal{S}_{\mathbb{X}}^{\text{out}} \circ [\mathbf{f}^{-1}] \cap \text{Id}\} \tag{7}$$

where *Id* is the identity contractor.

Remark. The separator defined by (6) corresponds to what we call the *transformation* of a separator by \mathbf{f} and we write $\mathcal{S}_{\mathbb{Y}} = \mathbf{f}(\mathcal{S}_{\mathbb{X}})$. As a consequence, thanks to the theorem, we can add to (5) the property

$$\text{(iv)} \quad \mathbf{f}(\mathbb{X}) \sim \mathbf{f}(\mathcal{S}_{\mathbb{X}}).$$

which will be used later for our state estimation problem.

Proof. The separator $\mathcal{S}_{\mathbb{Y}}$ is equivalent to $\mathbb{Y} = \mathbf{f}(\mathbb{X})$ if

$$\begin{cases} \mathcal{S}_{\mathbb{Y}}^{\text{out}}([\mathbf{y}] \cap \mathbb{Y}) = [\mathbf{y}] \cap \mathbb{Y} \\ \mathcal{S}_{\mathbb{Y}}^{\text{in}}([\mathbf{y}] \cap \overline{\mathbb{Y}}) = [\mathbf{y}] \cap \overline{\mathbb{Y}}. \end{cases} \tag{8}$$

Since $\mathcal{S}_{\mathbb{Y}}^{\text{out}}([\mathbf{y}]) \subset [\mathbf{y}]$ and $\mathcal{S}_{\mathbb{Y}}^{\text{in}}([\mathbf{y}]) \subset [\mathbf{y}]$, it suffices to prove that

$$\begin{cases} \text{(i)} \quad \mathcal{S}_{\mathbb{Y}}^{\text{out}}([\mathbf{y}]) \supset [\mathbf{y}] \cap \mathbb{Y} \\ \text{(ii)} \quad \mathcal{S}_{\mathbb{Y}}^{\text{in}}([\mathbf{y}]) \supset [\mathbf{y}] \cap \overline{\mathbb{Y}}. \end{cases} \tag{9}$$

Let us first prove (i). We have

$$\begin{aligned} [\mathbf{y}] \cap \mathbb{Y} &= \mathbf{f}(\mathbf{f}^{-1}([\mathbf{y}]) \cap \mathbf{f}^{-1}(\mathbb{Y})) & \mathbf{f} \text{ is bijective} \\ &= \mathbf{f}(\mathbf{f}^{-1}([\mathbf{y}]) \cap \mathbb{X}) & \mathbb{X} = \mathbf{f}^{-1}(\mathbb{Y}) \\ &\subset \mathbf{f}([\mathbf{f}^{-1}([\mathbf{y}]) \cap \mathbb{X}) & [\mathbf{f}^{-1}] \text{ is an inclusion function for } \mathbf{f}^{-1} \\ &\subset \mathbf{f}(\mathcal{S}_{\mathbb{X}}^{\text{out}}([\mathbf{f}^{-1}([\mathbf{y}])])) & \mathcal{S}_{\mathbb{X}}^{\text{out}} \text{ is a contractor for } \mathbb{X} \\ &\subset [\mathbf{f}] \circ \mathcal{S}_{\mathbb{X}}^{\text{out}} \circ [\mathbf{f}^{-1}]([\mathbf{y}]) & [\mathbf{f}] \text{ is an inclusion function for } \mathbf{f} \end{aligned} \tag{10}$$

Thus $[\mathbf{y}] \cap \mathbb{Y} \subset ([\mathbf{f}] \circ \mathcal{S}_{\mathbb{X}}^{\text{out}} \circ [\mathbf{f}^{-1}] ([\mathbf{y}]) \cap [\mathbf{y}]) = \mathcal{S}_{\mathbb{Y}}^{\text{out}}([\mathbf{y}])$. Let us now prove (ii). We have

$$\begin{aligned} [\mathbf{y}] \cap \overline{\mathbb{Y}} &= \mathbf{f} (\mathbf{f}^{-1} ([\mathbf{y}]) \cap \mathbf{f}^{-1} (\overline{\mathbb{Y}})) & \mathbf{f} \text{ is bijective} \\ &= \mathbf{f} (\mathbf{f}^{-1} ([\mathbf{y}]) \cap \overline{\mathbb{X}}) & \overline{\mathbb{X}} = \mathbf{f}^{-1} (\overline{\mathbb{Y}}) \\ &\subset \mathbf{f} ([\mathbf{f}^{-1}] ([\mathbf{y}]) \cap \overline{\mathbb{X}}) & [\mathbf{f}^{-1}] \text{ is an inclusion function for } \mathbf{f}^{-1} \\ &\subset \mathbf{f} (\mathcal{S}_{\overline{\mathbb{X}}}^{\text{in}} ([\mathbf{f}^{-1}] ([\mathbf{y}])) & \mathcal{S}_{\overline{\mathbb{X}}}^{\text{in}} \text{ is a contractor for } \overline{\mathbb{X}} \\ &\subset [\mathbf{f}] \circ \mathcal{S}_{\overline{\mathbb{X}}}^{\text{in}} \circ [\mathbf{f}^{-1}] ([\mathbf{y}]) & [\mathbf{f}] \text{ is an inclusion function for } \mathbf{f} \end{aligned} \quad (11)$$

Thus $[\mathbf{y}] \cap \mathbb{Y} \subset ([\mathbf{f}] \circ \mathcal{S}_{\overline{\mathbb{X}}}^{\text{out}} \circ [\mathbf{f}^{-1}] ([\mathbf{y}]) \cap [\mathbf{y}]) \cap \mathbb{Y} = \mathcal{S}_{\mathbb{Y}}^{\text{in}}([\mathbf{y}])$ which terminates the proof. ■

Example. Consider the constraint

$$\left\| \begin{pmatrix} 2 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} \cos \alpha & \sin \alpha \\ -\sin \alpha & \cos \alpha \end{pmatrix} \begin{pmatrix} y_1 - 1 \\ y_2 - 2 \end{pmatrix} \right\| \in [1, 3]. \quad (12)$$

If we apply an efficient forward-backward contractor in a paver, we get the contractions illustrated by the paving of Figure 1, left. Now, if we take

$$\begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} 2 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} \cos \alpha & \sin \alpha \\ -\sin \alpha & \cos \alpha \end{pmatrix} \begin{pmatrix} y_1 - 1 \\ y_2 - 2 \end{pmatrix} = \mathbf{f}^{-1}(\mathbf{y}) \quad (13)$$

or equivalently

$$\begin{pmatrix} y_1 \\ y_2 \end{pmatrix} = \begin{pmatrix} \cos \alpha & -\sin \alpha \\ \sin \alpha & \cos \alpha \end{pmatrix} \begin{pmatrix} \frac{1}{2} & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} + \begin{pmatrix} 1 \\ 2 \end{pmatrix} = \mathbf{f}(\mathbf{x}), \quad (14)$$

we get

$$\mathbf{y} = \mathbf{f}(\mathbf{x}), \text{ and } \|\mathbf{x}\| \in [1, 3]. \quad (15)$$

An optimal separator $\mathcal{S}_{\mathbb{X}}$ can be built for \mathbf{x} and the separator transform provides us a separator $\mathcal{S}_{\mathbb{Y}}$ for \mathbb{Y} . As illustrated by Figure 1, right, the resulting separator $\mathcal{S}_{\mathbb{Y}}$ is more efficient than the classical one based on forward-backward contractors. Note that in case we are not able to have an inner approximation for \mathbf{f}^{-1} , the problem of finding an inner approximation of the image of a set $\mathbf{f}(\mathbb{X})$ becomes much more difficult. See, *e.g.*, [VJVS05] [GJ10].

4 State estimation

If $\mathcal{S}_{\mathbb{X}(0)}$ is a separator for $\mathbb{X}(0)$ and if $\mathcal{S}_{\mathbb{Y}(k)}$ are separators for $\mathbb{Y}(k)$, then a separator for the set $\mathbb{X}(t)$ defined by (3) is

$$\mathcal{S}_{\mathbb{X}(t)} = \bigcap_{t_k \leq t} \varphi_{t_k, t} \circ \mathbf{g}^{-1}(\mathcal{S}_{\mathbb{Y}(k)}). \quad (16)$$

In this formula, $\mathbf{g}^{-1}(\mathcal{S}_{\mathbb{Y}(k)})$ is a separator. Due to the fact that $\varphi_{t_k, t}$ is bijective and that we are able to find an inclusion function for $\varphi_{t_k, t}$ and $\varphi_{t_k, t}^{-1}$ [RN11], the separator $\varphi_{t_k, t} \circ \mathbf{g}^{-1}(\mathcal{S}_{\mathbb{Y}(k)})$ is clearly defined using the separator transform. To illustrate the method, let us consider a robot described by

$$\begin{cases} \dot{\mathbf{x}}(t) = \begin{pmatrix} v(t) \cos \theta(t) \\ v(t) \sin \theta(t) \end{pmatrix} & \text{(evolution)} \\ \|\mathbf{x}(t_k)\| \in y(t_k) + [-0.3, 0.3], t_k = 0.1 \cdot k, k \in \mathbb{N} & \text{(observation)} \end{cases} \quad (17)$$




Fig. 1. Left. Contractions obtained using a classical forward-backward propagation; Right. Contractions obtained using the separator transform. The frame corresponds to the box $[-6, 6]^2$.

where $v(t)$ and $\theta(t)$ are measured with an accuracy of ± 0.03 . The observation equation is due to the fact that the robot measures every 0.1 sec its distance to the origin with an accuracy of ± 0.3 . The actual (but unknown) trajectory for the robot is

$$\mathbf{x}(t) = \begin{pmatrix} 2 + 3 \cos t \\ 2 \sin t \end{pmatrix}. \quad (18)$$

For $t \in 0.2 * k$, $k = 0, \dots, 7$, the sets $\mathbb{X}(t)$ obtained by our observer are represented on Figure 2. Black boxes are inside $\mathbb{X}(t)$, grey boxes are outside and the white boxes cover the boundary. For $t = 0$, $\mathbb{X}(t)$ is a ring which becomes a small set for $t = 1.4$ once the robot has moved sufficiently. The fact that the white area covering the boundary becomes thick is mainly due to the state errors inside the evolution equation.

References

- [CJ09] G. CHABERT AND L. JAULIN. Contractor Programming. *Artificial Intelligence* **173**, 1079–1100 (2009).
- [GJ10] A. GOLDSZTEJN AND L. JAULIN. Inner Approximation of the Range of Vector-Valued Functions. *Reliable Computing* pages 1–23 (2010).
- [GRMA13] A. GNING, B. RISTIC, L. MIHAYLOVA, AND F. ABDALLAH. An Introduction to Box Particle Filtering. *IEEE Signal Processing Magazine* **30**(1), 166–171 (2013).
- [Jau15] L. JAULIN. “Automation for Robotics”. ISTE editions (2015).
- [JD14] L. JAULIN AND B. DESROCHERS. Separators: a new interval tool to bracket solution sets; application to path planning. *Engineering Applications of Artificial Intelligence* **33**, 141–147 (2014).
- [JSD14] L. JAULIN, A. STANCU, AND B. DESROCHERS. Inner and outer approximations of probabilistic sets. In “ICVRAM 2014” (2014).
- [KJWM99] M. KIEFFER, L. JAULIN, E. WALTER, AND D. MEIZEL. Guaranteed mobile robot tracking using interval analysis. In “Proceedings of the MISC’99 Workshop on Applications of Interval Analysis to Systems and Control”, pages 347–359, Girona, Spain (1999).




Fig. 2. Inner and outer approximations of the set of all feasible state vectors $\mathbb{X}(t)$, for $t \in 0, 0.2, \dots, 1.4$. The frame boxes are $[-6, 6]^2$.

- [KK96] R. B. KEARFOTT AND V. KREINOVICH, editors. “Applications of Interval Computations”. Kluwer, Dordrecht, the Netherlands (1996).
- [Moo66] R. E. MOORE. “Interval Analysis”. Prentice-Hall, Englewood Cliffs, NJ (1966).
- [RN11] N. RAMDANI AND N. NEDIALKOV. Computing Reachable Sets for Uncertain Nonlinear Hybrid Systems using Interval Constraint Propagation Techniques. *Nonlinear Analysis: Hybrid Systems* **5**(2), 149–162 (2011).
- [VJVS05] P. HERRERO VINAS, L. JAULIN, J. VEHI, AND M. A. SAINZ. Inner and outer approximation of the polar diagram of a sailboat. In “workshop on Interval Analysis and Constraint Propagation for Applications (IntCP’05)”, Barcelona, Spain (2005).