



A radiative transfer based approach to merge SMOS and AMSR-e soil moisture retrievals into one consistent record

Robin van Der Schalie, Richard de Jeu, Yann H. Kerr, Jean-Pierre Wigneron, Nemesio Rodriguez-fernandez, Amen Al Yaari, Matthias Drusch, Susanne Mecklenburg, Han Dolman

► To cite this version:

Robin van Der Schalie, Richard de Jeu, Yann H. Kerr, Jean-Pierre Wigneron, Nemesio Rodriguez-fernandez, et al.. A radiative transfer based approach to merge SMOS and AMSR-e soil moisture retrievals into one consistent record. ESA-ESRIN Earth Observation for Water Cycle Science 2015, Oct 2015, Frascati, Italy. 2015. <hal-01237187>

HAL Id: hal-01237187

<https://hal.science/hal-01237187v1>

Submitted on 2 Dec 2015

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

A RADIATIVE TRANSFER BASED APPROACH TO MERGE SMOS AND AMSR-E SOIL MOISTURE INTO ONE CONSISTENT RECORD

Robin van der Schalie (r.vander.Schalie@vu.nl)^{1,2}, Richard de Jeu², Yann Kerr³, Jean Pierre Wigneron⁴, Nemesio Rodriguez Fernandez³, Amen Al-Yaari⁴, Matthias Drusch⁵, and Susanne Mecklenburg⁶

¹VU University Amsterdam NL; ²Transmissivity B.V. NL; ³CESBIO France; ⁴INRA France; ⁵ESA-ESTEC NL; ⁶ESA-ESRIN Italy

Abstract—Recently, a study funded by the European Space Agency (ESA), the SMOS Fusion project, was set up to provide guidelines for the development of a global soil moisture climate record with a special emphasis on the integration of SMOS. Three different data fusion approaches were designed and implemented on 10 year passive microwave data (2003-2013) from two different satellite sensors; the ESA Soil Moisture Ocean Salinity Mission (SMOS) and the NASA/JAXA Advanced Scanning Microwave Radiometer (AMSRE).

This study is part of the Fusion project and investigates a radiative transfer based approach to merge AMSRE and SMOS soil moisture retrievals using the Land Parameter Retrieval Model (LPRM). LPRM has already been successfully applied to SMOS observations in earlier studies and now we will focus on the merging strategy with AMSR-E. The first step in this project is a small update on the roughness parameterization in the SMOS LPRM. Then the merging of the AMSRE and SMOS LPRM is done by optimizing the AMSRE LPRM to best match the SMOS LPRM in soil moisture dynamics followed by a linear merging of the two datasets. The resulting AMSRE LPRM soil moisture retrievals are compared against ERA-Land, MERRA-Land and the International Soil Moisture Network (ISMN).

2

Roughness update

The new empirical method to estimate the roughness contains a correction for vegetation influences without using external sources of information, by first calculating SMOS LPRM as in the previous version, and using its retrieved Vegetation Optical Depth in the second run to correct for the vegetation influences. This has minimal effect on the high correlations as gotten with the previous version and improves the absolute values. Mean retrievals shown in Fig 1.

New roughness formula: $h = h_1 (A_v(1 - 2\theta) + B_v \tau_v)$




Fig. 1: Mean SMOS LPRM soil moisture after the improved roughness parameterization over the period of July 2010 to December 2013

3

Updated AMSRE LPRM

The AMSRE LPRM parameterization was updated for both the C-band and X-band frequencies and, in contrast to what it used to be, is now the same for both C- as X-band. The update includes a new roughness parameterization very similar to the one applied to SMOS LPRM and this is combined with an increase of the single scattering albedo from 0.05 and 0.06 to 0.075. These parameters were found during several optimization runs. This parameterization resulted in very high correlations between the AMSRE and SMOS LPRM products as can be seen in Fig 2. The updated soil moisture retrievals from AMSRE show very similar dynamics as that of SMOS LPRM over all areas that do not have dense vegetation or sandy deserts.




Fig. 2: Correlation coefficient of AMSRE LPRM vs SMOS LPRM using the updated parameterization




Fig. 3: Correlations of the updated C-band LPRM against MERRA-Land and ERA-Land

The long term dataset of AMSRE LPRM was tested for the period of 2003 to 2009 and compared against MERRA-Land and ERA-Land. Fig 3 shows the correlations that the new C-band AMSRE LPRM reaches against these two products. The results again show an overall good performance (>0.6) over all areas except for the densely vegetated, sandy desert and the boreal areas. Compared to the old method, the main places of improvement were the eastern part of the USA and Europe. For the unbiased root mean square error, the results are slightly improved compared to the old version, with an ubmse < 0.10 everywhere except for the boreal areas. This is expected due to difficulties defining the effects and filtering of permafrost, snow, open water and vegetation in these areas.

In Fig 4 you see the performance of C-band AMSRE LPRM against the available 560 in situ stations of the ISMN. The mean r of 0.49, ubmse of 0.084 and bias of -0.02 are all a slight improvement compared to the old LPRM version.

X-band retrievals against MERRA-Land, ERA-Land and the ISMN show a very similar behavior, only with a slightly higher noise level and quicker saturation over densely vegetated areas due to the higher frequency.




Fig. 4: Histograms of the comparison between C-band AMSRE LPRM and all the available ground observations from the international soil moisture network.

4

Conclusions

The results of this study show that the new roughness update for SMOS LPRM leads to a more natural distribution of soil moisture worldwide, while it keeps a similar high level of performance as before.

The update of AMSRE LPRM shows that the LPRM performs very similar for the different frequencies, leading to soil moisture retrievals that correlate very well with each other, modelled soil moisture from MERRA-Land and ERA-land and in situ measurements from the ISMN.

Acknowledgements

This project is funded by the ESA project "Passive Microwave Soil Moisture Data Fusion Study" (Contract # 4000110112).