

Interactions between polyphenols and polysaccharides: mechanisms and consequences in food processing and digestion


Catherine M.G.C. Renard, Aude A. Watrelot &
Carine Le Bourvellec



UMR408 Sécurité et Qualité des Produits d'Origine Végétale,
INRA, Université d'Avignon et des Pays du Vaucluse

EFFoST 2015
9-11 november 2015, Athens (Greece)





Why attempt to understand these interactions?

↖ Extraction of polyphenols

- Transfer from fruit to juice (wine, cider) or extraction to tea, coffee, chocolate...
- Major impact on colour and bitterness, astringency, « body »
- Need to understand the mechanisms for extraction vs. retention

↖ Health effects:

- Antioxidants: prevention of oxidation in the stomach
- Specific actions - interaction with receptors
 - Must be bioavailable - phenols or metabolites
 - Relevance of colonic fermentation metabolites


Free polyphenol → bioaccessible → None or poor bioavailability

Bound polyphenol → Not bioaccessible → Fermentation



Bioavailable metabolites





↖ **Introduction**

- Polyphenols & cell walls

↖ **Quantifying the interactions**


- Binding isotherms

↖ **An example of consequences**

- Modification of cider composition



↖ **Conclusion & perspectives**


- Colonic fermentation



Cell walls

- ↖ Surround the plant cells
- ↖ Structural in the cell walls
 - Protection and structure
- ↖ A complex assembly of macromolecules
 - Mainly polysaccharides
 - Cellulose microfibrils
 - Pectins
 - Hemicellulose
 - Minor amounts of glycoproteins
 - Polyphenols (FeA, lignin...)







Proanthocyanidins

↳ Proanthocyanidins =
« condensed tannins »

- Flavanol oligo and polymers
- Intracellular
- Characterized by:
 - Constitutive unit
 - Interflavanol binding
 - Degree of polymerization
- Abundant in apple and pear: procyanidins
 - Epicatechin (catechin)
 - B-type, primarily 4-8 binding
 - DP usually 3-8, up to a 100
 - Concentrations up to 5 g/ kg







Quantifying interactions by binding isotherms

↳ Model suspensions

Purified procyanidins




1-Incubation
2-Filtration



Bound procyanidins

- Freeze-dry and thiolysis




Free procyanidins

- OD 280 nm
- Freeze-dry and thiolysis

Cell walls
Devoid of phenolics

↳ What can be compared:

- Physico-chemical conditions (pH, T°, ionic strength, solvent...)
- Proanthocyanidins structure and size
- Cell walls: origin and pretreatment, drying...




Procyanidins adsorb spontaneously to cell walls


↳ Binding isotherms

$$PPf = \frac{N_{\max} \cdot K_L \cdot [PP_L]}{1 + K_L \cdot [PP_L]}$$

↳ Binding

- H-bonds and hydrophobic interactions
- Totally reversible using appropriate solvent (not water!)
- Binding increases with DP of procyanidins
- Amounts adsorbed vary with initial concentrations, cell wall mode of production and physico-chemical conditions
- High affinity for pectins, no impact of protein





Modifying cell wall affinity for procyanidins


↖ Extraction of pectins

- Chemical extraction or heat-treatment
- Decreased affinity

↖ Modifying surface properties by drying

- Mild drying (solvent-exchange) vs harsh drying

- Increase specific area and decrease porosity
- Decrease affinity and increase max absorption



Polyphenols and cider

↖ Organoleptic characters


- Colour, bitterness and astringency

↖ Evolution of phenolic composition during processing


- Polyphenol extraction from the fruit:
- Modification of structures (oxidation)


↖ Experiments with isolated procyanidins indicate an effect of temperature

- Extraction at different temperatures
- Apples with different average degrees of polymerisation
- Different levels of oxidation :
 - No oxidation for impact of non-covalent interaction



Phenolic composition changes from apple to juice





Extraction of procyanidins does vary with pressing temperature




Experiment:




- Three varieties
 - DM: DP = 3, GU: DP = 63, KE: DP = 3.5
- Four temperatures
 - 4°C, 11°C, 18°C, 25°C
- Inerted press and juice with NaF

Concentration (mg/L)



Extraction variation with temperature is validated

The model designed from experiments with isolated cell walls and procyanidins gives satisfactory previsions



Conclusions and perspectives

↖ Procyanidins bind spontaneously to cell walls

- This is very common artefact:
 - Procyanidins in all Rosaceae...
- And can be an artefact in cell wall preparation

↖ Consequences


- In food processing and for procyanidins health impact

↖ What about heating ?

- Evolution of non-covalent adducts ?
- Formation of covalent adducts ?

↖ Fate of non-covalent adducts in the gut ?

- Impact of non-covalent adducts vs procyanidins alone :
 - Short fatty acid production
 - Procyanidin colonic metabolites production
 - Microbiota



The main contributors

↳ Unité de Recherches Cidriques - Rennes

- A. Baron,
- S. Guyot,
- JM Le Queré,
- C. Le Bourvellec

↳ UMR SQPOV - Avignon

- C. Le Bourvellec
- A. Watrelot
- M. Kebe
- JF Maingonnat

↳ ESA Ronan Symoneaux


↳ CERMAV Anne Imberty

↳ U. Bordeaux S. Quideaux

↳ Pôle Agronomique Ouest

« Cidre et Polyphénols » 2000-2004

↳ FP6 FLAVO and ISAFRUIT




↳ ANR TEMPANTIOX

Agence Nationale de la Recherche



↳ FP7 DREAM



Thank you for your attention
Looking forward to welcoming you soon at:



2nd Euro-Mediterranean Symposium on Fruit and Vegetable Processing

An integrated view on sustainability and how it can interact with quality, safety and the consumers

Avignon, 4-6 April 2016

For more info:

<https://colloque.inra.fr/fruitvegprocessing2016>

