

Shapely monads and analytic functors

Richard Garner, Tom Hirschowitz

▶ To cite this version:

Richard Garner, Tom Hirschowitz. Shapely monads and analytic functors. 2015. hal-01246365v1

HAL Id: hal-01246365 https://hal.science/hal-01246365v1

Preprint submitted on 18 Dec 2015 (v1), last revised 10 Oct 2017 (v3)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

SHAPELY MONADS AND ANALYTIC FUNCTORS

RICHARD GARNER AND TOM HIRSCHOWITZ

1. INTRODUCTION

In mathematics and computer science, we often encounter structures which are faithfully encoded by a graphical calculus of the following sort. The basic data of the structure are depicted as certain diagrams; the basic operations of the structure act by glueing together these diagrams along certain parts of their boundaries; and the axioms of the structure are just those necessary to ensure that "every two ways of glueing a compound diagram together agree".

Commonly, such calculi depict structures wherein "functions", "arrows" or "processes" are wired together along input or output "ports". For instance, we have *multicategories* [21], whose arrows have many inputs but only one output; *polycategories* [29], whose arrows have multiple inputs and outputs, with composition subject to a linear wiring discipline; and *coloured properads* [31] and PROPS [24], which are like polycategories but allow for non-linear wirings.

Mathematical structures such as these are important in algebraic topology and homological algebra—encoding, for example, operations arising on infinite loop spaces [26] or on Hochschild cochains [27]—but also in logic and computer science. For example, polycategories encode the underlying semantics of a linear sequent calculus [22], while PROPs have recently been used as an algebraic foundation for notions of computational network such as signal flow graphs [3] and Bayesian networks [10]. Other kinds of graphical structures arising in computer science include *proof nets* [11, §2], *interaction nets* [20] and *bigraphs* [14].

There is an established approach to describing structures of the above kind using monads on presheaf categories. The presheaf category captures the essential topology of the underlying graphical calculus, while the monad encodes both the wiring operations of the structure and the axioms that they obey; the algebras for the monad are the instances of the structure. One aspect which this approach does not account for is that the axioms should be determined by the requirement that "every two ways of wiring a compound diagram together agree". The first main contribution of this paper is to rectify this: we explain the observed form of the axioms as a property of the associated monad—which we term *shapeliness*—stating that "every two operations of the same shape coincide".

Hirschowitz acknowledges the support of the French ANR projet blanc Récré ANR-11-BS02-0010; Garner acknowledges the support of the Australian Research Council Discovery Projects DP110102360 and DP130101969. This research was partially undertaken during a visit by Hirschowitz to Sydney funded by Macquarie University.

In fact, shapeliness gives not just a way of *characterising* the monads encoding graphical structures, but also a systematic way of *generating* them. For a given graphical calculus, it is typically easy to find a presheaf category encoding the basic diagram-shapes of the calculus, and an endofunctor thereon encoding the basic wiring operations; we then obtain the desired monad as the *free shapely* monad on the given endofunctor. The algebras for this monad can be seen as denotational models of the graphical calculus in question; and though we do not do this here, one can envisage this being used to attach workable denotational semantics to, for example, interaction nets; the syntactic part of bigraphs; or MLL proof nets without units.

Formalising the notion of shapely monad turns out to be a delicate task. In the end, we will define a monad on $\mathscr{PC} = [\mathscr{C}^{\mathrm{op}}, \mathbf{Set}]$ to be *shapely* just when it is a submonad of a *universal* shapely monad U with "exactly one operation of each shape". We will find U by seeking a terminal object in a suitable monoidal category of *endofunctors* of \mathscr{PC} ; once found, terminality will automatically endow this object with a monad structure, so giving the desired U.

This leaves the problem of choosing a suitable monoidal category of endofunctors. An obvious but wrong choice would be the whole functor category $[\mathscr{PC}, \mathscr{PC}]$: with this choice, U would be the monad constant at 1, and a general monad would be shapely just when it took values in subobjects of 1. This is manifestly not what we want; the problem is that terminality in the full functor category encodes the property of having "exactly one operation of each shape" for what are overly crude notions of "operation" and "operation shape".

Refining these notions, as we shall do, means looking for a terminal object in some smaller category of endofunctors of \mathscr{PC} . Choosing this category turns out to be an interesting design problem: some natural candidates have a terminal object, but are not closed under composition, while others are closed under composition, but fail to have a terminal object. Our eventual solution will triangulate between these failures, but we make no claims to its definitiveness; in fact, we consider the value of our work to lie as much in the exploration of the problem's design space as in the particular solution we adopt.

The technical foundation of our approach will be a theory of *analytic functors* between presheaf categories, and the second main contribution of this paper is to develop this theory. Analytic endofunctors of **Set** were introduced by Joyal in [16] as a categorical setting for constructions in enumerative combinatorics; their importance for computer science has been recognised in work such as [1, 9, 12]. An endofunctor F of **Set** is *analytic* when it can be written in the form:

$$FX = \sum_{i \in I} X^{\alpha_i} / G_i$$

for an *I*-indexed family of natural numbers α_i and subgroups $G_i \leq \mathfrak{S}_{\alpha_i}$; the quotients are by the permutation actions of G_i on the factors of X^{α_i} . Generalising this, we will call a functor $F: \mathscr{PD} \to \mathscr{PC}$ analytic if it takes the form:

(1.1)
$$FX(c) = \sum_{i \in I_c} \mathscr{P}\mathscr{D}(\alpha_i, X)_{/G_i}$$

 $\mathbf{2}$

for some family of presheaves $\alpha_i \in \mathscr{PD}$ and subgroups $G_i \leq \operatorname{Aut}(\alpha_i)$ of the automorphism group of each α_i . Just as in the case studied in [16], these generalised analytic functors have a strongly combinatorial flavour; moreover, all of the monads derived from graphical calculi that we will consider turn out to be analytic in this sense¹. It is therefore reasonable that we should look for a universal shapely monad among analytic endofunctors of a presheaf category. In the end, we are able to produce such a universal monad only in certain special cases; but along the way, we develop various results of independent interest, for example:

- (i) We give a combinatorial representation of the category of finitary analytic functors and transformations $\mathscr{PC} \to \mathscr{PD}$ (Proposition 4.16) and show that this category always has a terminal object (Proposition 4.20).
- (ii) We show that analytic functors between presheaf categories need *not* be closed under composition (Proposition 4.23); this is by contrast to analytic endofunctors of **Set**, which *are* composition-closed.
- (iii) We introduce a condition on analytic functors which we call *cellularity*, that is sufficient to ensure that they do compose (Proposition 5.15).
- (iv) We see that, unfortunately, the introduction of cellularity also destroys the terminal object among finitary analytic endofunctors (Proposition 5.20).

We conclude this introduction with a brief overview of the rest of the paper. We start in Section 2 by developing some motivating examples of structures expressible by graphical calculi: we discuss the *polycategories* of [28] and the richer structures of (coloured) *properads* [31] and PROPS [24], and in each case show how the structure at issue can be described as the algebras for a monad on a presheaf category. In Section 3, we begin our pursuit of the notion of universal shapely monad. We do not immediately consider the analytic functors discussed above, but rather the narrower class of *familially representable* or *familial* functors; these were introduced by Diers [7] and studied by *inter alia* Johnstone, Leinster and Weber [6, 33, 23], and are precisely the analytic functors whose expression (1.1) involves only *trivial* groups G_i . We recall basic aspects of the theory of familial functors, including closure under composition, but show that there is usually no terminal object among familial endofunctors, and hence no universal shapely monad among them.

In Section 4, we attempt to fix up the lack of a terminal object among familial endofunctors by passing to the more general analytic functors. As is visible from (i) and (ii) above, we succeed in doing this, but only at the cost of breaking the composability of familial endofunctors. Section 5 attempts to fix this new problem by introducing the more restricted class of *cellular* analytic functors; as in (iii) and (iv) above, this does indeed resolve the problem of composability but at the same time reintroduces the problem of the existence of a terminal object.

At this point, in Section 6, we declare ourselves unable to find a further refinement of the notion of cellularity that, in full generality, fixes both composability

¹This is why we do not make use of the various other generalisations of analyticity to presheaf categories in [1, 9, 8]; none of these generalisations can express the monads associated to our graphical examples.

and existence of a terminal object. However, in the presheaf categories relevant to the motivating examples of Section 2, we are able to impose an additional *ad hoc* condition on top of cellularity which is sufficient to ensure that the cellular functors in this class both compose *and* admit a terminal object: using this, we finally obtain the desired notion of shapely monad, and are able to exhibit the monads encoding the graphical structures of interest as free shapely monads on the basic wiring operations of the structure.

2. Motivating examples

Before developing our general theory of shapeliness, we describe in detail some of the examples we would like to capture, so that we can see as we go along how they fit into the framework. Our main example will be the *polycategories* of [28], though we also discuss *properads* [31] and PROPS [24]. In describing these examples, we first give an elementary presentation of the structure, and then show how it can be encoded by a suitable monad on a presheaf category.

2.1. **Polycategories.** Polycategories were introduced in [28] as a semantic model for a two-sided propositional sequent calculus; although originally used to model the classical Genzten calculus, it later became clear [22] that they are precisely the structure underlying the multiplicative fragment of linear logic.

Definition 2.1. A small (symmetric) polycategory \mathscr{C} comprises a set $ob(\mathscr{C})$ of *objects*; sets $\mathscr{C}(\vec{A}; \vec{B})$ of *morphisms* for each pair of lists $\vec{A} = (A_1, \ldots, A_n)$ and $\vec{B} = (B_1, \ldots, B_m)$ of objects; and the following further data:

- *Identity* morphisms $id_A \in \mathscr{C}(A; A)$ for each object A.
- Composition operations giving for each $f \in \mathscr{C}(\vec{A}; \vec{B})$ and $g \in \mathscr{C}(\vec{C}; \vec{D})$ and indices i, j with $B_i = C_j$, a morphism

$$g_{j} \circ_{i} f \in \mathscr{C}(\vec{C}_{< j}, \vec{A}, \vec{C}_{> j}; \vec{B}_{< i}, \vec{D}, \vec{B}_{> i}) ,$$

here we use comma to denote concatenation of lists, and write $C_{<j}$ for the list (C_1, \ldots, C_{j-1}) , and so on.

• Exchange operations giving for each $f \in \mathscr{C}(\vec{A}; \vec{B})$ and permutations $\varphi \in \mathfrak{S}_n$ and $\psi \in \mathfrak{S}_m$ an element

$$\psi \cdot f \cdot \varphi \in \mathscr{C}(A_{\varphi}; B_{\psi^{-1}})$$

where \vec{A}_{φ} denotes the list $(A_{\varphi(1)}, \ldots, A_{\varphi(n)})$ and likewise for $\vec{B}_{\psi^{-1}}$.

These data are required to satisfy various axioms, which we give in detail mainly to show how unpalatable they are when presented algebraically, and without any serious expectation that the reader should work through the details.

- The unit axioms $f_{i} \circ_{1} \operatorname{id}_{A_{i}} = f = \operatorname{id}_{B_{j-1}} \circ_{j} f$ for all $f \in \mathscr{C}(\vec{A}; \vec{B})$ and valid indices i, j.
- The associativity axiom for all $f \in \mathscr{C}(\vec{A}; \vec{B}), g \in \mathscr{C}(\vec{C}; \vec{D})$ and $h \in \mathscr{C}(\vec{E}; \vec{F})$ and indices i, j, k, ℓ with $B_i = C_j$ and $D_k = E_\ell$:

$$(h_{\ell} \circ_k g)_{i+\ell-1} \circ_i f = h_{\ell} \circ_{k+i-1} (g_{i} \circ_i f).$$

• The left interchange axiom for all $f \in \mathscr{C}(\vec{A}; \vec{B}), g \in \mathscr{C}(\vec{C}; \vec{D})$ and $h \in \mathscr{C}(\vec{E}; \vec{F})$ and indices i, j and $k_1 < k_2$ such that $B_i = E_{k_1}$ and $D_j = E_{k_2}$:

$$(h_{k_2} \circ_j g)_{k_1} \circ_i f = \psi \cdot ((h_{k_1} \circ_i f)_{k_2 + n - 1} \circ_j g) .$$

Here, n is the length of the list \vec{A} and ψ is the evident permutation for which $(\vec{B}_{<i}, \vec{D}_{<j}, \vec{F}, \vec{D}_{>j}, \vec{B}_{>i})_{\psi} = (\vec{D}_{<j}, \vec{B}_{<i}, \vec{F}, \vec{B}_{>i}, \vec{D}_{>j}).$

• The right interchange axiom for all f, g and h as above and indices $i_1 < i_2$ and j, k such that $B_{i_1} = C_j$ and $B_{i_2} = E_k$:

$$g_{j} \circ_{i_{1}} (h_{k} \circ_{i_{2}} f) = (h_{k} \circ_{i_{2}+m-1} (g_{j} \circ_{i_{1}} f)) \cdot \varphi .$$

This time, m is the length of the list \vec{D} and φ is the evident permutation for which $(\vec{E}_{< k}, \vec{C}_{< j}, \vec{A}, \vec{C}_{> j}, \vec{E}_{> k})_{\varphi} = (\vec{C}_{< j}, \vec{E}_{< k}, \vec{A}, \vec{E}_{> k}, \vec{C}_{> j}).$

- The usual action axioms for the exchange operation: $\mathrm{id}_m \cdot f \cdot \mathrm{id}_n = f$ and $(\psi_2 \psi_1) \cdot f \cdot (\varphi_1 \varphi_2) = \psi_2 \cdot (\psi_1 \cdot f \cdot \varphi_1) \cdot \varphi_2.$
- Compatibility of exchange and composition for all $f \in \mathscr{C}(\vec{A}; \vec{B}), g \in \mathscr{C}(\vec{C}; \vec{D})$, and suitable permutations $\varphi_1, \varphi_2, \psi_1$ and ψ_2 :

$$(\psi_2 \cdot g \cdot \varphi_2)_j \circ_i (\psi_1 \cdot f \cdot \varphi_1) = \bar{\psi} \cdot (g_{\varphi_2(j)} \circ_{\psi_1^{-1}(i)} f) \cdot \bar{\varphi}$$

where $\bar{\varphi}$ is determined by $(\vec{C}_{<\varphi_2(j)}, \vec{A}, \vec{C}_{>\varphi_2(j)})_{\bar{\varphi}} = ((\vec{C}_{\varphi_2})_{<j}, \vec{A}_{\varphi_1}, (\vec{C}_{\varphi_2})_{>j})$ and $\bar{\psi}$ by $(\vec{B}_{<\psi_1^{-1}(i)}, \vec{D}, \vec{B}_{>\psi_1^{-1}(i)})_{\bar{\psi}^{-1}} = ((\vec{B}_{\psi_1^{-1}})_{<i}, \vec{D}_{\psi_2^{-1}}, (\vec{B}_{\psi_1^{-1}})_{>i}).$

We write **Polycat** for the category of small polycategories and the structurepreserving maps between them.

The definition of polycategory is much more easily parsed by way of the following graphical presentation: a morphism in $\mathscr{C}(A_1, \ldots, A_n; B_1, \ldots, B_m)$ is depicted as a box with *n* wires going into it and *m* going out:



The actions by the symmetric groups are depicted by rearranging input or output wires; for example, if $f \in \mathscr{C}(A_1, A_2, A_3; B_1, B_2, B_3, B_4)$, then the element $(124) \cdot f \cdot (123)$ is depicted as left below. Now identities are depicted as bare wires, and the composition $g_j \circ_i f$ as the plugging of the *i*th output of f into the *j*th input

of g, as on the right in:



With this presentation, the axioms for a polycategory can be seen simply as asserting that various ways of building up the same compound diagram coincide. This has the advantage of allowing us to avoid the the careful index-tracking needed to state the axioms in the algebraic formulation; however it is not yet precise. As a first step in this direction, we show how the graphical presentation allows us to derive a description of polycategories as the algebras for a monad on a presheaf category.

Definition 2.2. Let P denote the category with objects $\{\star\} + \mathbb{N} \times \mathbb{N}$ and with non-identity maps $\sigma_1, \ldots, \sigma_n$ and $\tau_1, \ldots, \tau_m \colon \star \to (n, m)$. A presheaf $X \in \mathscr{P}\mathsf{P}$ is called a *polygraph*²; elements of $X(\star)$ are called *vertices*, elements of X(n, m)are called *edges*, and we write s_1, \ldots, s_n and t_1, \ldots, t_m for $X(\sigma_1), \ldots, X(\sigma_n)$ and $X(\tau_1), \ldots, X(\tau_m)$. The *sources* and *targets* of an edge *e* are its images under the maps s_1, \ldots, s_n , respectively t_1, \ldots, t_m .

The functor $U: \operatorname{Polycat} \to \mathscr{P}\mathsf{P}$ sending each polycategory to its underlying polygraph is easily seen to be monadic, so that we may identify polycategories with algebras for the induced monad on $\mathscr{P}\mathsf{P}$. To describe this monad explicitly, we exploit the correspondence between the diagrams drawn above and presheaves over P . For instance, the diagram (2.1) for a morphism with *n* inputs and *m* outputs corresponds to the representable presheaf $\mathsf{y}_{(n,m)} = \mathsf{P}(-,(n,m))$, while the diagram (2.2) for a composite $g_i \circ_i f$ corresponds to a pushout

²Our usage follows [2]; note that these polygraphs are completely unrelated to those of [4].

in $\mathscr{P}\mathsf{P}$. Writing $A = \mathsf{y}_{(p,q)} \mathsf{j} \bullet_i \mathsf{y}_{(n,m)}$, the further composite $h_{\ell} \circ_{k+i-1} (g_{\mathfrak{j}} \circ_i f)$ in the associativity axiom corresponds to the pushout in $\mathscr{P}\mathsf{P}$ as to the left in:



which clearly coincides to within isomorphism with the polygraph representing $(h_{\ell} \circ_k g)_{j+\ell-1} \circ_i f$, since both are colimits for the diagram above right. Iteratively taking pushouts of this kind yields the following class of polygraphs describing the compound wiring operations of a polycategory; eventually, in Section 6.3 below, we will be able to generate all of these shapes from those for the basic wiring operations, but for the moment we give a more hands-on construction.

Definition 2.3. A finite polygraph is one with finitely many vertices and edges. An (n,m)-labelling of a finite polygraph is given by choices of vertices ℓ_1, \ldots, ℓ_n and r_1, \ldots, r_m , called the *leaves* and roots respectively. An *isomorphism* of labelled polygraphs is one respecting the labellings. Let $\mathscr{L}(n,m)$ be a set of isomorphismclass representatives of (n,m)-labelled finite polygraphs; we write |X| for the underlying polygraph of $X \in \mathscr{L}(n,m)$ and ℓ^X and r^X for the labellings. Now:

- (a) Let $id \in \mathscr{L}(1,1)$ be y_{\star} labelled in the unique possible way;
- (b) Given $X \in \mathscr{L}(n,m)$, $Y \in \mathscr{L}(p,q)$ and indices $1 \leq i \leq m$ and $1 \leq j \leq p$, let $Y_j \bullet_i X \in \mathscr{L}(n+p-1,m+q-1)$ be such that there is a pushout of underlying polygraphs

(2.4)
$$\begin{array}{c} y_{\star} \xrightarrow{r_{i}^{X}} |X| \\ \ell_{j}^{Y} \downarrow \qquad \qquad \downarrow u \\ |Y| \xrightarrow{v} |Y_{j} \bullet_{i} X \end{array}$$

with the labelling of the leaves and roots given respectively by:

$$v\ell_1^Y, \dots, v\ell_{j-1}^Y, \ u\ell_1^X, \dots, u\ell_n^X, \ v\ell_{j+1}^Y, \dots, v\ell_p^Y$$

and $ur_1^X, \dots, ur_{i-1}^X, \ vr_1^Y, \dots, vr_q^Y, \ ur_{i+1}^X, \dots, ur_m^X$

- (c) For any $X \in \mathscr{L}(n,m)$, $\varphi \in \mathfrak{S}_n$ and $\psi \in \mathfrak{S}_m$, let $\psi \cdot X \cdot \varphi \in \mathscr{L}(n,m)$ be |X| labelled by $\ell_{\varphi(1)}, \ldots, \ell_{\varphi(n)}$ and $r_{\psi^{-1}(1)}, \ldots, r_{\psi^{-1}(m)}$.
- (d) For each $n, m \in \mathbb{N}$, let $\langle n, m \rangle \in \mathscr{L}(n, m)$ be $\mathsf{y}_{(n,m)}$ labelled by $\mathsf{y}_{\sigma_1}, \ldots, \mathsf{y}_{\sigma_n}$ and $\mathsf{y}_{\tau_1}, \ldots, \mathsf{y}_{\tau_m}$.

Now let $\mathscr{T}(n,m) \subset \mathscr{L}(n,m)$ be the subsets obtained by closing the elements $\langle n,m \rangle$ in (d) under the operations in (a)–(c).³

³Note that implicit in these definitions are the assumptions that the elements id and $\langle n, m \rangle$ of (a) and (d) are the chosen representatives of their isomorphism-classes, and that relabelling a representative $X \in \mathscr{L}(n,m)$ as in (c) yields another such; we are clearly at liberty to make these assumptions.

As noted above, the labelled polygraphs in the sets $\mathscr{T}(n,m)$ represent the compound wiring operations of a polycategory; following [18, 19], we may characterise them in a direct combinatorial manner.

Definition 2.4. For any polygraph X, we write G_X for the undirected multigraph obtained as follows: the nodes are the disjoint union of the sets of edges and of vertices of X, and there is an arc $v \frown e$ for each way that v is a source or a target of e. A polygraph X is called a *polygraphic tree* if it has finitely many edges and vertices, and moreover:

- Each vertex of X is a source of at most one edge;
- Each vertex of X is a target of at most one edge;
- G_X is acyclic (in particular without multiple edges) and connected (in particular non-empty).

A labelled polygraphic tree is a polygraphic tree equipped with an (n, m)-labelling for which ℓ_1, \ldots, ℓ_n enumerate the vertices which are not the targets of any edge, and r_1, \ldots, r_m enumerate the vertices which are not the source of any edge.

Lemma 2.5. $\mathcal{T}(n,m)$ is the set of isomorphism-class representatives of (n,m)labelled polygraphic trees.

Proof. It is easy to see that each $\langle n, m \rangle$ is a labelled polygraphic tree, and that the labelled polygraphic trees are closed under (a)–(c) above. Conversely, we may show that any labelled polygraphic tree T is in $\mathscr{T}(n,m)$ by an easy induction on the number of edges in T.

The labelled polygraphic trees may now be used to give a concrete description of the left adjoint F_{P} to $U: \mathsf{Polycat} \to \mathscr{P}_{\mathsf{P}}$. Given Lemma 2.5, the proof of the following result is straightforward, if tedious; the reader may reconstruct it by suitably adapting [25, Proposition 1.9.2] or [18, §2.2.7].

Proposition 2.6. The free polycategory F_PX on a polygraph $X \in \mathscr{P}P$ has object set $X(\star)$ and morphism sets $(F_PX)(v_1, \ldots, v_n; w_1, \ldots, w_m)$ given by

$$\sum_{T \in \mathscr{T}(n,m)} \{ f \colon |T| \to X \text{ in } \mathscr{P}\mathsf{P} \colon f(\ell_i^T) = v_i \text{ and } f(r_j^T) = w_j \}$$

The identity morphism in $(F_PX)(v;v)$ is the pair $(id \in \mathscr{T}(1,1), v: y_* \to X)$; the composite $(S, f)_j \circ_i (T, g)$ is given by $(S_j \bullet_i T, f_j \bullet_i g)$, where $f_j \bullet_i g$ is the unique map out of the pushout (2.4) induced by f and g; and the exchange operation is defined by $\psi \cdot (T, f) \cdot \varphi = (\psi \cdot T \cdot \varphi, f)$.

There is another way of presenting polycategories as the algebras for a monad on a presheaf category. Though it is further away from the graphical intution, it is quite a common approach in mathematical practice, and still fits into the general framework we will develop; it therefore seems to be worth describing here. The idea is to incorporate the exchange operations into the underlying presheaf:

8

Definition 2.7. Let P_s be the category obtained from P by adjoining arrows $\xi_{\varphi,\psi}: (n,m) \to (n,m)$ for each $\varphi \in \mathfrak{S}_n$ and $\psi \in \mathfrak{S}_m$, subject to the equations:

$$\begin{aligned} \xi_{\mathrm{id}_n,\mathrm{id}_m} &= \mathrm{id}_{(n,m)} & & \xi_{\varphi_1,\psi_1} \circ \xi_{\varphi_2,\psi_2} = \xi_{\varphi_1\varphi_2,\psi_2\psi_1} \\ \xi_{\varphi,\psi} \circ \sigma_i &= \sigma_{\varphi(i)} & & \xi_{\varphi,\psi} \circ \tau_{\psi(j)} = \tau_j . \end{aligned}$$

A presheaf $X \in \mathscr{P}\mathsf{P}_s$ is called a symmetric polygraph.

Like before, the forgetful $U: \operatorname{Polycat} \to \mathscr{P}_s$ is monadic, and its left adjoint can now be described using labelled symmetric polygraphic trees. Let $\mathscr{L}_s(n,m)$ and $\mathscr{T}_s(n,m)$ be defined exactly like $\mathscr{L}(n,m)$ and $\mathscr{T}(n,m)$ in Definition 2.3 but working over the category \mathscr{P}_s of symmetric polygraphs. Since each symmetric polygraph $\mathsf{y}_{(n,m)}$ has free action by $\mathfrak{S}_n \times \mathfrak{S}_m^{\mathrm{op}}$, the same will be true of the underlying symmetric polygraph of any $T \in \mathscr{T}_s(n,m)$; in fact, it is easy to see that these T's are precisely the images⁴ of the non-symmetric trees in $\mathscr{T}(n,m)$ under the free functor $\mathscr{P}_s \to \mathscr{P}_s$. The key difference is that a symmetric labelled tree $T \in \mathscr{T}_s(n,m)$ may admit a non-trivial group \mathfrak{S}_T of label-preserving automorphisms; the construction of the free polycategory must now quotient out by the action of these automorphisms.

Proposition 2.8. The free polycategory $F_{\mathsf{P}_s}X$ on $X \in \mathscr{P}_{\mathsf{P}_s}$ has object set $X(\star)$ and morphism sets $(F_{\mathsf{P}_s}X)(v_1, \ldots, v_n; w_1, \ldots, w_m)$ given by

$$\sum_{T \in \mathscr{T}_s(n,m)} \{ f \colon |T| \to X \text{ in } \mathscr{P}\mathsf{P}_s \colon f(\ell_i^T) = v_i, \ f(r_j^T) = w_j \}_{/\mathfrak{S}_T}$$

with remaining structure defined analogously to Proposition 2.6 above.

2.2. **Properads and PROPs.** Though polycategories will serve as our main running example, it is worth saying a few words about some variations on the same theme. A (coloured) properad [31] specialises a polycategory by allowing the plugging of several outputs of a morphism $f \in \mathscr{C}(\vec{A}; \vec{B})$ into a corresponding number of inputs of $g \in \mathscr{C}(\vec{C}; \vec{D})$. This means that, for any non-empty sequences of indices $I = \{i, \ldots, i+k\}$ and $J = \{j, \ldots, j+k\}$ such that $B_{i+\ell} = C_{j+\ell}$ for each $0 \leq \ell \leq k$, there is a composite morphism

(2.5)
$$g_{J} \circ_{I} f \in \mathscr{C}(\vec{C}_{< j}, \vec{A}, \vec{C}_{> j+k}; \vec{B}_{< i}, \vec{D}, \vec{B}_{> i+k})$$

While it would be possible to state the axioms for a properad in an algebraic manner paralleling Definition 2.1 above, it is simpler to proceed directly to a description of them as the algebras for a monad on $\mathscr{P}P$. For this, we first generate the class of labelled *properadic* trees: we do so as in Definition 2.3 but with clause (b) modified so as to allow for pushouts of the form

⁴Though note that non-isomorphic elements of $\mathscr{T}(n,m)$ may become isomorphic in $\mathscr{T}_{s}(n,m)$.

The monad for properads on $\mathscr{P}\mathsf{P}$ may now be defined as in Proposition 2.6 but with reference to the larger class of properadic trees. As in the polycategorical case, we can also view properads as monadic over symmetric polygraphs; a description of the free properad monad in this case is in [19].

Finally, the PROPs of [24] can be seen as a further specialisation of properads, in which morphisms may also be composed together along an *empty* set of wires; so we augment the data of a properad with composition operations assigning to each $f \in \mathscr{C}(\vec{A}; \vec{B})$ and $g \in \mathscr{C}(\vec{C}; \vec{D})$ a morphism

$$(2.6) g_{\varnothing \circ_{\varnothing}} f \in \mathscr{C}(A,C;B,D)$$

which, graphically, encodes the operation of placing two string diagrams next to each other; and a chosen element $0 \in \mathscr{C}(\ ; \)$, corresponding to the empty string diagram. From here we may proceed exactly as before to view PROPs as algebras for monads on both $\mathscr{P}P$ and $\mathscr{P}P_s$.

3. FAMILIAL FUNCTORS AND SHAPELINESS

Now that we have described various "graphically specified" structures as algebras for monads on presheaf categories, we begin our attempts to obtain these monads via a notion of shapeliness. As in the introduction, our approach will be to seek on the appropriate presheaf category a *universal* shapely monad U with "exactly one operation of each shape", and to generate the monad encoding the given structure as a suitable submonad of U. In this section, we look for U as a terminal object among *familially representable*, or more shortly *familial*, endofunctors—ones which pointwise are coproducts of representables. While this turns out not quite to work, the techniques we develop will be crucial to our subsequent efforts.

3.1. Linear operations and familial functors. The key concept underlying the notion of familial functor is that of a *linear operation*.

Definition 3.1. Given $F: \mathscr{A} \to \mathscr{B}$ and objects $A \in \mathscr{A}$ and $B \in \mathscr{B}$, an *F*-operation of input arity A at stage B is a map $t: B \to FA$. An F-operation t is *linear* if it is initial in its connected component of the comma category $B \downarrow F$.

An operation $t: B \to TA$ of a monad T on \mathscr{A} corresponds to a family of interpretation functions $\llbracket t \rrbracket: \mathscr{A}(A, X) \to \mathscr{A}(B, X)$, one for each T-algebra (X, x); maps of $B \downarrow T$ account for reindexing such T-operations so as to act only on part of their input arity, so that linearity expresses the idea of an operation which "consumes all its input arity".

Lemma 3.2. An operation $t: B \to FA$ is linear if and only if for every square of the following form, there is a unique $h: A \to A'$ with Fh.t = u; it then follows also that fh = g.

$$(3.1) \qquad \begin{array}{c} B \xrightarrow{u} FA' \\ \downarrow & Fh \\ FA \xrightarrow{Fg} FA'' \end{array}$$

10

Proof. This is [7, Proposition 0].

Now a *familial* functor is one whose operations are all reindexings of linear ones. In giving the definition, we say that Y covers X if there is a map $Y \to X$.

Definition 3.3. A functor $F \colon \mathscr{A} \to \mathscr{B}$ is familial at stage $B \in \mathscr{B}$ if each operation in $B \downarrow F$ is covered by a linear one; a transformation $\alpha \colon F \Rightarrow G$ is familial at stage B if F and G are so, and the induced functor $B \downarrow F \to B \downarrow G$ preserves linear operations. We write simply familial to mean "familial at every stage".

Familial functors were introduced by Diers [7]; his terminology is that familial functors are those "having a left multiadjoint". Our name is a shortening of the term "familially representable" used—for the special case $\mathscr{B} = \mathbf{Set}$ —in [15].

Lemma 3.4. A functor $F : \mathscr{A} \to \mathscr{B}$ is familial at stage $B \in \mathscr{B}$ if and only if the functor $\mathscr{B}(B, F) : \mathscr{A} \to \mathbf{Set}$ is a (possibly large) coproduct of representables.

Proof. For F to be familial at stage B is for $B \downarrow F$ to be a coproduct of categories with initial objects; since $B \downarrow F$ is the category of elements of $\mathscr{B}(B, F)$, this is to say that this latter functor is a coproduct of representables.

3.2. **Pointwise familiality.** We are interested in familial endofunctors of presheaf categories; later, we will need more general familial functors with *codomain* a presheaf category. The salient kind of familiality for these is as follows.

Definition 3.5. A functor $F: \mathscr{A} \to \mathscr{PC}$ or transformation $\alpha: F \Rightarrow G: \mathscr{A} \to \mathscr{PC}$ is *pointwise familial* if it is familial at all representable stages; F is called *small* if $\pi_0(\mathsf{y}_c \downarrow F)$ is a mere *set* for each $c \in \mathscr{C}$. We write $\mathbf{FAM}_0(\mathscr{A}, \mathscr{PC})$ for the category of small pointwise familial functors and pointwise familial transformations.

By Lemma 3.4, $F: \mathscr{A} \to \mathscr{PC}$ is (small) pointwise familial just when each $(F-)c \in [\mathscr{A}, \mathbf{Set}]$ is a small coproduct of representables. So, for example, the "free polycategory" endofunctor on the category $\mathscr{P}\mathsf{P}$ of polygraphs as in Proposition 2.6 is pointwise familial, but the corresponding endofunctor on the category $\mathscr{P}\mathsf{P}_s$ of symmetric polygraphs is *not* so, as it involves not just coproducts of representables but also quotients by group actions. We will be able to handle the latter example when we consider *analytic* functors in the following section.

Clearly, a small pointwise familial F is determined by the representable summands of each (F-)c and how these transform under maps $(F-)f: (F-)d \to (F-)c$. The following definition provides a compact way of representing these basic data. In giving it, we assume that any given $F \in \mathbf{FAM}_0(\mathscr{A}, \mathscr{PC})$ is equipped with a *choice* of linear operation in each connected component of $y_c \downarrow F$; for each $t \in y_c \downarrow F$ we write \tilde{t} for the chosen linear operation which covers it. We also write el X for the *category of elements* of a presheaf $X \in \mathscr{PC}$, whose object-set is $\sum_{c \in \mathscr{C}} Xc$ and whose morphisms from $s \in Xc$ to $t \in Xd$ are arrows $f: c \to d$ of \mathscr{C} with s = (Xf)(t).

Definition 3.6. The spectrum [7, Definition 3] of a small pointwise familial $F: \mathscr{A} \to \mathscr{PC}$ is the presheaf $S_F \in \mathscr{PC}$ given by:

 $S_F(c) = \{t \in \mathsf{y}_c \downarrow F : \tilde{t} = t\}$ and $S_F(f : d \to c) : t \mapsto \widetilde{t\mathsf{y}_f}$.

The canonical diagram of F is the functor $D_F \colon \operatorname{el} S_F \to \mathscr{PC} \downarrow F$ with:

$$D_F(t) = \bigvee_{FA}^{\mathbf{y}_c} \quad \text{and} \quad D_F(f: \widetilde{ty_f} \to t) = \bigvee_{t \in Y_f} \bigvee_{FA'} \bigvee_{Fu_f}^{\mathbf{y}_f} \downarrow_t \\ FA' \xrightarrow{Fu_f} FA ,$$

where the lower right map is uniquely induced by linearity of ty_f . The *exponent* [33, Definition 7.1] of F is the functor E_F : $\operatorname{el} S_F \to \mathscr{A}$ obtained by composing the canonical diagram with the second projection $\pi_2: \mathscr{PC} \downarrow F \to \mathscr{A}$.

As elements of $S_F(c)$ are in bijection with connected components of $y_c \downarrow F$, the presheaf S_F is equally the colimit of F; smallness is just what is needed to ensure this colimit exists. In particular, smallness is vacuous when either \mathscr{A} is small or \mathscr{A} has a terminal object, and in the latter case, we may take $S_F = F1$.

We now make precise the idea that a pointwise familial functor is completely determined by its spectrum and its exponent, by constructing an equivalence between $\mathbf{FAM}_0(\mathscr{A}, \mathscr{PC})$ and the following category:

Definition 3.7. For any \mathscr{A} and small \mathscr{C} , the category $\operatorname{el}_{\mathscr{C}} /\!\!/ \mathscr{A}$ has as objects, pairs of a presheaf $S \in \mathscr{PC}$ and a functor $E \colon \operatorname{el} S \to \mathscr{A}$, and as maps $(S, E) \to (T, D)$, pairs of a presheaf map $p \colon S \to T$ and a natural isomorphism φ of the form:

$$(3.2) \qquad \qquad \begin{array}{c} \operatorname{el} S & \xrightarrow{\operatorname{el} p} & \operatorname{el} T \\ \searrow & \swarrow & \swarrow \\ \swarrow & \swarrow & D \end{array}$$

This category is the " \mathscr{C} -Fam(\mathscr{A})" of [32, Definition 2.10], and the following result is a generalisation of Theorem 2.18 of *ibid*.

Proposition 3.8. The assignation $F \mapsto (S_F, E_F)$ is the action on objects of an equivalence of categories between $FAM_0(\mathscr{A}, \mathscr{PC})$ and $el_{\mathscr{C}} /\!\!/ \mathscr{A}$.

Proof. First let $\alpha: F \Rightarrow G$ in $\text{FAM}_0(\mathscr{A}, \mathscr{PC})$. To give the transformation α is to give transformations $\alpha_c: (F_{-})c \Rightarrow (G_{-})c$ naturally in c; since by Lemma 3.4 the functor $(F_{-})c$ is a coproduct of representables, giving each α_c is equivalent to giving the G-linear operation $\alpha_A.t: y_c \to GA$ obtained by acting α on each chosen F-linear operation $t: y_c \to FA$. But $\alpha_A.t$ factorises as on the left in:

$$(3.3) \qquad \begin{array}{c} p(t) := \widetilde{\alpha_A.t} & \forall c & el S_F \xrightarrow{el p} el S_G \\ GA' \xrightarrow{G\varphi_t} GA & & E_F \xrightarrow{\varphi} \swarrow E_G \end{array}$$

using the chosen linear operations of G, and linearity of $\alpha_A t$ is clearly equivalent to each φ_t being invertible. Thus, to give the pointwise familial α is equally to specify for each chosen linear $t \in \mathsf{y}_c \downarrow F$ a chosen linear $p(t) \in \mathsf{y}_c \downarrow G$ together with an isomorphism $\varphi_t \colon E_G(p(t)) \to E_F(t)$. All this must be done naturally in c so that to give α is equally to give a pair (p, φ) as right above with φ invertible.

This defines $\mathbf{FAM}_0(\mathscr{A}, \mathscr{PC}) \to \operatorname{el}_{\mathscr{C}} /\!\!/ \mathscr{A}$ on morphisms and simultaneously shows that it is fully faithful (functoriality is easily checked); it remains to

prove essential surjectivity. Given $S \in \mathscr{PC}$ and $E: \operatorname{el} S \to \mathscr{A}$, define a functor $F: \mathscr{A} \to \mathscr{PC}$ by taking $(F-)c = \sum_{t \in Sc} \mathscr{A}(Et, -)$ and taking $(F-)(f: d \to c)$ to be the unique natural transformation rendering commutative each diagram:

$$\begin{array}{c} \mathscr{A}(Et,-) \xrightarrow{\mathscr{A}(Ef,-)} \mathscr{A}(E(tf),-) \\ \iota \\ \downarrow \\ \sum_{t \in Sc} \mathscr{A}(Et,-) \xrightarrow{(F-)f} \sum_{u \in Sd} \mathscr{A}(Eu,-) \end{array}$$

By Lemma 3.4, F is pointwise familial, and is moreover small since the coproduct defining (F-)c is so; now by choosing the linear operations in $y_c \downarrow F$ to be those $\gamma_t : y_c \to FEt$ picking out the pairs $(t, 1_{Et})$, we have a bijection $S \to S_F$ sending t to γ_t , which, since $E_F(\gamma_t) = Et$, commutes *strictly* with the functors to \mathscr{A} . \Box

3.3. Composition of familial functors. In seeking a universal shapely monad among the class of familial endofunctors, we must consider both composability and existence of a terminal object; we start with composability. The following lemma gives the properties of linear operations necessary to establish our results.

Lemma 3.9. Let $F \colon \mathscr{A} \to \mathscr{B}$ and $G \colon \mathscr{B} \to \mathscr{C}$ be functors.

- (i) If $s: C \to GB$ is G-linear and $t: B \to FA$ is F-linear, then the composite $Gt.s: C \to GB \to GFA$ is GF-linear.
- (ii) The full subcategory $\operatorname{Lin}(\mathscr{B} \downarrow F) \subset \mathscr{B} \downarrow F$ on the linear operations is closed under pointwise colimits (ones created by the projection $\mathscr{B} \downarrow F \to \mathscr{B} \times \mathscr{A}$).

Proof. An easy exercise using Lemma 3.2.

Definition 3.10. Given
$$\mathscr{A}' \subset \mathscr{A}$$
 and $\mathscr{B}' \subset \mathscr{B}$ full replete subcategories, we say that $F: \mathscr{A} \to \mathscr{B}$ is $(\mathscr{A}', \mathscr{B}')$ -familial if it is familial at each stage $B \in \mathscr{B}'$ and each linear $t \in B \downarrow F$ has input arity in \mathscr{A}' . A transformation $\alpha: F \Rightarrow G$ between such functors is $(\mathscr{A}', \mathscr{B}')$ -familial if it is familial at every stage $B \in \mathscr{B}'$.

In this terminology, a familial functor $F: \mathscr{A} \to \mathscr{B}$ is equally $(\mathscr{A}, \mathscr{B})$ -familial, while a pointwise familial functor $\mathscr{A} \to \mathscr{PC}$ is equally an $(\mathscr{A}, \mathscr{YC})$ -familial one. The next result improves in very mild ways on [7, p. 985] and [33, Corollary 5.15].

Proposition 3.11. If $F: \mathscr{A} \to \mathscr{B}$ and $G: \mathscr{B} \to \mathscr{C}$ are $(\mathscr{A}', \mathscr{B}')$ - and $(\mathscr{B}', \mathscr{C}')$ -familial, then their composite is $(\mathscr{A}', \mathscr{C}')$ -familial, and has as linear operations at stage $C \in \mathscr{C}'$ precisely the composites $Gt.s: C \to GB \to GFA$ of G- and F-linear operations. The correspondingly familial transformations between these functors are likewise composable; in particular, there is a 2-category FAM of categories, familial functors and familial transformations.

Proof. To show that GF is $(\mathscr{A}', \mathscr{C}')$ -familial it suffices, by an easy argument, to show that any $s: C \to GFX$ with $C \in \mathscr{C}'$ is covered in $C \downarrow GF$ by some linear operation with input arity in \mathscr{A}' . But we have successive factorisations



with $B \in \mathscr{B}'$ and $A \in \mathscr{A}'$ by applying familiality of G to s and of F to t. By Lemma 3.9(i), the composite $G\tilde{t}.\tilde{s}$ is GF-linear so that $f: G\tilde{t}.\tilde{s} \to s$ is the required cover. The remaining parts of the result are now easy consequences. \Box

Since we are really interested in pointwise familial functors between presheaf categories, we should like to know that these, too, are closed under composition. The key to showing this is the following result.

Proposition 3.12. Let \mathscr{A} be cocomplete. For any functor $F : \mathscr{A} \to \mathscr{B}$ or transformation $\alpha : F \Rightarrow G$, the full subcategory $\mathscr{B}' \subset \mathscr{B}$ whose objects are those stages $B \in \mathscr{B}$ at which F (respectively α) is familial is closed in \mathscr{B} under colimits.

Proof. Suppose given $F: \mathscr{A} \to \mathscr{B}$, a diagram $D: \mathscr{I} \to \mathscr{B}$ such that F is familial at each $DI \in \mathscr{B}$, and a colimiting cocone $(p_I: DI \to V)_{I \in \mathscr{I}}$; we must show that F is also familial at V. So let $t: V \to FA$, and consider the diagram of linear operations $D_t: \mathscr{I} \to \mathscr{B} \downarrow F$ defined by:

$$\begin{array}{ccc} DI \\ D_t(I) = \begin{array}{c} DI \\ \widetilde{tp_I} \\ FA_I \end{array} \end{array} \qquad D_t(f: I \to J) = \begin{array}{c} DI \xrightarrow{Df} DJ \\ \widetilde{tp_I} \\ FA_I \xrightarrow{Fu_f} FA_I \end{array}$$

where the map u_f is the unique one induced by linearity of $\widetilde{tp_I}$. Since \mathscr{A} is cocomplete, the diagram D_t admits a pointwise colimit $u: V \to FW$, which by Lemma 3.9(ii) is itself linear. There is a cocone $D_t \Rightarrow \Delta t$ with components $(p_I, q_I): \widetilde{tp_I} \to t$ where the maps q_I are, again, induced by linearity of $\widetilde{tp_I}$, and this now induces a map $u \to t$ in $V \downarrow F$ providing the desired linear cover of t. This shows F is familial at V, and also that $t \in V \downarrow F$ is linear just when its cocone $(p, q): D_t \Rightarrow \Delta t$ is colimiting; using this last fact, the corresponding result for transformations α follows easily. \Box

As every presheaf is a colimit of representables, we immediately conclude from the preceding two results that:

Corollary 3.13. If \mathscr{A} is cocomplete, then each pointwise familial functor or transformation in $\operatorname{FAM}_0(\mathscr{A}, \mathscr{PC})$ is familial; whence there is a 2-category FAM_0 of presheaf categories and pointwise familial functors and transformations.

In the next section, size considerations will force us to bound the input arities of the pointwise familial functors we consider. As we would still like such functors to compose, we introduce the relevant notions and prove composability here.

Definition 3.14. We write $\mathscr{FC} \subset \mathscr{PC}$ for the full, replete subcategory of *finitely* presentable presheaves: those expressible as a finite colimit of representables. A pointwise familial functor $\mathscr{PC} \to \mathscr{PD}$ is called *finitary* if it is $(\mathscr{FC}, \mathsf{yD})$ -familial.

The modifier "finitary" typically refers to a functor which preserves filtered colimits; that our usage agrees with this follows from Lemma 3.4 and the fact that a representable $\mathscr{PC}(A, -): \mathscr{PC} \to \mathbf{Set}$ is finitary just when A is in \mathscr{FC} .

To see that finitary pointwise familial functors and transformations compose, we appeal to Proposition 3.11 and the following result:

Proposition 3.15. The pointwise familial $F: \mathscr{PC} \to \mathscr{PD}$ is finitary if and only if it is $(\mathscr{FC}, \mathscr{FD})$ -familial; whence there is a 2-category \mathbf{FAM}_0^{ω} of presheaf categories, finitary pointwise familial functors and pointwise familial transformations.

Proof. For the non-trivial direction, let $t: B \to FA$ with $B \in \mathscr{FD}$. On expressing B as a finite colimit of representables, the proof of Proposition 3.12 yields a cover of t by a linear operation $u: B \to FA$ obtained as a finite colimit in $\mathscr{B} \downarrow F$ of linear operations of the form $y_{d_I} \to FA_I$. By assumption, each A_I is in \mathscr{FC} , whence $A = \operatorname{colim}_I A_I$ is too.

3.4. Universal familial endofunctors. We now have all the ingredients we require for our first attempt at constructing a universal shapely monad U on \mathscr{PC} . As anticipated in the previous section, a naive attempt to construct it as a terminal object in the monoidal category $FAM_0(\mathscr{PC}, \mathscr{PC})$ fails for size reasons.

Proposition 3.16. If $\mathscr{C} \neq 0$ and the category \mathscr{A} has a proper class of nonisomorphic objects, then $\mathbf{FAM}_0(\mathscr{A}, \mathscr{PC})$ has no terminal object; in particular, if $\mathscr{C} \neq 0$ then $\mathbf{FAM}_0(\mathscr{PC}, \mathscr{PC})$ has no terminal object.

Proof. By Proposition 3.8, it suffices to show that $\operatorname{el}_{\mathscr{C}} /\!\!/ \mathscr{A}$ has no terminal object. Suppose that (S, E) were terminal; fixing some $c \in \mathscr{C}$, we would then have for each $A \in \mathscr{A}$ a unique map

$$\operatorname{el} \mathsf{y}_c \xrightarrow{\operatorname{el} t_A} \operatorname{el} S$$

$$\Delta A \xrightarrow{\varphi_A} \swarrow E$$

where ΔA is the constant functor at A. Note that $t_A \in Sc$ satisfies $Et_A \cong A$; since there are a proper class of non-isomorphic A's, there must be a proper class of distinct t_A 's, contradicting the fact that Sc is a set. \Box

What permits the above negative argument is the fact that a pointwise familial functor may have linear operations of arbitrarily large input arity; this suggests restricting attention to the *finitary* pointwise familial functors whose linear input arities lie in the essentially small⁵ \mathscr{FC} . We first note that:

Lemma 3.17. Precomposition with the inclusion $J : \mathscr{FC} \to \mathscr{PC}$ induces an equivalence between the categories $\operatorname{FAM}_0^{\omega}(\mathscr{PC}, \mathscr{PD})$ and $\operatorname{FAM}_0(\mathscr{FC}, \mathscr{PD})$.

Proof. Precomposing the equivalence $\mathbf{FAM}_0(\mathscr{FC}, \mathscr{PD}) \to \mathrm{el}_{\mathscr{D}} /\!\!/ \mathscr{FC}$ of Proposition 3.8 by $(-) \circ J : \mathbf{FAM}_0^{\omega}(\mathscr{PC}, \mathscr{PD}) \to \mathbf{FAM}_0(\mathscr{FC}, \mathscr{PD})$ evidently yields another equivalence; whence, by two-out-of-three, $(-) \circ J$ is an equivalence. \Box

Unfortunately, even with the finitariness restriction we are still unable to construct a strictly terminal familial endofunctor:

⁵A category is *essentially small* if it is equivalent to a small category.

Proposition 3.18. If $\mathscr{C} \neq 0$ and \mathscr{A} is essentially small, then $\operatorname{FAM}_0(\mathscr{A}, \mathscr{PC})$ always has a weakly terminal object, but has a terminal object if and only if \mathscr{A} has no non-identity automorphisms; consequently, if $\mathscr{C} \neq 0$, then $\operatorname{FAM}_0^{\omega}(\mathscr{PC}, \mathscr{PC})$ has a weakly terminal object, but not a terminal object.

Proof. By Proposition 3.8 we may prove the stated properties for the equivalent category $\operatorname{el}_{\mathscr{C}} /\!\!/ \mathscr{A}$; but as $\mathscr{A} \simeq \mathscr{A}'$ with \mathscr{A}' small and now $\operatorname{el}_{\mathscr{C}} /\!\!/ \mathscr{A} \simeq \operatorname{el}_{\mathscr{C}} /\!\!/ \mathscr{A}'$, we may assume without loss of generality that \mathscr{A} is itself small. We construct a weakly terminal (S, E) in $\operatorname{el}_{\mathscr{C}} /\!\!/ \mathscr{A}$ as follows. The presheaf $S \in \mathscr{P}_{\mathscr{C}}$ has:

$$Sc = \{ \text{functors } F \colon \mathscr{C}/c \to \mathscr{A} \}$$
 and $S(f \colon d \to c) \colon F \mapsto F(f \circ -) ,$

while $E: el S \to \mathscr{A}$ is given by $E(F: \mathscr{C}/c \to \mathscr{A}) = F(1_c)$ on objects, and by:

$$f: F(f \circ -) \to F \qquad \mapsto \qquad F(f: f \to 1_c): Ff \to F1_c$$

on morphisms. To see weak terminality of (S, E), consider some other (T, D) in $\operatorname{el}_{\mathscr{C}}/\!\!/ \mathscr{A}$. We define $p: T \to S$ in \mathscr{PC} by sending $t \in Tc$ to the element

$$p(t) \colon \mathscr{C}/c = \operatorname{el} \mathsf{y}_c \xrightarrow{\operatorname{el} t} \operatorname{el} T \xrightarrow{D} \mathscr{A}$$

of Sc. Naturality of p in c is easy; while from the equality $p(t)(1_c) = Dt$, we deduce $E \circ \operatorname{el} p = D$: $\operatorname{el} T \to \mathscr{A}$ and so $(p, 1_D): (T, D) \to (S, E)$ in $\operatorname{el}_{\mathscr{C}} /\!\!/ \mathscr{A}$.

Now let \mathscr{A} have no non-identity automorphisms; replacing it by its (equivalent) skeleton, we may assume that in fact it has no non-identity isomorphisms, and so that each map (3.2) of $\operatorname{el}_{\mathscr{C}}/\!\!/ \mathscr{A}$ has φ an *identity*. In this case, we claim the weakly terminal (S, E) given above is terminal. Indeed, if $(q, 1_D): (T, D) \to (S, E)$ is any map in $\operatorname{el}_{\mathscr{C}}/\!\!/ \mathscr{A}$, then for each $t \in Tc$, the functor $q(t): \mathscr{C}/c \to \mathscr{A}$ satisfies $q(t)(1_c) = Dt = p(t)(1_c)$; but then $q(t)(h) = q(th)(1_d) = p(t)(h)$ for all $h: d \to c$, whence q = p as required.

Next let \mathscr{A} admit the non-identity automorphism $a \in \mathscr{A}(A, A)$, and assume that there is a terminal object (T, D) in $\operatorname{el}_{\mathscr{C}} /\!\!/ \mathscr{A}$; we derive a contradiction. By terminality of (T, D), there is for any $c \in \mathscr{C}$ a unique pair as on the left in

where here ΔA is the constant functor at A. But now the triangle on the right also describes a morphism $(y_c, \Delta A) \rightarrow (T, D)$; so we must have $\varphi \Delta a = \varphi$ and so, by invertibility of φ , that $\Delta a = \mathrm{id}_{\Delta_A}$, contradicting $a \neq \mathrm{id}_A$.

For the final claim, note that we have $\mathbf{FAM}_0^{\omega}(\mathscr{PC}, \mathscr{PC}) \simeq \mathbf{FAM}_0(\mathscr{FC}, \mathscr{PC})$ by Lemma 3.17; now if $\mathscr{C} \neq 0$, then the essentially small \mathscr{FC} certainly contains non-identity automorphisms—for instance, the switch map $y_c + y_c \to y_c + y_c$ —and so $\mathbf{FAM}_0^{\omega}(\mathscr{PC}, \mathscr{PC})$ has a weakly terminal object, but no terminal object. \Box

4. Analytic functors and shapeliness

The underlying reason that there is no terminal object among finitary familial endofunctors of a presheaf category is that linear operations cannot be fixed by automorphisms of their input arities; this means that such automorphisms may be propagated up to the level of familial functors, so obstructing the existence of a terminal object. The next step in our pursuit of a universal shapely monad will attempt to resolve this problem by introducing *analytic* functors, whose generating operations can be fixed by input arity automorphisms.

4.1. Generic operations and analytic functors. The fundamental step in moving from familial to analytic functors is to generalise from linear to generic operations. In what follows, we write \mathfrak{S}_X for the automorphism group of any object $X \in \mathscr{C}$.

Definition 4.1. An object $X \in \mathscr{C}$ is *Galois* if for each $Y \in \mathscr{C}$, the composition action makes $\mathscr{C}(X,Y)$ into a connected \mathfrak{S}_X -set. An operation $t: B \to FA$ of a functor $F: \mathscr{A} \to \mathscr{B}$ is *generic* if it is Galois in its connected component of $B \downarrow F$.

An object X is Galois when it admits a map to every other object (weak initiality) and, for any pair of maps $f, f' \colon X \rightrightarrows Y$, there is an automorphism $\sigma \in \mathfrak{S}_X$ with $f' = f\sigma$ (transitivity); thus, Galois objects are initial "up to a group of automorphisms". In these terms, a generic F-operation can be understood as one which, like a linear operation, consumes all of its input arity, but which may now be invariant under certain automorphisms of that arity.

The next result identifies our generic operations with those of [33, Definition 5.2], which when $\mathscr{A} = \mathscr{B} = \mathbf{Set}$ and B = 1 are equally those of [16].

Lemma 4.2. An operation $t: B \to FA$ is generic if and only if for every square of the following form there exists some $\ell: A \to Y$ with $F\ell.t = u$ and $h\ell = k$:

$$(4.1) \qquad \begin{array}{c} B \xrightarrow{u} FY \\ t & Ft \\ FA \xrightarrow{Fk} FZ \end{array} \\ B \xrightarrow{Fk} FZ \end{array}$$

Proof. It is easy to see—as in [5, Remarks 2.4], for example—that an object $X \in \mathscr{C}$ is Galois just when it is weakly initial, and each cospan $X \to Z \leftarrow Y$ can be completed to a commuting triangle by some $X \to Y$. Now the condition on t above says that any cospan $t \to v \leftarrow u$ in $B \downarrow F$ can be completed to a commuting triangle, which thus says that t is Galois in its connected component.

Corollary 4.3. Any map $h: u \to t$ in $B \downarrow F$ with generic codomain is a split epimorphism; in particular, any map between generic operations is an isomorphism.

Proof. Take $k = 1_A$ in (4.1).

Replacing linear operations with generic ones in the definition of familial functor yields the notion of *analytic* functor.

Definition 4.4. A functor $F: \mathscr{A} \to \mathscr{B}$ is analytic at stage $B \in \mathscr{B}$ if each operation in $B \downarrow F$ is covered by a generic one; a transformation $\alpha: F \Rightarrow G$ is analytic at stage B if F and G are so, and the induced functor $B \downarrow F \rightarrow B \downarrow G$ preserves generic operations. We write simply *analytic* to mean "analytic at every stage".

Analytic endofunctors of **Set** and *weakly cartesian* transformations were introduced in [16]; by [33, Theorems 10.10 & 10.11], they are precisely the (filteredcolimit preserving) analytic endofunctors and transformations in our sense.

Just as familial functors are obtained from coproducts of representables, so analytic functors arise from coproducts of *near-representables* in the sense of [30]:

Definition 4.5. Let $A \in \mathscr{A}$ and $G \leq \mathfrak{S}_A$. A coinvariant for G is a joint coequaliser $q: A \to A_{/G}$ for the set of morphisms $\{\sigma: A \to A \mid \sigma \in G\}$; dually, an invariant for G is a joint equaliser $\iota: A^{\setminus G} \to A$ for the maps in G. A functor $F: \mathscr{A} \to \mathbf{Set}$ is near-representable if $F \cong \mathscr{A}(A, -)_{/G}$ for some $A \in \mathscr{A}$ and $G \leq \mathfrak{S}_A = \mathfrak{S}_{\mathscr{A}(A, -)}$.

Lemma 4.6. A functor $F: \mathscr{A} \to \mathscr{B}$ is analytic at stage $B \in \mathscr{B}$ if and only if the functor $\mathscr{B}(B, F): \mathscr{A} \to \mathbf{Set}$ is a (possibly large) coproduct of near-representables.

Proof. This will follow as in Lemma 3.4 once we have proved that: $F \in [\mathscr{A}, \mathbf{Set}]$ is near-representable just when $\operatorname{el} F$ contains a Galois object. For any (x, A) in $\operatorname{el} F$, let $G = \mathfrak{S}_{(x,A)} \leq \mathfrak{S}_A$; now $x \colon \mathsf{y}_A \to F$ coequalises $\mathsf{y}_\sigma \colon \mathsf{y}_A \to \mathsf{y}_A$ for each $\sigma \in G$, and so descends to a map $\bar{x} \colon \mathsf{y}_{A/G} \to F$. It suffices to show that \bar{x} is an isomorphism just when (x, A) is Galois. Surjectivity of \bar{x} clearly corresponds to weak initiality of (x, A); injectivity requires that, for any $f, f' \colon A \rightrightarrows B$ with $\bar{x}(f) = \bar{x}(f') = y$, we have $f' = f\sigma$ for some $\sigma \in G$, or in other words, that for any $f, f' \colon (x, A) \rightrightarrows (y, B)$ in $\operatorname{el} F$, there is some $\sigma \colon (x, A) \to (x, A)$ in G with $f' = f\sigma$: which is transitivity of (x, A). \Box

4.2. **Pointwise analyticity.** As before, when we consider endofunctors of presheaf categories, or more generally functors *into* a presheaf category, the most appropriate kind of analyticity is pointwise:

Definition 4.7. A functor $F: \mathscr{A} \to \mathscr{PC}$ or transformation $\alpha: F \Rightarrow G: \mathscr{A} \to \mathscr{PC}$ is *pointwise analytic* if it is analytic at all representable stages; F is called *small* if $\pi_0(\mathsf{y}_c \downarrow F)$ is a mere set for each $c \in \mathscr{C}$. We write $\mathbf{AN}_0(\mathscr{A}, \mathscr{PC})$ for the category of small pointwise analytic functors and pointwise analytic transformations.

In particular, by Lemma 4.6, a functor $F: \mathscr{A} \to \mathscr{PC}$ is small pointwise analytic just when each $(F_{-})c$ is a small coproduct of near-representables; so, for example, comparing with the formula of Proposition 2.8, we find—as promised above—that the "free polycategory" endofunctor on the category of *symmetric* polygraphs is pointwise analytic, though it is not pointwise familial.

As in the familial case, a small pointwise analytic F is determined by the nearrepresentable summands of each (F-)c and how these transform under maps $(F-)f: (F-)d \to (F-)c$. We wish to give a representation of these data analogous to Definition 3.6; the new aspect is that, in encoding a near-representable summand $\mathscr{A}(A,-)_{/G}$, we must record not just the arity A but also the group Gof automorphisms which fix it. We do this using the notion of *orbit category*. **Definition 4.8.** The orbit category $\mathscr{O}(\mathscr{A})$ of a category \mathscr{A} has as objects, pairs (A, G) where $A \in \mathscr{A}$ and $G \leq \mathfrak{S}_A$, and as morphisms $[f]: (A, G) \to (B, H)$, equivalence classes of maps $f: A \to B$ in \mathscr{A} with the property that

(4.2) for all
$$\tau \in H$$
, there exists $\sigma \in G$ with $\tau f = f\sigma$,

where [f] = [f'] when there exists $\sigma \in G$ with $f' = f\sigma$. We write $J : \mathscr{A} \to \mathscr{O}(\mathscr{A})$ for the full embedding sending A to (A, 1).

Intuitively, we regard the generating operations of $F \in \mathbf{AN}_0(\mathscr{A}, \mathscr{PC})$ as having input arities drawn not from \mathscr{A} but from $\mathscr{O}(\mathscr{A})$; we will make this precise by equating such functors F with pointwise familial ones $F' \in \mathbf{FAM}_0(\mathscr{O}(\mathscr{A}), \mathscr{PC})$. First we describe the passage between functors with domains \mathscr{A} and $\mathscr{O}(\mathscr{A})$.

Proposition 4.9. $\mathcal{O}(\mathscr{A})$ admits all group invariants, and for any category \mathscr{C} admitting group invariants, composition with J induces an equivalence

(4.3)
$$\mathbf{INVAR}(\mathscr{O}(\mathscr{A}),\mathscr{C}) \xrightarrow{(-)\circ J} \mathbf{CAT}(\mathscr{A},\mathscr{C})$$

with domain the category of invariant-preserving functors and transformations.

Proof. The statement says that $\mathscr{O}(\mathscr{A})$ is the free completion of \mathscr{A} under group invariants, and by [17, Theorem 5.35], this completion may be found as the full subcategory of $[\mathscr{A}, \mathbf{Set}]^{\mathrm{op}}$ obtained by closing the representables under group invariants. So it suffices to identify $\mathscr{O}(\mathscr{A})$ with this full subcategory. Direct calculation using the Yoneda lemma shows that maps in $\mathscr{O}(\mathscr{A})$ from (A, G) to (B, H) are in bijection with maps $\mathscr{A}(B, -)_{/H} \to \mathscr{A}(A, -)_{/G}$ in $[\mathscr{A}, \mathbf{Set}]$; so there is a fully faithful $K \colon \mathscr{O}(\mathscr{A}) \to [\mathscr{A}, \mathbf{Set}]^{\mathrm{op}}$ with $K(A, G) = \mathscr{A}(A, -)_{/G}$. Clearly, each $\mathscr{A}(A, -)_{/G}$ in the image of K lies in the closure of the representables in $[\mathscr{A}, \mathbf{Set}]^{\mathrm{op}}$ under group invariants, and so it suffices to show that this subcategory in fact has all group invariants—which is [30, Proposition 2.2].

Explicitly, if \mathscr{C} admits group invariants and $H: \mathscr{A} \to \mathscr{C}$, then the invariantpreserving extension $H': \mathscr{O}(\mathscr{A}) \to \mathscr{C}$ is defined by $H'(A, G) = HA^{\setminus HG}$, where here $HG = \{H\sigma : \sigma \in G\} \leq \mathfrak{S}_{HA}$. In particular, if $F: \mathscr{A} \to \mathscr{B}$ is any functor between categories, then applying this construction to $JF: \mathscr{A} \to \mathscr{O}(\mathscr{B})$ yields an invariant-preserving $\mathscr{O}(F): \mathscr{O}(\mathscr{A}) \to \mathscr{O}(\mathscr{B})$ given by $\mathscr{O}(F)(A, G) = (FA, FG)$.

We will now show that, when $\mathscr{B} = \mathscr{PC}$, the equivalence (4.3) restricts back to one between pointwise analytic functors out of \mathscr{A} and pointwise familial ones out of $\mathscr{O}(\mathscr{A})$. However, under this equivalence, pointwise analytic transformations correspond not to familial ones but to *near-familial* ones in the following sense:

Definition 4.10. A morphism $[f]: (A, G) \to (B, H)$ in $\mathcal{O}(\mathscr{A})$ is called *vertical* if the underlying map $f: A \to B$ is invertible in \mathscr{A} . For any $F': \mathcal{O}(\mathscr{A}) \to \mathscr{B}$, an operation t in $B \downarrow F'$ is called *near-linear* if it admits a vertical map from a linear operation. If $F, G: \mathcal{O}(\mathscr{A}) \to \mathscr{B}$ are familial at stage B, then a transformation $\alpha: F \Rightarrow G$ is *near-familial* at stage B if it preserves near-linear operations.

We now give our equivalence result, together with a preparatory lemma; in the statement of the lemma, we call an object of an orbit category $\mathscr{O}(\mathscr{A})$ near-initial if it admits a vertical map from an initial object.

- **Lemma 4.11.** (i) \mathscr{A} has a Galois object if and only if $\mathscr{O}(\mathscr{A})$ has an initial one. $F: \mathscr{A} \to \mathscr{B}$ preserves Galois objects if and only if $\mathscr{O}(F): \mathscr{O}(\mathscr{A}) \to \mathscr{O}(\mathscr{B})$ preserves near-initial objects.
- (ii) Let \mathscr{B} admit group invariants and let $F \colon \mathscr{A} \to \mathscr{B}$ have invariant-preserving extension $F' \colon \mathscr{O}(\mathscr{A}) \to \mathscr{B}$. We have $B \downarrow F' \cong \mathscr{O}(B \downarrow F)$ naturally in F.

Proof. For (i), $A \in \mathscr{A}$ is Galois just when each $\mathscr{A}(A, B)$ is a connected \mathfrak{S}_A -set. This is easily equivalent to the existence of a unique $[u_B]: (A, \mathfrak{S}_A) \to (B, 1)$ in $\mathscr{O}(\mathscr{A})$ for each $B \in \mathscr{A}$; as $[u_B]$ clearly factors through each $[1]: (B, H) \to (B, 1)$, this is equivalent to (A, \mathfrak{S}_A) being initial in $\mathscr{O}(\mathscr{A})$. The second claim is immediate on observing that $(A, G) \in \mathscr{O}(\mathscr{A})$ is near-initial if and only if $(A, \mathfrak{S}_A) \in \mathscr{O}(\mathscr{A})$ is initial, if and only if $A \in \mathscr{A}$ is Galois.

For (ii), an object of $B \downarrow F'$ comprises $(A, G) \in \mathscr{O}(\mathscr{A})$ and $u: B \to FA^{\setminus FG}$ in \mathscr{B} . Now, to give u is equally to give a map $t: B \to FA$ satisfying $F\sigma.t = t$ for all $\sigma \in G$. This condition says that each $\sigma \in G$ lies in $\mathfrak{S}_t \leq \mathfrak{S}_A$, and so an object of $B \downarrow F'$ is equally a pair $(t \in \mathscr{B} \downarrow F, G \leq \mathfrak{S}_t)$. Arguing similarly on morphisms, we conclude that $B \downarrow F' \cong \mathscr{O}(B \downarrow F)$; naturality in F is straightforward. \Box

Proposition 4.12. Let \mathscr{B} admit group invariants. Under the equivalence (4.3), functors and transformations $\mathscr{A} \to \mathscr{B}$ which are analytic at stage B correspond to functors and transformations $\mathscr{O}(\mathscr{A}) \to \mathscr{B}$ which are familial, respectively near-familial at stage B. When $\mathscr{B} = \mathscr{PC}$, the equivalence (4.3) restricts to one

(4.4)
$$\operatorname{NFAM}_{0}(\mathscr{O}(\mathscr{A}), \mathscr{PC}) \xrightarrow{(-)^{\circ}} \operatorname{AN}_{0}(\mathscr{A}, \mathscr{PC})$$

with as domain the category of small pointwise familial functors and pointwise near-familial transformations $\mathcal{O}(\mathcal{A}) \to \mathcal{PC}$.

Proof. Let $F: \mathscr{A} \to \mathscr{B}$ have invariant-preserving extension $F': \mathscr{O}(\mathscr{A}) \to \mathscr{B}$. Easily $\mathscr{O}(-)$ preserves connected components, and so by Lemma 4.11, each $B \downarrow F$ is a coproduct of categories with Galois objects just when each $\mathscr{O}(B \downarrow F) \cong B \downarrow$ F' is a coproduct of categories with initial objects. Moreover, if $\alpha: F \Rightarrow G$ is a transformation between functors analytic at stage B, with invariant-preserving extension $\alpha': F' \Rightarrow G'$, then by Lemma 4.11, each functor $B \downarrow \alpha: B \downarrow F \to$ $B \downarrow G$ preserves Galois objects just when each $\mathscr{O}(B \downarrow \alpha) \cong B \downarrow \alpha'$ preserves near-initial objects. This proves the first claim.

Now suppose that $\mathscr{B} = \mathscr{PC}$. It is immediate that smallness is preserved under the preceding equivalences, and so the only additional point to verify is that $\mathbf{NFAM}_0(\mathscr{O}(\mathscr{A}), \mathscr{PC}) \subset \mathbf{INVAR}(\mathscr{O}(\mathscr{A}), \mathscr{PC})$. But if $G: \mathscr{O}(\mathscr{A}) \to \mathscr{PC}$ is pointwise familial, then each $(G)c: \mathscr{O}(\mathscr{A}) \to \mathbf{Set}$, being a coproduct of representables, preserves connected limits and in particular group invariants; whence G preserves group invariants, as limits in \mathscr{PC} are pointwise. \Box

Using this result, we may now give the promised analytic analogue of Definition 3.6, describing each small pointwise analytic $F: \mathscr{A} \to \mathscr{PC}$ in terms of the near-representable summands of each (F)c.

Definition 4.13. Let $F: \mathscr{A} \to \mathscr{PC}$ be small pointwise analytic. The *spectrum* $S_F \in \mathscr{PC}$ and *exponent* E_F : el $S_F \to \mathscr{O}(\mathscr{A})$ of F are the spectrum and exponent of the small pointwise familial $F': \mathscr{O}(\mathscr{A}) \to \mathscr{PC}$ corresponding to F under (4.4).

Remark 4.14. Let us unpack this definition. Given $F \in \mathbf{AN}_0(\mathscr{A}, \mathscr{PC})$, we choose like before a generic operation in each connected component of $y_c \downarrow F$, and write \tilde{t} for the chosen generic cover of $t \in y_c \downarrow F$. The spectrum of F is now exactly as in Definition 3.6, while the exponent E_F : el $S_F \to \mathscr{O}(\mathscr{A})$ is given on objects by $E_F(t: y_c \to TA) = (A, \mathfrak{S}_t)$, where \mathfrak{S}_t is the automorphism group of $t \in y_c \downarrow T$, or equally the set of all $\sigma \in \mathfrak{S}_A$ such that $(T\sigma)(t) = t$. To define E_F on a map $f: \tilde{ty}_f \to t$ of el S_F , we form the square

$$\begin{array}{c} \mathsf{y}_c \xrightarrow{\mathsf{y}_f} \mathsf{y}_d \\ t' = t \widetilde{\mathsf{y}}_f \bigcup_{f \in I} f A' \xrightarrow{Fu_f} FA \end{array}$$

whose lower edge is any map induced by weak initiality of \widetilde{ty}_f in $y_c \downarrow F$, and take $E_F(f) = [u_f]: (A', \mathfrak{S}_{t'}) \to (A, \mathfrak{S}_t)$. Note that the mapping $f \mapsto u_f$ is only functorial "up to automorphism groups", so that E_F may not exist as a functor $el S_F \to \mathscr{A}$.

Just as in the familial case, a small pointwise analytic $\mathscr{A} \to \mathscr{PC}$ can be recovered from its spectrum and exponent. We express this in terms of an equivalence between $\mathbf{AN}_0(\mathscr{A}, \mathscr{PC})$ and the following category:

Definition 4.15. For any \mathscr{A} and small \mathscr{C} , the category $\operatorname{el}_{\mathscr{C}}/\!\!/_{v} \mathscr{O}(\mathscr{A})$ has as objects, pairs $(S \in \mathscr{PC}, E: \operatorname{el} S \to \mathscr{O}(\mathscr{A}))$, and as maps $(S, E) \to (T, D)$, pairs of a presheaf map $p: S \to T$ and a pointwise vertical transformation φ of the form:

(4.5)
$$\operatorname{el} S \xrightarrow{\operatorname{el} p} \operatorname{el} T$$
$$\xrightarrow{\mathcal{G}} \mathcal{G}(\mathscr{A}) .$$

Proposition 4.16. The assignation $F \mapsto (S_F, E_F)$ is the action on objects of an equivalence of categories between $AN_0(\mathscr{A}, \mathscr{PC})$ and $el_{\mathscr{C}} /\!\!/_{\mathcal{V}} \mathscr{O}(\mathscr{A})$.

Proof. By Proposition 4.12, it suffices to show that $F' \mapsto (S_{F'}, E_{F'})$ underlies an equivalence of categories $\operatorname{NFAM}_0(\mathscr{O}(\mathscr{A}), \mathscr{PC}) \to \operatorname{el}_{\mathscr{C}} /\!\!/_v \mathscr{O}(\mathscr{A})$. This is almost exactly as in Proposition 3.8, with the only difference arising on morphisms. It is easy to see that for a transformation $\alpha' \colon F' \Rightarrow G'$ between pointwise familial $F' \colon \mathscr{O}(\mathscr{A}) \to \mathscr{PC}$, near-familiality is equivalent to the requirement that each triangle as to the left of (3.3) should have φ_t vertical, rather than invertible: this accounts for the differing 2-cell data between (3.2) and (4.5).

4.3. Universal analytic endofunctors. Now in seeking a universal shapely monad among analytic endofunctors, we must as before consider both composability and existence of a terminal object. This time we deal with terminality first. As in Proposition 3.16, there is a size obstruction to constructing a terminal object of the category $\mathbf{AN}_0(\mathcal{PC}, \mathcal{PC})$, and so we immediately restrict our attention to finitary ones:

Definition 4.17. If $\mathscr{A}' \subset \mathscr{A}$ and $\mathscr{B}' \subset \mathscr{B}$ are full replete subcategories, we say that $F: \mathscr{A} \to \mathscr{B}$ is $(\mathscr{A}', \mathscr{B}')$ -analytic it is analytic at each $B \in \mathscr{B}'$, and each generic $t \in B \downarrow F$ has input arity in \mathscr{A}' . A transformation $\alpha: F \Rightarrow G$ between such functors is $(\mathscr{A}', \mathscr{B}')$ -analytic if it is analytic at every stage $B \in \mathscr{B}'$.

Like before, the finitary analytic functors of the following definition are equally those which preserve filtered colimits; this now follows using Lemma 4.6.

Definition 4.18. A pointwise analytic $F: \mathscr{PC} \to \mathscr{PD}$ is called *finitary* if it is $(\mathscr{FC}, \mathscr{YD})$ -analytic. We write $\mathbf{AN}_0^{\omega}(\mathscr{PC}, \mathscr{PD})$ for the category of finitary pointwise analytic functors and pointwise analytic transformations.

This restriction is in fact enough: $\mathbf{AN}_0^{\omega}(\mathscr{PC}, \mathscr{PC})$ —and more generally, $\mathbf{AN}_0^{\omega}(\mathscr{PD}, \mathscr{PC})$ does have a terminal object. To see this, we first argue as in Lemma 3.17 to establish an equivalence between the categories $\mathbf{AN}_0^{\omega}(\mathscr{PD}, \mathscr{PC})$ and $\mathbf{AN}_0(\mathscr{FD}, \mathscr{PC})$; the claim will now follow once we show more generally that $\mathbf{AN}_0(\mathscr{A}, \mathscr{PC})$ has a terminal object whenever \mathscr{A} is essentially small. The key to proving this is the following lemma; in it, for any categories \mathscr{I} and \mathscr{A} , we write $[\mathscr{I}, \mathscr{O}(\mathscr{A})]_v$ for the category of functors $\mathscr{I} \to \mathscr{O}(\mathscr{A})$ and pointwise vertical transformations.

Lemma 4.19. Each connected component of $[\mathscr{I}, \mathscr{O}(\mathscr{A})]_v$ has an initial object.

Proof. Let $(T,G): \mathscr{I} \to \mathscr{O}(\mathscr{A})$ be given on objects by $I \mapsto (TI,G_I)$ and on morphisms by $f \mapsto [Tf]: (TI,G_I) \to (TJ,G_J)$. Note that the family of subgroups $(G_I \leq \mathfrak{S}_{TI})_{I \in \mathscr{I}}$ satisfy the condition that

(*) for all $f: I \to J$ and $\sigma \in G_J$, there exists $\tau \in G_I$ with $Tf \circ \tau = \sigma \circ Tf$.

Call a family of subgroups $H = (H_I \leq \mathfrak{S}_{TI})_{I \in \mathscr{I}}$ suitable if it satisfies (\star) with H_I and H_J in place of G_I and G_J . We claim that, if H^x is a suitable family of subgroups for each $x \in X$, then the family of subgroups $\bigvee_x H^x = (\bigvee_x H_I^x)_{I \in \mathscr{I}}$ is again suitable (here the join $\bigvee_x H_I^x$ is taken in the lattice of subgroups of \mathfrak{S}_{TI}). Indeed, if $f: I \to J$ in \mathscr{I} and $\sigma \in \bigvee_x H_J^x$, then $\sigma = \sigma_1 \cdots \sigma_n$ for some $\sigma_i \in H_J^{x_i}$; now by suitability of each H^{x_i} , there are elements $\tau_i \in H_I^{x_i}$ with $Tf.\tau_i = \sigma_i.Tf$ for each i, and so $\tau = \tau_1 \cdots \tau_n \in \bigvee_x H_I^x$ is an element with $Tf.\tau = \sigma.Tf$.

It follows that there is a largest suitable family of subgroups given by:

 $\tilde{G} = \bigvee \{ H : H \text{ is a suitable family of subgroups} \}$.

By suitability, $[Tf]: (TI, \tilde{G}_I) \to (TJ, \tilde{G}_J)$ is well-defined for each $f: I \to J$; as $G_I \leq \tilde{G}_I$, this assignation is functorial in f and so we obtain $(T, \tilde{G}): \mathscr{I} \to \mathscr{O}(\mathscr{A})$ and a vertical transformation $\xi: (T, \tilde{G}) \to (T, G)$ with components $\xi_I = [1_{TI}]$. We claim that (T, \tilde{G}) is in fact initial in its connected component.

First we show that any pair of vertical transformations $\alpha, \beta \colon (T,G) \Rightarrow (S,H)$ are equal. Each component α_I or β_I is an equivalence class of maps $TI \to SI$, and so we may consider the family of subgroups

$$(K_I = \langle a^{-1}b \mid a \in \alpha_I, b \in \beta_I \rangle \leqslant \mathfrak{S}_{TI})_{I \in \mathscr{I}} .$$

We claim this family is suitable: for then $K_I \leq \tilde{G}_I$ so that $a^{-1}b \in \tilde{G}_I$ for all $a \in \alpha_I$ and $b \in \beta_I$, whence $\alpha_I = \beta_I$ as required. For suitability, it suffices to show that, if $(a,b) \in \alpha_J \times \beta_J$ and $f: I \to J$, then there exists $(c,d) \in \alpha_I \times \beta_I$ with $Tf.c^{-1}d = a^{-1}b.Tf$. For any $c \in \alpha_I$ we have by naturality of α that

22

[a.Tf] = [Sf.c]; but then $a.Tf = Sf.(c\sigma)$ for some $\sigma \in \tilde{G}_I$, and so on replacing c by $c\sigma \in \alpha_I$ we may take it that in fact a.Tf = Sf.c. Similarly, we can find $d \in \beta_I$ such that b.Tf = Sf.d, and now $Tf.c^{-1}d = a^{-1}b.Tf$ as required.

To show initiality of (T, G) in its connected component, it now suffices to show that, for all cospans $\alpha \colon (T, \tilde{G}) \to (S, H) \leftarrow (R, K) \colon \beta$ in $[\mathscr{I}, \mathscr{O}(\mathscr{A})]_v$ there is some map $\gamma \colon (T, \tilde{G}) \to (R, K)$. To this end, consider the family of subgroups

$$(L_I = \langle a^{-1}bc^{-1}d \mid a, d \in \alpha_I, b, c \in \beta_I \rangle \leqslant \mathfrak{S}_{TI})_{I \in \mathscr{I}}$$

Repeating the above argument shows this family is suitable, and so $L_I \leq \tilde{G}_I$ for all $I \in \mathscr{I}$. Now, choosing any $a \in \alpha_I$ and $b \in \beta_I$, we have for each $I \in \mathscr{I}$ a welldefined map $\gamma_I = [b^{-1}a]: (TI, \tilde{G}_I) \to (RI, K_I)$. Indeed, if $\sigma \in K_I$ then $b\sigma \in \beta_I$ and so $\tau = a^{-1}b\sigma b^{-1}a \in L_I \leq \tilde{G}_I$ satisfies $b^{-1}a.\tau = \sigma b^{-1}a$ as required. Clearly $\beta_I.\gamma_I = \alpha_I$ for each I; since each β_I is vertical, hence monic in $\mathscr{O}(\mathscr{A})$, we conclude by naturality of α that $\gamma: (T, \tilde{G}) \to (R, K)$ is also natural as required. \Box

Using this, we are finally able to prove:

Proposition 4.20. If \mathscr{A} is essentially small, then the category $\mathbf{AN}_0(\mathscr{A}, \mathscr{PC})$ has a terminal object; in particular, any $\mathbf{AN}_0^{\omega}(\mathscr{PQ}, \mathscr{PC})$ has a terminal object.

Proof. It suffices by Proposition 4.16 to show that the equivalent category $el_{\mathscr{C}} /\!\!/_v \mathscr{O}(\mathscr{A})$ has a terminal object, and as before, we may assume without loss of generality that \mathscr{A} is in fact small. For any $c \in \mathscr{C}$, we know by Lemma 4.19 that each connected component of $[\mathscr{C}/c, \mathscr{O}(\mathscr{A})]_v$ has an initial object; make a choice of such, and for each $F \in [\mathscr{C}/c, \mathscr{O}(\mathscr{A})]_v$, write \tilde{F} for the chosen initial object in its connected component, and $u: \tilde{F} \to F$ for the unique vertical transformation. The required terminal $(S, E) \in el_{\mathscr{C}} /\!\!/_v \mathscr{O}(\mathscr{A})$ now has:

 $Sc = \{F \in [\mathscr{C}/c, \mathscr{O}(\mathscr{A})]_v : \tilde{F} = F\}$ and $S(f : d \to c) : F \mapsto \widetilde{F(f \circ -)}$, and has $E : \text{el} S \to \mathscr{O}(\mathscr{A})$ given by $E(F, c) = F(1_c)$ on objects, and

$$f: (\widetilde{F(f \circ -)}, d) \to (F, c) \quad \mapsto \quad \widetilde{F(f \circ -)}(1_d) \xrightarrow{u_{1_d}} F(f \circ -)(1_d) = Ff \xrightarrow{Ff} F1_c$$

on morphisms. To see terminality of (S, E), let (T, D) be another object of $\operatorname{el}_{\mathscr{C}}/\!\!/_{v} \mathscr{O}(\mathscr{A})$. To define a map $f: T \to S$, we form for each $t \in Tc$ the composite (4.6) $F_t: \mathscr{C}/c = \operatorname{el} \mathsf{y}_c \xrightarrow{\operatorname{el} t} \operatorname{el} T \xrightarrow{D} \mathscr{O}(\mathscr{A})$

and now define $p(t) = F_t \in Sc$. This is easily seen to be natural in c; moreover, we have a pointwise vertical transformation

$$\operatorname{el} T \xrightarrow{\operatorname{el} p} \operatorname{el} S$$

$$\xrightarrow{\varphi} \swarrow E$$

$$\mathcal{O}(\mathscr{A})$$

whose component at $t \in Tc$ is the map $u_{1_c} \colon E(p(t)) = \widetilde{F}_t(1_c) \to F_t(1_c) = Dt$. So we have a map $(p, \varphi) \colon (T, D) \to (S, E)$ and to conclude the proof, we must show that any $(q, \psi) \colon (T, D) \to (S, E)$ is equal to (p, φ) . For each $t \in Tc$, consider the composite functor

$$G_t \colon \mathscr{C}/c = \operatorname{el} \mathsf{y}_c \xrightarrow{\operatorname{el} t} \operatorname{el} T \xrightarrow{\operatorname{el} q} \operatorname{el} S \xrightarrow{E} \mathscr{O}(\mathscr{A}) \ .$$

By naturality of q, this functor sends $f: d \to c$ to $q(t)(f \circ -)(1_d) \in \mathscr{O}(\mathscr{A})$, and there is now a vertical transformation $\xi: G_t \Rightarrow q(t): \mathscr{C}/c \to \mathscr{O}(\mathscr{A})$ with component

$$G_t(f) = q(\widetilde{t})(\widetilde{f} \circ -)(1_d) \xrightarrow{u_{1_d}} q(t)(f \circ -)(1_d) = q(t)(f)$$

at $f: d \to c \in \mathscr{C}/c$. Since q(t) is a (chosen) initial object in its connected component of $[\mathscr{C}/c, \mathscr{O}(\mathscr{A})]_v$, the map $\xi: G_t \Rightarrow q(t)$ must be a split epimorphism; since every map of $[\mathscr{C}/c, \mathscr{O}(\mathscr{A})]_v$ is (pointwise monomorphic and hence) monomorphic, ξ is thus invertible, so that $G_t \cong q(t)$. Since the composite vertical transformation

(4.7)
$$\begin{array}{c} \mathscr{C}/c \xrightarrow{\operatorname{el} qt} \operatorname{el} S \\ \downarrow \psi \cdot \operatorname{el} t \\ F_{t}=D \cdot \operatorname{el} t \\ \mathscr{O}(\mathscr{A}) \end{array}$$

exhibits G_t (the upper composite) as connected to F_t in $[\mathscr{C}/c, \mathscr{O}(\mathscr{A})]_v$, this determines q(t) uniquely as being $\tilde{F}_t = p(t)$; since this holds for all $t \in \operatorname{el} T$, we conclude that p = q. Moreover, as $G_t \cong q(t)$ is initial in its connected component of $[\mathscr{C}/c, \mathscr{O}(\mathscr{A})]_v$, the 2-cell in (4.7) must be equal to $\varphi \cdot \operatorname{el} t \colon G_t \Rightarrow F_t$; as this holds for all $t \in \operatorname{el} T$, we have $\varphi = \psi$ as required.

Remark 4.21. For any small category \mathscr{C} , we may view the terminal object U of $\mathbf{AN}_0^{\omega}(\mathscr{PC},\mathscr{PC})$ as an object in $\mathbf{AN}_0(\mathscr{PC},\mathscr{PC})$. From its construction above, it is easy to see that any $F \in \mathbf{AN}_0(\mathscr{PC},\mathscr{PC})$ which admits a map to U must itself be finitary, so that the map $F \to U$ is unique if it exists. In other words, U is a *subterminal* object in $\mathbf{AN}_0(\mathscr{PC},\mathscr{PC})$; it follows that we can identify $\mathbf{AN}_0^{\omega}(\mathscr{PC},\mathscr{PC})$ with the slice category $\mathbf{AN}_0(\mathscr{PC},\mathscr{PC})/U$. We will revisit this point in Remark 6.3 below.

4.4. **Composition of analytic functors.** The passage from familial to analytic functors has thus fixed the problem we had previously, namely the lack of a terminal object among such functors. However, we are not in the clear yet, as we must still show that pointwise analytic functors compose. By modifying Lemma 3.9(i) to use Lemma 4.2 in place of Lemma 3.2 we may show that generic morphisms compose; now arguing as in Proposition 3.11 yields:

Proposition 4.22. If $F: \mathscr{A} \to \mathscr{B}$ and $G: \mathscr{B} \to \mathscr{C}$ are $(\mathscr{A}', \mathscr{B}')$ - and $(\mathscr{B}', \mathscr{C}')$ analytic, then their composite is $(\mathscr{A}', \mathscr{C}')$ -analytic, and has as generic operations at stage $C \in \mathscr{C}'$ precisely the composites $Gt.s: C \to GB \to GFA$ of Gand F-generic operations. The correspondingly analytic transformations between these functors are likewise composable; in particular, there is a 2-category **AN** of categories, analytic functors and analytic transformations.

However, this does not imply that pointwise analytic functors between presheaf categories are composable, since we do not know that pointwise analytic functors are necessarily analytic. In fact, this is not true, by virtue of:

Proposition 4.23. Pointwise analytic functors between presheaf categories are not closed under composition.

Proof. Consider the following two functors:

(4.8)
$$F: \mathbf{Set} \to \mathbf{Set}^2 \qquad G: \mathbf{Set}^2 \to \mathbf{Set} \\ X \mapsto (X^2 \to X^2/\mathfrak{S}_2) \qquad (A \to B) \mapsto A \times_B A$$

G is representable at $W = (2 \rightarrow 1)$, and so pointwise analytic; *F* is pointwise analytic with spectrum $1 \in \mathbf{Set}^2$ and exponent $\mathrm{el} 1 = \mathbf{2} \rightarrow \mathscr{O}(\mathbf{Set})$ picking out the arrow $(2, \mathfrak{S}_2) \rightarrow (2, 1)$. The composite $GF: \mathbf{Set} \rightarrow \mathbf{Set}$ sends a set *X* to

$$X^2 \times_{X^2/\mathfrak{S}_2} X^2 = \{(a, b, c, d) \in X^4 : (a, b) = (c, d) \text{ or } (a, b) = (d, c)\}.$$

Now, no operation $(a, a, a, a): 1 \to GFX$ can be generic, because the square left below has no filler; while if $a \neq b \in X$, then no (a, b, a, b) or $(a, b, b, a): 1 \to GFX$ can be generic because the square below right has no filler in either direction.

$$\begin{array}{ccc} 1 & \overbrace{(a,a,a,a,a)}^{(0,1,0,1)} GF\{0,1\} & 1 & \overbrace{(a,b,b,a)}^{(a,b,b,a)} GFX \\ \downarrow & \downarrow GF! & (a,b,a,b) & \downarrow & \downarrow GF! \\ GFX & \xrightarrow{GF!} GF1 & GFX & \xrightarrow{GF!} GF1 \end{array}$$

So GF is not pointwise analytic, as there are no generic operations in $1 \downarrow GF$. \Box

Corollary 4.24. $\mathbf{AN}_0^{\omega}(\mathscr{PC}, \mathscr{PC})$ need not be monoidal under composition.

Proof. If F and G are as in the preceding proof, then $F\pi_1$ and ΔG are easily seen to lie in $\mathbf{AN}_0^{\omega}(\mathbf{Set}^2, \mathbf{Set}^2)$. But if their composite $\Delta GF\pi_1$ were pointwise analytic, then so too would be $\pi_1(\Delta GF\pi_1)\Delta = GF$.

Remark 4.25. The preceding argument does not rule out the possibility that the composition-powers of the terminal finitary analytic endofunctor U of a presheaf category happen to be again analytic—which would allow for the construction of a monad structure on U. However, at least for the presheaf categories of our examples, it is easy to adapt the preceding argument to show that this is not so.

5. Cellular functors and shapeliness

We have now failed to construct a universal shapely monad on a presheaf category \mathscr{PC} in two different ways: there was no universal *familial* monad due to the lack of a terminal familial endofunctor, while there was no universal *analytic* monad due to the failure of pointwise analytic functors to be composition-closed.

Our next attempt to produce a universal shapely monad will focus on a special class of pointwise analytic functors, which we term *cellular*, that are closed under composition. This is achieved by way of an additional condition which allows their pointwise analyticity to be built up to analyticity at more complex stages, so that Proposition 4.22 can then be applied.

Building up this analyticity will require an analogue of Proposition 3.12, which showed that the stages of familiality of a functor $\mathscr{A} \to \mathscr{B}$ are closed under colimits. The reason this does not carry over unchanged to the analytic setting is that the analogue of Lemma 3.9(ii) fails to hold: generic operations are not closed under arbitrary colimits in $\mathscr{B} \downarrow F$. However, we will see in this section that there are certain kinds of colimit under which generic operations *are* closed, and for these, we do have an analogue of Proposition 3.12. With this result in place, we will be in a position to introduce the notion of cellular functor, this being a pointwise analytic functor whose generic operations have input arities that can be constructed out of the well-behaved kinds of colimit under which stages of analyticity are closed. This then allows us to show that cellular functors are closed under composition.

5.1. Arrow-genericity and arrow-analyticity. It is easy to see from the definitions that generic operations are closed under coproducts in $\mathscr{B} \downarrow F$, and at first this may appear to be all that we can salvage from Proposition 3.12 in the analytic case. However, there is in fact a class of morphisms in $\mathscr{B} \downarrow F$ along which generic operations are closed under pushout; we now introduce this class.

Definition 5.1. Let $F: \mathscr{A} \to \mathscr{B}$. We say that a map $(b, a): t_1 \to t_2$ in $\mathscr{B} \downarrow F$ as below is *arrow-generic* if t_1 and t_2 are *F*-generic and $(t_1, t_2): b \to F^2 a$ is generic for the functor $F^2: \mathscr{A}^2 \to \mathscr{B}^2$.

(5.1)
$$\begin{array}{c} B_1 \xrightarrow{b} B_2 \\ t_1 \downarrow \qquad \qquad \downarrow t_2 \\ FA_1 \xrightarrow{Fa} FA_2 \end{array}$$

Just as with linear and generic operations, there is a characterisation of arrowgenericity as a diagonal filling property:

Lemma 5.2. Let t_1, t_2 be F-generic operations. A map (b, a): $t_1 \rightarrow t_2$ in $\mathscr{B} \downarrow F$ is arrow-generic just when for every commuting diagram as below (with hj = ka), there exists ℓ as shown with $h\ell = k$ and $F\ell \cdot t_2 = u$ and $\ell a = j$.

(5.2)
$$\begin{array}{c} B_{1} \xrightarrow{b} B_{2} \xrightarrow{u} FY \\ \downarrow & \downarrow \\ t_{1} \downarrow & f_{2} \downarrow \\ Fj & \downarrow \\ FA_{1} \xrightarrow{F_{a}} FA_{2} \xrightarrow{Fk} FZ \end{array}$$

It follows that the class of arrow-generic maps in $\mathscr{B} \downarrow F$ contains the isomorphisms and is composition-closed.

Proof. First assume the condition in the statement. We must show that for any cube as below left, there are diagonal fillers $j_1: A_1 \to Y_1$ and $j_2: A_2 \to Y_2$ with $h_i j_i = k_i$ and $F j_i.t_i = u_i$ and $j_2 a = y j_1$. Applying genericity of t_1 to the front face yields the required j_1 ; now the left and back faces give the solid part of a diagram as in (5.2) with the composite $y j_1$ as its diagonal, and applying the stated condition to this yields a filler j_2 for the back face satisfying the required

equations.



Suppose conversely that $(b, a): t_1 \to t_2$ is arrow-generic. Given a diagram as in the solid part of (5.2), we apply arrow-genericity to the cube above right to obtain fillers for the front and back faces making everything commute. The front filler is necessarily j, and so the back filler is the $\ell: A_2 \to Y$ required for (5.2).

As mentioned above, the reason for introducing arrow-generic maps is that generic operations in $\mathscr{B} \downarrow F$ are closed under pushout along them; we show this in the next section, but first let us introduce the associated notion of analyticity.

Definition 5.3. A functor $F: \mathscr{A} \to \mathscr{B}$ is arrow-analytic at stage $b \in \mathscr{B}(B_1, B_2)$ if F is analytic at stages $B_1, B_2 \in \mathscr{B}$ and F^2 is analytic at stage $b \in \mathscr{B}^2$; we define arrow-analyticity of a transformation $\alpha: F \Rightarrow G$ correspondingly.

It should not yet be clear whether arrow-analyticity is a property that will be fulfilled in examples of interest. We will see that this is the case in Lemma 5.18 below, where we characterise arrow-generic morphisms $t_1 \rightarrow t_2$ in terms of an easily-satisfied relation between the automorphism groups of $t_1 \in B_1 \downarrow F$ and $t_2 \in B_2 \downarrow F$. Combining this with the following lemma will allow us to find many examples of arrow-analytic functors and transformations.

Lemma 5.4. A functor $F: \mathscr{A} \to \mathscr{B}$ is arrow-analytic at $b \in \mathscr{B}(B_1, B_2)$ if and only if it is analytic at $B_1, B_2 \in \mathscr{B}$ and every $(b, a): t_1 \to t_2$ in $\mathscr{B} \downarrow F$ between generic operations is arrow-generic. A transformation $\alpha: F \Rightarrow G$ between two such functors is arrow-analytic at b if and only if it is analytic at $B_1, B_2 \in \mathscr{B}$.

Proof. If F is analytic at stages $B_1, B_2 \in \mathscr{B}$, then every square as to the front of the diagram below left can be factorised through the back faces as displayed:



Here, $u_1: \tilde{t}_1 \to t_1$ and $u_2: \tilde{t}_2 \to t_2$ are generic covers obtained from analyticity at B_1 and B_2 , and a is induced by applying Lemma 4.2 to the generic \tilde{t}_1 . Now if the hypotheses in the statement hold, then $(\tilde{t}_1, \tilde{t}_2)$ is F^2 -generic and so each (t_1, t_2) in $b \downarrow F^2$ admits a generic cover, as required for F^2 to be analytic at b.

Suppose conversely that F^2 is analytic at b, and consider a square as right above with generic sides; we must show that it is arrow-generic. So construct covers

$$(\widetilde{t}_1, \widetilde{t}_2) \xrightarrow{(u_1, u_2)} (t_1, t_2) \xrightarrow{(v_1, v_2)} (s_1, s_2)$$

in $b \downarrow F^2$, where (t_1, t_2) is an F^2 -generic cover, and where $(\tilde{t}_1, \tilde{t}_2)$ is obtained as above left using analyticity of F at B_1, B_2 . Since \tilde{t}_1 and s_1 are both F-generic operations at stage B_1 , v_1u_1 is invertible by Corollary 4.3 and so u_1 is a split monomorphism; similarly u_2 is split monic. On the other hand, since (t_1, t_2) is F^2 -generic, the map (u_1, u_2) must—by Corollary 4.3 again—be a pointwise split epimorphism: whence u_1, u_2, v_1 and v_2 are invertible, so that (s_1, s_2) , like (t_1, t_2) , is arrow-generic as required. It follows that, if F is arrow-analytic at b, then the generic operations in $b \downarrow F^2$ are precisely the squares with generic sides; the statement about arrow-analytic transformations follows directly from this. \Box

In the sequel, we will make use of this characterisation of arrow-genericity without further comment. We conclude this section by recording the analogue of Definitions 3.10 and 4.17 for arrow-analytic functors:

Definition 5.5. If $\mathscr{I} \subset \mathscr{A}^2$ and $\mathscr{J} \subset \mathscr{B}^2$ are full replete subcategories, we say that $F: \mathscr{A} \to \mathscr{B}$ is $(\mathscr{I}, \mathscr{J})$ -arrow-analytic if it is arrow-analytic at each $b \in \mathscr{J}$, and each arrow-generic $(b, a): t_1 \to t_2$ as in (5.1) with $b \in \mathscr{J}$ has $a \in \mathscr{I}$. A transformation $\alpha: F \Rightarrow G$ between such functors is $(\mathscr{I}, \mathscr{J})$ -arrow-analytic if it is arrow-analytic at every stage $b \in \mathscr{J}$.

Proposition 5.6. If $F: \mathscr{A} \to \mathscr{B}$ and $G: \mathscr{B} \to \mathscr{C}$ are $(\mathscr{I}, \mathscr{J})$ - and $(\mathscr{J}, \mathscr{K})$ -arrow-analytic, then their composite is $(\mathscr{I}, \mathscr{K})$ -arrow-analytic, and correspondingly for the transformations between such functors.

Proof. Direct from the definitions and Proposition 4.22.

5.2. Building up stages of analyticity. We are now ready to see what the purpose of arrow-genericity and arrow-analyticity really is. We begin with a lemma which provides an analytic analogue of Lemma 3.9(ii) above.

Lemma 5.7. Let $F: \mathscr{A} \to \mathscr{B}$ and consider a pointwise pushout square in $\mathscr{B} \downarrow F$ as below. If s_1 , s_2 and t_1 are generic and (d, c) is arrow-generic, then t_2 is also generic and (b, a) is also arrow-generic.

(5.3)
$$s_1 \xrightarrow{(g,f)} t_1$$
$$(d,c) \downarrow \qquad \qquad \downarrow^{(b,a)}$$
$$s_2 \xrightarrow{(m,n)} t_2$$

Proof. We will show that every diagram as in the solid part of (5.2) admits a dotted filler; since t_1 is generic, this immediately implies that t_2 is generic, and so by Lemma 5.2 that (b, a) is arrow-generic. To prove the claim, observe that the stated filling condition can be described as a *left lifting property*: it says that, for each $h: Y \to Z$ in \mathscr{A} , each square in $\mathscr{B} \downarrow F$ as left below has a diagonal

filler.

(5.4)
$$\begin{array}{c} t_1 \xrightarrow{(ub,j)} 1_{FY} & s_1 \xrightarrow{(ubg,jf)} 1_{FY} \\ (b,a) \downarrow & \downarrow (u,\ell) \\ t_2 \xrightarrow{(u,k)} Fh & s_2 \xrightarrow{(udg,\ell)} Fh \end{array}$$

Pasting the given square with the pushout (5.3) gives a square as right above; since (d, c) is arrow-generic, we induce a filler for this square as indicated and so by the universal property of pushout the required filler (u, ℓ) as left above. \Box

In fact, we can do better than this: the characterisation of arrow-generic maps by a left lifting property allows us to show that they are also closed under *transfinite composition* [13, Definition 2.1.1]. As we do not need this further fact, we leave its verification to the interested reader.

We now use the preceding lemma to give the promised analytic analogue of Proposition 3.12.

Proposition 5.8. Let \mathscr{A} be cocomplete and let $F \colon \mathscr{A} \to \mathscr{B}$.

- (i) If F is analytic at B, then it is arrow-analytic at 1_B ; if F is arrow-analytic at composable maps b and c, then it is also arrow-analytic at cb.
- (ii) For any pushout as below in \mathscr{B} , if F is analytic at B_1 , B_2 and C_1 and arrow-analytic at b, then it is also analytic at C_2 and arrow-analytic at c.

$$(5.5) \qquad \begin{array}{c} B_1 \xrightarrow{f_1} C_1 \\ b \\ B_2 \xrightarrow{f_2} C_2 \end{array}$$

The analogous results hold for natural transformations $\alpha \colon F \Rightarrow G \colon \mathscr{A} \to \mathscr{B}$.

Again, we could add an additional clause to this proposition showing closure of stages of arrow-analyticity under transfinite composition, but we refrain from doing so as we have no use for it in what follows.

Proof. Let F be analytic at B. Any map $(1_B, a): t_1 \to t_2$ between generic operations in $\mathscr{B} \downarrow F$ is a map in $B \downarrow F$, so that f is invertible by Corollary 4.3 and $(1_B, f)$ is arrow-generic by Lemma 5.2. This shows that B is arrow-analytic at 1_B . Suppose next that F is arrow-analytic at $b: B_1 \to B_2$ and $c: B_2 \to B_3$. Given a square as below left with t_1 and t_3 generic, let $e: t_2 \to t_3 c$ be a generic cover in $B_2 \downarrow F$ and let d be induced by genericity as centre below; this yields a factorisation of the left square as to the far right.

$$\begin{array}{cccc} B_1 & \xrightarrow{cb} B_3 & B_1 & \xrightarrow{t_2b} FA_2 & B_1 & \xrightarrow{b} B_2 & \xrightarrow{c} B_3 \\ t_1 & \downarrow & \downarrow t_3 & t_1 & \downarrow & \overbrace{Fd}^{\times} \downarrow Fe & t_1 & \downarrow & t_2 & \downarrow & \downarrow t_3 \\ FA_1 & \xrightarrow{Fa} FA_3 & FA_1 & \xrightarrow{Fa} FA_3 & FA_1 & \xrightarrow{Fd} FA_2 & \xrightarrow{Fe} FA_3 \end{array}$$

By arrow-analyticity at b and c, both small squares are arrow-generic, whence also their composite by Lemma 5.2; this shows that B is arrow-analytic at cb as required for (i). Now suppose the hypotheses of (ii). We first show that F is analytic at C_2 . Let $z: C_2 \to FZ$, and consider the left cube in:



The front, left and right faces arise from generic covers $m: t_2 \to zf_2$, $a: t_1 \to t_2b$ and $n: u_1 \to zc$, while the map g_1 across the back face is obtained as in (i) using genericity of t_1 . Since the top face is a pushout, and \mathscr{A} is cocomplete, the back and left faces admit a pushout in $\mathscr{B} \downarrow F$ which may be taken to be as right above. Since t_1 and t_2 are generic and F is arrow-analytic at b, the map $(b, a): t_1 \to t_2$ is arrow-generic; since u_1 is also generic, we conclude by Lemma 5.7 that u_2 is generic and $(c, y): u_1 \to u_2$ is arrow-generic.

Now taking $w: Y_2 \to Z$ to be the unique map with wy = n and $wg_2 = m$, we see that $w: u_2 \to z$ provides a generic cover of z in $C_2 \downarrow F$, so that F is analytic at C_2 as required. For arrow-analyticity at c, suppose that $(c, n): u_1 \to z$ is a map between generic operations in $\mathscr{B} \downarrow F$. We may complete this to a cube as left above and form the generic pushout u_2 as to the right. Now since z is generic, the induced map $u_2 \to z$ in $C_2 \downarrow F$ is invertible by Corollary 4.3. So the left cube above is also a pushout; as $(b, a): t_1 \to t_2$ is arrow-generic, so too is $(c, n): u_1 \to z$ by Lemma 5.7.

5.3. Cellular analytic functors. By using Proposition 5.8, we can now build up the analyticity of a pointwise analytic functor between presheaf categories to analyticity at more complex stages by assuming suitable instances of arrow-analyticity. In order to specify what these more complex stages are, we borrow some ideas from algebraic topology, in particular the theory of *cell complexes* in model categories; see [13, §2.1.2], for example.

Definition 5.9. Let *I* be a class of maps in a category \mathscr{C} with an initial object. A map $f: X \to Y$ is called a *finite relative I-complex* if either it is an isomorphism, or it can be written as a finite composite

(5.7)
$$X = X_0 \xrightarrow{f_1} X_1 \to \dots \xrightarrow{f_m} X_m = Y$$

where each f_i is a pushout of a map in I. An object $Y \in \mathscr{C}$ is called a *finite* I-complex if the unique map $0 \to Y$ is a finite relative I-complex. We write $\mathbf{Cx}(I) \subset \mathscr{C}$ for the full subcategory on the finite I-complexes, and $\mathbf{Cx}^2(I) \subset \mathscr{C}^2$ for the full subcategory on the relative finite I-complexes between finite I-complexes.

The modifier "finite" here comes from the fact that in (5.7) we allow only finite compositions; the general notion of cell complex in topology also allows for *transfinite* ones, and everything that follows could be adapted to this greater generality; however, like before, we refrain from giving this as we will not need it.

Definition 5.10. Let \mathscr{C} be a small category. A *bordage* on $\mathscr{P}\mathscr{C}$ is a filtered family $\mathscr{Q} = I_0 \subset I_1 \subset \cdots \subset \bigcup_n I_n = I$ of maps in $\mathscr{P}\mathscr{C}$ such that:

(i) Each $g \in I_{n+1}$ has representable codomain and domain a finite I_n -complex;

(ii) Each representable is a finite *I*-complex.

Note that condition (i) for a bordage ensures that each map in I has domain a finite I-complex and, as such, is an object of $\mathbf{Cx}^2(I)$; this is something which need not be true for a general class of maps I. Condition (ii) is much less important than (i) and will only play a role in Proposition 5.15 below.

- **Examples 5.11.** (i) Any presheaf category \mathscr{PC} has a bordage given by $I = I_1 = \{ 0 \rightarrow y_c : c \in \mathscr{C} \}$. The finite *I*-complexes are the finite coproducts of representables, and the finite relative *I*-complexes are coproduct injections with complement a finite *I*-complex.
 - (ii) Let **2** be the arrow category $f: 0 \to 1$. The presheaf category $\mathscr{P}\mathbf{2}$ has a bordage given by $I_1 = \{0 \to y_0\}$ and $I \setminus I_1 = \{y_f: y_0 \to y_1\}$. The finite *I*-complexes are all finitely presentable presheaves, and the finite relative *I*-complexes are the monomorphisms with cofinite image.
- (iii) Let \mathscr{G} be the category $s, t: 0 \Rightarrow 1$. The presheaf category $\mathscr{P}\mathscr{G}$ has a bordage given by $I_1 = \{0 \to y_0\}$ and $I \setminus I_1 = \{\langle y_s, y_t \rangle : y_0 + y_0 \to y_1\}$, whose finite *I*-complexes and finite relative *I*-complexes are as in (ii).
- (iv) Changing $I \setminus I_1$ in the preceding example to be $\{y_t : y_0 \to y_1\}$ yields another bordage on \mathscr{PG} whose finite *I*-complexes are now finite *forests* whose edges are all directed towards the roots. Changing $I \setminus I_1$ to be $\{y_s : y_0 \to y_1\}$ yields finite forests with edges directed away from the roots, while taking $I \setminus I_1$ to be $\{y_s, y_t\}$ yields finite forests whose edges may be oriented arbitrarily.

We will see further examples of bordages when we revisit the motivating examples of polycategories, properads and PROPs in Section 6 below.

Definition 5.12. Let I be a bordage on \mathscr{PC} . A pointwise analytic $F : \mathscr{A} \to \mathscr{PC}$ is *I*-cellular if any square as below with $b \in I$ and t_1, t_2 generic is arrow-generic.

(5.8)
$$\begin{array}{c} S \xrightarrow{b} y_c \\ t_1 \downarrow \qquad \qquad \downarrow t_2 \\ FA_1 \xrightarrow{Fa} FA_2 \end{array}$$

We will see in the following section that the cellularity condition is very easy to check in practice. Note that cellularity *almost* says that F is arrow-analytic at $b: S \to y_c$ for each $b \in I$, except that we do not assume that F is analytic at the domain object S. In fact, this is true by virtue of:

Proposition 5.13. Let I be a bordage on \mathscr{PC} and let \mathscr{A} be a cocomplete category. Any I-cellular $F: \mathscr{A} \to \mathscr{PC}$ is analytic at all $B \in \mathbf{Cx}(I)$ and arrow-analytic at all $b \in \mathbf{Cx}^2(I)$. The same holds for pointwise analytic transformations $\alpha: F \Rightarrow G$ between I-cellular functors.

Proof. Let F be *I*-cellular. We prove by induction on n that F is analytic at every $B \in \mathbf{Cx}(I_n)$ and $b \in \mathbf{Cx}^2(I_n)$. For the base case n = 0, every $B \in \mathbf{Cx}(I_0)$ is initial: thus $B \downarrow F \cong \mathscr{A}$, and so as \mathscr{A} has an initial object, F is analytic at B. Since any $b \in \mathbf{Cx}^2(I_0)$ is invertible, F is analytic at b by Proposition 5.8.

Now assume the result for n. Each map $b: S \to y_c$ in I_{n+1} has domain in $\mathbf{Cx}(I_n)$, and so F is analytic at S; thus I-cellularity implies that F is arrowanalytic at every $b \in I_{n+1}$. Applying Proposition 5.8 finitely many times shows that, if $f: X \to Y$ is a finite relative I_{n+1} -cell complex for which F is analytic at X, then F is also analytic at Y and arrow-analytic at f. Taking X to be initial and using the base case, shows that F is analytic at every $B \in \mathbf{Cx}(I_{n+1})$; while taking X to be an arbitrary finite I_{n+1} -complex shows that F is arrowanalytic at every $b \in \mathbf{Cx}^2(I_{n+1})$. The case of transformations is similar, and so omitted. \Box

The preceding proposition shows us that the pointwise analyticity of functors $F: \mathscr{PC} \to \mathscr{PD}$ and $G: \mathscr{PD} \to \mathscr{PE}$ is stable under composition if there is a bordage I on \mathscr{PD} such that F is I-cellular and the input arities of G's generic operations are I-cell complexes. However, GF need not then satisfy any cellularity conditions allowing it to compose further; the following definition ensures this.

Definition 5.14. Let I and J be bordages on \mathscr{PC} and \mathscr{PD} . A pointwise analytic functor $F: \mathscr{PC} \to \mathscr{PD}$ is called (I, J)-cellular if every square (5.8) with $b \in J$ and t_1, t_2 generic is arrow-generic and has a a finite relative I-complex. We write $\mathbf{CELL}((\mathscr{C}, I), (\mathscr{D}, J))$ for the category of (I, J)-cellular functors and pointwise analytic transformations.

Proposition 5.15. Let I and J be bordages on \mathscr{PC} and \mathscr{PD} . A functor $F: \mathscr{PC} \to \mathscr{PD}$ is (I, J)-cellular if and only if it is $(\mathbf{Cx}^2(I), \mathbf{Cx}^2(J))$ -arrow-analytic. In particular, there is a 2-category **CELL** of presheaf categories equipped with bordages, cellular analytic functors, and pointwise analytic transformations.

Proof. By condition (ii) for a bordage, each representable in \mathscr{PD} is a finite J-complex, whence any $(\mathbf{Cx}^2(I), \mathbf{Cx}^2(J))$ -arrow-analytic functor is pointwise analytic. By condition (i), we have $J \subset \mathbf{Cx}^2(J)$, and so any $(\mathbf{Cx}^2(I), \mathbf{Cx}^2(J))$ -arrow-analytic functor is (I, J)-cellular. This proves the "if" direction. For the "only if", if F is (I, J)-cellular, then it is certainly J-cellular, and so arrow-analytic at each $b \in \mathbf{Cx}^2(J)$; it remains to show that each square (5.1) with $b \in \mathbf{Cx}^2(J)$ and t_1, t_2 generic has $a \in \mathbf{Cx}^2(I)$. Let K denote the collection of all arrows $b \in \mathbf{Cx}^2(J)$ for which each square (5.1) with t_1, t_2 generic has a a finite relative I-complex. By assumption $J \subset K$, and by examining the proof of Proposition 5.8 we see that K is stable under pushout and closed under composition, and so must comprise all of $\mathbf{Cx}^2(J)$. So each (5.1) with $b \in \mathbf{Cx}^2(J)$ and t_1, t_2 generic has a a relative finite I-complex, and it remains to show that the domain and codomain

of a are in fact finite *I*-complexes. But whenever $B \in \mathbf{Cx}(J)$ and $t: B \to FA$ is generic, the following square has top edge in $\mathbf{Cx}^2(J)$ and generic sides:



and so has bottom map a finite relative *I*-complex; whence $A \in \mathbf{Cx}(I)$ as required. The existence of the 2-category **CELL** now follows from this together with Proposition 5.6.

5.4. A combinatorial characterisation of arrow-genericity. We now know that cellular pointwise analytic functors are closed under composition; what we have not yet seen is that there *are* any cellular functors. In this section, we give a result which will allow us to verify that a given pointwise analytic functor is indeed cellular. The key concept required is that of a *minimal extension*.

Definition 5.16. Let $f: A \to B$ and let $\sigma \in \mathfrak{S}_A$. An extension of σ along f is some $\tau \in \mathfrak{S}_B$ for which $\tau f = f\sigma$. An extension is called *minimal* if whenever $g: B \to C$ satisfies $gf\sigma = gf$, also $g\tau = \tau$; equivalently, if $f, f\sigma: A \rightrightarrows B$ admit a coequaliser q, then τ is minimal just when $q\tau = q$.

The name is motivated by the case of $f: A \to B$ an injection in **Set**; for such an f, any $\sigma \in \mathfrak{S}_A$ is easily seen to have a unique minimal extension $\tau \in \mathfrak{S}_B$ with

(5.9)
$$\tau(x) = \begin{cases} \sigma(x) & \text{if } x \in \text{Im}(f); \\ x & \text{otherwise.} \end{cases}$$

So τ is minimal among extensions of σ in that it permutes the smallest possible part of B. This intuition works for monomorphisms in any presheaf category:

Proposition 5.17. If $f: A \rightarrow B$ is monic in \mathscr{PC} then each $\sigma \in \mathfrak{S}_A$ admits at most one minimal extension along f; this extension exists just when

(5.10)
$$x \notin f(A) \text{ and } xh \in f(A) \implies xh \in f(A^{\setminus \sigma})$$

for all $x \in Bc$ and $h: d \to c$, and is then given componentwise as in (5.9).

Proof. If q coequalises f and $f\sigma$, then $\tau \in \mathfrak{S}_B$ is a minimal extension of σ just when $q\tau = \tau$, just when $q_c\tau_c = \tau_c$ for all c, just when each τ_c is a minimal extension of σ_c (as colimits in \mathscr{PC} are pointwise). So any minimal extension of σ must be given componentwise by (5.9), with (5.10) being just what is needed to ensure naturality of this definition in c.

We now use the notion of minimal extension to give a combinatorial characterisation of arrow-generic morphisms. As in Remark 4.14, we use \mathfrak{S}_{t_1} and \mathfrak{S}_{t_2} to denote the automorphism groups of $t_1 \in B_1 \downarrow F$ and $t_2 \in B_2 \downarrow F$.

Lemma 5.18. Let \mathscr{A} have coequalisers and $F : \mathscr{A} \to \mathscr{B}$. A map $(b, a) : t_1 \to t_2$ between generic operations in $\mathscr{B} \downarrow F$ is arrow-generic if and only if each $\sigma \in \mathfrak{S}_{t_1}$ admits a minimal extension $\tau \in \mathfrak{S}_{t_2}$ along $a : A_1 \to A_2$. Proof. We use the alternate characterisation of arrow-genericity of Lemma 5.2. For the "if" direction, suppose given a diagram as in the solid part of (5.2). As t_2 is generic, there exists a map $g: A_2 \to Y$ with hg = k and $Fg.t_2 = u$; now both ga and j are maps $t_1 \to ub$ in $B_1 \downarrow F$ and so by genericity of t_1 there is some $\sigma \in \mathfrak{S}_{t_1}$ with $j = ga\sigma$. By the assumption on a, there is some minimal $\tau \in \mathfrak{S}_{t_2}$ with $\tau a = a\sigma$; letting $\ell = g\tau$, we have $\ell a = g\tau a = ga\sigma = j$ and $F\ell.t_2 = Fg.F\tau.t_2 = Fg.t_2 = u$. Now $ka\sigma = hga\sigma = hj = ka$, whence $k\tau = k$ by minimality of τ , and so $h\ell = hg\tau = k\tau = k$ as required for ℓ to be a filler.

For the "only if" direction, let $(b, a): t_1 \to t_2$ be arrow-generic and let $\sigma \in \mathfrak{S}_{t_1}$; we must find a minimal extension $\tau \in \mathfrak{S}_{t_2}$ along a. Let $q: A_2 \to Q$ be a coequaliser of a and $a\sigma$, and consider the diagram



The solid part clearly commutes, and so we induce a map τ as displayed making everything commute. Since t_2 is generic, τ is invertible by Corollary 4.3; since $F\tau t_2 = t_2$, we have $\tau \in \mathfrak{S}_{t_2}$. Moreover, $\tau a = a\sigma$, so τ is an extension of σ ; while $q\tau = q$ so that τ is minimal.

This result allows us to check in a concrete fashion the (I, J)-cellularity of a pointwise analytic $F: \mathscr{PC} \to \mathscr{PD}$. For this, it suffices to check (I, J_n) cellularity for each n. This is trivial for n = 0 since $J_0 = \emptyset$; so suppose now that we have verified it up to n. To check (I, J_{n+1}) -cellularity, we must show that, for any $b: S \to y_d$ in $J_{n+1} \setminus J_n$, each square (5.1) with generic sides is arrow-generic with a a finite relative I-complex. We can do this using the previous result so long as we can compute all such squares (5.1).

Now, as F is (I, J_n) -cellular, it is by Proposition 5.13 analytic at the finite J_n complex S. Thus, each square (5.1) is obtained from a generic $t_2: y_d \to FA_2$ —
which we can classify by pointwise analyticity of F—upon forming a generic
cover $a: t_1 \to t_2 b$ in $S \downarrow F$, which can be calculated explicitly by applying the
algorithm of Proposition 5.8 to some presentation of S as a finite J_n -complex.

Example 5.19. Let J be the bordage of Examples 5.11(ii) on $\mathscr{P}\mathbf{2}$, let I be any bordage on \mathscr{PC} , and let $F: \mathscr{PC} \to \mathscr{P}\mathbf{2}$ be pointwise analytic. Recalling that $J_1 = \{0 \to y_0\}$, the condition for F to be (I, J_1) -cellular is that, for each generic $t: y_0 \to FA$, the induced square



with generic sides is arrow-generic with a a finite relative *I*-complex. Arrow-genericity is trivial in this case, and so the condition is simply that A is a

finite *I*-complex. Now, since $J_2 \setminus J_1 = \{y_f : y_0 \to y_1\}$, we see that *F* will be $(I, J_2) = (I, J)$ -cellular when, for each generic $t_2 : y_0 \to FB$, the induced square

$$\begin{array}{c} y_0 \xrightarrow{y_f} y_1 \\ t_1 \downarrow & \downarrow t_2 \\ FA \xrightarrow{Fa} FB \end{array}$$

with generic sides is arrow-generic and has a a finite relative *J*-complex. Let us see what this says relative to an explicit presentation of F as a pointwise coproduct of near-representables:

$$F \qquad = \qquad \sum_{u \in U} \mathscr{PC}(B_u, -)_{/G_u} \xrightarrow{\alpha} \sum_{v \in V} \mathscr{PC}(A_v, -)_{/H_v} \; .$$

Here, the map α is determined by a function $h: U \to V$ together with natural transformations $\alpha_u: \mathscr{PC}(B_u, -)_{/G_u} \to \mathscr{PC}(A_{hu}, -)_{/H_{hu}}$ —which, as in the proof of Proposition 4.9, correspond to maps $[a_u]: (A_{hu}, H_{hu}) \to (B_u, G_u)$ in $\mathscr{O}(\mathscr{PC})$ (note that this is really just an explicit description of the exponent of F). In these terms, the necessary conditions for F to be (I, J)-cellular are that: each A_v should be a finite I-complex; each $a_u: A_{hu} \to B_u$ should be a finite relative I-complex; and each $\sigma \in H_{hu}$ should have a minimal extension $\tau \in G_u$ along a_u .

5.5. Universal cellular functors. We have now achieved what we set out to do in this section, by exhibiting a class of pointwise analytic functors which is closed under composition. Our broader objective, recall, is to construct a universal shapely monad as a terminal object among a suitable class of composable endo-functors, and it may appear that we have now achieved this. However, we are not yet done, as we must still check the existence of a terminal object among cellular functors. Unfortunately, we have:

Proposition 5.20. CELL($(\mathscr{C}, I), (\mathscr{D}, J)$) need not admit a terminal object.

Proof. Let $\mathscr{C} = \mathscr{D} = \mathbf{2}$ and let both I and J be the bordage $\{0 \to \mathsf{y}_0, \mathsf{y}_0 \to \mathsf{y}_1\}$ of Examples 5.11(ii). Consider the endofunctor $F: \mathscr{P}\mathbf{2} \to \mathscr{P}\mathbf{2}$ sending $f: A \to B$ to $f \times f: A \times A \to B \times B$; this is pointwise analytic, with explicit presentation

$$F \qquad = \qquad \mathscr{P} \mathbf{2}(\mathsf{y}_1 + \mathsf{y}_1, -)_{/1} \xrightarrow{\mathscr{P} \mathbf{2}(\mathsf{y}_f + \mathsf{y}_f, -)} \mathscr{P} \mathbf{2}(\mathsf{y}_0 + \mathsf{y}_0, -)_{/1} \; .$$

To see that F is (I, I)-cellular, we observe that $y_0 + y_0$ is a finite I-complex (= finitely presentable presheaf), that $y_f + y_f$ is a relative finite I-complex (= monomorphism with cofinite image), and that the minimal extension condition is trivially satisfied: this verifies the three conditions of Example 5.19, as required.

In the terminology of Definition 4.13, F has spectrum $S_F = 1 \in \mathscr{P}\mathbf{2}$ —so that el $S_F \cong \mathbf{2}$ —and exponent $E_F \colon \mathbf{2} \to \mathscr{O}(\mathscr{P}\mathbf{2})$ picking out the arrow

$$[y_f + y_f]: (y_0 + y_0, 1) \to (y_1 + y_1, 1)$$

of $\mathscr{O}(\mathscr{P}\mathbf{2})$. It follows using Proposition 4.16 that for each $G \in \mathbf{AN}_0(\mathscr{P}\mathbf{2}, \mathscr{P}\mathbf{2})$, pointwise analytic transformations $\alpha \colon F \Rightarrow G$ correspond bijectively with squares

(5.11)
$$\begin{array}{c} y_0 \xrightarrow{y_f} y_1 \\ t_1 \downarrow & \downarrow t_2 \\ G(y_0 + y_0) \xrightarrow{G(y_f + y_f)} G(y_1 + y_1) \end{array}$$

in $\mathscr{P}\mathbf{2}$ with generic sides. We claim that whenever G is (I, I)-cellular, there are always two distinct such squares, so that G cannot be terminal in $\mathbf{CELL}((\mathbf{2}, I), (\mathbf{2}, I))$. Since G was arbitrary, this concludes the proof. Now, if G is (I, I)-cellular, then any square as displayed above must be arrow-generic; by Lemma 5.18, this means that each $\sigma \in \mathfrak{S}_{t_1}$ admits a minimal extension $\tau \in \mathfrak{S}_{t_2}$ along $y_f + y_f$. Using Proposition 5.17, we see that the switch isomorphism $\sigma_{12} \colon y_0 + y_0 \to y_0 + y_0$ has no minimal extension along $y_f + y_f$, and so we must have $\mathfrak{S}_{t_1} = 1$. Since $[y_f + y_f] \colon (y_0 + y_0, \mathfrak{S}_{t_1}) \to (y_1 + y_1, \mathfrak{S}_{t_2})$ in $\mathscr{O}(\mathscr{P}\mathbf{2})$, it follows that $\mathfrak{S}_{t_2} = 1$ too; whence the square

$$\begin{array}{c} \mathsf{y}_{0} \xrightarrow{\mathsf{y}_{f}} \mathsf{y}_{1} \\ G\sigma_{12}.t_{1} \downarrow \qquad \qquad \downarrow G\sigma_{12}.t_{2} \\ G(\mathsf{y}_{0} + \mathsf{y}_{0}) \xrightarrow{G(\mathsf{y}_{f} + \mathsf{y}_{f})} G(\mathsf{y}_{1} + \mathsf{y}_{1}) \end{array}$$

is a second, distinct instance of (5.11). This proves the claim.

6. Shapeliness in context

We have now failed for a third time to exhibit a notion of universal shapely monad: the imposition of cellularity, which fixed the failure of general analytic functors to compose, did so at the cost of destroying the terminal object existing among them. At this point, we prefer to leave for future work the problem of finding a general notion of shapeliness, and concentrate instead on giving a solution for the particular motivating examples from Section 2.

For these examples, the notion of cellularity turns out to be almost sufficient: some simple *ad hoc* additional conditions will be enough to obtain the desired universal shapely monad U. With this in place, we can define a general shapely monad to be any pointwise analytic submonad of U, and then provide an inductive construction of the free shapely monad on a generating set of operations. Finally, we apply this construction to our motivating examples, and thereby realise the main goal of this paper by exhibiting the monads for polycategories, properads and PROPs as free shapely monads on the basic wiring operations.

6.1. Universal shapely monads on (symmetric) polygraphs. Our first goal is to construct universal shapely monads on the presheaf categories of polygraphs and symmetric polygraphs from Section 2 which are suitable for analysing polycategories, properads and PROPs. We will find these universal monads among the class of cellular analytic endofunctors of the previous section; but in order to neutralise the counterargument of Proposition 5.20, we must further restrict the functors under consideration. We build on the observation that the monads in our examples act trivially on the set of objects of a (symmetric) polygraph.

Definition 6.1. An endofunctor F of either $\mathscr{P}\mathsf{P}$ or $\mathscr{P}\mathsf{P}_s$ is called *framed* if $FX(\star) \cong A \times X(\star)$ for some fixed set A.

Of course, "acting trivially" on objects is only the special case A = 1 of this definition. The reason for allowing the more general definition is to ensure that any pointwise analytic functor admitting a pointwise analytic transformation to a framed one is itself framed; see Remark 6.3 below.

Since framed functors are clearly stable under composition, we have for any bordage I on $\mathscr{P}\mathsf{P}$ or $\mathscr{P}\mathsf{P}_s$ a monoidal category of framed (I, I)-cellular endofunctors. In both the symmetric and non-symmetric cases, we are free to choose I in any way which ensures that the monads for polycategories, properads and PROPs are all in fact (I, I)-cellular; we now describe suitable such choices, and check by hand that corresponding monoidal categories of framed (I, I)-cellular endofunctors have a terminal object—so giving the desired universal shapely monads. Let us begin with the case of $\mathscr{P}\mathsf{P}$. The bordage I_P we take has $(I_\mathsf{P})_1 = \{0 \to y_\star\}$ and $I_\mathsf{P} \setminus (I_\mathsf{P})_1$ the set of the following maps for all $n, m \in \mathbb{N}$:

$$\langle \mathsf{y}_{\sigma_1}, \ldots, \mathsf{y}_{\sigma_n} \rangle \colon \mathsf{y}_{\star} + \cdots + \mathsf{y}_{\star} \to \mathsf{y}_{(n,m)} \text{ and } \langle \mathsf{y}_{\tau_1}, \ldots, \mathsf{y}_{\tau_m} \rangle \colon \mathsf{y}_{\star} + \cdots + \mathsf{y}_{\star} \to \mathsf{y}_{(n,m)}$$
.

We now show that there is a universal framed $(I_{\mathsf{P}}, I_{\mathsf{P}})$ -cellular endofunctor. We make use of the sets $\mathscr{L}(n,m)$ of (n,m)-labelled finite polygraphs from Definition 2.3 above. We will call $X \in \mathscr{L}(n,m)$ well-labelled if the maps

$$\langle \ell_1^X, \dots, \ell_n^X \rangle \colon \mathsf{y}_{\star} + \dots + \mathsf{y}_{\star} \to |X| \quad \text{and} \quad \langle r_1^X, \dots, r_m^X \rangle \colon \mathsf{y}_{\star} + \dots + \mathsf{y}_{\star} \to |X|$$

are both relative finite I_{P} -complexes; and, extending the notation of Proposition 2.8, we write \mathfrak{S}_X for the set of label-preserving automorphisms of |X|.

Proposition 6.2. The monoidal category of framed $(I_{\mathsf{P}}, I_{\mathsf{P}})$ -cellular endofunctors of $\mathscr{P}\mathsf{P}$ has a terminal object U_{P} , which thus underlies a monad on $\mathscr{P}\mathsf{P}$, the universal shapely monad. The spectrum $S \in \mathscr{P}\mathsf{P}$ of U_{P} may be taken to be:

$$S(\star) = \{u\}$$
 and $S(n,m) = \{X \in \mathcal{L}(n,m) : X \text{ is well-labelled}\}$

and the exponent $E \colon \operatorname{el} S \to \mathscr{O}(\mathscr{P}\mathsf{P})$ to have $E(u) = \mathsf{y}_{\star}$ and

$$E(X) = (|X|, \mathfrak{S}_X), \qquad E(\sigma_i \colon u \to X) = [\ell_i^X], \qquad E(\tau_j \colon u \to X) = [r_i^X].$$

Proof. By Proposition 4.16, we have $\mathbf{AN}_0(\mathscr{P}\mathsf{P}, \mathscr{P}\mathsf{P}) \simeq \operatorname{el}_{\mathsf{P}} /\!\!/_v \mathscr{O}(\mathscr{P}\mathsf{P})$; if we define $(\operatorname{el}_{\mathsf{P}} /\!\!/_v \mathscr{O}(\mathscr{P}\mathsf{P}))' \subset \operatorname{el}_{\mathsf{P}} /\!\!/_v \mathscr{O}(\mathscr{P}\mathsf{P})$ to be the full subcategory corresponding under this equivalence to the full subcategory of framed $(I_{\mathsf{P}}, I_{\mathsf{P}})$ -cellular endofunctors, then it suffices to show that (S, E) as defined above is terminal in this category.

First, let us call a functor $\mathsf{P}/\star \cong 1 \to \mathscr{O}(\mathscr{P}\mathsf{P})$ acceptable if it picks out the object $(\mathsf{y}_{\star}, 1)$, and a functor $\mathsf{P}/(n, m) \to \mathscr{O}(\mathscr{P}\mathsf{P})$ acceptable if it takes the form



with (X, ℓ, r) a well-labelled polygraph. By an argument like Example 5.19 above, a pointwise analytic $F: \mathscr{P}\mathsf{P} \to \mathscr{P}\mathsf{P}$ is framed $(I_\mathsf{P}, I_\mathsf{P})$ -cellular just when, for each element $t \in S_F(x)$ of its spectrum, the composite

$$F_t \colon \mathsf{P}/x = \operatorname{el} \mathsf{y}_x \xrightarrow{\operatorname{el} t} \operatorname{el} S_F \xrightarrow{E_F} \mathscr{O}(\mathscr{P}\mathsf{P})$$

is acceptable. Noting that this F_t is the same as (4.6) appearing in the proof of Proposition 4.20, we thus continue by emulating the rest of that proof.

Recall the key Lemma 4.19 stating that, for each $x \in \mathsf{P}$, the connected component of any $F \in [\mathsf{P}/x, \mathscr{O}(\mathscr{P}\mathsf{P})]_v$ contains an initial object \tilde{F} . We claim that, if F is acceptable, then so too is \tilde{F} . This is trivial when $x = \star$, while if x = (n, m), then \tilde{F} is obtained from F as in (6.1) simply by changing its value at $\mathrm{id}_{(n,m)}$ from (X, G) to (X, \mathfrak{S}_X) . So the analogue of Lemma 4.19 holds for acceptable functors; it follows that we can define a terminal object (S', E') for $(\mathrm{el}_{\mathsf{P}} /\!\!/_v \mathscr{O}(\mathscr{P}\mathsf{P}))'$ by taking

(6.2)
$$S'(x) = \{ F \in [\mathsf{P}/x, \mathscr{O}(\mathscr{P}\mathsf{P})]_v : \tilde{F} = F \text{ is acceptable} \}$$

with the remaining data defined exactly as in Proposition 4.20 above. All that remains is to identify this (S', E') with the (S, E) in the statement. Once again, this is trivial at stage \star , while at stage (n, m), any acceptable F by definition has the form (6.1); but the further requirement that $F = \tilde{F}$ means that $G = \mathfrak{S}_X$, so that F determines and is determined by the well-labelled polygraph (X, ℓ, r) . \Box

Remark 6.3. As in Remark 4.21, if we view the terminal framed cellular endofunctor of $\mathscr{P}P$ as an object $U \in \mathbf{AN}_0(\mathscr{P}P, \mathscr{P}P)$, then any pointwise analytic Fwhich admits a map to U in this category must itself be framed (I_P, I_P) -cellular. So U is subterminal in $\mathbf{AN}_0(\mathscr{P}P, \mathscr{P}P)$, and the slice category $\mathbf{AN}_0(\mathscr{P}P, \mathscr{P}P)/U$ may be identified with the monoidal category of framed (I_P, I_P) -cellular endofunctors.

The case of the presheaf category $\mathscr{P}\mathsf{P}_s$ of symmetric polygraphs is very similar: the maps in the bordage I_{P_s} are identical in form to those of I_P —though now living on a different category—and we now obtain:

Proposition 6.4. The monoidal category of framed $(I_{\mathsf{P}_s}, I_{\mathsf{P}_s})$ -cellular endofunctors of $\mathscr{P}_{\mathsf{P}_s}$ has a terminal object U_{P_s} , which thus underlies a monad on $\mathscr{P}_{\mathsf{P}_s}$, the universal shapely monad. The spectrum $S \in \mathscr{P}_{\mathsf{P}_s}$ of U_{P_s} may be taken to be:

 $S(\star) = \{u\}$ and $S(n,m) = \{X \in \mathscr{L}_s(n,m) : X \text{ is well-labelled}\}$,

with symmetric actions on S(n,m) given by $X \mapsto \psi \cdot X \cdot \varphi$ as in Definition 2.3(c); the exponent $E: \text{ el } S \to \mathscr{O}(\mathscr{P}\mathsf{P}_s)$ now has $E(u) = \mathsf{y}_{\star}, E(X) = (|X|, \mathfrak{S}_X)$ and

$$E(\sigma_i \colon u \to X) = [\ell_i^X], \quad E(\tau_j \colon u \to X) = [r_i^X], \quad E(\xi_{\varphi,\psi} \colon \psi \cdot X \cdot \varphi \to X) = [1_{|X|}]$$

6.2. Free shapely monads. Now that we have universal shapely monads on the presheaf categories of polygraphs and symmetric polygraphs, we are finally in a position to define more general shapely monads. It will be convenient to abstract away from the particularities of our examples as follows.

Definition 6.5. Let $U \in \mathbf{AN}_0(\mathscr{PC}, \mathscr{PC})$ be subterminal. We write \mathbf{AN}_U for the full subcategory of $\mathbf{AN}_0(\mathscr{PC}, \mathscr{PC})$ on the *U*-analytic endofunctors: those

1

admitting a map to U. We call U nice if \mathbf{AN}_U is closed in $\mathbf{CAT}(\mathscr{PC}, \mathscr{PC})$ under the composition monoidal structure, and in this case we write \mathbf{MND}_U for the category of U-analytic monoids: monoids in \mathbf{AN}_U .

Clearly, the subterminal U in $\mathbf{AN}_0(\mathscr{PC}, \mathscr{PC})$ becomes terminal in \mathbf{AN}_U ; when U is nice, this terminal object has a unique monoid structure making it into a terminal object U in \mathbf{MND}_U . The universal shapely monads of Propositions 6.2 and 6.4 arise in this way from the nice subterminal objects U_P and U_{P_s} in the categories of pointwise analytic endofunctors of \mathscr{PP} and \mathscr{PP}_s ; here " U_P -analytic" means "framed (I_P, I_P) -cellular" and likewise for P_s .

Definition 6.6. Let $U \in \mathbf{AN}_0(\mathscr{PC}, \mathscr{PC})$ be subterminal. A *U*-analytic endofunctor is *shapely* if the unique pointwise analytic $F \to U$ is pointwise monic. If *U* is nice, then a *U*-analytic monad is called *shapely* if its underlying endofunctor is so. We write $\mathbf{SH}_U \subset \mathbf{AN}_U$ and $\mathbf{SHM}_U \subset \mathbf{MND}_U$ for the full subcategories on the shapely endofunctors and monads.

By the free shapely monad on a shapely endofunctor F, we mean the value at F of a left adjoint to the forgetful functor $\mathbf{SHM}_U \to \mathbf{SH}_U$. To construct free shapely monads we will first need to analyse more closely the structure of shapely endofunctors. The following two results are the key to doing so.

Proposition 6.7. For any \mathscr{A} and \mathscr{C} (with \mathscr{C} small), the category $AN_0(\mathscr{A}, \mathscr{PC})$ admits a factorisation system (pointwise epi, pointwise mono).

Proof. First we show that pointwise epimorphic and pointwise monomorphic transformations are orthogonal in $AN_0(\mathscr{A}, \mathscr{PC})$: this says that any square



in $\mathbf{AN}_0(\mathscr{A}, \mathscr{PC})$ with α pointwise epimorphic and β pointwise monomorphic admits a unique diagonal filler ε as displayed. As pointwise epimorphic and monomorphic transformations are orthogonal in $\mathbf{CAT}(\mathscr{A}, \mathscr{PC})$, there is certainly a unique transformation $\varepsilon \colon G \Rightarrow H$; we must show it is pointwise analytic. For each $c \in \mathscr{C}$ we have the factorisation

$$\mathbf{y}_c \downarrow \delta \qquad = \qquad \mathbf{y}_c \downarrow G \xrightarrow{\mathbf{y}_c \downarrow \varepsilon} \mathbf{y}_c \downarrow H \xrightarrow{\mathbf{y}_c \downarrow \beta} \mathbf{y}_c \downarrow K \; .$$

Now since β is pointwise monomorphic, $y_c \downarrow \beta$ is fully faithful and so reflects generic operations; since $y_c \downarrow \delta$ preserves them, we conclude that $y_c \downarrow \varepsilon$ preserves generics, whence ε is pointwise analytic as required.

It remains to show that any $\delta: G \Rightarrow K$ in $\operatorname{AN}_0(\mathscr{PC}, \mathscr{PC})$ has a pointwise (epi, mono) factorisation. Let $\delta = \beta \varepsilon: G \Rightarrow H \Rightarrow K$ be such a factorisation in $\operatorname{CAT}(\mathscr{A}, \mathscr{PC})$; we must show that H, β and ε are pointwise analytic. We argue as before to see that each $y_c \downarrow \varepsilon$ preserves generics, but since ε is pointwise epimorphic, $y_c \downarrow \varepsilon$ is also surjective on objects. It follows that each $t \in y_c \downarrow H$ has a generic cover obtained by lifting along $y_c \downarrow \varepsilon$, taking a generic cover there, and then applying the generic-preserving $y_c \downarrow \varepsilon$. This shows that H and ε are pointwise analytic; moreover, it follows easily that each generic operation in $y_c \downarrow H$ is the image of a generic operation in $y_c \downarrow G$. Since $y_c \downarrow \delta$ preserves generics, it follows that $y_c \downarrow \beta$ does so too, so that β is also pointwise analytic. \Box

Proposition 6.8. The (pointwise epi, pointwise mono) factorisation system on $\mathbf{AN}_0(\mathscr{A}, \mathscr{PC})$ corresponds under Proposition 4.16 to the factorisation system $(\mathscr{E}, \mathscr{M})$ on $\mathrm{el}_{\mathscr{C}}/\!\!/_{v} \mathscr{O}(\mathscr{A})$ for which \mathscr{E} and \mathscr{M} comprise those maps (p, φ) as in (4.5) for which p is epimorphic, respectively p is monic and φ is invertible.

Proof. As (pointwise) epimorphic and monomorphic maps form a factorisation system on \mathscr{PC} , it is not hard to show that $(\mathscr{E}, \mathscr{M})$ is indeed a factorisation system on $\mathrm{el}_{\mathscr{C}}/\!\!/_{v} \mathscr{O}(\mathscr{A})$. It is moreover easy to see that under the equivalence $\mathbf{AN}_{0}(\mathscr{A}, \mathscr{PC}) \simeq \mathrm{el}_{\mathscr{C}}/\!\!/_{v} \mathscr{O}(\mathscr{A})$, pointwise epimorphic maps correspond to maps in \mathscr{E} ; it follows that pointwise monomorphic maps correspond to ones in \mathscr{M} . \Box

Corollary 6.9. Let $U \in \mathbf{AN}_0(\mathscr{PC}, \mathscr{PC})$ be subterminal. The category \mathbf{SH}_U of shapely U-analytic functors is equivalent to the poset of subfunctors of U's spectrum $S_U \in \mathscr{PC}$; in particular, \mathbf{SH}_U is a complete preorder, whose joins are given by unions of subfunctors of $U \in \mathbf{CAT}(\mathscr{A}, \mathscr{PC})$.

Proof. By the preceding result, the shapely U-analytic endofunctors correspond under Proposition 4.16 to the \mathscr{M} -subobjects of (S_U, E_U) in $\operatorname{el}_{\mathscr{C}}/\!\!/_v \mathscr{O}(\mathscr{PC})$. Any such subobject is easily seen to have a unique representative of the form $(p, 1): (S, E_U. \operatorname{el} p) \to$ (S_U, E_U) for $p: S \hookrightarrow S_U$ a subfunctor inclusion. This proves the first claim; the stated form of joins in SH_U follows by transporting across the equivalence $\operatorname{el}_{\mathscr{C}}/\!\!/_v \mathscr{O}(\mathscr{PC}) \simeq \operatorname{AN}_0(\mathscr{PC}, \mathscr{PC})$. \Box

Example 6.10. Consider the subterminal $U_{\mathsf{P}} \in \mathbf{AN}_0(\mathscr{P}\mathsf{P}, \mathscr{P}\mathsf{P})$ which classifies framed $(I_{\mathsf{P}}, I_{\mathsf{P}})$ -cellular endofunctors. By Proposition 6.2, the spectrum S of U_{P} has $S(\star) = \{u\}$ and S(n, m) the set of well-labelled elements in $\mathscr{L}(n, m)$. We will say that a subpresheaf of S is *non-degenerate* if it contains $u \in S(\star)$. Clearly, a non-degenerate subpresheaf is given by selecting arbitrary subsets $\mathscr{F}(n, m)$ of well-labelled elements from each $\mathscr{L}(n, m)$; the corresponding shapely endofunctor $F: \mathscr{P}\mathsf{P} \to \mathscr{P}\mathsf{P}$ —which we also call non-degenerate—satisfies $FA(\star) = A(\star)$ and

$$FA(n,m) = \sum_{X \in \mathscr{F}(n,m)} \mathscr{P}\mathsf{P}(|X|,A)_{\mathfrak{S}_X}$$

We may express this subsequently by saying that the non-degenerate F contains the well-labelled polygraphs in each $\mathscr{F}(n,m)$. For example, it is easy to see that the identity endofunctor of $\mathscr{P}\mathsf{P}$ contains precisely each of the well-labelled polygraphs $\langle n, m \rangle$ of Definition 2.3(d).

Returning to the general situation, when U is a nice subterminal object of $\mathbf{AN}_0(\mathscr{PC}, \mathscr{PC})$, the composition monoidal structure on \mathbf{AN}_U induces by way of Proposition 6.7 the following binary operation on \mathbf{SH}_U .

Definition 6.11. Let $U \in \mathbf{AN}_0(\mathscr{PC}, \mathscr{PC})$ be nice. For any $F, G \in \mathbf{SH}_U$, we let $F \cdot G \in \mathbf{SH}_U$ be the pointwise monic image of the unique $u: FG \to U$ in \mathbf{AN}_U :



The following lemma describes the basic properties of this operation.

Lemma 6.12. Let $U \in \mathbf{AN}_0(\mathscr{PC}, \mathscr{PC})$ be nice. The assignation $F, G \mapsto F \cdot G$ defines a monotone map $\mathbf{SH}_U \times \mathbf{SH}_U \to \mathbf{SH}_U$ which satisfies:

$$F \cdot \mathrm{id} \cong F$$
 $\mathrm{id} \cdot G \cong G$ and $(F \cdot G) \cdot H \leqslant F \cdot (G \cdot H)$.

Moreover, each $(-) \cdot G \colon \mathbf{SH}_U \to \mathbf{SH}_U$ preserves joins, and if U is finitary, then each $F \cdot (-)$ preserves directed joins.

Proof. Monotonicity and the first two displayed equations are obvious. For the third, consider the hexagon left below in $\mathbf{AN}_0(\mathscr{PC}, \mathscr{PC})$; the indicated arrows are pointwise epimorphic or pointwise monomorphic, whence by orthogonality there is a filler as displayed.



Next we show that $(-) \cdot G$ preserves joins of shapely functors. Let $\bigvee_i F_i$ be any such join; since it is computed as a union of subfunctors of U, the induced transformation $\Sigma_i F_i \to \bigvee_i F_i$ in **CAT**($\mathscr{PC}, \mathscr{PC}$) is epimorphic, whence also its precomposition $\Sigma_i F_i G \to (\bigvee_i F_i)G$. Thus in the hexagon right above, each edge is pointwise epi or mono as indicated, so that by orthogonality we induce an isomorphism $(\bigvee_i F_i) \cdot G \cong \bigvee_i (F_i \cdot G)$ as indicated.

Suppose now that U is finitary; by Remark 4.21, any $F \in \mathbf{SH}_U$ is then also finitary. Now any directed join $\bigvee_i G_i$ in \mathbf{SH}_U , being a union of subfunctors of U, may be computed as the colimit in $\mathbf{CAT}(\mathscr{PC}, \mathscr{PC})$ of the filtered diagram of subfunctor inclusions. Because any $F \in \mathbf{SH}_U$ is finitary, it will preserve this colimit, so that the induced map $\Sigma_i FG_i \to F(\bigvee_i G_i)$ in $\mathbf{CAT}(\mathscr{PC}, \mathscr{PC})$ is pointwise epimorphic. The argument of the previous paragraph now carries over *mutatis mutandis* to show that $F \cdot (\bigvee_i G_i) \cong \bigvee_i (F \cdot G_i)$ as required. \Box

Proposition 6.13. Let $U \in \mathbf{AN}_0(\mathscr{PC}, \mathscr{PC})$ be finitary and nice. The forgetful $\mathbf{SHM}_U \to \mathbf{SH}_U$ is a reflective inclusion of preorders, whose image comprises those $F \in \mathbf{SH}_U$ with $\mathrm{id} \leq F$ and $F \cdot F \leq F$. The left adjoint, giving the free shapely monad on $F \in \mathbf{SH}_U$ is defined by:

$$F \qquad \mapsto \qquad \bar{F} = \bigvee_{n \in \mathbb{N}} (\mathrm{id} \lor F)^{\cdot n}$$

where here $F^{\cdot 0} = \text{id}$ and $F^{\cdot n+1} = F \cdot F^{\cdot n}$.

Proof. The only non-trivial point is the verification that \overline{F} is indeed a reflection of F into \mathbf{SHM}_U . First, we have $\mathrm{id} = (\mathrm{id} \vee F)^{\cdot 0} \leq \overline{F}$ and $\overline{F} \cdot \overline{F} \leq \overline{F}$, since

$$\bar{F} \cdot \bar{F} \cong \bigvee_{n} (\mathrm{id} \vee F)^{\cdot n} \cdot \bar{F} \cong \bigvee_{n,m} (\mathrm{id} \vee F)^{\cdot n} \cdot (\mathrm{id} \vee F)^{\cdot m} \leqslant \bigvee_{n,m} (\mathrm{id} \vee F)^{\cdot (n+m)} \leqslant \bar{F}$$

where the first two equalities use cocontinuity of $(-) \cdot \overline{F}$ and directed cocontinuity of each $(\mathrm{id} \vee F)^{\cdot n} \cdot (-)$ (noting that the join defining \overline{F} is indeed directed) and the third inequality uses repeatedly $(F \cdot G) \cdot H \leq F \cdot (G \cdot H)$. So $\overline{F} \in \mathbf{SHM}_U$; moreover, if $G \in \mathbf{SHM}_U$ satisfies $F \leq G$, then since $\mathrm{id} \leq G$ we have $(1 \vee F) \leq G$; furthermore, if $(1 \vee F)^{\cdot n} \leq G$, then

$$(1 \lor F)^{\cdot (n+1)} = (1 \lor F) \cdot (1 \lor F)^{\cdot n} \le G \cdot G \le G$$

so that by induction on n we have $(1 \vee F)^{\cdot n} \leq G$ for all n and so, finally, that $\overline{F} \leq G$. This proves that \overline{F} is a reflection of F into **SHM**_U as desired. \Box

6.3. Polycategories, properads and PROPs. We are now ready to apply the preceding theory to our motivating examples. We concentrate on exhibiting the "free polycategory" monads on $\mathscr{P}\mathsf{P}$ and $\mathscr{P}\mathsf{P}_s$ as free shapely monads, but also indicate how this extends to the cases of properads and PROPs.

We begin in the non-symmetric case $\mathscr{P}\mathsf{P}$ by describing a non-trivial shapely endofunctor Σ_P which encodes the basic polycategorical wiring operations; for this, it suffices by Example 6.10 to describe which well-labelled polygraphs Σ_P will contain. We make use of the operations on polygraphs of Definition 2.3 above. It is easy to see that the elements id $\in \mathscr{L}(1,1)$ and $\langle n,m \rangle \in \mathscr{L}(n,m)$ in parts (a) and (d) of this definition are well-labelled, and that the operations $(-)_j \bullet_i (-)$ and $\psi \cdot (-) \cdot \varphi$ of parts (b) and (c) preserve well-labelledness; so it makes sense to give:

Definition 6.14. Let Σ_{P} be the non-degenerate shapely U_{P} -analytic endofunctor of $\mathscr{P}\mathsf{P}$ which contains the following well-labelled polygraphs:

- (i) id $\in \mathscr{L}(1,1)$;
- (ii) $\psi \cdot \langle n, m \rangle \cdot \varphi$ for each n, m and permutations $\varphi \in \mathfrak{S}_n$ and $\psi \in \mathfrak{S}_m$;
- (iii) $\langle p,q \rangle_j \bullet_i \langle n,m \rangle \in \mathscr{L}(n+p-1,m+q-1)$ for all n,m,p,q and all suitable indices i,j.

Theorem 6.15. The free shapely monad on the shapely U_{P} -analytic endofunctor Σ_{P} is the "free polycategory" monad on \mathscr{P}_{P} .

Proof. Since Σ_{P} contains each of the shapes $\langle n, m \rangle$, we have by Example 6.10 that id $\subset \Sigma_{\mathsf{P}}$; so by the formula of Proposition 6.13, the free shapely monad on Σ_{P} is given by $\bigvee_{n} (\Sigma_{\mathsf{P}})^{\cdot n}$. To compute this, we first calculate for any non-degenerate shapely U_{P} -analytic endofunctor F the composite $\Sigma_{\mathsf{P}} \cdot F$. Since F is non-degenerate, it is by Example 6.10 specified by families of well-labelled polygraphs $\mathscr{F}(n,m) \subset \mathscr{L}(n,m)$; it is easy to see that $\Sigma_{\mathsf{P}} \cdot F$ is then also non-degenerate, and so it will suffice to determine the well-labelled polygraphs which it contains. These polygraphs correspond to generic operations of $\Sigma_{\mathsf{P}} \cdot F$ at stage $y_{(n,m)}$, and by Definition 6.11 and Proposition 6.8, such operations are precisely

the images of the generic operations of $\Sigma_{\mathsf{P}}F$ at stage $\mathsf{y}_{(n,m)}$ under the unique pointwise analytic $\Sigma_{\mathsf{P}}F \to U_{\mathsf{P}}$; so it will suffice to compute these.

Now, by Proposition 4.22, any generic operation $v: y_{(n,m)} \to \Sigma_{\mathsf{P}} FA$ is the composite of a Σ_{P} -generic operation $s: y_{(n,m)} \to \Sigma_{\mathsf{P}} B$ and an *F*-generic operation $t: B \to FA$. The first possibility is that

$$v = \mathsf{y}_{(1,1)} \xrightarrow{s} \Sigma_{\mathsf{P}}(\mathsf{y}_{\star}) \xrightarrow{\Sigma_{\mathsf{P}}(t)} \Sigma_{\mathsf{P}}F(\mathsf{y}_{\star})$$

where s corresponds to $\mathrm{id} \in \mathscr{L}(1,1)$. This v is sent by $\Sigma_{\mathsf{P}}F \to U_{\mathsf{P}}$ to a welllabelled polygraph $X \in \mathscr{L}(1,1)$ with $|X| = \mathsf{y}_{\star}$, which clearly forces $X = \mathrm{id}$. The next possibility is that

$$v = \mathsf{y}_{(n,m)} \xrightarrow{s} \Sigma_{\mathsf{P}}(\mathsf{y}_{(n,m)}) \xrightarrow{\Sigma_{\mathsf{P}}(t)} \Sigma_{\mathsf{P}}F(|X|)$$

where s corresponds to $\psi \cdot \langle n, m \rangle \cdot \varphi \in \mathscr{L}(n, m)$, and t corresponds to some welllabelled $X \in \mathscr{F}(n, m)$. The composite v is sent by $\Sigma_{\mathsf{P}}F \to U_{\mathsf{P}}$ to a well-labelled $Y \in \mathscr{L}(n, m)$ with underlying polygraph |X|; to calculate the leaf labellings $\ell_1^Y, \ldots, \ell_n^Y$, we apply (4.6) and Remark 4.14, which tell us that they arise by taking generic covers as to the left in:

$$\begin{array}{cccc} & & & & & & & & \\ y_{\star} & \xrightarrow{y_{\sigma_{i}}} & & & & \\ y_{\star} & \xrightarrow{y_{\sigma_{i}}} & & & & \\ y_{\star} & \xrightarrow{y_{\sigma_{i}}} & & & \\ \hline y_{\tau_{i}} & & & & \\ \hline y_{\tau_{i}} & & \\ \hline$$

But from the given forms of s and t, we have generic covers as to the right, and so must have that $\ell_i^Y = \ell_{\varphi(i)}^X$. The same argument shows that $r_j^Y = r_{\psi^{-1}(j)}^X$, and so we conclude that in fact $Y = \psi \cdot X \cdot \varphi$. The final possibility is that

$$v = \mathsf{y}_{(n+p-1,m+q-1)} \xrightarrow{s} \Sigma_{\mathsf{P}}(\mathsf{y}_{(p,q)} \ j \bullet_i \mathsf{y}_{(n,m)}) \xrightarrow{\Sigma_{\mathsf{P}}(t)} \Sigma_{\mathsf{P}} FA$$

where s corresponds to $\langle p,q \rangle_{j} \bullet_i \langle n,m \rangle \in \mathscr{L}(n+p-1,m+q-1)$. As for the *F*-generic t, by virtue of the pushout (2.3) and Proposition 5.8, this must arise from a pointwise pushout in $\mathscr{P}\mathsf{P} \downarrow F$ of the form:



where t_0 is the unique *F*-generic operation at stage y_* , and t_1 and t_2 are *F*-generic operations corresponding to well-labelled polygraphs $X \in \mathscr{F}(n,m)$ and $Y \in \mathscr{F}(p,q)$. Since the bottom face is a pushout, we conclude that the generic v must correspond to a well-labelled polygraph $Z \in \mathscr{L}(n+p-1,m+q-1)$ with

 $|Z| = |Y_j \bullet_i X|$; now a similar calculation to before shows that the labellings of Z are such that, in fact, we have $Z = Y_j \bullet_i X$.

In sum, we have now shown that, for any non-trivial shapely U_{P} -analytic endofunctor F containing the well-labelled polygraphs $\mathscr{F}(n,m)$, the well-labelled polygraphs contained in the shapely composite $\Sigma_{\mathsf{P}} \cdot F$ are given by:

- (i) id $\in \mathscr{L}(1,1)$;
- (ii) $\psi \cdot X \cdot \varphi \in \mathscr{L}(n,m)$ for all $X \in \mathscr{F}(n,m)$, $\varphi \in \mathfrak{S}_n$ and $\psi \in \mathfrak{S}_m$;
- (iii) $Y_{j} \bullet_{i} X \in \mathscr{L}(n+p-1, m+q-1)$ for all $X \in \mathscr{F}(n,m), Y \in \mathscr{F}(p,q)$ and suitable indices i, j.

Consequently, the well-labelled polygraphs contained in the free shapely monad $\bigvee_n (\Sigma_{\mathsf{P}})^{\cdot n}$ are those obtained by closing the $\langle n, m \rangle$'s under the operations (a)–(c) of Definition 2.3, and by definition, these are precisely the finite labelled polygraphic trees $\mathscr{T}(n,m)$. It follows from Example 6.10 that the free shapely monad T on Σ_{P} is given by $TX(\star) = X(\star)$ and

$$TX(n,m) = \sum_{T \in \mathscr{T}(n,m)} \mathscr{P}\mathsf{P}(|T|,X)_{/\mathfrak{S}_T} = \sum_{T \in \mathscr{T}(n,m)} \mathscr{P}\mathsf{P}(|T|,X)$$

where the second equality follows from the easy observation that a labelled polygraphic tree has *no* non-trivial label-preserving automorphisms. Comparing this with the formula of Proposition 2.6 gives the result. \Box

The argument just given for the free polycategory monad on $\mathscr{P}P$ applies equally well to the free polycategory monad on $\mathscr{P}P_s$. By adapting Example 6.10, we see that the non-trivial shapely endofunctors of $\mathscr{P}P_s$ are specified by giving subsets $\mathscr{F}_s(n,m) \subset \mathscr{L}_s(n,m)$ of well-labelled finite symmetric trees; so we can define a shapely endofunctor Σ_{P_s} by requiring it to contain id $\in \mathscr{L}_s(1,1)$ and each $\langle p,q \rangle_j \bullet_i \langle n,m \rangle \in \mathscr{L}_s(n+p-1,m+q-1)$. Now following the precise same argument as in Theorem 6.15 gives:

Theorem 6.16. The free shapely monad on the shapely U_{P_s} -analytic endofunctor Σ_{P_s} is the "free polycategory" monad on $\mathscr{P}_{\mathsf{P}_s}$.

Finally, let us say a few brief words about the passage from polycategories to properads and to PROPs. This is almost completely trivial in our framework; all we need to do is to replace the closure operations of Definition 2.3 which defined the class of polycategorical trees with the corresponding closure operations defining the properadic trees or the trees for PROPs. Thus, for example, the monad for properads on the category $\mathscr{P}P$ arises as the free shapely monad on the shapely endofunctor specified by the well-labelled polygraphs

id,
$$\psi \cdot \langle n, m \rangle \cdot \varphi$$
 and $\langle p, q \rangle_I \bullet_J \langle n, m \rangle$.

The remaining cases proceed similarly.

References

- ABBOTT, M., ALTENKIRCH, T., GHANI, N., AND MCBRIDE, C. Constructing polymorphic programs with quotient types. In *Mathematics of program construction*, vol. 3125 of *Lecture Notes in Computer Science*. Springer, 2004, pp. 2–15.
- [2] BLUTE, R. F., COCKETT, J. R. B., SEELY, R. A. G., AND TRIMBLE, T. H. Natural deduction and coherence for weakly distributive categories. *Journal of Pure and Applied Algebra 113* (1996), 229–296.

- [3] BONCHI, F., SOBOCIŃSKI, P., AND ZANASI, F. Full abstraction for signal flow graphs. In Principles of Programming Languages, POPL'15. (2015).
- [4] BURRONI, A. Higher-dimensional word problems with applications to equational logic. *Theoretical Computer Science* 115, 1 (1993), 43–62.
- [5] CARAMELLO, O. Fraïssé's construction from a topos-theoretic perspective. Logica Universalis 8, 2 (2014), 261–281.
- [6] CARBONI, A., AND JOHNSTONE, P. Connected limits, familial representability and Artin glueing. Mathematical Structures in Computer Science 5, 4 (1995), 441–459.
- [7] DIERS, Y. Spectres et localisations relatifs à un foncteur. Comptes rendus hebdomadaires des séances de l'Académie des sciences 287, 15 (1978), 985–988.
- [8] FIORE, M. Analytic functors between presheaf categories over groupoids. Theoretical Computer Science 546 (2014), 120–131.
- FIORE, M., GAMBINO, N., HYLAND, M., AND WINSKEL, G. The cartesian closed bicategory of generalised species of structures. *Journal of the London Mathematical Society* 77, 1 (2008), 203–220.
- [10] FONG, B. Causal theories: A categorical perspective on bayesian networks. Master's thesis, University of Oxford, 2012.
- [11] GIRARD, J.-Y. Linear logic. Theoretical Computer Science 50, 1 (1987), 1–101.
- [12] HASEGAWA, R. Two applications of analytic functors. *Theoretical Computer Science 272*, 1-2 (2002), 113–175.
- [13] HOVEY, M. Model categories, vol. 63 of Mathematical Surveys and Monographs. American Mathematical Society, 1999.
- [14] JENSEN, O. H., AND MILNER, R. Bigraphs and mobile processes (revised). Tech. Rep. TR580, University of Cambridge, 2004.
- [15] JOHNSON, M., AND WALTERS, R. F. C. Algebra objects and algebra families for finite limit theories. Journal of Pure and Applied Algebra 83, 3 (1992), 283–293.
- [16] JOYAL, A. Foncteurs analytiques et espèces de structures. In Combinatoire énumérative (Montreal, 1985), vol. 1234 of Lecture Notes in Mathematics. Springer, 1986, pp. 126–159.
- [17] KELLY, G. M. Basic concepts of enriched category theory, vol. 64 of London Mathematical Society Lecture Note Series. Cambridge University Press, 1982. Republished as: Reprints in Theory and Applications of Categories 10 (2005).
- [18] KOCK, J. Polynomial functors and trees. International Mathematics Research Notices, 3 (2011), 609–673.
- [19] KOCK, J. Graphs, hypergraphs, and properads. Preprint, available as http://arxiv.org/abs/1407.3744, 2014.
- [20] LAFONT, Y. Interaction nets. In POPL (1990), ACM.
- [21] LAMBEK, J. Deductive systems and categories. II. Standard constructions and closed categories. In *Category Theory, Homology Theory and their Applications, I*, no. 42 in Lecture Notes in Mathematics. Springer, 1969, pp. 76–122.
- [22] LAMBEK, J. From categorial grammar to bilinear logic. In Substructural logics (Tübingen, 1990), vol. 2 of Studies in Logic and Computation. Oxford University Press, 1993, pp. 207– 237.
- [23] LEINSTER, T. Higher operads, higher categories, vol. 298 of London Mathematical Society Lecture Note Series. Cambridge University Press, 2004.
- [24] MAC LANE, S. Categorical algebra. Bulletin of the American Mathematical Society 71 (1965), 40–106.
- [25] MARKL, M., SHNIDER, S., AND STASHEFF, J. Operads in algebra, topology and physics, vol. 96 of Mathematical Surveys and Monographs. American Mathematical Society, Providence, RI, 2002.
- [26] MAY, J. P. The geometry of iterated loop spaces, vol. 271 of Lecture Notes in Mathematics. Springer, 1972.
- [27] MCCLURE, J. E., AND SMITH, J. H. A solution of Deligne's Hochschild cohomology conjecture. In *Recent progress in homotopy theory (Baltimore, 2000)*, vol. 293 of *Contemporary Mathematics*. American Mathematical Society, 2002, pp. 153–193.

R. GARNER AND T. HIRSCHOWITZ

- [28] SZABO, M. E. Polycategories. Communications in Algebra 3, 8 (1975), 663–689.
- [29] SZABO, M. E. The logic of closed categories. Notre Dame Journal of Formal Logic 18, 3 (1977), 441–457.
- [30] TAMBARA, D. Finite categories with pushouts. Theory and Applications of Categories 30 (2015), 1017–1031.
- [31] VALLETTE, B. A Koszul duality for PROPs. Transactions of the American Mathematical Society 359, 10 (2007), 4865–4943.
- [32] WEBER, M. Symmetric Operads for Globular Sets. PhD thesis, Macquarie University, 2001.
- [33] WEBER, M. Generic morphisms, parametric representations and weakly Cartesian monads. Theory and Applications of Categories 13 (2004), 191–234.

Department of Mathematics, Macquarie University, NSW 2109, Australia E-mail address: richard.garner@mq.edu.au

CNRS, UNIVERSITÉ SAVOIE MONT-BLANC *E-mail address:* tom.hirschowitz@univ-smb.fr

46