



HAL
open science

THE AMPLIFICATION METHOD IN THE $GL(3)$ HECKE ALGEBRA

Roman Holowinsky, Guillaume Ricotta, Emmanuel Royer

► **To cite this version:**

Roman Holowinsky, Guillaume Ricotta, Emmanuel Royer. THE AMPLIFICATION METHOD IN THE $GL(3)$ HECKE ALGEBRA. Publications Mathématiques de Besançon : Algèbre et Théorie des Nombres, 2016, pp.13-40. hal-01247675

HAL Id: hal-01247675

<https://hal.science/hal-01247675>

Submitted on 22 Dec 2015

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

THE AMPLIFICATION METHOD IN THE $GL(3)$ HECKE ALGEBRA

ROMAN HOLOWINSKY, GUILLAUME RICOTTA, AND EMMANUEL ROYER

ABSTRACT. This article contains all of the technical ingredients required to implement an effective, explicit and unconditional amplifier in the context of $GL(3)$ automorphic forms. In particular, several coset decomposition computations in the $GL(3)$ Hecke algebra are explicitly done.

CONTENTS

| | |
|---|----|
| 1. Introduction and statement of the results | 1 |
| 1.1. Motivation: sup-norms of $GL(n)$ Hecke-Maaß cusp forms | 1 |
| 1.2. The $GL(2)$ and $GL(3)$ amplifier | 2 |
| 1.3. Statement of the results | 4 |
| 1.4. Organization of the paper | 6 |
| 2. Background on the $GL(3)$ Hecke algebra | 6 |
| 3. Proof of the linearizations given in Theorem A | 8 |
| 3.1. Linearization of $\Lambda \text{diag}(1, 1, p)\Lambda * \Lambda \text{diag}(1, p, p)\Lambda$ | 8 |
| 3.2. Linearization of $\Lambda \text{diag}(1, p, p^2)\Lambda * \Lambda \text{diag}(1, p, p^2)\Lambda$ | 11 |
| Appendix A. Decomposition of Λ -double cosets into Λ -cosets | 15 |
| A.1. Decomposition and degree of $\Lambda \text{diag}(1, 1, p)\Lambda$ | 16 |
| A.2. Decomposition and degree of $\Lambda \text{diag}(1, p, p)\Lambda$ | 17 |
| A.3. Decomposition and degree of $\Lambda \text{diag}(1, p, p^2)\Lambda$ | 17 |
| A.4. Degree of $\Lambda \text{diag}(1, p^2, p^4)\Lambda$ | 19 |
| A.5. Degree of $\Lambda \text{diag}(1, p^3, p^3)\Lambda$ | 22 |
| A.6. Degree of $\Lambda \text{diag}(1, 1, p^3)\Lambda$ | 24 |
| References | 26 |

1. INTRODUCTION AND STATEMENT OF THE RESULTS

1.1. **Motivation: sup-norms of $GL(n)$ Hecke-Maaß cusp forms.** Let f be a $GL(n)$ Maaß cusp form and K be a fixed compact subset of $SL_n(\mathbb{R})/SO_n(\mathbb{R})$ (see [Gol06]). The generic or local bound for the sup-norm of f restricted to K is given by

$$\|f|_K\|_\infty \ll \lambda_f^{n(n-1)/8}$$

where λ_f is the Laplace eigenvalue of f (see [Sar]). Note that F. Brumley and N. Templier noticed in [BT] that the previous bound does not hold when $n \geq 6$ if f is not restricted to a compact.

Date: Version of March 10, 2015.

1991 Mathematics Subject Classification. Primary 11F99, 11F60; Secondary 11F55.

Key words and phrases. Amplification method, Hecke operators, Hecke algebras.

If f is an eigenform of the Hecke algebra, however, then the generic bound is not expected to be the correct order of magnitude for the sup-norm of the restriction of f to a fixed compact. This is essentially due to the fact that the Hecke operators are additional symmetries on the ambient space. In other words, we expect there to exist an absolute positive constant $\delta_n > 0$ such that

$$\|f|_K\|_\infty \ll \lambda_f^{n(n-1)/8-\delta_n}. \quad (1.1)$$

H. Iwaniec and P. Sarnak proved the bound given in (1.1) in [IS95] when $n = 2$ for $\delta_2 = 1/24$. Note that improving this constant δ_2 seems to be a very delicate open problem. The case $n = 3$ was completed by the authors in [HRR] with $\delta_3 = 1/124$. The general case was done, without an explicit value for δ_n , in a series of recent impressive works by V. Blomer and P. Maga in [BMb] and in [BMa].

All of the above achievements (and much more) were made possible thanks to generalizations of the amplification method developed by W. Duke, J. Friedlander and H. Iwaniec for $GL(1)$ and $GL(2)$ (see [FI92], [Iwa92] and [DFI94] for example). In particular, the proof of (1.1) for $n = 3$ with $\delta_3 = 1/124$ relies on Theorem B of this article which was stated without proof in [HRR] as Proposition 4.1¹. For the sake of completeness and future use, we provide the full details of the proof of Theorem B, including computations, here in this article.

1.2. The $GL(2)$ and $GL(3)$ amplifier. The general principle behind the construction of an amplifier, is the existence of an identity which allows one to write a non-zero constant as a finite sum of Hecke eigenvalues. In the most basic context of $GL(2)$ automorphic forms, this identity is

$$\lambda_f(p)^2 - \lambda_f(p^2) = 1 \quad (1.2)$$

where p is any prime and $\lambda_f(n)$ is the n -th Hecke eigenvalue of a Hecke-Maaß cusp form f of full level, i.e. $T_n f = \lambda_f(n)f$ where

$$(T_n f)(z) = \frac{1}{\sqrt{n}} \sum_{ad=n} \sum_{b=1}^d f\left(\frac{az+b}{d}\right).$$

One may interpret the above identity as the fact that the Rankin-Selberg convolution factors as the product of the adjoint square and the Riemann zeta function and therefore has a pole at $s = 1$.

From the identity (1.2), one constructs an amplifier

$$A_f := \left| \sum_{\ell} \alpha_{\ell} \lambda_f(\ell) \right|^2$$

with

$$\alpha_{\ell} := \begin{cases} \lambda_{f_0}(\ell) & \text{if } \ell \leq \sqrt{L} \text{ is a prime number,} \\ -1 & \text{if } \ell \leq L \text{ is the square of a prime number,} \\ 0 & \text{otherwise} \end{cases}$$

¹Theorem B and Proposition 4.1 in [HRR] are not entirely identical. Since releasing our first article [HRR], we have noticed a simplification in the construction of the amplifier. Therefore, Theorem B only contains the identities in Proposition 4.1 of [HRR] which are necessary for the amplification method. The implied power saving in the Laplace eigenvalue for the sup-norm bound remains the same.

for some fixed form f_0 . The advantage to constructing such an amplifier is that it is expected to be small² for general forms f while satisfying the lower bound $A_f \gg_\varepsilon L^{1-\varepsilon}$ when $f = f_0$.

Reinterpreting (1.2) in terms of Hecke operators, we may write

$$T_p \circ T_p - T_{p^2} = Id.$$

In application to the sup-norm problem for $GL(2)$ via a pre-trace formula argument, this translates into a need to geometrically understand the behavior of the following collection of operators on an automorphic kernel:

$$T_p, T_p \circ T_q^*, T_p \circ T_{q^2}^* \text{ and } T_{p^2} \circ T_{q^2}^*$$

both in the cases of primes $p = q$ and $p \neq q$. Since the Hecke operators T_n in $GL(2)$ are self-adjoint and computationally pleasant to work with due to their relatively simple composition law, one quickly computes that the above collection of Hecke operators may be reduced to the study of

$$T_p, T_{pq}, T_{pq^2} \text{ and } T_{p^2q^2}.$$

In truth, one has an opportunity to further reduce the collection of necessary Hecke operators through the simple inequality

$$A_f \leq 2 \left| \sum_p \alpha_p \lambda_f(p) \right|^2 + 2 \left| \sum_p \alpha_{p^2} \lambda_f(p^2) \right|^2. \quad (1.3)$$

Indeed, an appropriate application of (1.3) (see for example [BHM]) allows one to further restrict the set of necessary Hecke operators to

$$T_{pq} \text{ and } T_{p^2q^2}$$

both in the cases of primes $p = q$ and $p \neq q$.

The case of $GL(3)$ is much more computationally involved due to the lack of self-adjointness of the Hecke operators and their multiplication law. Instead of looking at identities involving Hecke eigenvalues, we start immediately with the Hecke operators themselves (see §2 for definitions). Our fundamental identity now will be

$$T_{\text{diag}(1,p,p)} \circ T_{\text{diag}(1,1,p)} - T_{\text{diag}(1,p,p^2)} = (p^2 + p + 1)Id.$$

We set $c_f(p) = a_f(p, 1)$, $c_f(p)^* = \overline{a_f(p, 1)}$ and $c_f(p^2)$ to be the eigenvalues of $p^{-1}T_{\text{diag}(1,1,p)} = T_p$, $p^{-1}T_{\text{diag}(1,p,p)} = T_p^*$ and $p^{-2}T_{\text{diag}(1,p,p^2)}$ respectively when acting on a form f . See (2.2) and (2.3) for the precise definitions. We construct the amplifier

$$A_f := \left| \sum_\ell \alpha_\ell c_f(\ell) \right|^2$$

with³

$$\alpha_\ell := \begin{cases} c_{f_0}(\ell)^* & \text{if } \ell \leq \sqrt{L} \text{ is a prime number,} \\ -1 & \text{if } \ell \leq L \text{ is the square of a prime number,} \\ 0 & \text{otherwise.} \end{cases}$$

²This follows, at least conditionally, from a suitable version of the Generalized Riemann Hypothesis. However, we do not need to use this fact.

³One may also choose a variant, in the spirit of [Ven10], which involves the signs of $c_{f_0}(\ell)$.

As in the case of $GL(2)$, this amplifier will satisfy $A_{f_0} \gg_\varepsilon L^{1-\varepsilon}$ and A_f is otherwise expected to be small for $f \neq f_0$.

Applying the inequality

$$A_f \leq 2 \left| \sum_p \alpha_p c_f(p) \right|^2 + 2 \left| \sum_p \alpha_{p^2} c_f(p^2) \right|^2,$$

one is reduced to understanding the actions of

$$T_{\text{diag}(1,1,p)} \circ T_{\text{diag}(1,q,q)} \text{ and } T_{\text{diag}(1,p,p^2)} \circ T_{\text{diag}(1,q,q^2)}$$

both in the cases of primes $p = q$ and $p \neq q$ on the relevant automorphic kernel. In the following sections, we compute the above compositions as linear combinations of other Hecke operators and state our main result as Theorem B. In the end, we shall see that the following operators are the relevant ones for our application

$$T_{\text{diag}(1,p,pq)}, T_{\text{diag}(1,pq,p^2q^2)}, T_{\text{diag}(1,p^3,p^3)} \text{ and } T_{\text{diag}(1,1,p^3)}$$

for primes $p = q$ and $p \neq q$.

1.3. Statement of the results.

Theorem A—Let p be a prime number and $\Lambda = GL_3(\mathbb{Z})$.

- The set $R_{1,1,p}$ (respectively $R_{1,p,p}$, R_{1,p,p^2}) defined in Proposition A.1 (respectively Proposition A.2, Proposition A.3) is a complete system of representatives for the distinct Λ -right cosets in the Λ -double coset of $\text{diag}(1, 1, p)$ (respectively $\text{diag}(1, 1, p)$, $\text{diag}(1, p, p^2)$) modulo Λ .
- The following formulas for the degrees of Λ -double cosets hold.

$$\begin{aligned} \deg(\text{diag}(1, 1, p)) &= p^2 + p + 1, \\ \deg(\text{diag}(1, p, p)) &= p^2 + p + 1, \\ \deg(\text{diag}(1, p, p^2)) &= p(p+1)(p^2 + p + 1), \\ \deg(\text{diag}(p, p, p)) &= 1, \\ \deg(\text{diag}(1, p^2, p^4)) &= p^5(p+1)(p^2 + p + 1), \\ \deg(\text{diag}(1, p^3, p^3)) &= p^4(p^2 + p + 1), \\ \deg(\text{diag}(p, p, p^4)) &= p^4(p^2 + p + 1), \\ \deg(\text{diag}(p, p^2, p^3)) &= p(p+1)(p^2 + p + 1), \\ \deg(\text{diag}(p^2, p^2, p^2)) &= 1. \end{aligned}$$

- Finally,

$$\Lambda \text{diag}(1, 1, p) \Lambda * \Lambda \text{diag}(1, p, p) \Lambda = \Lambda \text{diag}(1, p, p^2) \Lambda + (p^2 + p + 1) \Lambda \text{diag}(p, p, p) \Lambda. \quad (1.4)$$

and

$$\begin{aligned} \Lambda \text{diag}(1, p, p^2) \Lambda * \Lambda \text{diag}(1, p, p^2) \Lambda &= \Lambda \text{diag}(1, p^2, p^4) \Lambda + (p+1) \Lambda \text{diag}(1, p^3, p^3) \Lambda \\ &+ (p+1) \Lambda \text{diag}(p, p, p^4) \Lambda + (p+1)(2p-1) \Lambda \text{diag}(p, p^2, p^3) \Lambda \\ &+ p(p+1)(p^2 + p + 1) \Lambda \text{diag}(p^2, p^2, p^2) \Lambda. \quad (1.5) \end{aligned}$$

Remark 1.1–In [Kod67], T. Kodama explicitly computed the product of other double cosets in the slightly harder case of the Hecke ring for the symplectic group. The results stated in the previous theorem are similar in spirit.

Remark 1.2–It is well-known that a Λ -double coset can be identified with its characteristic function χ . Under this identification, the multiplication law between Λ -double cosets is the classical convolution between functions. If $\mu = (\mu_1, \mu_2, \mu_3)$ with $\mu_1 \geq \mu_2 \geq \mu_3 \geq 0$ and $\nu = (\nu_1, \nu_2, \nu_3)$ with $\nu_1 \geq \nu_2 \geq \nu_3 \geq 0$ are two partitions of length less than n , then

$$\chi_{\Lambda \text{diag}(p^{\mu_1}, p^{\mu_2}, p^{\mu_3})\Lambda} * \chi_{\Lambda \text{diag}(p^{\nu_1}, p^{\nu_2}, p^{\nu_3})\Lambda} = \sum_{\lambda} g_{\mu, \nu}^{\lambda}(p) \chi_{\Lambda \text{diag}(p^{\lambda_1}, p^{\lambda_2}, p^{\lambda_3})\Lambda}$$

for any prime number p where λ ranges over the partitions of length less than n and the $g_{\mu, \nu}^{\lambda}(p)$ are the Hall polynomials (see [Mac95, Equation (2.6) Page 295]). The sum on the right-hand side of the above equality is finite since only a finite number of the Hall polynomials are non-zero. However, determining which Hall polynomials vanish is not straightforward (see [Mac95, Equation (4.3) Page 188]). Using Sage, one can check that

$$\begin{aligned} g_{(2,1,0), (2,1,0)}^{(4,2,0)}(p) &= 1, \\ g_{(2,1,0), (2,1,0)}^{(3,3,0)}(p) &= p+1, \\ g_{(2,1,0), (2,1,0)}^{(4,1,1)}(p) &= p+1, \\ g_{(2,1,0), (2,1,0)}^{(3,2,1)}(p) &= (p+1)(2p-1), \\ g_{(2,1,0), (2,1,0)}^{(2,2,2)}(p) &= p(p+1)(p^2+p+1) \end{aligned}$$

and one can recover the coefficients occurring in (1.5). We prefer to give a different proof, which has the advantage of producing explicit systems of representatives for the Λ -right cosets and formulas for the degrees.

Corollary B–If p and q are two prime numbers then

$$T_{\text{diag}(1,p,p)} \circ T_{\text{diag}(1,1,q)} = T_{\text{diag}(1,p,pq)} + \delta_{p=q}(p^2+p+1)Id \quad (1.6)$$

and

$$\begin{aligned} T_{\text{diag}(1,p,p^2)} \circ T_{\text{diag}(1,q,q^2)} &= T_{\text{diag}(1,pq,p^2q^2)} + \delta_{p=q}(p+1)(T_{\text{diag}(1,p^3,p^3)} + T_{\text{diag}(1,1,p^3)}) \\ &+ \delta_{p=q}(p+1)(2p-1)T_{\text{diag}(1,p,p^2)} + \delta_{p=q}p(p+1)(p^2+p+1)Id. \end{aligned} \quad (1.7)$$

When $p \neq q$, the previous corollary immediately follows from (2.6) and (2.7). When $p = q$, it is a consequence of the previous theorem and of (2.6).

Remark 1.3–As observed by L. Silberman and A. Venkatesh in [SA] and used by V. Blomer and P. Maga in [BMb] and in [BMa], the precise formulas for the Hall polynomials occurring in (1.6) and in (1.7) are not really needed for the purpose of the amplification method, since the Hall polynomials are easily well approximated for p and q large by the much easier Schur polynomials. Nevertheless, the precise list of the Hecke operators relevant for the amplification method, namely occurring in (1.6) and in (1.7), seems to be crucial in order to obtain the best possible explicit result. For instance, G. Harcos and N. Templier used such a list in order to prove the best known subconvexity exponent for the sup-norm of $GL(2)$ automorphic forms in the level aspect in [HT13].

1.4. Organization of the paper. The general background on $GL(3)$ Maaß cusp forms and on the $GL(3)$ Hecke algebra is given in Section 2. The linearizations involved in Theorem A are detailed in Section 3. The proof requires decompositions of Λ -double cosets into Λ -left and right cosets and computations of degrees as done in Appendix A.

Notations— Λ stands for the group $GL_3(\mathbb{Z})$ of 3×3 invertible matrices with integer coefficients. If g is a 3×3 matrix with real coefficients then ${}^t g$ stands for its transpose. For $g \in GL_3(\mathbb{Q})$ we let T_g denote the Hecke operator associated to g (see §2). If a, b and c are three rational numbers then

- $\text{diag}(a, b, c)$ denotes the diagonal 3×3 matrix with a, b and c as diagonal coefficients;
- $L_{a,b,c}$ (respectively $R_{a,b,c}$) stands for a system of representatives for the decomposition of the Λ -double coset $\Lambda \text{diag}(a, b, c) \Lambda$ into Λ -left (respectively right) cosets.

Acknowledgements— The authors would like to thank the referees for a very careful reading of the manuscript.

They also want to thank J. Cogdell, É. Fouvry, E. Kowalski and P. Michel for fruitful discussions related to this work.

This article was worked out at several places: Université Blaise Pascal (Clermont-Ferrand), the Department of Mathematics of The Ohio State University (Columbus), École Polytechnique Fédérale de Lausanne, Eidgenössische Technische Hochschule (Zurich), Forschungsinstitut für Mathematik (Zurich). The authors would like to thank all of these institutions for their hospitality and inspiring working conditions.

The research of R. Holowinsky was partially supported by the Sloan fellowship BR2011-083 and the NSF grant DMS-1068043. In-person collaboration with the other authors was made possible through this funding.

The research of G. Ricotta was partially supported by a Marie Curie Intra European Fellowship within the 7th European Community Framework Programme. The grant agreement number of this project, whose acronym is ANERAUTOHI, is PIEF-GA-2009-25271. He would like to thank ETH and its entire staff for the excellent working conditions. Last but not least, the second author would like to express his gratitude to K. Belabas for his crucial but isolated support for Analytic Number Theory among the Number Theory research team A2X (Institut de Mathématiques de Bordeaux, Université de Bordeaux).

2. BACKGROUND ON THE $GL(3)$ HECKE ALGEBRA

Convenient references for this section include [AZ95], [Gol06], [New72] and [Shi94].

Let f be a $GL(3)$ Maaß cusp form of full level. Such f admits a Fourier expansion

$$f(g) = \sum_{\gamma \in U_2(\mathbb{Z}) \backslash SL_2(\mathbb{Z})} \sum_{\substack{m_1 \geq 1 \\ m_2 \in \mathbb{Z} \setminus \{0\}}} \frac{a_f(m_1, m_2)}{m_1 |m_2|} W_{\text{Ja}} \left(\begin{pmatrix} m_1 |m_2| & & \\ & m_1 & \\ & & 1 \end{pmatrix} \begin{pmatrix} \gamma & \\ & 1 \end{pmatrix} g, \nu_f, \psi_{1, \frac{m_2}{|m_2|}} \right)$$

for $g \in GL_3(\mathbb{R})$ (see [Gol06, Equation (6.2.1)]). Here $U_2(\mathbb{Z})$ stands for the \mathbb{Z} -points of the group of upper-triangular unipotent 2×2 matrices. $\nu_f \in \mathbb{C}^2$ is the type

of f , whose components are complex numbers characterized by the property that, for every invariant differential operator D in the center of the universal enveloping algebra of $GL_3(\mathbb{R})$, the cusp form f is an eigenfunction of D with the same eigenvalue as the power function I_{ν_f} , which is defined in [Gol06, Equation (5.1.1)]. $\psi_{1,\pm 1}$ is the character of the group of upper-triangular unipotent real 3×3 matrices defined by

$$\psi_{1,\pm 1} \left(\begin{pmatrix} 1 & u_{1,2} & u_{1,3} \\ & 1 & u_{2,3} \\ & & 1 \end{pmatrix} \right) = e^{2i\pi(u_{2,3} \pm u_{1,2})}.$$

$W_{Ja}(*, \nu_f, \psi_{1,\pm 1})$ stands for the $GL(3)$ Jacquet Whittaker function of type ν_f and character $\psi_{1,\pm 1}$ defined in [Gol06, Equation 6.1.2]. The complex number $a_f(m_1, m_2)$ is the (m_1, m_2) -th *Fourier coefficient* of f for m_1 a positive integer and m_2 a non-vanishing integer.

For $g \in GL_3(\mathbb{Q})$, the *Hecke operator* T_g is defined by

$$T_g(f)(h) = \sum_{\delta \in \Lambda \setminus \Lambda g \Lambda} f(\delta h)$$

for $h \in GL_3(\mathbb{R})$ (see [AZ95, Chapter 3, Sections 1.1 and 1.5]). The *degree* of g or T_g defined by

$$\deg(g) = \deg(T_g) = \text{card}(\Lambda \setminus \Lambda g \Lambda)$$

is scaling invariant, in the sense that

$$\deg(rg) = \deg(g) \tag{2.1}$$

for $r \in \mathbb{Q}^\times$. The adjoint of T_g for the Petersson inner product is $T_{g^{-1}}$. The algebra of Hecke operators \mathbb{T} is the ring of endomorphisms generated by all the T_g 's with $g \in GL_3(\mathbb{Q})$, a commutative algebra of normal endomorphisms (see [Gol06, Theorem 6.4.6]), which contains the m -th normalized Hecke operator

$$T_m = \frac{1}{m} \sum_{\substack{g = \text{diag}(y_1, y_2, y_3) \\ y_1 | y_2 | y_3 \\ y_1 y_2 y_3 = m}} T_g \tag{2.2}$$

for all positive integers m . A *Hecke-Maaß cusp form* f of full level is a Maaß cusp form of full level, which is an eigenfunction of \mathbb{T} . In particular, it satisfies

$$T_m(f) = a_f(m, 1)f \text{ and } T_m^*(f) = a_f(1, m)f = \overline{a_f(m, 1)}f \tag{2.3}$$

according to [Gol06, Theorem 6.4.11].

The algebra \mathbb{T} is isomorphic to the *absolute Hecke algebra*, the free \mathbb{Z} -module generated by the double cosets $\Lambda g \Lambda$ where g ranges over $\Lambda \setminus GL_3(\mathbb{Q}) / \Lambda$ and endowed with the following multiplication law. If g_1 and g_2 belong to $GL_3(\mathbb{Q})$ and

$$\Lambda g_1 \Lambda = \bigcup_{i=1}^{\deg(g_1)} \Lambda \alpha_i \text{ and } \Lambda g_2 \Lambda = \bigcup_{j=1}^{\deg(g_2)} \Lambda \beta_j$$

then

$$\Lambda g_1 \Lambda * \Lambda g_2 \Lambda = \sum_{\Lambda h \Lambda \subset \Lambda g_1 \Lambda g_2 \Lambda} m(g_1, g_2; h) \Lambda h \Lambda \tag{2.4}$$

where $h \in GL_3(\mathbb{Q})$ ranges over a system of representatives of the Λ -double cosets contained in the set $\Lambda g_1 \Lambda g_2 \Lambda$ and

$$\begin{aligned} m(g_1, g_2; h) &= \text{card}(\{(i, j) \in \{1, \dots, \deg(g_1)\} \times \{1, \dots, \deg(g_2)\}, \alpha_i \beta_j \in \Lambda h\}), \quad (2.5) \\ &= \frac{1}{\deg(h)} \text{card}(\{(i, j) \in \{1, \dots, \deg(g_1)\} \times \{1, \dots, \deg(g_2)\}, \alpha_i \beta_j \in \Lambda h\}), \\ &= \frac{\deg(g_2)}{\deg(h)} \text{card}(\{i \in \{1, \dots, \deg(g_1)\}, \alpha_i g_2 \in \Lambda h\}). \end{aligned}$$

Confer [AZ95, Lemma 1.5 Page 96]. In particular,

$$\Lambda \text{diag}(r, r, r) \Lambda * \Lambda g \Lambda = \Lambda r g \Lambda \quad (2.6)$$

for $g \in GL_3(\mathbb{Q})$ and $r \in \mathbb{Q}^\times$ ([AZ95, Lemma 2.4 Page 107]). In addition, for p and q two distinct prime numbers,

$$\Lambda \text{diag}(1, p^{\alpha_1}, p^{\alpha_2}) \Lambda * \Lambda \text{diag}(1, q^{\beta_1}, q^{\beta_2}) \Lambda = \Lambda \text{diag}(1, p^{\alpha_1} q^{\beta_1}, p^{\alpha_2} q^{\beta_2}) \Lambda \quad (2.7)$$

where $\alpha_1, \alpha_2, \beta_1, \beta_2$ are non-negative integers by [AZ95, Proposition 2.5 Page 107].

Every double coset $\Lambda g \Lambda$ with g in $GL_3(\mathbb{Q})$ contains a unique representative of the form

$$r \text{diag}(1, s_1(g), s_2(g)) \quad (2.8)$$

where $r \in \mathbb{Q}^*$ and $s_1(g), s_2(g)$ are some positive integers satisfying $s_1(g) \mid s_2(g)$ (see [AZ95, Lemma 2.2]).

Finally, let $g = [g_{i,j}]_{1 \leq i, j \leq 3}$ be a 3×3 matrix with integer coefficients. Its *determinantal divisors* are the non-negative integers given by

$$\begin{aligned} d_1(g) &= \text{gcd}(\{g_{i,j}, 1 \leq i, j \leq 3\}), \\ d_2(g) &= \text{gcd}(\{\text{determinants of } 2 \times 2 \text{ submatrices of } g\}), \\ d_3(g) &= |\det(g)|. \end{aligned}$$

and its *determinantal vector* is $\mathbf{d}(g) = (d_1(g), d_2(g), d_3(g))$. The determinantal divisors turn out to be useful since if h is another 3×3 matrix with integer coefficients then h belongs to $\Lambda g \Lambda$ if and only if $d_k(h) = d_k(g)$ for $1 \leq k \leq 3$ (see [New72]).

3. PROOF OF THE LINEARIZATIONS GIVEN IN THEOREM A

3.1. Linearization of $\Lambda \text{diag}(1, 1, p) \Lambda * \Lambda \text{diag}(1, p, p) \Lambda$. This section contains the proof of (1.4).

By (2.4), the product of these double cosets equals

$$\sum_{\Lambda h \Lambda \subset \Lambda \text{diag}(1, 1, p) \Lambda \text{diag}(1, p, p) \Lambda} m \left(\begin{pmatrix} 1 & & \\ & 1 & \\ & & p \end{pmatrix}, \begin{pmatrix} 1 & & \\ & p & \\ & & p \end{pmatrix}; h \right) \Lambda h \Lambda$$

where $h \in GL_3(\mathbb{Q})$ ranges over a system of representatives of the Λ -double cosets contained in the set

$$\Lambda \begin{pmatrix} 1 & & \\ & 1 & \\ & & p \end{pmatrix} \Lambda \begin{pmatrix} 1 & & \\ & p & \\ & & p \end{pmatrix} \Lambda.$$

Let us determine the matrices h occurring in this sum. Let h in $GL_3(\mathbb{Q})$ be such that $\Lambda h \Lambda$ is included in the previous set. By (2.8), one has uniquely

$$\Lambda h \Lambda = \Lambda \varepsilon \frac{\lambda_1}{\lambda_2} \text{diag}(1, s_1, s_2) \Lambda$$

with $\varepsilon = \pm 1$, $\lambda_1, \lambda_2 > 0$, $(\lambda_1, \lambda_2) = 1$, $s_1, s_2 > 0$, $s_1 \mid s_2$. The inclusion is equivalent to

$$\Lambda \varepsilon \lambda_1 \text{diag}(1, s_1, s_2) \Lambda = \Lambda \lambda_2 \delta_1 \delta_2 \Lambda$$

for some matrices $\delta_1 \in R_{1,1,p}$ and $\delta_2 \in L_{1,p,p}$ by (A.3) and (A.4). So, both matrices have the same determinantal divisors ie

$$\begin{aligned} \varepsilon \lambda_1 &= \lambda_2 d_1(\delta_1 \delta_2), \\ \lambda_1^2 s_1 &= \lambda_2^2 d_2(\delta_1 \delta_2), \\ \varepsilon \lambda_1^3 s_1 s_2 &= \lambda_2^3 d_3(\delta_1 \delta_2) = \lambda_2^3 p^3. \end{aligned}$$

One can check that the set

$$\{\delta_1 \delta_2, (\delta_1, \delta_2) \in R_{1,1,p} \times L_{1,p,p}\}$$

is made exactly of the matrices

$$\begin{aligned} \begin{pmatrix} p & & \\ & p & \\ & & p \end{pmatrix} &\rightsquigarrow \mathbf{d}(\delta_1 \delta_2) = (p, p^2, p^3), \\ \begin{pmatrix} p & d_1 + D_1 & \\ & p & \\ & & p \end{pmatrix} &\rightsquigarrow \mathbf{d}(\delta_1 \delta_2) = ((p, d_1 + D_1), p(p, d_1 + D_1), p^3), \\ \begin{pmatrix} p & e_1 + E_1 & \\ & p & f_1 + F_1 \\ & & p \end{pmatrix} &\rightsquigarrow \mathbf{d}(\delta_1 \delta_2) = ((p, e_1 + E_1, f_1 + F_1), p(p, e_1 + E_1, f_1 + F_1), p^3) \end{aligned}$$

and

$$\begin{aligned} \begin{pmatrix} p^2 & & pE_1 \\ & p & F_1 \\ & & 1 \end{pmatrix} &\rightsquigarrow \mathbf{d}(\delta_1 \delta_2) = (1, p, p^3), \\ \begin{pmatrix} p^2 & pD_1 & \\ & 1 & \\ & & p \end{pmatrix} &\rightsquigarrow \mathbf{d}(\delta_1 \delta_2) = (1, p, p^3), \\ \begin{pmatrix} 1 & pd_1 & \\ & p^2 & \\ & & p \end{pmatrix} &\rightsquigarrow \mathbf{d}(\delta_1 \delta_2) = (1, p, p^3), \\ \begin{pmatrix} p & pd_1 & d_1 F_1 + E_1 \\ & p^2 & pF_1 \\ & & 1 \end{pmatrix} &\rightsquigarrow \mathbf{d}(\delta_1 \delta_2) = (1, p, p^3), \\ \begin{pmatrix} 1 & pe_1 & \\ & p & pf_1 \\ & & p^2 \end{pmatrix} &\rightsquigarrow \mathbf{d}(\delta_1 \delta_2) = (1, p, p^3), \\ \begin{pmatrix} p & d_2 & pe_1 \\ & 1 & pf_1 \\ & & p^2 \end{pmatrix} &\rightsquigarrow \mathbf{d}(\delta_1 \delta_2) = (1, p, p^3) \end{aligned}$$

with $0 \leq d_1, e_1, f_1, D_1, E_1, F_1 < p$. As a consequence, only two cases can occur since

$$\mathbf{d}(\delta_1 \delta_2) \in \{(1, p, p^3), (p, p^2, p^3)\}.$$

First case: $\mathbf{d}(\delta_1 \delta_2) = (1, p, p^3)$.

$$\begin{aligned} \varepsilon \lambda_1 &= \lambda_2, \\ \lambda_1^2 s_1 &= \lambda_2^2 p, \\ \varepsilon \lambda_1^3 s_1 s_2 &= \lambda_2^3 p^3. \end{aligned}$$

The first equation gives $\varepsilon = \lambda_1 = \lambda_2 = 1$ by the coprimality of λ_1 and λ_2 . The second equation gives $s_1 = p$. The third equation gives $s_2 = p^2$. Thus,

$$\Lambda h \Lambda = \Lambda \text{diag}(1, p, p^2) \Lambda.$$

Second case: $\mathbf{d}(\delta_1 \delta_2) = (p, p^2, p^3)$.

$$\begin{aligned} \varepsilon \lambda_1 &= \lambda_2 p, \\ \lambda_1^2 s_1 &= \lambda_2^2 p^2, \\ \varepsilon \lambda_1^3 s_1 s_2 &= \lambda_2^3 p^3. \end{aligned}$$

The first equation gives $\varepsilon = \lambda_2 = 1$ and $\lambda_1 = p$ by the coprimality of λ_1 and λ_2 . The second equation gives $s_1 = 1$. The third equation gives $s_2 = 1$. Thus,

$$\Lambda h \Lambda = \Lambda \text{diag}(p, p, p) \Lambda.$$

As a consequence,

$$\Lambda \text{diag}(1, 1, p) \Lambda * \Lambda \text{diag}(1, p, p) \Lambda = m_1 \Lambda \begin{pmatrix} 1 & & \\ & p & \\ & & p^2 \end{pmatrix} \Lambda + m_2 \Lambda \begin{pmatrix} p & & \\ & p & \\ & & p \end{pmatrix} \Lambda$$

where

$$\begin{aligned} m_1 &:= m \left(\begin{pmatrix} 1 & & \\ & 1 & \\ & & p \end{pmatrix}, \begin{pmatrix} 1 & & \\ & p & \\ & & p \end{pmatrix}; \begin{pmatrix} 1 & & \\ & p & \\ & & p^2 \end{pmatrix} \right), \\ m_2 &:= m \left(\begin{pmatrix} 1 & & \\ & 1 & \\ & & p \end{pmatrix}, \begin{pmatrix} 1 & & \\ & p & \\ & & p \end{pmatrix}; \begin{pmatrix} p & & \\ & p & \\ & & p \end{pmatrix} \right). \end{aligned}$$

Let us compute the value of m_2 first. By (2.5), (A.5) and (2.1),

$$m_2 = (p^2 + p + 1) \left| \left\{ \delta_1 \in R_{1,1,p}, \delta_1 \begin{pmatrix} 1 & & \\ & p & \\ & & p \end{pmatrix} \in \Lambda \begin{pmatrix} p & & \\ & p & \\ & & p \end{pmatrix} \Lambda \right\} \right|.$$

Let us compute the remaining cardinality. One can check that the set

$$\left\{ \delta_1 \begin{pmatrix} 1 & & \\ & p & \\ & & p \end{pmatrix}, \delta_1 \in R_{1,1,p} \right\}$$

is exactly made of the matrices

$$\begin{aligned} \begin{pmatrix} p & & \\ & p & \\ & & p \end{pmatrix} &\rightsquigarrow (d_1, d_2, d_3) = (p, p^2, p^3), \\ \begin{pmatrix} 1 & pd_1 & \\ & p^2 & \\ & & p \end{pmatrix} &\rightsquigarrow (d_1, d_2, d_3) = (1, p, p^3), \\ \begin{pmatrix} 1 & 0 & pe_1 \\ & p & pf_1 \\ & & p^2 \end{pmatrix} &\rightsquigarrow (d_1, d_2, d_3) = (1, p, p^3) \end{aligned}$$

with $0 \leq d_1, e_1, f_1 < p$. The fact that the determinantal vector of $\text{diag}(p, p, p)$ is $\text{diag}(p, p^2, p^3)$ implies that

$$\left\{ \delta_1 \in R_{1,1,p}, \delta_1 \begin{pmatrix} 1 & & \\ & p & \\ & & p \end{pmatrix} \in \Lambda \begin{pmatrix} p & & \\ & p & \\ & & p \end{pmatrix} \Lambda \right\} = \left\{ \begin{pmatrix} p & & \\ & p & \\ & & p \end{pmatrix} \right\}$$

and is of cardinality 1 such that $m_2 = p^2 + p + 1$.

Now, let us compute the value of m_1 . By (2.5), (A.5) and (A.7),

$$m_1 = \frac{1}{p(p+1)} \left| \left\{ \delta_1 \in R_{1,1,p}, \delta_1 \begin{pmatrix} 1 & & \\ & p & \\ & & p \end{pmatrix} \in \Lambda \begin{pmatrix} 1 & & \\ & p & \\ & & p^2 \end{pmatrix} \Lambda \right\} \right|.$$

Let us compute the remaining cardinality. Both the analysis done for m_2 and the fact that the determinantal vector of $\text{diag}(1, p, p^2)$ is $(1, p, p^2)$ imply that

$$\begin{aligned} \left\{ \delta_1 \in R_{1,1,p}, \delta_1 \begin{pmatrix} 1 & & \\ & p & \\ & & p \end{pmatrix} \in \Lambda \begin{pmatrix} 1 & & \\ & p & \\ & & p^2 \end{pmatrix} \Lambda \right\} &= \bigcup_{0 \leq d_1 < p} \left\{ \begin{pmatrix} 1 & d_1 & \\ & p & \\ & & 1 \end{pmatrix} \right\} \\ &\quad \bigcup_{0 \leq e_1, f_1 < p} \left\{ \begin{pmatrix} 1 & e_1 & \\ & 1 & f_1 \\ & & p \end{pmatrix} \right\} \end{aligned}$$

which is of cardinality $p(p+1)$ such that $m_1 = 1$.

3.2. Linearization of $\Lambda \text{diag}(1, p, p^2) \Lambda * \Lambda \text{diag}(1, p, p^2) \Lambda$. This section contains the proof of (1.5).

By (2.4), the product of these double cosets equals

$$\sum_{\Lambda h \Lambda \subset \Lambda \text{diag}(1, p, p^2) \Lambda \text{diag}(1, p, p^2) \Lambda} m \left(\begin{pmatrix} 1 & & \\ & p & \\ & & p^2 \end{pmatrix}, \begin{pmatrix} 1 & & \\ & p & \\ & & p^2 \end{pmatrix}; h \right) \Lambda h \Lambda$$

where $h \in GL_3(\mathbb{Q})$ ranges over a system of representatives of the Λ -double cosets contained in the set

$$\Lambda \begin{pmatrix} 1 & & \\ & p & \\ & & p^2 \end{pmatrix} \Lambda \begin{pmatrix} 1 & & \\ & p & \\ & & p^2 \end{pmatrix} \Lambda.$$

Let us determine the relevant matrices h occurring in this sum. Let h in $GL_3(\mathbb{Q})$ be such that $\Lambda h \Lambda$ is included in the previous set. By (2.8), one has uniquely

$$\Lambda h \Lambda = \Lambda \varepsilon \frac{\lambda_1}{\lambda_2} \text{diag}(1, s_1, s_2) \Lambda$$

with $\varepsilon = \pm 1$, $\lambda_1, \lambda_2 > 0$, $(\lambda_1, \lambda_2) = 1$, $s_1, s_2 > 0$, $s_1 \mid s_2$. The inclusion is equivalent to

$$\Lambda \varepsilon \lambda_1 \text{diag}(1, s_1, s_2) \Lambda = \Lambda \lambda_2 \delta_1 \delta_2 \Lambda$$

for some matrices $\delta_1 \in R_{1,p,p^2}$ and $\delta_2 \in L_{1,p,p^2}$ by (A.6). So, both matrices have the same determinantal divisors ie

$$\begin{aligned} \varepsilon \lambda_1 &= \lambda_2 d_1(\delta_1 \delta_2), \\ \lambda_1^2 s_1 &= \lambda_2^2 d_2(\delta_1 \delta_2), \\ \varepsilon \lambda_1^3 s_1 s_2 &= \lambda_2^3 d_3(\delta_1 \delta_2) = \lambda_2^3 p^6. \end{aligned}$$

By (A.6), a straightforward but tedious computation ensures that the set

$$\{\mathbf{d}(\delta_1 \delta_2), (\delta_1, \delta_2) \in R_{1,p,p^2} \times L_{1,p,p^2}\}$$

is a subset of

$$\{(1, p^2, p^6), (1, p^3, p^6), (p, p^2, p^6), (p, p^3, p^6), (p^2, p^4, p^6)\}.$$

Case 1: $(d_1, d_2, d_3) = (1, p^2, p^6)$.

$$\begin{aligned} \varepsilon \lambda_1 &= \lambda_2, \\ \lambda_1^2 s_1 &= \lambda_2^2 p^2, \\ \varepsilon \lambda_1^3 s_1 s_2 &= \lambda_2^3 p^6. \end{aligned}$$

The first equation gives $\varepsilon = \lambda_1 = \lambda_2 = 1$ by the coprimality of λ_1 and λ_2 . The second equation gives $s_1 = p^2$. The third equation gives $s_2 = p^4$. Thus,

$$\Lambda h \Lambda = \Lambda \text{diag}(1, p^2, p^4) \Lambda.$$

Case 2: $(d_1, d_2, d_3) = (1, p^3, p^6)$.

$$\begin{aligned} \varepsilon \lambda_1 &= \lambda_2, \\ \lambda_1^2 s_1 &= \lambda_2^2 p^3, \\ \varepsilon \lambda_1^3 s_1 s_2 &= \lambda_2^3 p^6. \end{aligned}$$

The first equation gives $\varepsilon = \lambda_1 = \lambda_2 = 1$ by the coprimality of λ_1 and λ_2 . The second equation gives $s_1 = p^3$. The third equation gives $s_2 = p^3$. Thus,

$$\Lambda h \Lambda = \Lambda \text{diag}(1, p^3, p^3) \Lambda.$$

Case 3: $(d_1, d_2, d_3) = (p, p^2, p^6)$.

$$\begin{aligned} \varepsilon \lambda_1 &= \lambda_2 p, \\ \lambda_1^2 s_1 &= \lambda_2^2 p^2, \\ \varepsilon \lambda_1^3 s_1 s_2 &= \lambda_2^3 p^6. \end{aligned}$$

The first equation gives $\varepsilon = \lambda_2 = 1$ and $\lambda_1 = p$ by the coprimality of λ_1 and λ_2 . The second equation gives $s_1 = 1$. The third equation gives $s_2 = p^3$. Thus,

$$\Lambda h \Lambda = \Lambda \text{diag}(p, p, p^4) \Lambda.$$

Case 4: $(d_1, d_2, d_3) = (p, p^3, p^6)$.

$$\begin{aligned}\varepsilon\lambda_1 &= \lambda_2 p, \\ \lambda_1^2 s_1 &= \lambda_2^2 p^3, \\ \varepsilon\lambda_1^3 s_1 s_2 &= \lambda_2^3 p^6.\end{aligned}$$

The first equation gives $\varepsilon = \lambda_2 = 1$ and $\lambda_1 = p$ by the coprimality of λ_1 and λ_2 . The second equation gives $s_1 = p$. The third equation gives $s_2 = p^2$. Thus,

$$\Lambda h \Lambda = \Lambda \text{diag}(p, p^2, p^3) \Lambda.$$

Case 5: $(d_1, d_2, d_3) = (p^2, p^4, p^6)$.

$$\begin{aligned}\varepsilon\lambda_1 &= \lambda_2 p^2, \\ \lambda_1^2 s_1 &= \lambda_2^2 p^4, \\ \varepsilon\lambda_1^3 s_1 s_2 &= \lambda_2^3 p^6.\end{aligned}$$

The first equation gives $\varepsilon = \lambda_2 = 1$ and $\lambda_1 = p^2$ by the coprimality of λ_1 and λ_2 . The second equation gives $s_1 = 1$. The third equation gives $s_2 = 1$. Thus,

$$\Lambda h \Lambda = \Lambda \text{diag}(p^2, p^2, p^2) \Lambda.$$

As a consequence,

$$\begin{aligned}\Lambda \text{diag}(1, p, p^2) \Lambda * \Lambda \text{diag}(1, p, p^2) \Lambda &= m_1 \Lambda \begin{pmatrix} 1 & & \\ & p^2 & \\ & & p^4 \end{pmatrix} \Lambda + m_2 \Lambda \begin{pmatrix} 1 & & \\ & p^3 & \\ & & p^3 \end{pmatrix} \Lambda \\ &+ m_3 \Lambda \begin{pmatrix} p & & \\ & p & \\ & & p^4 \end{pmatrix} \Lambda + m_4 \Lambda \begin{pmatrix} p & & \\ & p^2 & \\ & & p^3 \end{pmatrix} \Lambda + m_5 \Lambda \begin{pmatrix} p^2 & & \\ & p^2 & \\ & & p^2 \end{pmatrix} \Lambda\end{aligned}$$

where

$$\begin{aligned}m_1 &:= m \left(\begin{pmatrix} 1 & & \\ & p & \\ & & p^2 \end{pmatrix}, \begin{pmatrix} 1 & & \\ & p & \\ & & p^2 \end{pmatrix}; \begin{pmatrix} 1 & & \\ & p^2 & \\ & & p^4 \end{pmatrix} \right), \\ m_2 &:= m \left(\begin{pmatrix} 1 & & \\ & p & \\ & & p^2 \end{pmatrix}, \begin{pmatrix} 1 & & \\ & p & \\ & & p^2 \end{pmatrix}; \begin{pmatrix} 1 & & \\ & p^3 & \\ & & p^3 \end{pmatrix} \right), \\ m_3 &:= m \left(\begin{pmatrix} 1 & & \\ & p & \\ & & p^2 \end{pmatrix}, \begin{pmatrix} 1 & & \\ & p & \\ & & p^2 \end{pmatrix}; \begin{pmatrix} p & & \\ & p & \\ & & p^4 \end{pmatrix} \right), \\ m_4 &:= m \left(\begin{pmatrix} 1 & & \\ & p & \\ & & p^2 \end{pmatrix}, \begin{pmatrix} 1 & & \\ & p & \\ & & p^2 \end{pmatrix}; \begin{pmatrix} p & & \\ & p^2 & \\ & & p^3 \end{pmatrix} \right), \\ m_5 &:= m \left(\begin{pmatrix} 1 & & \\ & p & \\ & & p^2 \end{pmatrix}, \begin{pmatrix} 1 & & \\ & p & \\ & & p^2 \end{pmatrix}; \begin{pmatrix} p^2 & & \\ & p^2 & \\ & & p^2 \end{pmatrix} \right).\end{aligned}$$

Let us compute the value of m_1 . By (2.5), (A.7) and (A.8),

$$m_1 = \frac{1}{p^4} \left| \left\{ \delta_1 \in R_{1,p,p^2}, \delta_1 \begin{pmatrix} 1 & & \\ & p & \\ & & p^2 \end{pmatrix} \in \Lambda \begin{pmatrix} 1 & & \\ & p^2 & \\ & & p^4 \end{pmatrix} \Lambda \right\} \right|.$$

Let us compute the remaining cardinality. One can check that the set

$$\left\{ \delta_1 \begin{pmatrix} 1 & & \\ & p & \\ & & p^2 \end{pmatrix}, \delta_1 \in R_{1,p,p^2} \right\}$$

is exactly made of the matrices

$$\begin{pmatrix} p^2 & & \\ & p & p^2 f_1 \\ & & p^3 \end{pmatrix} \rightsquigarrow (d_1, d_2, d_3) = (p, p^3, p^6),$$

$$\begin{pmatrix} p & p d_2 & \\ & p^3 & \\ & & p^2 \end{pmatrix} (p \mid d_2) \rightsquigarrow (d_1, d_2, d_3) = (p, p^3, p^6),$$

$$\begin{pmatrix} p & p d_1 & p^2 e_1 \\ & p^2 & p^2 f_1 \\ & & p^3 \end{pmatrix} (d_1 f_1 = 0, (d_1, e_1, f_1) \neq (0, 0, 0)) \rightsquigarrow (d_1, d_2, d_3) = (p, p^3, p^6)$$

and

$$\begin{pmatrix} 1 & p d_1 & p^2 e_2 \\ & p^2 & p^2 f_2 \\ & & p^4 \end{pmatrix} (p \mid f_2) \rightsquigarrow (d_1, d_2, d_3) = (1, p^2, p^6)$$

and

$$\begin{pmatrix} 1 & p d_2 & p^2 e_1 \\ & p^3 & \\ & & p^3 \end{pmatrix} \rightsquigarrow (d_1, d_2, d_3) = (1, p^3, p^6)$$

and

$$\begin{pmatrix} p & p^2 e_2 \\ & p & p^2 f_2 \\ & & p^4 \end{pmatrix} (p \mid e_2) \rightsquigarrow (d_1, d_2, d_3) = (p, p^2, p^6)$$

and

$$\begin{pmatrix} p^2 & & \\ & p^2 & \\ & & p^2 \end{pmatrix} \rightsquigarrow (d_1, d_2, d_3) = (p^2, p^4, p^6)$$

where $0 \leq d_1, e_1, f_1 < p$ and $0 \leq d_2, e_2, f_2 < p^2$. The fact that the determinantal vector of $\text{diag}(1, p^2, p^4)$ is $(1, p^2, p^6)$ implies that $m_1 = 1$.

Let us compute the value of m_2 . By (2.5), (A.7) and (A.9),

$$m_2 = \frac{p+1}{p^3} \left| \left\{ \delta_1 \in R_{1,p,p^2}, \delta_1 \begin{pmatrix} 1 & & \\ & p & \\ & & p^2 \end{pmatrix} \in \Lambda \begin{pmatrix} 1 & & \\ & p^3 & \\ & & p^3 \end{pmatrix} \Lambda \right\} \right|.$$

Both the analysis done for m_1 and the fact that the determinantal vector of $\text{diag}(1, p^3, p^3)$ is $(1, p^3, p^6)$ imply that $m_2 = p+1$.

Let us compute the value of m_3 . By (2.5), (A.7), (2.1) and (A.10),

$$m_3 = \frac{p+1}{p^3} \left| \left\{ \delta_1 \in R_{1,p,p^2}, \delta_1 \begin{pmatrix} 1 & & \\ & p & \\ & & p^2 \end{pmatrix} \in \Lambda \begin{pmatrix} p & & \\ & p & \\ & & p^4 \end{pmatrix} \Lambda \right\} \right|.$$

Both the analysis done for m_1 and the fact that the determinantal vector of $\text{diag}(p, p, p^4)$ is (p, p^2, p^6) imply that $m_3 = p + 1$.

Let us compute the value of m_4 . By (2.5), (A.7), (2.1) and (A.7),

$$m_4 = \frac{p+1}{p^3} \left| \left\{ \delta_1 \in R_{1,p,p^2}, \delta_1 \begin{pmatrix} 1 & & \\ & p & \\ & & p^2 \end{pmatrix} \in \Lambda \begin{pmatrix} p & & \\ & p^2 & \\ & & p^3 \end{pmatrix} \Lambda \right\} \right|.$$

Both the analysis done for m_1 and the fact that the determinantal vector of $\text{diag}(p, p^2, p^3)$ is (p, p^3, p^6) imply that $m_4 = (p+1)(2p-1)$.

Let us compute the value of m_5 . By (2.5), (A.7) and (2.1),

$$m_5 = p(p+1)(p^2+p+1) \left| \left\{ \delta_1 \in R_{1,p,p^2}, \delta_1 \begin{pmatrix} 1 & & \\ & p & \\ & & p^2 \end{pmatrix} \in \Lambda \begin{pmatrix} p^2 & & \\ & p^2 & \\ & & p^2 \end{pmatrix} \Lambda \right\} \right|.$$

Both the analysis done for m_1 and the fact that the determinantal vector of $\text{diag}(p^2, p^2, p^2)$ is (p^2, p^4, p^6) imply that $m_5 = p(p+1)(p^2+p+1)$.

APPENDIX A. DECOMPOSITION OF Λ -DOUBLE COSETS INTO Λ -COSETS

By [AZ95, Lemma 1.2 Page 94 and Lemma 2.1 Page 105], we know that every Λ -double coset $\Lambda g \Lambda$ with g in $GL_3(\mathbb{Q})$ with integer coefficients is both a finite union of Λ -left cosets and Λ -right cosets. In addition, every Λ -right coset Λg contains a unique upper-triangular column reduced representative, namely

$$\Lambda g = \Lambda \begin{pmatrix} a & d & e \\ & b & f \\ & & c \end{pmatrix}$$

where $0 \leq d < b$ and $0 \leq e, f < c$ by [AZ95, Lemma 2.7 Page 109].

As a consequence, every Λ -left coset $g \Lambda$ contains a unique upper-triangular row reduced representative, namely

$$g \Lambda = \begin{pmatrix} a & d & e \\ & b & f \\ & & c \end{pmatrix} \Lambda$$

where $0 \leq d, e < a$ and $0 \leq f < b$. More explicitly, if $UW^t g W = H$ is the upper-triangular column reduced representative of the Λ -right coset $\Lambda W^t g W$ with W the anti-diagonal matrix with 1's on the anti-diagonal then $g W^t U W = W^t H W$ is the upper-triangular row reduced representative of the Λ -left coset $g \Lambda$.

The previous fact also entails that

$$\Lambda g \Lambda = \bigcup_{\delta \in R_g} \Lambda \delta \Rightarrow \Lambda g \Lambda = \bigcup_{\delta \in W^t R_g W} \delta \Lambda \quad (\text{A.1})$$

since

$$\Lambda g \Lambda = W \Lambda g \Lambda = W^t (\Lambda g \Lambda) = W \bigcup_{\delta \in R_g} {}^t \delta \Lambda = \bigcup_{\delta \in R_g} W^t \delta W \Lambda.$$

Let us finish with a useful elementary practical remark for the computations done in the following sections of the appendix. If H is an upper-triangular column

reduced matrix in a Λ -double coset $\Lambda \text{diag}(p^{\alpha_1}, p^{\alpha_2}, p^{\alpha_3}) \Lambda$ where p is a prime number and α_1, α_2 and α_3 are non-negative integers then

$$H = \begin{pmatrix} p^{\delta_1} & * & * \\ & p^{\delta_2} & * \\ & & p^{\delta_3} \end{pmatrix}, \sum_{j=1}^3 (\alpha_j - \delta_j) = 0, \forall j \in \{1, 2, 3\}, 0 \leq \delta_j \leq \max_{1 \leq k \leq 3} \alpha_k. \quad (\text{A.2})$$

The fact that the diagonal coefficients of H are powers of p comes from the determinant equation. The condition on the exponents of these diagonal coefficients follows from the fact that $p^{\max\{\alpha_k, 1 \leq k \leq 3\}} H^{-1}$ has integer coefficients.

A.1. Decomposition and degree of $\Lambda \text{diag}(1, 1, p) \Lambda$.

Proposition A.1—One has

$$\Lambda \text{diag}(1, 1, p) \Lambda = \bigcup_{\delta \in R_{1,1,p}} \Lambda \delta = \bigcup_{\delta \in L_{1,1,p}} \delta \Lambda \quad (\text{A.3})$$

where

$$R_{1,1,p} = \{\text{diag}(p, 1, 1)\} \bigcup_{0 \leq d_1 < p} \left\{ \begin{pmatrix} 1 & d_1 & \\ & p & \\ & & 1 \end{pmatrix} \right\} \bigcup_{0 \leq e_1, f_1 < p} \left\{ \begin{pmatrix} 1 & 0 & e_1 \\ & 1 & f_1 \\ & & p \end{pmatrix} \right\}$$

and

$$L_{1,1,p} = \{\text{diag}(1, 1, p)\} \bigcup_{0 \leq f_1 < p} \left\{ \begin{pmatrix} 1 & & \\ & p & f_1 \\ & & 1 \end{pmatrix} \right\} \bigcup_{0 \leq d_1, e_1 < p} \left\{ \begin{pmatrix} p & d_1 & e_1 \\ & 1 & \\ & & 1 \end{pmatrix} \right\}.$$

In particular,

$$\deg(\text{diag}(1, 1, p)) = p^2 + p + 1.$$

Proof of Proposition A.1. The decomposition into Λ -right cosets implies the decomposition into Λ -left cosets by (A.1). The possible upper-triangular column reduced matrices δ that can occur in the decomposition into Λ -right cosets are

$$\begin{aligned} \begin{pmatrix} 1 & 0 & e_1 \\ & 1 & f_1 \\ & & p \end{pmatrix} &\rightsquigarrow \mathbf{d}(\delta) = (1, 1, p), \\ \begin{pmatrix} 1 & d_1 & 0 \\ & p & 0 \\ & & 1 \end{pmatrix} &\rightsquigarrow \mathbf{d}(\delta) = (1, 1, p), \\ \begin{pmatrix} p & 0 & 0 \\ & 1 & 0 \\ & & 1 \end{pmatrix} &\rightsquigarrow \mathbf{d}(\delta) = (1, 1, p) \end{aligned}$$

where $0 \leq d_1, e_1, f_1 < p$. The fact that the determinantal vector of $\text{diag}(1, 1, p)$ is $(1, 1, p)$ implies the decomposition into Λ -left cosets given in (A.3) and the computation of the degree too. \square

A.2. Decomposition and degree of $\Lambda \text{diag}(1, p, p)\Lambda$.

Proposition A.2– One has

$$\Lambda \text{diag}(1, p, p)\Lambda = \cup_{\delta \in L_{1,p,p}} \delta \Lambda \quad (\text{A.4})$$

where

$$L_{1,p,p} = \{\text{diag}(1, p, p)\} \cup_{0 \leq e_1, f_1 < p} \left\{ \begin{pmatrix} p & e_1 \\ & p & f_1 \\ & & 1 \end{pmatrix} \right\} \cup_{0 \leq d_1 < p} \left\{ \begin{pmatrix} p & d_1 \\ & 1 \\ & & p \end{pmatrix} \right\}$$

In particular,

$$\deg(\text{diag}(1, p, p)) = p^2 + p + 1. \quad (\text{A.5})$$

Proof of Proposition A.2. By (A.2), the possible upper-triangular row reduced matrices δ that can occur in the decomposition into Λ -left cosets are

$$\begin{aligned} \begin{pmatrix} p & d_1 & e_1 \\ & p & f_1 \\ & & 1 \end{pmatrix} &\rightsquigarrow \mathbf{d}(\delta) = (1, (p, d_1, d_1 f_1), p^2), \\ \begin{pmatrix} p & d_1 & e_1 \\ & 1 \\ & & p \end{pmatrix} &\rightsquigarrow \mathbf{d}(\delta) = (1, (p, e_1), p^2), \\ \begin{pmatrix} 1 & & \\ & p & f_1 \\ & & p \end{pmatrix} &\rightsquigarrow \mathbf{d}(\delta) = (1, (p, f_1), p^2) \end{aligned}$$

where $0 \leq d_1, e_1, f_1 < p$. The fact that the determinantal vector of $\text{diag}(1, p, p)$ is $(1, p, p^2)$ implies the decomposition into Λ -left cosets given in (A.4) and the computation of the degree too. \square

A.3. Decomposition and degree of $\Lambda \text{diag}(1, p, p^2)\Lambda$.

Proposition A.3– One has

$$\Lambda \text{diag}(1, p, p^2)\Lambda = \cup_{\delta \in R_{1,p,p^2}} \Lambda \delta = \cup_{\delta \in L_{1,p,p^2}} \delta \Lambda \quad (\text{A.6})$$

where

$$\begin{aligned} R_{1,p,p^2} = & \cup_{\substack{0 \leq d_1 < p \\ 0 \leq e_2, f_2 < p^2 \\ p \mid f_2}} \left\{ \begin{pmatrix} 1 & d_1 & e_2 \\ & p & f_2 \\ & & p^2 \end{pmatrix} \right\} \cup_{\substack{0 \leq e_1 < p \\ 0 \leq d_2 < p^2}} \left\{ \begin{pmatrix} 1 & d_2 & e_1 \\ & p^2 \\ & & p \end{pmatrix} \right\} \\ & \cup_{\substack{0 \leq e_2, f_2 < p^2 \\ p \mid e_2}} \left\{ \begin{pmatrix} p & & e_2 \\ & 1 & f_2 \\ & & p^2 \end{pmatrix} \right\} \cup_{0 \leq f_1 < p} \left\{ \begin{pmatrix} p^2 & & \\ & 1 & f_1 \\ & & p \end{pmatrix} \right\} \cup_{\substack{0 \leq d_2 < p^2 \\ p \mid d_2}} \left\{ \begin{pmatrix} p & d_2 \\ & p^2 \\ & & 1 \end{pmatrix} \right\} \\ & \cup_{\substack{0 \leq d_1, e_1, f_1 < p \\ d_1 f_1 = 0 \\ (d_1, e_1, f_1) \neq (0,0,0)}} \left\{ \begin{pmatrix} p & d_1 & e_1 \\ & p & f_1 \\ & & p \end{pmatrix} \right\} \cup \left\{ \begin{pmatrix} p^2 & & \\ & p & \\ & & 1 \end{pmatrix} \right\} \end{aligned}$$

and

$$\begin{aligned}
L_{1,p,p^2} = & \bigcup_{\substack{0 \leq f_1 < p \\ 0 \leq d_2, e_2 < p^2 \\ p | d_2}} \left\{ \begin{pmatrix} p^2 & d_2 & e_2 \\ & p & f_1 \\ & & 1 \end{pmatrix} \right\} \bigcup_{\substack{0 \leq e_1 < p \\ 0 \leq f_2 < p^2}} \left\{ \begin{pmatrix} p & & e_1 \\ & p^2 & f_2 \\ & & 1 \end{pmatrix} \right\} \\
& \bigcup_{\substack{0 \leq d_2, e_2 < p^2 \\ p | e_2}} \left\{ \begin{pmatrix} p^2 & d_2 & e_2 \\ & 1 & \\ & & p \end{pmatrix} \right\} \bigcup_{0 \leq d_1 < p} \left\{ \begin{pmatrix} p & d_1 & \\ & 1 & \\ & & p^2 \end{pmatrix} \right\} \bigcup_{\substack{0 \leq f_2 < p^2 \\ p | f_2}} \left\{ \begin{pmatrix} 1 & & f_2 \\ & p^2 & \\ & & p \end{pmatrix} \right\} \\
& \bigcup_{\substack{0 \leq d_1, e_1, f_1 < p \\ d_1 f_1 = 0 \\ (d_1, e_1, f_1) \neq (0,0,0)}} \left\{ \begin{pmatrix} p & d_1 & e_1 \\ & p & f_1 \\ & & p \end{pmatrix} \right\} \bigcup \left\{ \begin{pmatrix} p^2 & & \\ & p & \\ & & 1 \end{pmatrix} \right\}.
\end{aligned}$$

In particular,

$$\deg(\text{diag}(1, p, p^2)) = p(p+1)(1+p+p^2). \quad (\text{A.7})$$

Proof of Proposition A.3. The decomposition into Λ -right cosets implies the decomposition into Λ -left cosets by (A.1). By (A.2), the possible upper-triangular column reduced matrices δ that can occur in the decomposition into Λ -right cosets are

$$\text{Type 1: } \begin{pmatrix} p & d_1 & e_1 \\ & p & f_1 \\ & & p \end{pmatrix} \rightsquigarrow \mathbf{d}(\delta) = ((p, d_1, e_1, f_1), (p^2, p d_1, p f_1, d_1 f_1 - p e_1), p^3)$$

and

$$\text{Type 2: } \begin{pmatrix} 1 & d_1 & e_2 \\ & p & f_2 \\ & & p^2 \end{pmatrix} \rightsquigarrow \mathbf{d}(\delta) = (1, (p, f_2), p^3),$$

$$\text{Type 3: } \begin{pmatrix} 1 & d_2 & e_1 \\ & p^2 & f_1 \\ & & p \end{pmatrix} \rightsquigarrow \mathbf{d}(\delta) = (1, (p, f_1), p^3),$$

$$\text{Type 4: } \begin{pmatrix} p & & e_2 \\ & 1 & f_2 \\ & & p^2 \end{pmatrix} \rightsquigarrow \mathbf{d}(\delta) = (1, (p, e_2), p^3),$$

$$\text{Type 5: } \begin{pmatrix} p^2 & & e_1 \\ & 1 & f_1 \\ & & p \end{pmatrix} \rightsquigarrow \mathbf{d}(\delta) = (1, (p, e_1), p^3),$$

$$\text{Type 6: } \begin{pmatrix} p & d_2 & \\ & p^2 & \\ & & 1 \end{pmatrix} \rightsquigarrow \mathbf{d}(\delta) = (1, (p, d_2), p^3),$$

$$\text{Type 7: } \begin{pmatrix} p^2 & d_1 & \\ & p & \\ & & 1 \end{pmatrix} \rightsquigarrow \mathbf{d}(\delta) = (1, (p, d_1), p^3)$$

where $0 \leq d_1, e_1, f_1 < p$ and $0 \leq d_2, e_2, f_2 < p^2$. Let us count the matrices among the previous ones, whose determinantal vector is the same as the one of $\text{diag}(1, p, p^2)$, namely $(1, p, p^3)$.

Let us consider the matrices of type 1. The condition on $d_2(\delta)$ implies $d_1 \neq 0$. The condition on $d_1(\delta)$ implies that $(d_1, e_1, f_1) \neq (0, 0, 0)$. The condition on $d_2(\delta)$ implies $p \mid d_1 f_1$ such that $p \mid d_1$ or $p \mid f_1$, namely $d_1 = 0$ or $f_1 = 0$. There are $(p-1)(2p+1)$ such matrices of type 1.

Let us consider the matrices of type 2. The condition on $d_2(\delta)$ implies $p \mid f_2$. There are p^4 such matrices of type 2.

Let us consider the matrices of type 3. The condition on $d_2(\delta)$ implies $f_1 = 0$. There are p^3 such matrices of type 3.

Let us consider the matrices of type 4. The condition on $d_2(\delta)$ implies $p \mid e_2$. There are p^3 such matrices of type 4.

Let us consider the matrices of type 5. The condition on $d_2(\delta)$ implies $e_1 = 0$. There are p such matrices of type 5.

Let us consider the matrices of type 6. The condition on $d_2(\delta)$ implies $p \mid d_2$. There are p such matrices of type 6.

Let us consider the matrices of type 7. The condition on $d_2(\delta)$ implies $d_1 = 0$. There is 1 such matrix of type 7.

One can recover the decomposition in Λ -right cosets given in (A.6) and the value of the degree given in (A.7) by summing all the contributions in the previous paragraphs. \square

A.4. Degree of $\Lambda \text{diag}(1, p^2, p^4) \Lambda$.

Proposition A.4– One has

$$\deg(\text{diag}(1, p^2, p^4)) = p^5(p+1)(p^2+p+1). \quad (\text{A.8})$$

Proof of Proposition A.4. By (A.2), the possible upper-triangular column reduced matrices δ that can occur in the decomposition into Λ -right cosets are

$$\begin{aligned} \text{Type 1: } & \begin{pmatrix} p^4 & d_2 & \\ & p^2 & \\ & & 1 \end{pmatrix} \rightsquigarrow \mathbf{d}(\delta) = (1, (p^2, d_2), p^6), \\ \text{Type 2: } & \begin{pmatrix} p^4 & & e_2 \\ & 1 & f_2 \\ & & p^2 \end{pmatrix} \rightsquigarrow \mathbf{d}(\delta) = (1, (p^2, e_2), p^6), \\ \text{Type 3: } & \begin{pmatrix} p^2 & d_4 & \\ & p^4 & \\ & & 1 \end{pmatrix} \rightsquigarrow \mathbf{d}(\delta) = (1, (p^2, d_4), p^6), \\ \text{Type 4: } & \begin{pmatrix} p^2 & & e_4 \\ & 1 & f_4 \\ & & p^4 \end{pmatrix} \rightsquigarrow \mathbf{d}(\delta) = (1, (p^2, e_4), p^6), \\ \text{Type 5: } & \begin{pmatrix} 1 & d_4 & e_2 \\ & p^4 & f_2 \\ & & p^2 \end{pmatrix} \rightsquigarrow \mathbf{d}(\delta) = (1, (p^2, f_2, d_4 f_2), p^6), \\ \text{Type 6: } & \begin{pmatrix} 1 & d_2 & e_4 \\ & p^2 & f_4 \\ & & p^4 \end{pmatrix} \rightsquigarrow \mathbf{d}(\delta) = (1, (p^2, f_4, d_2 f_4), p^6) \end{aligned}$$

and

$$\text{Type 7: } \begin{pmatrix} p^4 & d_1 & e_1 \\ & p & f_1 \\ & & p \end{pmatrix} \rightsquigarrow \mathbf{d}(\delta) = ((p, d_1, e_1, f_1), (p^2, pd_1, d_1f_1 - pe_1), p^6),$$

$$\text{Type 8: } \begin{pmatrix} p & d_4 & e_1 \\ & p^4 & f_1 \\ & & p \end{pmatrix} \rightsquigarrow \mathbf{d}(\delta) = ((p, d_4, e_1, f_1), (p^2, pf_1, pd_4, d_4f_1), p^6),$$

$$\text{Type 9: } \begin{pmatrix} p & d_1 & e_4 \\ & p & f_4 \\ & & p^4 \end{pmatrix} \rightsquigarrow \mathbf{d}(\delta) = ((p, d_1, e_4, f_4), (p^2, pf_4, d_1f_4 - pe_4), p^6)$$

and

$$\text{Type 10: } \begin{pmatrix} p^3 & d_3 & \\ & p^3 & \\ & & 1 \end{pmatrix} \rightsquigarrow \mathbf{d}(\delta) = (1, (p^3, d_3), p^6),$$

$$\text{Type 11: } \begin{pmatrix} p^3 & & e_3 \\ & 1 & f_3 \\ & & p^3 \end{pmatrix} \rightsquigarrow \mathbf{d}(\delta) = (1, (p^3, e_3), p^6),$$

$$\text{Type 12: } \begin{pmatrix} 1 & d_3 & e_3 \\ & p^3 & f_3 \\ & & p^3 \end{pmatrix} \rightsquigarrow \mathbf{d}(\delta) = (1, (p^3, f_3, d_3f_3), p^6)$$

and

$$\text{Type 13: } \begin{pmatrix} p^3 & d_2 & e_1 \\ & p^2 & f_1 \\ & & p \end{pmatrix} \rightsquigarrow \mathbf{d}(\delta) = ((p, d_2, e_1, f_1), (p^3, pd_2, d_2f_1 - p^2e_1), p^6)$$

$$\text{Type 14: } \begin{pmatrix} p^3 & d_1 & e_2 \\ & p & f_2 \\ & & p^2 \end{pmatrix} \rightsquigarrow \mathbf{d}(\delta) = ((p, d_1, e_2, f_2), (p^3, p^2d_1, d_1f_2 - pe_2), p^6)$$

$$\text{Type 15: } \begin{pmatrix} p^2 & d_3 & e_1 \\ & p^3 & f_1 \\ & & p \end{pmatrix} \rightsquigarrow \mathbf{d}(\delta) = ((p, d_3, e_1, f_1), (p^3, pd_3, p^2f_1, d_3f_1), p^6)$$

$$\text{Type 16: } \begin{pmatrix} p^2 & d_1 & e_3 \\ & p & f_3 \\ & & p^3 \end{pmatrix} \rightsquigarrow \mathbf{d}(\delta) = ((p, d_1, e_3, f_3), (p^3, d_1f_3 - pe_3, p^2f_3), p^6)$$

and

$$\text{Type 17: } \begin{pmatrix} p & d_3 & e_2 \\ & p^3 & f_2 \\ & & p^2 \end{pmatrix} \rightsquigarrow \mathbf{d}(\delta) = ((p, d_3, e_2, f_2), (p^3, pf_2, p^2d_3, d_3f_2), p^6)$$

$$\text{Type 18: } \begin{pmatrix} p & d_2 & e_3 \\ & p^2 & f_3 \\ & & p^3 \end{pmatrix} \rightsquigarrow \mathbf{d}(\delta) = ((p, d_2, e_3, f_3), (p^3, pf_3, d_2f_3 - p^2e_3), p^6)$$

and

$$\text{Type 19: } \begin{pmatrix} p^2 & d_2 & e_2 \\ & p^2 & f_2 \\ & & p^2 \end{pmatrix} \rightsquigarrow \mathbf{d}(\delta) = ((p^2, d_2, e_2, f_2), (p^4, p^2d_2, p^2f_2, d_2f_2 - p^2e_2), p^6)$$

where $0 \leq d_j, e_j, f_j < p^j$ for $j = 1, 2, 3, 4$. Let us count the matrices among the previous ones, whose determinantal vector is the same as the one of $\text{diag}(1, p^2, p^4)$, namely $(1, p^2, p^6)$.

Let us consider the matrices of type 1. The condition on $d_2(\delta)$ implies $d_2 = 0$. There is 1 relevant matrix of type 1.

Let us consider the matrices of type 2. The condition on $d_2(\delta)$ implies $e_2 = 0$. There are p^2 relevant matrices of type 2.

Let us consider the matrices of type 3. The condition on $d_2(\delta)$ implies $p^2 \mid d_4$. There are p^2 relevant matrices of type 3.

Let us consider the matrices of type 4. The condition on $d_2(\delta)$ implies $p^2 \mid e_4$. There are p^6 relevant matrices of type 4.

Let us consider the matrices of type 5. The condition on $d_2(\delta)$ implies $e_2 = 0$. There are p^6 relevant matrices of type 5.

Let us consider the matrices of type 6. The condition on $d_2(\delta)$ implies $p^2 \mid f_4$. There are p^8 relevant matrices of type 6.

Let us consider the matrices of type 7. The condition on $d_2(\delta)$ implies $d_1 = e_1 = 0$ and the condition on $d_1(\delta)$ implies $f_1 \neq 0$. There are $p - 1$ relevant matrices of type 7.

Let us consider the matrices of type 8. The condition on $d_2(\delta)$ implies $f_1 = 0$ and $p \mid d_4$. The condition on $d_1(\delta)$ implies $e_1 \neq 0$. There are $p^3(p - 1)$ relevant matrices of type 8.

Let us consider the matrices of type 9. The condition on $d_2(\delta)$ implies $p \mid f_4$ and $p \mid d_1 f_4 / p - e_4$. One has $d_1 \neq 0$ since otherwise $d_1(\delta) = 1 = (p, e_4)$ and $d_2(\delta) = p(p, e_4) = p \neq p^2$. Thus, d_1 is invertible modulo p and $f_4 / p \equiv e_4 \overline{d_1} \pmod{p}$ such that f_4 / p can take p^2 values. There are $(p - 1)p^6$ relevant matrices of type 9.

Let us consider the matrices of type 10. The condition on $d_2(\delta)$ implies $p^2 \parallel d_3$. There are $p - 1$ relevant matrices of type 10.

Let us consider the matrices of type 11. The condition on $d_2(\delta)$ implies $p^2 \parallel e_3$. There are $(p - 1)p^3$ relevant matrices of type 11.

Let us consider the matrices of type 12. The condition on $d_2(\delta)$ implies $p^2 \parallel f_3$. There are $(p - 1)p^6$ relevant matrices of type 12.

Let us consider the matrices of type 13. Note that $(e_1, f_1) \neq (0, 0)$ since otherwise $d_1(\delta) = 1 = (p, d_2)$, which implies that $d_2(\delta) = (pd_2, p^3) = p \neq p^2$. As a consequence, $d_1(\delta) = 1 = (p, d_2, e_1, f_1)$. The fact that $d_2(\delta) = p^2$ implies that $p \mid d_2$ and $p \mid f_1 d_2 / p$, namely $f_1 = 0$ or $d_2 = 0$. If $d_2 = 0$ then $d_2(\delta) = p^2 = (p^3, p^2 e_1)$ such that $e_1 \neq 0$. There are $p(p - 1)$ such matrices. If $d_2 \neq 0$ then $f_1 = 0$, $d_2(\delta) = p^2(p, d_2 / p, e_1) = p^2$ since d_2 / p is coprime with p and $d_1(\delta) = 1 = (p, e_1)$ such that $e_1 \neq 0$. There are $(p - 1)^2$ such matrices. Finally, there are $(p - 1)(2p - 1)$ relevant matrices of type 13.

Let us consider the matrices of type 14. The fact that $d_2(\delta) = p^2$ implies that $p^2 \mid d_1 f_2 - p e_2$. If $d_1 = 0$ then $p \mid e_2$ and $d_2(\delta) = p^2 = (p^3, p^2 e_2 / p)$ if $e_2 \neq 0$. $d_1(\delta) = 1 = (p, f_2)$ implies that $p \nmid f_2$. There are $(p - 1)(p^2 - p)$ such matrices. If $d_1 \neq 0$ then the value of f_2 is fixed by $f_2 \equiv p e_2 \overline{d_1} \pmod{p^2}$ and $d_1(\delta) = (p, d_1) = 1$. There are $p^2(p - 1)$ such matrices. Finally, there are $(p - 1)(2p^2 - p)$ relevant matrices of type 14.

Let us consider the matrices of type 15. The condition $d_2(\delta) = p^2$ implies that $p \mid d_3$ and $p \mid f_1 d_3 / p$. If $f_1 = 0$ then $d_2(\delta) = p^2 = p^2(p, d_3 / p)$ such that $p \parallel d_3$. The condition $d_1(\delta) = 1 = (p, e_1)$ implies that $e_1 \neq 0$. There are $(p^2 - p)(p - 1)$ such

matrices. If $f_1 \neq 0$ then $p^2 \mid d_3$ and $d_1(\delta) = 1$. There are $p^2(p-1)$ such matrices. Finally, there are $(p-1)(2p^2-p)$ relevant matrices of type 15.

Let us consider the matrices of type 16. The condition $d_2(\delta) = p^2$ implies that $p^2 \mid d_1 f_3 - p e_3$. If $p \mid e_3$ then $p^2 \mid d_1 f_3$. If $p \mid e_3$ and $p \mid d_1$ then $d_1 = 0$ and the condition $d_1(\delta) = 1 = (p, f_3)$ implies that $p \nmid f_3$ and $d_2(\delta) = p^2$. There are $p^2(p^3 - p^2)$ such matrices. If $p \mid e_3$ and $p \nmid d_1$ then $p^2 \mid f_3$ then $d_2(\delta) = p^2(p, d_1 f_3 / p^2 - e_3 / p) \neq p^2$ if and only if $f_3 / p^2 \equiv \overline{d_1} e_3 / p \pmod{p}$, which given d_1 and e_3 / p can happen for only one value of f_3 / p^2 . There are $(p-1)p^2(p-1)$ such matrices. If $p \nmid e_3$ then $d_1(\delta) = 1 = (p, d_1, e_3, f_3)$. The condition $p^2 \mid d_1 f_3 - p e_3$ implies that $p^2 \nmid d_1 f_3$ and $p \nmid d_1$ but $p \mid f_3$. The condition $d_2(\delta)$ implies that $p \parallel d_1 f_3 / p - e_3$. Given d_1 and e_3 , there are p choices for f_3 / p given by $f_3 / p \equiv \overline{d_1} e_3 \pmod{p}$ but one has to remove the value satisfying $f_3 / p \equiv \overline{d_1} e_3 \pmod{p^2}$. There are $(p-1)(p^3 - p^2)(p-1)$ such matrices. Finally, there are $p^3(p-1)(2p-1)$ relevant matrices of type 16.

Let us consider the matrices of type 17. The condition $d_2(\delta) = p^2$ implies that $p \mid f_2$. If $f_2 = 0$ then $d_2(\delta) = p^2 = p^2(p, d_3)$ such that $p \nmid d_3$, which implies $d_1(\delta) = 1$. There are $(p^3 - p^2)p^2$ such matrices. If $f_2 \neq 0$ then $p \mid d_3$ since $p \mid d_3 f_2 / p$, in which case $d_2(\delta) = p^2$. The condition $d_1(\delta) = 1$ implies that $p \nmid e_2$. There are $p^2(p^2 - p)(p-1)$ such matrices. Finally, there are $p^3(p-1)(2p-1)$ relevant matrices of type 17.

Let us consider the matrices of type 18. The condition $d_2(\delta) = p^2$ implies that $p \mid f_3$ and $p \mid d_2 f_3 / p$. If $p^2 \mid f_3$ then $d_2(\delta) = p^2 = p^2(p, d_2 f_3 / p^2 - e_3)$. One has to remove the p^2 values of e_3 satisfying $e_3 \equiv d_2 f_3 / p^2 \pmod{p}$. In this case, one has $d_1(\delta) = (p, d_2, e_3) = 1$ since if $p \mid (d_2, e_3)$ then $(p, d_2 f_3 / p^2 - e_3) \neq 1$. There are $p^2(p^3 - p^2)p$ such matrices. If $p^2 \nmid f_3$ then $p \mid d_2$ and the conditions on $d_1(\delta)$ and $d_2(\delta)$ are satisfied. There are $p(p^3 - p^2)(p^2 - p)$ such matrices. Finally, there are $p^4(p-1)(2p-1)$ relevant matrices of type 18.

Let us consider the matrices of type 19. The condition on $d_2(\delta)$ implies that $p^2 \mid d_2 f_2$. If $d_2 = 0$ then $d_2(\delta) = p^2 = p^2(p^4, e_2, f_2)$ and $d_1(\delta) = 1 = (p^2, e_2, f_2)$. One has to remove the couples (e_2, f_2) satisfying $p \mid e_2$ and $p \mid f_2$, namely p^2 couples. There are $p^4 - p^2$ such matrices. If $d_2 \neq 0$ and $f_2 = 0$ then $d_2(\delta) = p^2 = p^2(p^2, d_2, e_2)$ and $d_1(\delta) = 1 = (p^2, d_2, e_2)$. One has to remove the couples (d_2, e_2) satisfying $p \mid d_2$ and $p \mid e_2$, namely $(p-1)p$ couples. There are $(p^2 - 1)p^2 - (p-1)p$ such matrices. If $d_2 \neq 0$ and $f_2 \neq 0$ then $d_2(\delta) = p^2 = p^2(p^2, p d_2 / p, p f_2 / p, d_2 f_2 / p^2 - e_2)$ and $d_1(\delta) = 1 = (p^2, e_2)$. Thus, $p \nmid e_2$ and $p \nmid d_2 f_2 / p^2 - e_2$. Among the $p^2 - p$ values of e_2 satisfying $p \nmid e_2$, one has to remove these satisfying $e_2 \equiv d_2 f_2 / p^2 \pmod{p}$ of cardinal p . There are $(p-1)(p^2 - 2p)(p-1)$ such matrices. Finally, there are $p(p-1)(3p^2 - p + 1)$ relevant matrices of type 19.

One can recover the value of the degree given in (A.8) by summing all the contributions in the previous paragraphs. \square

A.5. Degree of $\Lambda \text{diag}(1, p^3, p^3) \Lambda$.

Proposition A.5—One has

$$\deg(\text{diag}(1, p^3, p^3)) = p^4(p^2 + p + 1). \quad (\text{A.9})$$

Proof of Proposition A.5. By (A.2), the possible upper-triangular column reduced matrices δ that can occur in the decomposition into Λ -right cosets are

$$\text{Type 1: } \begin{pmatrix} p^3 & d_3 & \\ & p^3 & \\ & & 1 \end{pmatrix} \rightsquigarrow \mathbf{d}(\delta) = (1, (p^3, d_3), p^6),$$

$$\text{Type 2: } \begin{pmatrix} p^3 & & e_3 \\ & 1 & f_3 \\ & & p^3 \end{pmatrix} \rightsquigarrow \mathbf{d}(\delta) = (1, (p^3, e_3), p^6),$$

$$\text{Type 3: } \begin{pmatrix} 1 & d_3 & e_3 \\ & p^3 & f_3 \\ & & p^3 \end{pmatrix} \rightsquigarrow \mathbf{d}(\delta) = (1, (p^3, f_3, d_3 f_3), p^6)$$

and

$$\text{Type 4: } \begin{pmatrix} p^3 & d_2 & e_1 \\ & p^2 & f_1 \\ & & p \end{pmatrix} \rightsquigarrow \mathbf{d}(\delta) = ((p, d_2, e_1, f_1), (p^3, p d_2, d_2 f_1 - p^2 e_1), p^6),$$

$$\text{Type 5: } \begin{pmatrix} p^3 & d_1 & e_2 \\ & p & f_2 \\ & & p^2 \end{pmatrix} \rightsquigarrow \mathbf{d}(\delta) = ((p, d_1, e_2, f_2), (p^3, p^2 d_1, d_1 f_2 - p e_2), p^6),$$

$$\text{Type 6: } \begin{pmatrix} p^2 & d_3 & e_1 \\ & p^3 & f_1 \\ & & p \end{pmatrix} \rightsquigarrow \mathbf{d}(\delta) = ((p, d_3, e_1, f_1), (p^3, p d_3, p^2 f_1, d_3 f_1), p^6),$$

$$\text{Type 7: } \begin{pmatrix} p^2 & d_1 & e_3 \\ & p & f_3 \\ & & p^3 \end{pmatrix} \rightsquigarrow \mathbf{d}(\delta) = ((p, d_1, e_3, f_3), (p^3, p^2 f_1, d_1 f_3 - p e_3), p^6),$$

$$\text{Type 8: } \begin{pmatrix} p & d_3 & e_2 \\ & p^3 & f_2 \\ & & p^2 \end{pmatrix} \rightsquigarrow \mathbf{d}(\delta) = ((p, d_3, e_2, f_2), (p^3, p^2 d_3, p f_2, d_3 f_2), p^6),$$

$$\text{Type 9: } \begin{pmatrix} p & d_1 & e_3 \\ & p^2 & f_3 \\ & & p^3 \end{pmatrix} \rightsquigarrow \mathbf{d}(\delta) = ((p, d_1, e_4, f_4), (p^3, p f_3, d_2 f_3 - p^2 e_3), p^6)$$

and

$$\text{Type 10: } \begin{pmatrix} p^2 & d_2 & e_2 \\ & p^2 & f_2 \\ & & p^2 \end{pmatrix} \rightsquigarrow \mathbf{d}(\delta) = ((p^2, d_2, e_2, f_2), (p^4, p^2 d_2, p^2 f_2, d_2 f_2 - p^2 e_2), p^6)$$

where $0 \leq d_j, e_j, f_j < p^j$ for $j = 1, 2, 3$. Let us count the matrices among the previous ones, whose determinantal vector is the same as the one of $\text{diag}(1, p^3, p^3)$, namely $(1, p^3, p^6)$.

Let us consider the matrices of type 1. The condition on $d_2(\delta)$ implies $d_3 = 0$. There is 1 relevant matrix of type 1.

Let us consider the matrices of type 2. The condition on $d_2(\delta)$ implies $e_3 = 0$. There are p^3 relevant matrices of type 2.

Let us consider the matrices of type 3. The condition on $d_2(\delta)$ implies $f_3 = 0$. There are p^6 relevant matrices of type 3.

Let us consider the matrices of type 4. The condition on $d_2(\delta)$ implies $d_2 = 0$ and $e_1 = 0$. Then, $d_1(\delta) = 1 = (p, f_1)$ such that $f_1 \neq 0$. There are $p - 1$ relevant matrices of type 4.

Let us consider the matrices of type 5. The condition on $d_2(\delta)$ implies $d_1 = 0$ and $e_2 = 0$. Then, $d_1(\delta) = 1 = (p, f_2)$ such that $p \nmid f_2$. There are $p^2 - p$ relevant matrices of type 5.

Let us consider the matrices of type 6. The condition on $d_2(\delta)$ implies $f_1 = 0$ and $p^2 \mid d_3$. Then, $d_1(\delta) = 1 = (p, e_1)$ such that $e_1 \neq 0$. There are $p(p - 1)$ relevant matrices of type 6.

Let us consider the matrices of type 7. The condition on $d_2(\delta)$ implies $p \mid f_3$ and $p \mid d_1 f_3 / p - e_3$. One has $d_1 \neq 0$ since otherwise $p^2 \mid e_2$ by the condition on $d_2(\delta)$ such that $d_1(\delta) = p \neq 1$. Thus, d_1 is invertible modulo p and $f_3 / p \equiv e_3 \overline{d_1} \pmod{p^2}$ is fixed. There are $(p - 1)p^3$ relevant matrices of type 7.

Let us consider the matrices of type 8. The condition on $d_2(\delta)$ implies $p \mid d_3$ and $f_2 = 0$. Then, $d_1(\delta) = 1 = (p, e_2)$ such that $p \nmid e_2$. There are $p^2(p^2 - p)$ relevant matrices of type 8.

Let us consider the matrices of type 9. The condition on $d_2(\delta)$ implies $p^2 \mid f_3$ and $p \mid d_2 f_3 / p - e_3$. If $f_3 = 0$ then $p \mid e_3$ and $d_1(\delta) = 1 = (p, d_2)$ such that $p \nmid d_2$. There are $(p^2 - p)p^2$ such matrices. If $f_3 \neq 0$ then $d_2 \equiv e_3 \overline{f_3 / p^2} \pmod{p}$ can take p values. Then, $d_1(\delta) = 1 = (p, e_3)$ such that $p \nmid e_3$. There are $p(p^3 - p^2)(p - 1)$ such matrices. Finally, there are $p^4(p - 1)$ relevant matrices of type 9.

Let us consider the matrices of type 10. The condition on $d_2(\delta)$ implies $p \mid d_2$, $p \mid f_2$ and $p \mid d_2 f_2 / p^2 - e_2$. One has $d_2 \neq 0$ since otherwise $d_2(\delta) = p^2(p^2, e_2) = p^2 \overline{d_1(\delta)} = p^2 \neq p^3$. Thus, d_2 / p is invertible modulo p and f_2 is fixed by $f_2 / p \equiv e_2 \overline{d_2 / p} \pmod{p}$. Then, $d_1(\delta) = 1 = (p, e_2)$ such that $p \nmid e_2$, $p \nmid f_2 / p$ and $d_2(\delta) = p^3$. There are $(p - 1)(p^2 - p)$ relevant matrices of type 10.

One can recover the value of the degree given in (A.9) by summing all the contributions in the previous paragraphs. \square

A.6. Degree of $\Lambda \text{diag}(1, 1, p^3) \Lambda$.

Proposition A.6– One has

$$\deg(\text{diag}(1, 1, p^3)) = p^4(p^2 + p + 1). \quad (\text{A.10})$$

Proof of Proposition A.6. By (A.2), the possible upper-triangular column reduced matrices δ that can occur in the decomposition of the Λ -double coset $\Lambda \text{diag}(1, 1, p^3) \Lambda$ into Λ -right cosets are

$$\begin{aligned} \text{Type 1: } & \begin{pmatrix} p^3 & & \\ & 1 & \\ & & 1 \end{pmatrix} \rightsquigarrow \mathbf{d}(\delta) = (1, 1, p^3), \\ \text{Type 2: } & \begin{pmatrix} 1 & d_3 & \\ & p^3 & \\ & & 1 \end{pmatrix} \rightsquigarrow \mathbf{d}(\delta) = (1, 1, p^3), \\ \text{Type 3: } & \begin{pmatrix} 1 & e_3 & \\ & 1 & f_3 \\ & & p^3 \end{pmatrix} \rightsquigarrow \mathbf{d}(\delta) = (1, 1, p^3) \end{aligned}$$

and

$$\text{Type 4: } \begin{pmatrix} 1 & d_1 & e_2 \\ & p & f_2 \\ & & p^2 \end{pmatrix} \rightsquigarrow \mathbf{d}(\delta) = (1, (p, f_2), p^3),$$

$$\text{Type 5: } \begin{pmatrix} 1 & d_2 & e_1 \\ & p^2 & f_1 \\ & & p \end{pmatrix} \rightsquigarrow \mathbf{d}(\delta) = (1, (p, f_1), p^3),$$

$$\text{Type 6: } \begin{pmatrix} p & e_2 \\ & 1 & f_2 \\ & & p^2 \end{pmatrix} \rightsquigarrow \mathbf{d}(\delta) = (1, (p, e_2), p^3),$$

$$\text{Type 7: } \begin{pmatrix} p^2 & e_1 \\ & 1 & f_1 \\ & & p \end{pmatrix} \rightsquigarrow \mathbf{d}(\delta) = (1, (p, e_1), p^3),$$

$$\text{Type 8: } \begin{pmatrix} p & d_2 \\ & p^2 & \\ & & 1 \end{pmatrix} \rightsquigarrow \mathbf{d}(\delta) = (1, (p, d_2), p^3),$$

$$\text{Type 9: } \begin{pmatrix} p^2 & d_1 \\ & p & \\ & & 1 \end{pmatrix} \rightsquigarrow \mathbf{d}(\delta) = (1, (p, d_1), p^3)$$

and

$$\text{Type 10: } \begin{pmatrix} p & d_1 & e_1 \\ & p & f_1 \\ & & p \end{pmatrix} \rightsquigarrow \mathbf{d}(\delta) = ((p, d_1, e_1, f_1), (p^2, pd_1, pf_1, d_1f_1 - pe_1), p^3)$$

where $0 \leq d_j, e_j, f_j < p^j$ for $j = 1, 2, 3$. Let us count the matrices among the previous ones, whose determinantal vector is the same as the one of $\text{diag}(1, 1, p^3)$, namely $(1, 1, p^3)$.

Let us consider the matrices of type 1. There is 1 relevant matrix of type 1.

Let us consider the matrices of type 2. There are p^3 relevant matrices of type 2.

Let us consider the matrices of type 3. There are p^6 relevant matrices of type 3.

Let us consider the matrices of type 4. The condition on $d_2(\delta)$ implies $p \nmid f_2$.

There are $p^3(p^2 - p)$ relevant matrices of type 4.

Let us consider the matrices of type 5. The condition on $d_2(\delta)$ implies $f_1 \neq 0$.

There are $p^3(p - 1)$ relevant matrices of type 5.

Let us consider the matrices of type 6. The condition on $d_2(\delta)$ implies $p \nmid e_2$.

There are $p^2(p^2 - p)$ relevant matrices of type 6.

Let us consider the matrices of type 7. The condition on $d_2(\delta)$ implies $e_1 \neq 0$.

There are $p(p - 1)$ relevant matrices of type 7.

Let us consider the matrices of type 8. The condition on $d_2(\delta)$ implies $p \nmid d_2$.

There are $p^2 - p$ relevant matrices of type 8.

Let us consider the matrices of type 9. The condition on $d_2(\delta)$ implies $d_1 \neq 0$.

There are $p - 1$ relevant matrices of type 9.

Let us consider the matrices of type 10. One has $d_1 \neq 0$ since otherwise $p \mid d_2(\delta)$. Thus, $d_1(\delta) = 1$. In addition, $f_1 \neq 0$ since otherwise $p \mid d_2(\delta)$. There are $p(p - 1)^2$ relevant matrices of type 10.

One can recover the value of the degree given in (A.10) by summing all the contributions in the previous paragraphs. \square

REFERENCES

- [AZ95] A. N. Andrianov and V. G. Zhuravljëv. *Modular forms and Hecke operators*, volume 145 of *Translations of Mathematical Monographs*. American Mathematical Society, Providence, RI, 1995. Translated from the 1990 Russian original by Neal Koblitz.
- [BHM] V. Blomer, G. Harcos, and D. Milićević. Bounds for eigenforms on arithmetic hyperbolic 3-manifolds. available at <http://arxiv.org/abs/1401.5154>.
- [BMa] Valentin Blomer and Péter Maga. Subconvexity for sup-norms of automorphic forms on $PGL(n)$. available at <http://arxiv.org/pdf/1405.6691.pdf>.
- [BMb] Valentin Blomer and Péter Maga. The sup-norm problem for $PGL(4)$. available at <http://arxiv.org/pdf/1404.4331.pdf>.
- [BT] Farrell Brumley and Nicolas Templier. Large values of cusp forms on $GL(n)$. available at <http://arxiv.org/abs/1411.4317>.
- [DFI94] W. Duke, J. B. Friedlander, and H. Iwaniec. Bounds for automorphic L -functions. II. *Invent. Math.*, 115(2):219–239, 1994.
- [FI92] J. Friedlander and H. Iwaniec. A mean-value theorem for character sums. *Michigan Math. J.*, 39(1):153–159, 1992.
- [Gol06] Dorian Goldfeld. *Automorphic forms and L -functions for the group $GL(n, \mathbb{R})$* , volume 99 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 2006. With an appendix by Kevin A. Broughan.
- [HRR] Roman Holowinsky, Guillaume Ricotta, and Emmanuel Royer. On the sup-norm of $SL(3)$ Hecke-Maass cusp form. available at <http://arxiv.org/abs/1411.4317>.
- [HT13] Gergely Harcos and Nicolas Templier. On the sup-norm of Maass cusp forms of large level. III. *Math. Ann.*, 356(1):209–216, 2013.
- [IS95] H. Iwaniec and P. Sarnak. L^∞ norms of eigenfunctions of arithmetic surfaces. *Ann. of Math. (2)*, 141(2):301–320, 1995.
- [Iwa92] Henryk Iwaniec. The spectral growth of automorphic L -functions. *J. Reine Angew. Math.*, 428:139–159, 1992.
- [Kod67] Tetsuo Kodama. On the law of product in the Hecke ring for the symplectic group. *Mem. Fac. Sci. Kyushu Univ. Ser. A*, 21:108–121, 1967.
- [Mac95] I. G. Macdonald. *Symmetric functions and Hall polynomials*. Oxford Mathematical Monographs. The Clarendon Press, Oxford University Press, New York, second edition, 1995. With contributions by A. Zelevinsky, Oxford Science Publications.
- [New72] Morris Newman. *Integral matrices*. Academic Press, New York, 1972. Pure and Applied Mathematics, Vol. 45.
- [SA] L. Silberman and Venkatesh A. Entropy bounds for Hecke eigenfunctions on division algebras. preprint available at <http://www.math.ubc.ca/~lior/work/>.
- [Sar] Peter Sarnak. Letter to Morawetz. available at <http://www.math.princeton.edu/sarnak>.
- [Shi94] Goro Shimura. *Introduction to the arithmetic theory of automorphic functions*, volume 11 of *Publications of the Mathematical Society of Japan*. Princeton University Press, Princeton, NJ, 1994. Reprint of the 1971 original, Kanô Memorial Lectures, 1.
- [Ven10] Akshay Venkatesh. Sparse equidistribution problems, period bounds and subconvexity. *Ann. of Math. (2)*, 172(2):989–1094, 2010.

DEPARTMENT OF MATHEMATICS, THE OHIO STATE UNIVERSITY, 100 MATH TOWER, 231 WEST 18TH AVENUE, COLUMBUS, OH 43210-1174
E-mail address: holowinsky.1@osu.edu

UNIVERSITÉ DE BORDEAUX, IMB, 351 COURS DE LA LIBÉRATION, 33405 TALENCE CEDEX, FRANCE
E-mail address: Guillaume.Ricotta@math.u-bordeaux1.fr

UNIVERSITÉ BLAISE PASCAL, LABORATOIRE DE MATHÉMATIQUES, LES CÉZEAUX, BP 80026, F-63171 AUBIÈRE CEDEX, FRANCE
E-mail address: emmanuel.royer@math.univ-bpclermont.fr