



**HAL**  
open science

**Does the study of nano-microzooplankton community  
size structure effectively define their dynamics ?  
Investigation in the Bay of Biscay (France).**

E Marquis, Nathalie Niquil, Christine Dupuy

► **To cite this version:**

E Marquis, Nathalie Niquil, Christine Dupuy. Does the study of nano-microzooplankton community size structure effectively define their dynamics ? Investigation in the Bay of Biscay (France).. Journal of Plankton Research, 2011. hal-01248041

**HAL Id: hal-01248041**

**<https://hal.science/hal-01248041>**

Submitted on 26 Dec 2016

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



**Does the study of nano-microzooplankton community size structure effectively define their dynamics ? Investigation in the Bay of Biscay (France).**

Journal:	<i>Journal of Plankton Research</i>
Manuscript ID:	Draft
Manuscript Type:	Original Article
Date Submitted by the Author:	n/a
Complete List of Authors:	Marquis, Elise; National Taiwan University, Institute of Oceanography Niquil, Nathalie; Université de La Rochelle, UMR 6250, Laboratoire LIENSs Dupuy, Christine; Université de La Rochelle, UMR 6250, Laboratoire LIENSs
Keywords:	nano-microzooplankton, size structure, dynamics, taxonomy, Bay of Biscay

SCHOLARONE™  
Manuscripts

1  
2  
3 1 Does the study of nano-microzooplankton community size  
4  
5 2 structure effectively define their dynamics ? Investigation  
6  
7  
8 3 in the Bay of Biscay (France).  
9  
10  
11 4  
12  
13  
14 5

15 6 **Elise MARQUIS \***  
16  
17 7

18  
19 8 Laboratoire LIENSs, UMR 6250, Université de La Rochelle, Bâtiment ILE, 2 rue Olympe de Gouges, 17000  
20  
21 9 La Rochelle, France

22 10 E-mail address : [emarquis@me.com](mailto:emarquis@me.com)  
23  
24

25 11 Present address : Institute of Oceanography, National Taiwan University, No. 1, Section 4, Roosevelt Road,  
26  
27 12 Taipei, 10617, Taiwan

28 13 Tel: 00886(0)983341751  
29

30 14 Fax: 00886(0)233669746  
31  
32 15

33 16 **Nathalie NIQUIL**  
34  
35 17

36 18 Laboratoire LIENSs, UMR 6250, Université de La Rochelle, Bâtiment ILE, 2 rue Olympe de Gouges, 17000  
37  
38 19 La Rochelle, France

39  
40 20 E-mail address : [niquil@univ-lr.fr](mailto:niquil@univ-lr.fr)  
41  
42 21

43 22 **Christine DUPUY**  
44  
45 23

46  
47 24 Laboratoire LIENSs, UMR 6250, Université de La Rochelle, Bâtiment ILE, 2 rue Olympe de Gouges, 17000  
48  
49 25 La Rochelle, France

50 26 E-mail address : [cdupuy@univ-lr.fr](mailto:cdupuy@univ-lr.fr)  
51  
52  
53 27  
54  
55 28  
56  
57

58 29 \* Corresponding author  
59  
60 30

**ABSTRACT**

Studies of nano-microzooplankton dynamics based on the analysis of community taxonomic composition are highly time consuming and manpower demanding. Recent developments in automatic plankton counting technology are offering a new way to consider plankton dynamics with straightforward analysis of community size structures. The present study aimed to ensure that size is a good descriptor of nano-microzooplankton dynamics. The dynamics of the nano-microzooplankton community of the Bay of Biscay (France) over three sites and four sampling periods was analyzed using three different parameters of classification: taxon, body size ( $\text{ngC cell}^{-1}$ ) and equivalent spherical diameter ( $\mu\text{m}$ ). A Mantel test revealed that there was no difference in the characteristics of nano-microzooplankton dynamics when studying either community size structure or taxonomic composition. Moreover, a BEST test confirmed that the biotic and abiotic factors selected for impacting nano-microzooplankton community dynamics were the same among the three classification types. Considering those results, it was argued that nano-microplankton dynamics is well defined by the study of the community size structure. While focusing on nano-microzooplankton size structure seems promising, homogenizing the size descriptor used among studies would be needed in order to make worldwide data comparisons and ESD/biovolume should be favored.

**KEYWORDS:** nano-microzooplankton, size structure, dynamics, taxonomy, Bay of Biscay

## 1 INTRODUCTION

2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Open ocean photosynthesis is dominated by pico-nanophytoplanktonic production (0.2 to 20 m diameter, Sieburth et al., 1978), with most of the active primary producers smaller than 3  $\mu\text{m}$  (Waterbury et al., 1979; Fenchel, 1988). A complex assemblage of viruses, bacterioplankton and protozoans coexist with and support these small oceanic primary producers through their role in nutrient regeneration (Azam et al., 1983; Ducklow and Carlson, 1992; Sherr and Sherr, 2000). Larger phytoplankton such as diatoms and dinoflagellates show greater variability than pico-nanophytoplankton and so do their predators, i.e. larger protists and metazoans (Fenchel, 1988). Primary production is transferred to higher levels through two main pathways: classical food chain and microbial food web, according to the size of the main primary producers (Azam et al., 1983; Sherr et al., 1986; Sommaruga, 1995; Thingstad and Rassoulzadegan, 1999). Therefore, depending on which pathways is dominating the planktonic food web, the amount of energy lost (mainly through respiration), matter recycled and organic carbon available to plankton predators (mainly fish) will vary. For those reasons, our understanding of plankton dynamics and plankton food web functioning is intimately related to our appreciation of global biogeochemical fluxes. Despite the actual need for information about these issues when studying global warming concerns, the current amount of knowledge of the identity and the functional role of the major plankton groups as well as their dynamics prevent us evolving from the current simple plankton models towards more detailed and adaptive ecosystem modeling (Ducklow, 2003).

Since the paper of Azam et al. (1983), a high number of studies have focused on the dynamics and the role of nano-microzooplankton in oceanic, coastal and halieutic environments. Nano-microzooplankton are trophic intermediaries in pelagic food webs, permitting the transfer of carbon from pico- and nanoplankton to the metazoans (Sherr et al., 1986; Pierce and Turner, 1992). They occupy an essential trophic node in microbial food webs and their dynamics may be either controlled by predation or by resource availability as well as hydrography (e.g. Sanders, 1987; Cowlshaw, 2004). Despite such an essential role in the aquatic ecosystems, their taxonomical diversity is so high that it is clear that they are not perfectly known. In the early nineties, Sleigh (1991) discussed about the taxonomy of heterotroph protists, which are the main components of the nano-microzooplankton community, and stated “many species, and possibly phyla, remain to be described” (Sleigh, 1991). Today, almost twenty years later, and despite progress in DNA sequencing, protozoan taxonomy is still subject to study, discussion and controversy (e.g. Finlay, 2004; Adl et al., 2005).

Traditional studies based on taxonomic compositions of plankton communities are difficult to implement efficiently due to the limitation of resources, time and manpower to process samples

(Culverhouse et al., 2006). However, size is another possible way to classify plankton organisms, and the terminology of Sieburth (Sieburth et al., 1978) is useful and has become widely accepted. Recent development in metabolic theory (Brown et al., 2004) confirms older studies (e.g. Moloney and Field, 1989) on suggesting that size determines many biological properties of organisms such as respiration, nutrient uptake, production, ontogenetic growth, etc... (e.g. Zeuthen, 1970; Gillooly et al., 2001; West et al., 2003; Lupez-Urrutia et al., 2006). Moreover, in aquatic ecosystems, size usually determines predator-prey interactions (e.g. Frost, 1972; Peters and Downing, 1984; Caparroy et al., 2000). As such, size distribution of plankton may be used as indicators of ecosystem and trophic status. Additionally, recent studies have shown that one of the aquatic ecological responses to global warming is the reduction of body size (Daufresne et al., 2009; Moran et al., 2010). Regarding all these issues, studying planktonic and especially nano-microzooplanktonic dynamics based on the community size structure seems to be an interesting and promising research focus.

New technologies for automatic plankton counting are now available to plankton ecologists. Such instruments, e.g. the Flowcam technology (Sieracki et al., 1998), are excelling in size classification and identification of planktonic cells. Therefore, they would be very useful in studying the dynamics of planktonic community size structures. However, before commencing such studies, it is essential to ensure that size is a sufficient plankton community descriptor on its own. In other words, it is essential to understand if the dynamics resulting from those size-based analyses are showing the same pattern and the same spatio-temporal differences as the dynamics resulting from the taxonomy-based analysis.

This study aims to investigate whether the seasonal and spatial dynamics of the nano-microzooplankton may be revealed looking at the size structure of the community. Extensive data on planktonic communities collected in 2004 in the Bay of Biscay were used to answer this question. Size-Abundance spectra are used to understand the overall changes in the nano-microzooplankton composition and a comparison between taxon-based and size-based classifications helps to evaluate the relevance of the size structure analysis.

## MATERIAL AND METHODS

### Study site and hydrography

The continental shelf of the Bay of Biscay (Fig. 1) is up to 200 km wide with a surface area of 223,000 km<sup>2</sup>. Its hydrological structure is principally influenced by the seasonal dynamics of the

1  
2 1 Loire and the Gironde river plumes (Lazure and Jegou, 1998) and the shelf ecology shows a strong  
3  
4 2 variability in time related to temperate zone climatic fluctuations (Koutsikopoulos et al., 1998). At  
5  
6 3 four periods in 2004 (08 to 10 February, 23 to 25 April, 09 to 11 June, and 30 September to 02  
7  
8 4 October), three stations were sampled, located on the continental shelf, on an estuary-coast-offshore  
9  
10 5 triangle: “Gironde” (01°30W, 45°30N), “Coast” (01°30W, 45°01N) and “Offshore” (2°20W,  
11  
12 6 45°10N) (Fig. 1). Salinity and temperature profiles were measured using a CTD (Conductivity-  
13  
14 7 Temperature-Density) probe (Sea-Bird SBE 9). Concurrently, irradiance and fluorescence profiles  
15  
16 8 of the water column were measured in order to measure the depths of the photic zone and maximum  
17  
18 9 fluorescence in order to adapt the plankton sampling accordingly.

## 18 10 19 11 20 11 21 12 **Water and plankton sampling and analysis**

22 12  
23 13  
24 13  
25 14 At each station and sampling period, we collected water samples with 12-L Niskin bottles at  
26  
27 15 three different depths: subsurface, maximum fluorescence and bottom of the photic zone.

28  
29 16 Samples for dissolved inorganic nutrients analyses: nitrate (NO<sub>3</sub>), nitrite (NO<sub>2</sub>), silicate  
30  
31 17 (Si(OH)<sub>4</sub>) and phosphate (PO<sub>4</sub>), were immediately filtered through Whatman GF/F filters so that the  
32  
33 18 filtrate could be stored at -20°C until its analysis at the laboratory with a Skalar autoanalyzer  
34  
35 19 (Strickland and Parsons, 1972).

36 20 Samples for bacteria and picophytoplankton counting were fixed with formaldehyde (final  
37  
38 21 concentration 2%), frozen in liquid N<sub>2</sub> and enumerated using a FACSCan flow cytometer (Bd-  
39  
40 22 Bioscience) (Marie et al., 2000). Nanoflagellates were fixed with buffered paraformaldehyde (final  
41  
42 23 concentration 1%) then stained with DAPI and counted on 0.8 µm black polycarbonate filters by  
43  
44 24 epifluorescence microscopy (Sherr et al., 1994). Heterotrophic nanoflagellates (HNF) were  
45  
46 25 distinguished from pigmented (autotrophic) nanoflagellates (ANF) by the absence of chlorophyll  
47  
48 26 fluorescence. Microphytoplankton (diatoms and dinoflagellates) were fixed with formaldehyde  
49  
50 27 (final concentration 1%) plus alkaline lugol (final concentration 1%), enumerated and measured by  
51  
52 28 inverse microscopy (Utermöhl, 1958). Heterotrophic and mixotrophic dinoflagellates (HDF) were  
53  
54 29 determined from morphologic species recognition and relevant literature (e.g. Lessard and Swift,  
55  
56 30 1986). Ciliates were stained with alkaline lugol (1% final concentration), counted and measured by  
57  
58 31 inverse fluorescence microscopy. Ciliate samples from surface and bottom of the photic zone  
59  
60 32 collected in February at the Gironde station were not analyzed due to poor preservation. Diatom,  
33  
34 33 dinoflagellate and ciliate biovolumes were calculating by transforming the cell’s shapes into  
35  
36 34 geometric figures. Samples of metazoan microplankton were obtained by gently filtering 10 liters  
37  
38 35 of collected seawater through a 63µm size mesh. The retained organisms were then diluted in



1 filtered (< 63  $\mu\text{m}$ ) seawater and preserved in buffered formaldehyde (final concentration 2%). They  
2 were counted under a binocular microscope. All the conversion factors and equations used to  
3 convert the abundance of pico-, nano- and microplankton into biomass were obtained from the  
4 literature (Table 1). The equivalent spherical diameter (ESD) of each cell was obtained by  
5 calculating the diameter ( $\mu\text{m}$ ) of a hypothetical sphere of equivalent volume. The two conversion  
6 from biovolume to biomass and from biovolume to ESD resulted in an exponential relationships  
7 between the body mass and ESD of nano-microzooplankton cells in this study (Figure 2).

8 Mesozooplankton was collected from vertical tows through the entire photic zone using a  
9 200 $\mu\text{m}$ -mesh WP2 net, preserved in buffered formaldehyde (final concentration 2%) and counted  
10 under a binocular microscope. A second replicate was used to measure the mesozooplankton dry  
11 weight. Mesozooplankton carbon biomass was determined by multiplying dry weights of each  
12 sample by a factor of 0.38 (Bode et al., 1998).

#### 13 14 **Data analysis**

15 Size-abundance spectra (SAS) were obtained by presenting the  $\log_{10}$  of the abundances  
16 (cells  $\text{mL}^{-1}$ ) versus the  $\log_{10}$  of the geometric mean of the respective size class ( $\mu\text{m}^3$ ).

17 The nano-microzooplankton community was then classified under 3 different classification  
18 types:

- 19 1- Taxons: Heterotrophic nanoflagellates (HNF); Unarmoured Heterotrophic  
20 dinoflagellates (U-HDF); Armoured Heterotrophic dinoflagellates (A-  
21 HDF); Naked Ciliates; Tintinnid Ciliates; *Myrionecta rubra* ciliates;  
22 Microplanktonic metazoans.
- 23 2- Body Sizes: <100  $\text{ngC cell}^{-1}$ ; 100 to 1,000  $\text{ngC cell}^{-1}$ ; 1,000 to 5,000  $\text{ngC cell}^{-1}$ ;  
24 5,000 to 10,000  $\text{ngC cell}^{-1}$ ; 10,000 to 50,000  $\text{ngC cell}^{-1}$ ; 50,000 to  
25 100,000  $\text{ngC cell}^{-1}$ ; >100,000  $\text{ngC cell}^{-1}$ .
- 26 3- ESD (Equivalent Spherical Diameter): <20  $\mu\text{m}$ ; 20 to 30  $\mu\text{m}$ ; 30 to 40  $\mu\text{m}$ ; 40  
27 to 50  $\mu\text{m}$ ; 50 to 60  $\mu\text{m}$ ; 60 to 70  $\mu\text{m}$ ; >70  $\mu\text{m}$ .

28 All data were normalized prior to statistical analyses; using a double-root-transformation on  
29 the abundance and biomass data and a log-transformation on the environmental data (Legendre and  
30 Legendre, 1998). Two types of analyses were applied to the data sets: Mantel test (Legendre and  
31 Legendre, 1998) to analyze whether the different ways of classifying nano-microzooplankton lead  
32 to different description of its spatial/temporal dynamics, and BEST test, to analyze the biotic (i.e.  
33 trophic) and abiotic (i.e. environmental) factors impacting nano-microzooplankton community  
34 dynamics and to compare the factors selected for each classification types.



1  
2 1 First, we constructed 3 matrices using nano-microzooplankton abundances of the 3  
3 classification types (Taxons, Body Sizes and ESD). We used the average abundance values  
4 2 throughout the water column. The 3 matrices have 12 rows (3 sites and 4 sampling periods) and 7  
5 3 columns (7 different classes, for each classification). These matrices were used to calculate  
6 4 dissimilarity matrices in order to model the resemblance between the sampled sites/months by the  
7 5 mean values of community composition for each of the 3 nano-microzooplankton classification  
8 6 types. Bray and Curtis distance measure was used because it's particularly suitable for quantitative  
9 7 data (Legendre and Legendre, 1998). We then applied the Mantel test to see if there was a  
10 8 correlation between the different distances matrices (Legendre and Legendre, 1998). In a simple  
11 9 Mantel test, two matrices were compared. First a Pearson correlation ( $r$ ) was calculated between the  
12 10 matrices and after this, the second matrix was shuffled 10,000 times and the correlation  
13 11 recalculated. This permutation procedure was done in order to construct a law of distribution of  
14 12 correlation coefficients, in absence of statistically significant relation between the two matrices. The  
15 13 original coefficient was then compared with this distribution in order to determine its statistical  
16 14 significance ( $p$  value). The Mantel statistics were calculated for the 3 pairs formed by the different  
17 15 nano-microzooplankton classifications: Taxons/Body Sizes, Taxons/ESD, Body Sizes/ESD.

18 16 The second analysis was a BEST test (Clarke and Warwick, 2001) done using the software  
19 17 package PRIMER<sup>®</sup>6. The BEST analysis selected trophic and environmental variables that best  
20 18 explain patterns in the nano-microzooplankton assemblage. The test was conducted by maximizing  
21 19 a Spearman rank correlation between the resemblance matrices of environmental and trophic  
22 20 variables (Euclidean distance) and community abundances (Bray-Curtis distance). The significance  
23 21 of these results was tested using permutation tests. The environmental parameters analysed were:  
24 22 Temperature ( $^{\circ}\text{C}$ ), Salinity (PSU),  $\text{NO}_2$  concentration ( $\mu\text{mol l}^{-1}$ ),  $\text{NO}_3$  concentration ( $\mu\text{mol l}^{-1}$ ),  $\text{PO}_4$   
25 23 concentration ( $\mu\text{mol l}^{-1}$ ) and N/P ratio. The trophic parameters tested were the biomasses ( $\mu\text{gC l}^{-1}$ )  
26 24 of the possible nano-microzooplankton's preys as well as their possible predators: Heterotrophic  
27 25 Bacteria, Cyanobacteria, Picoeukaryotes, Autotrophic Nanoflagellates, Autotrophic Dinoflagellates  
28 26  $<20 \mu\text{m}$ , Autotrophic Dinoflagellates  $>20 \mu\text{m}$ , Diatoms  $<20 \mu\text{m}$ , Diatoms  $>20 \mu\text{m}$ , and  
29 27 Mesozooplankton.

## 30 28 **RESULTS**

### 31 29 **Seasonal and spatial dynamics of environmental conditions.**

1  
2 1 At the Gironde and Coast sites (Fig. 3), the water column varied from halostratified in  
3  
4 2 winter and spring to thermostratified in summer. In June at these two coastal sites, the water column  
5  
6 3 showed both types of stratification (Fig. 3). For example, at the Gironde site, changes in salinity and  
7  
8 4 temperature between the surface to a depth of 10 meters were 31.8 PSU to 34.3 PSU, and 17.2°C to  
9  
10 5 14.7°C. At the Offshore site, the water column showed a moderate halostratification in June with  
11  
12 6 only a surface salinity of 34.5 PSU (Fig. 3). The water column at the Offshore site was not stratified  
13  
14 7 in winter and early spring (February and April, Fig. 3) but became progressively thermostratified  
15  
16 8 from June to October; in late summer the water's temperature stays around 20.1°C from the surface  
17  
18 9 to a depth of 30 m subsequently dropping lower than 18°C below a depth of 40 m (Fig. 3).

18 10 At the three sites, the highest average concentrations of total NO (NO<sub>2</sub> and NO<sub>3</sub>) along the  
19  
20 11 photic zone were observed in February with, for example,  $12.1 \pm 1.6 \mu\text{mol l}^{-1}$  at the Gironde site  
21  
22 12 (Fig. 4). In April, June and September, NO average concentrations varied from  $5.8 \pm 0.6$  to  $0.2 \pm$   
23  
24 13  $0.1 \mu\text{mol l}^{-1}$  along the photic zones of the 3 studied sites (Fig. 4). Considering all three sites and all  
25  
26 14 4 studied periods, the average Si(OH)<sub>4</sub> concentrations along the photic zone were always lower than  
27  
28 15  $6 \mu\text{mol l}^{-1}$  and the average PO<sub>4</sub> concentrations along the photic zone never exceeded  $1 \mu\text{mol l}^{-1}$  (Fig.  
29  
30 16 4). The N/P ratio was highest in April at the two coastal sites (77 at Gironde and 173 at Coast) and  
31  
32 17 lowest in September (below the Redfield ratio with 10 at Gironde and 11 at Coast). The highest N/P  
33  
34 18 ratio at Offshore site was found in September but didn't exceed 13.

### 34 19 **Seasonal and spatial variations in phytoplankton biomass and composition**

35  
36 20  
37 21 The average autotrophic biomass along the photic zone decreased from February to October  
38  
39 22 at the Gironde site ( $376.6 \pm 459.3 \mu\text{gC l}^{-1}$  to  $10.9 \pm 2.4 \mu\text{gC l}^{-1}$ , Fig. 5). At the Coast and Offshore  
40  
41 23 sites, the average autotrophic biomass along the photic zone first increased between February and  
42  
43 24 April, reaching, respectively,  $116.2 \pm 15.6 \mu\text{gC l}^{-1}$  and  $155.8 \pm 40.1 \mu\text{gC l}^{-1}$  (Fig. 5a) and then  
44  
45 25 decreased in June and October. Large diatoms were responsible for more than 70% of the total  
46  
47 26 biomass in February, April and June at the Gironde site and in February and April at both Coast and  
48  
49 27 Offshore sites (Fig. 5b). In February, diatoms were dominated by large chain forming cells (e.g.  
50  
51 28 *Thalassiosira sp.*) at the two coastal sites. Smaller diatoms (i.e. diatoms <20 μm) such as  
52  
53 29 *Leptocylindrus minumus* became more abundant in April at Coast site (Fig. 5b). The biomass of  
54  
55 30 smaller autotrophs (ANF, cyanobacteria and picoeukaryotes) proportionally increased from April to  
56  
57 31 October at all three stations and dominated in October (> 60% of average biomass, Fig. 5b).

### 57 32 **Seasonal and spatial variations of nano-microzooplankton biomass and taxonomic composition**

58  
59 33  
60 34 At the three sites, average biomasses along the photic zone of nano-microzooplankton  
35 peaked in April at Gironde, Coast and Offshore with  $27.9 \pm 13.2 \mu\text{gC l}^{-1}$ ,  $57.1 \pm 32.8 \mu\text{gC l}^{-1}$  and

1  
2 1  $27.2 \pm 13.6 \mu\text{gC l}^{-1}$ , respectively (Fig. 6a). The average biomass was still relatively high in June at  
3  
4 2 the Gironde and Coast sites ( $> 19.5 \mu\text{gC l}^{-1}$ , Fig. 6a) but never exceeded  $8.2 \mu\text{gC l}^{-1}$  in February and  
5  
6 3 October. The Offshore site always had the lowest average biomasses of the three sites (Fig. 6a).

7 4 Ciliates, especially naked ciliates, counted for a large part ( $> 70\%$ , Fig. 6b) of the total  
8  
9 5 average nano-microzooplankton biomass at Coast site in February and April as well as in April at  
10  
11 6 Offshore site. The relative proportion of ciliates in the average biomass of nano-microzooplankton  
12  
13 7 was always the highest in April at the three studied sites (Fig 6b). The relative proportion of nauplii  
14  
15 8 biomass was higher in February and October compare to April and June at all sites (Fig. 6b) despite  
16  
17 9 small range of variations of their absolute abundances. The rest of the nano-microzooplankton  
18  
19 10 biomass was largely due to the unarmoured dinoflagellates (up to 61.8% of the total biomass, in  
20  
21 11 June at Offshore site, Fig. 6b). Even though the HNF had very high abundances (Table 2), their  
22  
23 12 contribution to the total biomass of the nano-microzooplankton was very low, overall less than 8%  
24  
25 13 (Fig. 6b).

#### 25 14 **Seasonal and spatial dynamics of nano-microzooplankton size structure.**

26  
27 15 Analysis of the size-abundance spectra (SAS) showed typical linear relationships at every  
28  
29 16 sites and every sampling period with regression coefficients comprised between 0.7 and 0.9 (Fig. 7).  
30  
31 17 Linear regression slope of the SAS at the Offshore site presented the highest temporal variations  
32  
33 18 with a decrease from -1.02 in February to -0.68 in April (Fig. 8). Both SAS slopes at the Offshore  
34  
35 19 and Coast sites decreased from February to April and then increased from April to October (Fig. 8).  
36  
37 20 At the Gironde site, the SAS slope varied slightly along the first three sampling periods but its value  
38  
39 21 increased in October and became very similar to the slopes of the two other sites (-0.85, -0.87 and -  
40  
41 22 0.89 respectively at Gironde, Coast and Offshore sites; Fig. 8).

42 23 Despite large variations in their abundances (Table 2), the relative proportion of most of the  
43  
44 24 small ESD size classes to the total biomass did not vary remarkably and remained low during the 4  
45  
46 25 sampling periods (Fig. 9a). Cells measuring less than  $40 \mu\text{m}$  ESD never counted for more than 16%  
47  
48 26 of the total nano-microzooplankton biomass at any of the sites (Fig. 9a). The relative proportion of  
49  
50 27 the  $40\text{-}50\mu\text{m}$  ESD-size class was at all times much higher with values above 16% (Fig. 9a).  
51  
52 28 Furthermore, it dominated the community in June at Gironde and Coast sites, contributing for  
53  
54 29 72.1% and 68.2% of the total biomass, respectively, (Fig. 9a) with cell abundances of several  
55  
56 30 thousand per liter (Table 2). With the exception of April, the nano-microzooplankton biomass at  
57  
58 31 Offshore site was always dominated by this  $40\text{-}50 \mu\text{m}$  ESD-size class with a relative proportion  
59  
60 32 greater than 40% (Fig. 9a). Large microzooplankton ( $> 60\mu\text{m}$  ESD) contributing substantially to the  
33  
34 34 total community biomass in February and October at Gironde and Coast, with a relative proportion  
greater than 46% and in April at Offshore site with a relative proportion of 52.8% (Fig. 9a).

1  
2 1 The dynamics of the nano-microzooplankton body size structure followed the same pattern  
3  
4 2 as the ESD size structure (Fig. 9b). The small body size classes ( $< 1 \text{ ngC cell}^{-1}$ ) didn't contribute  
5  
6 3 significantly to the final community biomass ( $< 24.5\%$ , Fig. 9b) but had very high abundance (Table  
7  
8 4 2). The relative proportion of the intermediate body size class ( $1-5 \text{ ngC cell}^{-1}$ ) was the highest in  
9  
10 5 June at the three sites and this size class dominated the final community biomass in April and June  
11 6 at Gironde, in June at Coast and in February and June at Offshore (Fig. 9b). The largest body size  
12  
13 7 classes ( $> 10 \text{ ngC cell}^{-1}$ ) dominated the nano-microzooplankton biomass in February and October at  
14  
15 8 Gironde and Coast sites and in April at Coast and Offshore sites (Fig. 9b) but never dominated the  
16  
17 9 total community abundance (Table 2).

### 18 10 **Comparing the three classification types.**

19  
20 11 The Mantel test revealed a positive and significant correlation between the three  
21  
22 12 microzooplankton classification types (Table 3). The level of similarities between each of the 12  
23  
24 13 data sets (3 sites and 4 sampling periods) in terms of nano-microzooplankton composition was  
25  
26 14 comparable with every classification we used ( $r > 0.8$  and  $p < 0.0001$ ). In other words, there was no  
27  
28 15 difference in the characteristics of nano-microzooplankton dynamics when studying either  
29  
30 16 community size structure or taxonomic composition.

### 31 17 **Linking nano-microzooplankton dynamics and environmental conditions.**

32 18 Rank correlation of physical variables to the nano-microzooplankton abundances indicated  
33  
34 19 that the best physical variables driving community patterns were temperature and  $\text{NO}_3$   
35  
36 20 concentration whatever classification types we were using (Table 4). BEST analysis also showed  
37  
38 21 that the nano-microzooplankton dynamic was associated with bacterial and cyanobacterial  
39  
40 22 dynamics with every classification types were used. These associations were also observed with  
41  
42 23 ANF dynamics when using ESD and Taxons classifications only (Table 4). It was finally noted that  
43  
44 24 the correlation coefficient obtained with the BEST analyses, while significant, were not in any case  
45  
46 25 very high ( $< 0.5$ , Table 4).

## 50 27 **DISCUSSION**

51  
52 28  
53 29 In temperate marine environments, seasonal variations of the physico-chemical parameters  
54  
55 30 of the water column have a strong influence on the structure of phytoplanktonic communities  
56  
57 31 (Margalef, 1958; Fenchel, 1988; Breton et al., 2000; Chang et al., 2003). In summer, vertical  
58  
59 32 stratification is generally associated with nutrient limitation and regenerated production based on  
60  
33 the uptake of ammonium by small-celled phytoplankton populations (Margalef, 1958; Valiela,  
34 1995; Chang et al., 2003). On the other hand, winter/spring vertical mixing is usually associated

1  
2 1 with nitrate availability and new production by large-celled phytoplankton populations (Margalef,  
3 1958; Valiela, 1995; Bury et al., 2001; Irigoien et al., 2005). In addition, coastal areas receiving  
4 2 river runoffs may see that latter increased due to the accumulation of terrestrial nutrients in late  
5 3 winter/spring (Domingues et al., 2005; Alvarez et al., 2009). Although the Bay of Biscay makes no  
6 4 exception with highest thermostratification of the water column in late summer at the three studied  
7 5 sites and high concentrations of nitrate in late winter at the site closest to the Gironde estuary, areas  
8 6 under the influence of the Gironde runoff showed high halostratification of the water column as  
9 7 soon as February. As a consequence, phytoplankton biomass and composition present strong spatio-  
10 8 temporal variations with large diatoms dominating the community in winter/spring and small  
11 9 autotrophic cells dominating in late summer. Nano-microzooplankton is a direct predator of  
12 10 phytoplankton cells (Capriulo, 1991; Capriulo et al., 1991; Calbet and Landry, 2004). As a bottom-  
13 11 up effect, we can assume that changes in nano-microzooplankton community will follow changes in  
14 12 phytoplankton community structure. Indeed, heterotrophic protist abundances have been described  
15 13 as a function of food availability in many previous studies (e.g. Dolan and Coats, 1990). Following  
16 14 this argument, the lack of suitable resources in September (dominance of cyanobacteria) and  
17 15 February (large diatoms) might be the principal factor responsible for the low summer and winter  
18 16 nano-microzooplankton biomasses in the Bay of Biscay. Verity (1986) and Lynn and Montagnes  
19 17 (1991) show that the distribution of ciliates generally present a close association with  
20 18 nanophytoplankton in temperate coastal ecosystems. In this study, ciliate biomasses seem more  
21 19 closely related to microphytoplankton than nanophytoplankton but this might be because those  
22 20 kinds of biomass composition snapshots are not powerful in showing the real level of produced  
23 21 biomass of highly grazed prey. However, the BEST analysis revealed that ANF biomasses were one  
24 22 of the main factors explaining the changes in the nano-microzooplankton taxonomic structure. In  
25 23 late summer, the proportion of nauplii in the nano-microzooplankton community increases along  
26 24 the coast of the Bay of Biscay since most of the reproduction of adult copepods occurs during  
27 25 summer (Sautour and Castel, 1998).

28 26  
29 27 Most of the trophic relationships within planktonic food webs are based on size, i.e. the  
30 28 larger consuming the smaller (Platt and Denman, 1977; Moloney and Field, 1991; Caparroy et al.,  
31 29 2000; Stock et al., 2008). Therefore, the size structure of nano-microzooplankton should reflect the  
32 30 phytoplanktonic composition with large grazers more abundant when large diatoms are the main  
33 31 primary producers. When the community is described with SAS, this situation would then  
34 32 correspond to flatter slopes, i.e. lower absolute values (Platt and Denman, 1977; Rodriguez and  
35 33 Mullin, 1986). This is what was observed with this data. The slopes of the linear regression, when  
36 34 fitted to the SAS, are steepest (i.e. highest absolute value) in October when bacteria are very  
37 35 abundant and when picophytoplankton is the principal primary producer. The three study sites show



1  
2 1 time differences in the phytoplanktonic dynamics with a diatom bloom occurring as soon as  
3  
4 2 February at Gironde but only in April at Coast and Offshore. This difference is visible when  
5  
6 3 analyzing the nano-microzooplankton SAS slope changes at the three sampling sites. The estuarine  
7  
8 4 site, Gironde, has a flatter slope in February due to the presence of bigger cells in the nano-  
9  
10 5 microzooplankton community. The SAS slopes at Coast and Offshore decrease in April showing an  
11  
12 6 increase of large cells in the nano-microzooplankton when the bloom of diatoms is peaking at those  
13  
14 7 two sites.

15 8 In the literature, most of the studies on plankton size structure are focused on phytoplankton  
16  
17 9 (e.g. Cermeno and Figueiras, 2008) or the modeling of the whole planktonic community (e.g. Baird  
18  
19 10 and Suthers, 2007). However, with the increasing interest on nano-microzooplankton community as  
20  
21 11 a major player of the planktonic food web (Azam et al., 1983), numerous studies are describing the  
22  
23 12 nano-microzooplankton dynamics in very different environments (e.g. Shinada et al., 2003; Gaedke  
24  
25 13 and Wickham, 2004; Fileman and Leakey, 2005). Identification of nano-microzooplankton cells  
26  
27 14 through classical microscopic methods is highly time consuming and new technologies such as  
28  
29 15 automatic plankton counters are quickly extending their attraction among ecologists (see review in  
30  
31 16 Benfield et al., 2007). But those technologies, despite being very efficient at counting cells, do not  
32  
33 17 have the same capabilities in cell identification as an expert human eye. A good understanding of  
34  
35 18 the community dynamics depends on a good description of its composition and structure. Size  
36  
37 19 parameters of nano-microzooplankton cells are easy to compile but the question is whether it is a  
38  
39 20 sufficient descriptor on its own.

40 21 The first problem occurs when choosing the best size parameter to use. Each study  
41  
42 22 corresponds to a different parameter: length, diameter, equivalent spherical diameter, biovolume,  
43  
44 23 cell carbon content, etc...(e.g. Kimmel et al., 2006; Cermeno and Figueiras, 2008; Basedow et al.,  
45  
46 24 2009). Despite the will to describe the same characteristic of a cell and its size, this variety may be  
47  
48 25 an obstacle to those who would like to compare data from different studies. In this study, the three  
49  
50 26 most widely used parameters were purposely used: biovolume, ESD and body mass. Since ESD  
51  
52 27 and biovolume are directly related to one another, one can be used to compare the structure of the  
53  
54 28 other without any difficulties. But depending on the conversion factor used, the body mass may not  
55  
56 29 be directly proportional to the first two parameters. In this case, the general relationship between  
57  
58 30 ESD and body mass matches an exponential slope. Although ESD is not representing the trophic  
59  
60 31 value of the nano-microzooplanktonic cell, it is an easier parameter to measure or estimate than the  
32  
33 32 body mass. Moreover, since the definition of ESD is the same for cells of every size and every sea,  
34  
35 33 it would be more reliable as a comparable and universal parameter. Therefore, we recommend on  
36  
37 34 using ESD as a proxy of size when studying nano-microzooplanktonic size structure.

1  
2 1 The second problem, and without doubt the most important, concerns the efficiency of size  
3  
4 2 descriptors to define nano-microzooplankton dynamics as clearly as the taxonomy. Before  
5  
6 3 answering that problem, the definition of “dynamics” of community needs to be agreed upon. As  
7  
8 4 proposed in Lindeman’s paper on tropho-dynamics (1942), the dynamics of a community is the  
9  
10 5 pattern of its changes in size and composition over time and space resulting from the coactions  
11  
12 6 between the living members of the community and reactions of those members with the non-living  
13  
14 7 environment. Following that definition, the nano-microzooplankton dynamics portrayed with size  
15  
16 8 descriptors should show the same pattern of changes and of differences between seasons and sites  
17  
18 9 as the dynamics portrayed with taxonomy. The Mantel test used in this study focused on comparing  
19  
20 10 the patterns of the spatio-temporal differences observed in the size and composition of the nano-  
21  
22 11 microplankton community with three different descriptors (taxonomy and two size parameters).  
23  
24 12 Statistics didn’t divulge any differences between the three dynamics. Therefore, we can argue that  
25  
26 13 using the size structure to characterize the dynamics of nano-microplankton lead to revealing the  
27  
28 14 same spatio-temporal patterns when using a taxonomic structure.

29 15 Finally, the ultimate goal of community dynamics studies is the understanding of the  
30  
31 16 relationships of a community with its environment and more precisely with its trophic environment  
32  
33 17 (i.e. Lindeman, 1942). Knowledge of community dynamics is necessary in food webs and  
34  
35 18 ecosystem process studies as it provides a way to analyze the transfer rates of energy matter and the  
36  
37 19 impact of environmental disturbances (de Ruiter et al., 2005). The historical way to examine  
38  
39 20 community dynamics was by changes in its taxonomic structure and the trophic relationships  
40  
41 21 between taxonomical groups (e.g. Andersen and Sorensen, 1986; Uitto et al., 1997). We have  
42  
43 22 shown with this study that size descriptors were efficiently defining nano-microzooplankton  
44  
45 23 dynamics, but are they also good in revealing biotic and abiotic factors controlling that dynamics?  
46  
47 24 Although correlations do not necessary mean direct causal relationships such as trophic links, they  
48  
49 25 might give information on the physical and biological parameters influencing the nano-  
50  
51 26 microzooplankton dynamics (Legendre and Legendre, 1998). In our study, the interest of such an  
52  
53 27 analysis was to compare the results obtained with the three different types of classification. The  
54  
55 28 results of the BEST analyses were the same for the three types of classification compared in our  
56  
57 29 study with very low but similar Spearman rank correlation coefficients. The only difference was  
58  
59 30 found with the Body size classification, for which only two biological parameters were significantly  
60  
61 31 explaining the nano-microzooplanktonic dynamics, compared to the three found for the two other  
62  
63 32 classifications.

64 33 In conclusion, considering the results of our study, we argue that nano-microplankton  
65  
66 34 dynamics are well defined by the study of their size structure. The use of automatic plankton  
67  
68 35 counters such as the Flowcam technology (Sieracki et al., 1998) is therefore relevant despite the



1  
2 1 lack of precise taxonomical identification. However, homogenizing the size descriptor used among  
3  
4 2 studies is needed in order to make worldwide data comparisons and ESD/biovolume should be  
5  
6 3 favored. We also recommend further investigations on size-based trophic relationships and size  
7  
8 4 structure dynamics in order to confirm the results found in the Bay of Biscay.  
9  
10 5

## 11 6 **ACKNOWLEDGEMENTS**

12 7

13  
14 8 This study was supported by the French “Programme National Environment Côtier” – Bay  
15  
16 9 of Biscay working site – and Ifremer. The authors wish to thank captains and crews of R/V Thalia  
17  
18 10 for support activities, C. Courties (Observatoire Océanologique de Banyuls, France) for the  
19  
20 11 flowcytometry analyses, F. Mornet (IFREMER, FREDD, France) and V. Huet (LIENSs, France)  
21  
22 12 for nutrient and chlorophyll measurements, P. Malterre and M-J Capdeville (students of University  
23  
24 13 of La Rochelle, France) for metazoan counts.  
25 14  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2 1 **BIBLIOGRAPHY**  
3  
4 2  
5  
6  
7 3

- 8 4 Adl S.M., Simpson, A.G.B., Farmer, M.A., Andersen, R.A., Anderson, O.R., Barta, J.R., Bowser,  
9 5 S.S., Brugerolle, G., Fensome, R.A., Fredericq, S., James, T.Y., Karpov, S., Kugrens, P.,  
10 6 Krug, J., Lane, C.E., Lewis, L.A., Lodge, J., Lynn, D.H., Mann, D.G., Mccourt, R.M.,  
11 7 Mendoza, L., Moestrup, O., Mozley-Standridge, S.E., Nerad, T.A., Shearer, C.A., Smirnov,  
12 8 A.V., Spiegel, F.W., Taylor, M.F.J.R. (2005) The new higher level classification of  
13 9 eukaryotes with emphasis on the taxonomy of protists. *Journal of Eukaryotic*  
14 10 *Microbiology*, **52**, 399-451  
15 11  
16 12 Alvarez E., Nogueira, E., Acuna, J.L., Lopez-Alvarez, M., Sostres, J.A. (2009) Short-term  
17 13 dynamics of late-winter phytoplankton blooms in a temperate ecosystem (Central  
18 14 Cantabrian Sea, Southern Bay of Biscay). *J Plankton Res*, **31**, 601-617  
19 15  
20 16 Andersen P., Sorensen, H.M. (1986) Population dynamics and trophic coupling in pelagic  
21 17 microorganisms in eutrophic coastal waters. *Mar Ecol Prog Ser*, **33**, 99-109  
22 18  
23 19 Azam F., Fenchel, T., Field, J.G., Gray, J.S., Meyer-Reil, L.A., Thingstad, F. (1983) The ecological  
24 20 role of water-column microbes in the sea. *Mar Ecol Prog Ser*, **10**, 257-263  
25 21  
26 22 Baird M.E., Suthers, I.M. (2007) A size-resolved pelagic ecosystem model. *Ecol Model*, **203**, 185-  
27 23 203  
28 24  
29 25 Basedow S.L., Tande, K.S., Zhou, M. (2009) Biovolume spectrum theories applied: spatial patterns  
30 26 of trophic levels within a mesozooplankton community at the polar front. *J Plankton Res*,  
31 27 fbp110  
32 28  
33 29 Benfield M.C., Grosjean, P., Culverhouse, P., Irigoien, X., Sieracki, M.E., Lopez-Urrutia, A., Dam,  
34 30 H.G., Hu, H.G., Davis, Q., Hansen, A., Pilskaln, C.H., Riseman, E., Schultz, H., Utgoff,  
35 31 P.E., Gorsky, G. (2007) RAPID - Research on Automated Plankton Identification.  
36 32 *Oceanography*, **20**, 12-26  
37 33  
38 34 Blanchot J., Rodier, M. (1996) Picophytoplankton abundance and biomass in the western tropical  
39 35 Pacific Ocean during the 1992 El Nino year: results from flow cytometry. *Deep-Sea Res Pt*  
40 36 *I*, **43**, 877-895  
41 37  
42 38 Bode A., Alvarez-Ossorio, M., Gonzalez, N. (1998) Estimation of mesozooplankton biomass in a  
43 39 coastal upwelling area off NW Spain. *J Plankton Res*, **20**, 1005-1014  
44 40  
45 41 Breton E., Brunet, C., Sautour, B., Brylinski, J.-M. (2000) Annual variations of phytoplankton  
46 42 biomass in the Eastern English Channel: comparison by pigment signatures and microscopic  
47 43 counts. *J Plankton Res*, **22**, 1423-1440  
48 44  
49 45  
50 46  
51 47  
52 48  
53 49  
54 50  
55 51  
56 52  
57 53  
58 54  
59 55  
60 56

- 1  
2 1 Brown J.H., Gillooly, J.F., Allen, A.P., Savage, V.M., West, G.B. (2004) Toward a metabolic  
3 theory of ecology *Ecology*, **85**, 1771-1789  
4 2  
5 3 Bury S.J., Boyd, P.W., Preston, T., Savidge, G., Owens, N.J.P. (2001) Size-fractionated primary  
6 production and nitrogen uptake during a North Atlantic phytoplankton bloom: implications  
7 for carbon export estimates. *Deep-Sea Res Pt I*, **48**, 689-720  
8 4  
9 5 Calbet A., Landry, M.R. (2004) Phytoplankton growth, microzooplankton grazing, and carbon  
10 cycling in marine systems. *Limnol Oceanogr*, **49**, 51-57  
11 6  
12 7 Caparroy P., Thygesen, U.H., Visser, A.W. (2000) Modelling the attack success of planktonic  
13 predators: patterns and mechanisms of prey size selectivity. *J Plankton Res*, **22**, 1871  
14 8  
15 9 Capriulo G.M. (1991) Community grazing in heterotrophic marine protista - session summary. In:  
16 Reid P.C., Turley, C.M. and Burkill, P.H. (eds) *Protozoa and their role in marine processes*.  
17 Springer-Verlag, Berlin, pp. 205-218  
18 10  
19 11 Capriulo G.M., Sherr, E.B., Sherr, B.F. (1991) Trophic behaviour and related community feeding  
20 activities of heterotrophic marine protists. In: Reid P.C., Turley, C.M. and Burkill, P.H.  
21 (eds) *Protozoa and their role in marine processes*. Springer-Verlag, Berlin, pp. 219-265  
22 12  
23 13 Cermeno P., Figueiras, F.G. (2008) Species richness and cell-size distribution: size structure of  
24 phytoplankton communities. *Mar Ecol Prog Ser*, **357**, 79-85  
25 14  
26 15 Chang F.H., Zeldis, J., Gall, M., Hall, J. (2003) Seasonal and spatial variation of phytoplankton  
27 assemblages, biomass and cell size from spring to summer across the north-eastern New  
28 Zealand continental shelf. *J Plankton Res*, **25**, 737-758  
29 16  
30 17 Clarke K.R., Warwick, R.M. (2001) *Changes in marine communities: an approach to statistical  
31 analysis and interpretation (PRIMER-E)*, Plymouth Marine Laboratory, Plymouth, UK  
32 18  
33 19 Cowlshaw R.J. (2004) Seasonal coupling between ciliate and phytoplankton standing stocks in the  
34 South Slough of Coos Bay, Oregon. *Estuaries*, **27**, 539-550  
35 20  
36 21 Culverhouse P., Williams, R., Benfield, M.C., Flood, P.R., Sell, A.F., Mazzocchi, M.G., Buttino, I.,  
37 Sieracki, M. (2006) Automatic image analysis of plankton: future perspectives. *Mar Ecol  
38 Prog Ser*, **312**, 297-309  
39 22  
40 23 Daufresne M., Lengfellner, K., Sommer, U. (2009) Global warming benefits the small in aquatic  
41 ecosystems. *P Nat Acad Sci USA*, **106**, 12788-12793  
42 24  
43 25 de Ruiter P.C., Wolters, V., Moore, J.C. (2005) Dynamic food webs. In: de Ruiter P.C., Wolters, V.  
44 and Moore, J.C. (eds) *Dynamic food webs*. Academic Press, pp. 3-9  
45 26  
46 27 Dolan J.R., Coats, D.W. (1990) Seasonal abundances of planktonic ciliates and microflagellates in  
47 mesohaline Chesapeake Bay Waters. *Est Coast Shelf S*, **31**, 157-175  
48 28  
49 29 Domingues R.B., Barbosa, A., Galvao, H. (2005) Nutrients, light and phytoplankton succession in a  
50 temperate estuary (the Guadiana, south-western Iberia). *Est Coast Shelf S*, **64**, 249-260  
51 30  
52 31  
53 32  
54 33  
55 34  
56 35  
57  
58  
59  
60

- 1  
2 1 Ducklow H.W. (2003) Biogeochemical Provinces: towards a JGOFS synthesis. In: Fasham M.J.R.  
3 (ed) *Ocean Biogeochemistry - The role of the ocean carbon cycle in Global Change*.  
4 2 Springer, Berlin, pp. 3-17  
5 3  
6 4 Ducklow H.W., Carlson, C.A. (1992) Oceanic Bacterial Production. *Adv Microb Ecol*, **12**, 113-181  
7 5  
8 6 Fenchel T. (1988) Marine plankton food chains. *Annu Rev Ecol Syst*, **19**, 19-38  
9 7  
10 8 Fileman E.S., Leakey, R.J.G. (2005) Microzooplankton dynamics during the development of the  
11 6 spring bloom in the north-east Atlantic. *J Mar Biol Ass UK*, **85**, 741-753  
12 7  
13 8 Finlay B.J. (2004) Protist taxonomy: an ecological perspective. *Philosophical Transactions of the*  
14 8 *Royal Society of London. Series B: Biological Sciences*, **359**, 599-610  
15 9  
16 9 Frost B.W. (1972) Effects of size and concentration of food particles on the feeding behavior of the  
17 10 marine planktonic copepod *Calanus pacificus*. *Limnol Oceanogr*, **17**, 805-815  
18 10  
19 11 Gaedke U., Wickham, S.A. (2004) Ciliate dynamics in response to changing biotic and abiotic  
20 11 conditions in a large, deep lake (Lake Constance). *Aquat Microb Ecol*, **34**, 247-261  
21 12  
22 13 Gillooly J.F., Brown, J.H., West, G.B., Savage, V.M., Charnov, E.L. (2001) Effects of Size and  
23 13 Temperature on Metabolic Rate. *Science*, **293**, 2248-2251  
24 14  
25 14 Gowing M.M., Garrison, D.L., Wishner, K.F., Gelfman, C. (2003) Mesopelagic microplankton of  
26 15 the Arabian Sea. *Deep-Sea Res Pt I*, **50**, 1205-1234  
27 15  
28 16 Irigoien X., Flynn, K.J., Harris, R.P. (2005) Phytoplankton blooms: a 'loophole' in  
29 16 microzooplankton grazing impact? *J Plankton Res*, **27**, 313-321  
30 17  
31 18 Kimmel D.G., Roman, M.R., Zhang, X.S. (2006) Spatial and temporal variability in factors  
32 18 affecting mesozooplankton dynamics in Chesapeake Bay: Evidence from biomass size  
33 19 spectra. *Limnol Oceanogr*, **51**, 131-141  
34 19  
35 20 Koutsikopoulos C., Beilois, P., Leroy, C., Taillefer, F. (1998) Temporal trends and spatial  
36 20 structures of the sea surface temperature in the Bay of Biscay. *Oceanol Acta*, **21**, 335-344  
37 21  
38 22 Labry C., Herbland, A., Delmas, D. (2002) The role of phosphorus on planktonic production of the  
39 22 Gironde plume waters in the Bay of Biscay. *J Plankton Res*, **24**, 97-117  
40 22  
41 23 Lazure P., Jegou, A.-M. (1998) 3D modelling of seasonal evolution of Loire and Gironde plumes on  
42 23 Biscay Bay continental shelf. *Oceanol Acta*, **21**, 165-177  
43 24  
44 25 Legendre P., Legendre, L. (1998) *Numerical ecology*, Elsevier, Amsterdam  
45 25  
46 26 Lessard E.J., Swift, E. (1986) Dinoflagellates from the North Atlantic classified as phototrophic or  
47 26 heterotrophic by epifluorescence microscopy. *J Plankton Res*, **8**, 1209-1215  
48 27  
49 28 Lindeman R.L. (1942) The Trophic-Dynamic Aspect of Ecology. *Ecology*, **23**, 399-417  
50 28  
51 29 Lupez-Urrutia Å., San Martin, E., Harris, R.P., Irigoien, X. (2006) Scaling the metabolic balance of  
52 29 the oceans. *P Nat Acad Sci USA*, **103**, 8739-8744  
53 30  
54 31  
55 32  
56 33  
57 34  
58  
59  
60

- 1  
2 1 Lynn D.H., Montagnes, D.J.S. (1991) Global production of heterotrophic marine planktonic ciliates.  
3  
4 2 In: Reid P.C., Turley, C.M. and Burkill, P.H. (eds) *Protozoa and their role in marine*  
5  
6 3 *processes*. Springer-Verlag, Berlin, pp. 281-307
- 7 4 Margalef R. (1958) Temporal succession and spatial heterogeneity in phytoplankton. In: Buzzati-  
8  
9 5 Traverso A.A. (ed) *Perspective in marine biology*. University of California Press, Berkeley  
10 6 and Los Angeles, pp. 323-349
- 11 7 Marie D., Partensky, F., Simon, N., Guillou, L., Vaultot, D. (2000) Flow cytometry analysis of  
12 8 marine picoplankton. In: Diamond R.A. and DeMaggio, S. (eds) *In living colors: Protocols*  
13 9 *in flow cytometry and cell sorting*. Springer-Verlag, Berlin, pp. 421-454
- 14 10 Menden-Deuer S., Lessard, E.J. (2000) Carbon to volume relationships for dinoflagellates, diatoms,  
15 11 and other protist plankton. *Limnol Oceanogr*, **45**, 569-579
- 16 12 Moloney C.L., Field, J.G. (1989) Generic allometric equations for rates of nutrient uptake,  
17 13 ingestion, and respiration in plankton organisms. *Limnol Oceanogr*, **34**, 1290-1299
- 18 14 Moloney C.L., Field, J.G. (1991) The size-based dynamics of plankton food webs. I. A simulation  
19 15 model of carbon and nitrogen flows. *J Plankton Res*, **13**, 1003-1038
- 20 16 Moran X.A.G., Lopez-Urrutia, A., Calvo-Diaz, A., Li, W.K.W. (2010) Increasing importance of  
21 17 small phytoplankton in a warmer ocean. *Global Change Biol*, **16**, 1137-1144
- 22 18 Pelegri S.P., Dolan, J.R., Rassoulzadegan, F. (1999) Use of high temperature catalytic oxidation  
23 19 (HTCO) to measure carbon content of microorganisms. *Aquat Microb Ecol*, **16**, 273-280
- 24 20 Peters R.H., Downing, J.A. (1984) Empirical Analysis of Zooplankton Filtering and Feeding Rates.  
25 21 *Limnol Oceanogr*, **29**, 763-784
- 26 22 Pierce R.W., Turner, J.T. (1992) Ecology of planktonic ciliates in marine food webs. *Rev Aquat S*,  
27 23 **6**, 139-181
- 28 24 Platt T., Denman, K.L. (1977) Organisation in the pelagic ecosystem. *Helgolander Meeresun*, **30**,  
29 25 575-581
- 30 26 Putt M., Stoecker, D.K. (1989) An experimentally determined carbon: volume ratio for marine  
31 27 "oligotrichous" ciliates from estuarine and coastal waters. *Limnol Oceanogr*, **34**, 1097-1103
- 32 28 Rodriguez J., Mullin, M.M. (1986) Relation between biomass and body weight of plankton in a  
33 29 steady state oceanic ecosystem. *Limnol Oceanogr*, **31**, 361-370
- 34 30 Sanders R.W. (1987) Tintinnids and other microzooplankton - seasonal distributions and  
35 31 relationships to resources and hydrography in a Maine estuary. *J Plankton Res*, **9**, 65-77
- 32 32 Sautour B., Castel, J. (1998) Importance of microzooplanktonic crustaceans in the coastal food  
33 33 chain: Bay of Marennes-Oleron, France. *Oceanologica Acta*, **21**, 105-112
- 34 34 Sherr E., Sherr, B.F., Paffenhofer, G.-A. (1986) Phagotrophic protozoa as food for metazoans : a  
35 35 "missing" trophic link in marine pelagic food webs? *Mar Microb Food Webs*, **1**, 61-80



- 1  
2 1 Sherr E.B., Caron, D.A., Sherr, B.F. (1994) Staining of heterotrophic protists for visualisation via  
3  
4 2 epifluorescence microscopy. In: Kemp P.F., Sherr, B.F., Sherr, E.B. and Coll, J.J. (eds)  
5  
6 3 *Handbook of methods in aquatic microbial ecology*. Lewis Publishers, Boca Raton, pp. 213-  
7  
8 4 227
- 9 5 Sherr E.B., Sherr, B.F. (2000) Marine microbes: an overview. In: Kirchman D.L. (ed) *Microbial*  
10  
11 6 *ecology of the oceans*. Wiley-Liss, New York, pp. 13-46
- 12 7 Shinada A., Ban, S., Ikeda, T. (2003) Seasonal changes in nano/micro-zooplankton herbivory and  
13  
14 8 heterotrophic nano-flagellates bacterivory off Cape Esan, Southwestern Hokkaido, Japan. *J*  
15  
16 9 *Oceanogr*, **59**, 609-618
- 17  
18 10 Shinada A., Ban, S., Yamada, Y., Ikeda, T. (2005) Seasonal variations of plankton food web  
19  
20 11 structure in the coastal water off Usujiri Southwestern Hokkaido, Japan. *J Oceanogr*, **61**,  
21  
22 12 645-654
- 23 13 Sieburth J.M., Smetacek, V., Lenz, J. (1978) Pelagic Ecosystem Structure: Heterotrophic  
24  
25 14 Compartments of the Plankton and Their Relationship to Plankton Size Fractions. *Limnol*  
26  
27 15 *Oceanogr*, **23**, 1256-1263
- 28  
29 16 Sieracki C.K., Sieracki, M.E., Yentsch, C.S. (1998) An imaging-in-flow system for automated  
30  
31 17 analysis of marine microplankton. *Mar Ecol Prog Ser*, **168**, 285-296
- 32 18 Sleight M.A. (1991) A taxonomic review of heterotrophic protists important in marine ecology. In:  
33  
34 19 Reid P.C., Turley, C.M. and Burkill, P.H. (eds) *Protozoa and their role in marine processes*.  
35  
36 20 Springer-Verlag, Berlin, pp. 9-38
- 37 21 Sommaruga R. (1995) Microbial and classical food webs: A visit to a hypertrophic lake. *FEMS*  
38  
39 22 *Microbiol Ecol*, **17**, 257-270
- 40  
41 23 Stock C.A., Powell, T.M., Levin, S.A. (2008) Bottom-up and top-down forcing in a simple size-  
42  
43 24 structured plankton dynamics model. *J Mar Syst*, **74**, 134-152
- 44 25 Strickland J.D.H., Parsons, T.R. (1972) A practical handbook of seawater analysis. *Bull Fish Res*  
45  
46 26 *Board Can*, **167**, 47-89
- 47  
48 27 Thingstad T.F., Rassoulzadegan, F. (1999) Conceptual models for the biogeochemical role of the  
49  
50 28 photic zone microbial food web, with particular reference to the Mediterranean Sea. *Prog*  
51  
52 29 *Oceanogr*, **44**, 271-286
- 53 30 Uitto A., Heiskanen, A.-S., Lignell, R., Autio, R., Pajuniemi, R. (1997) Summer dynamics of the  
54  
55 31 coastal planktonic food web in the northern Baltic Sea. *Mar Ecol Prog Ser*, **151**, 27-41
- 56  
57 32 Utermöhl H. (1958) Zur Vervollkommung der quantitativen phytoplankton-methodik. *Mitteilungen*  
58  
59 33 *Internationale Vereinigung für Theoretische und Angewandte Limnologie*, **9**, 1-38
- 60 34 Valiela I. (1995) *Marine ecological processes*, Springer-Verlag, New York

- 1  
2 1 Verity P.G., Langdon, C. (1984) Relationships between Lorica volume, carbon, nitrogen, and ATP  
3 content of tintinnids in Narragansett Bay. *J Plankton Res*, **6**, 859-868  
4 2  
5 3 Verity P.G. (1986) Growth rates of natural tintinnid populations in Narragansett Bay. *Mar Ecol*  
6 *Prog Ser*, **29**, 117-126  
7 4  
8 5 Waterbury J.B., Watson, S.W., Guillard, R.R.L., Brand, L.E. (1979) Widespread occurrence of a  
9 unicellular, marine, planktonic, cyanobacterium. *Nature*, **277**, 293-294  
10 6  
11 7 West G.B., Savage, V.M., Gillooly, J., Enquist, B.J., Woodruff, W.H., Brown, J.H. (2003)  
12 Physiology (communication arising): Why does metabolic rate scale with body size? *Nature*,  
13 **421**, 713-713  
14 8  
15 9  
16 10 Zeuthen E. (1970) Rate of living as related to body size of organisms. *Polskie Arch Hydrobiol*, **17**,  
17 21-30  
18 11  
19 12  
20 13  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## 1 FIGURE CAPTIONS

2 **Figure 1:** Map of the Bay of Biscay with the location of the three study sites. Dashed line is a  
3 schematic representation of the continental shelf limits.

4 **Figure 2:** Variations of the Body Size (ngC) of nano-microzooplankton cells depending on their  
5 ESD ( $\mu\text{m}$ ).

6 **Figure 3:** Temperature ( $^{\circ}\text{C}$ ) and salinity (PSU) profiles of the water column at the three stations  
7 and the four sampling periods.

8 **Figure 4:** Seasonal variations of nutrient concentration averages ( $\mu\text{M}$ ) over the photic zone of the  
9 three stations: Nitrites and Nitrates (N tot), Phosphates ( $\text{PO}_4$ ) and Silicates (Si).

10 **Figure 5:** Seasonal variations of the average autotrophic biomass (a) and its relative composition  
11 (b) over the photic zone at the three stations. Diat $>20$  for diatoms  $> 20\mu\text{m}$ , Diat $<20$  for  
12 diatoms  $< 20\mu\text{m}$ , ADF $>20$  for autotrophic dinoflagellates  $> 20\mu\text{m}$ , ADF $<20$  for autotrophic  
13 dinoflagellates  $< 20\mu\text{m}$ , ANF for autotrophic nanoflagellates, Picoeuk for eukaryotic  
14 picophytoplankton, Cyanobac for cyanobacteria.

15 **Figure 6:** Seasonal variations of average biomass of nano-microzooplankton (a) and its relative  
16 composition (b) over the photic zone at the three stations. HNF for Heteronano-flagellates; U-  
17 HDF for Unarmoured Hetero-dinoflagellates; A-HDF for Armoured Hetero-dinoflagellates;  
18 N-Cil for Naked Ciliates; Tint for Tintinnid Ciliates; Mr-Cil for Myrionecta rubra ciliates;  
19 Meta for Metazoans.

20 **Figure 7:** Size-abundance spectra and corresponding fitted linear regression ( $r^2>0.7$ ) of nano-  
21 microzooplankton at the 3 sites and for the 4 sampling periods. Size is represented by cell  
22 biovolumes ( $\mu\text{m}^3$ ).

23 **Figure 8:** Temporal and spatial variations of the linear regression slope of the size-abundance  
24 spectra as presented in figure 6.

25 **Figure 9:** Seasonal variations of the nano-microzooplankton community structure over the photic  
26 zone at the three stations in terms of relative biomasses of (a) ESD size ( $\mu\text{m}$ ) classes and (b)  
27 body size ( $\text{ngC cell}^{-1}$ ) classes.

29

1  
2 1 **TABLE CAPTIONS**  
3

4 2 **Table 1:** Factors and formulae with their reference used to convert biovolume to carbon mass of  
5 3 each plankton organisms.  
6  
7

8 4 **Table 2:** Seasonal abundances (cells  $l^{-1}$ ) with their standard deviations, of the different nano-  
9 5 microzooplankton classes from the 3 classification types (Taxons, ESD and Body Size) in average  
10 6 over the photic zone at the three stations.  
11  
12

13 7 **Table 3:** Statistical details of the Mantel tests run to compare similarities between the dynamics of  
14 8 nano-microzooplankton community using three different classification types: Taxons, Body Size  
15 9 and ESD.  $r$  refers to Pearson correlation coefficient. Statistical significance ( $p$ ) was estimated using  
16 10 10,000 permutations.  
17  
18

19 11 **Table 4.** Statistical details of the BEST analysis run to link nano-microzooplankton dynamics and  
20 12 environmental conditions (biological and physical conditions).  $r$  refers to Spearman rank correlation  
21 13 coefficient. Statistical significance ( $p$ ) was estimated with permutations and is  $< 0.01$ .  
22  
23  
24  
25  
26  
27  
28  
29  
30 14  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Table 1

<i>Plankton organisms</i>	<i>Conversion factor or formulae</i>	<i>References</i>
<b>Bacteria</b>	0.016 pgC cell <sup>-1</sup>	Labry et al. (2002)
<b>Cyanobacteria</b>	0.104 pgC cell <sup>-1</sup>	Blanchot and Rodier (1996)
<b>Eucaryotic picophytoplankton</b>	0.22 pgC cell <sup>-1</sup>	Shinada et al. (2005)
<b>Nanoflagellates</b>	0.125 pgC μm <sup>-3</sup> = 3.14 pgC cell <sup>-1</sup> (with mean biovolume of 25.2 μm <sup>3</sup> )	Pelegri et al. (1999) and our data of biovolumes
<b>Dinoflagellates</b>	Log <sub>10</sub> C (in pgC cell <sup>-1</sup> ) = -0.353 + 0.864 * log <sub>10</sub> V	Menden-Deuer & Lessard (2000)
<b>Diatoms</b>	Log <sub>10</sub> C (in pgC cell <sup>-1</sup> ) = -0.541 + 0.811 * log <sub>10</sub> V Log <sub>10</sub> C (in pgC cell <sup>-1</sup> ) = -0.933 + 0.881 * log <sub>10</sub> V	< 3000 μm <sup>3</sup> > 3000 μm <sup>3</sup> Menden-Deuer & Lessard (2000)
<b>Naked Ciliates</b>	0.19 pgC μm <sup>-3</sup>	Putt & Stoecker (1989)
<b>Ciliate Tintinnids</b>	C (in pgC cell <sup>-1</sup> ) = 444.5 + 0.053 * LV	Verity & Langdon (1984)
<b>Copepod Nauplii</b>	pgC ind <sup>-3</sup> = 0.08 × V	Gowing et al. (2003)

V: biovolume (in μm<sup>3</sup>) and LV: Lorica volume (in μm<sup>3</sup>)

Table 2

Classification	Class	Gironde				Coast				Offshore			
		February	April	June	October	February	April	June	October	February	April	June	October
Taxons	HNF ( $10^4$ )	12.7 ± 1.9	5.5 ± 2.6	4.4 ± 4.2	16.2 ± 14.2	14.6 ± 8.5	3.0 ± 0.8	1.7 ± 0.9	2.5 ± 2.4	15.4 ± 4.2	1.2 ± 1.0	1.0 ± 1.7	0.6 ± 0.6
	Mr-Cil	0.0	543.4	64.6	0.0	68.4	2810.1	55.1	11.4	108.3	963.3	3.8	19.0
		± 0.0	± 413.3	± 47.5	± 0.0	± 118.5	± 2517.3	± 64.1	± 19.7	± 114.4	± 1123.6	± 6.6	± 11.9
	Nak-Cil	53.2	4115.4	2976.4	596.6	1411.4	6627.2	3345.9	471.2	2471.9	2346.5	463.6	469.8
		± 92.1	± 2161.6	± 4479.3	± 638.2	± 1272.4	± 4211.9	± 4582.3	± 668.7	± 1587.9	± 1482.4	± 125.1	± 387.8
	Tint	7.6	7.6	9.5	38.0	318.6	0.0	17.1	22.8	19.0	1.9	0.0	0.0
		± 13.2	± 13.2	± 8.7	± 8.7	± 503.3	± 0.0	± 17.1	± 26.1	± 3.3	± 3.3	± 0.0	± 0.0
	U-HDF	1040.0	18373.3	5985.8	332.0	277.3	12133.3	8437.9	1061.9	7522.7	11856.0	5477.3	2617.3
	± 750.0	± 13130.9	± 10188.1	± 394.8	± 317.7	± 8002.8	± 7677.3	± 961.3	± 2422.0	± 4632.4	± 4993.4	± 2858.7	
	A-HDF	485.3	1109.3	2357.3	100.8	69.3	485.3	1079.3	434.7	520.0	485.3	225.3	28.9
		± 240.1	± 1067.4	± 2582.1	± 104.1	± 120.1	± 317.7	± 237.8	± 344.9	± 375.0	± 120.1	± 346.2	± 26.5
	Meta	43.1	44.3	31.2	47.6	20.4	39.6	17.5	24.3	30.2	12.4	12.1	2.5
		± 23.0	± 4.5	± 16.9	± 28.9	± 26.5	± 5.5	± 28.2	± 23.8	± 13.0	± 7.0	± 10.9	± 2.3
ESD ( $\mu\text{m}$ )	<20 ( $10^4$ )	12.7 ± 1.9	5.5 ± 2.6	4.4 ± 4.2	16.2 ± 14.2	14.6 ± 8.5	3.0 ± 0.8	1.7 ± 0.9	2.5 ± 2.4	15.4 ± 4.2	1.2 ± 1.0	1.0 ± 1.7	0.6 ± 0.6
	20-30	208.0	383.8	340.1	19.0	269.8	442.7	1448.2	140.6	1025.0	231.8	382.8	85.0
		± 360.3	± 399.9	± 564.5	± 18.3	± 373.0	± 327.5	± 2258.4	± 204.4	± 895.8	± 171.0	± 337.7	± 68.8
	30-40	208.0	10931.1	695.3	224.4	387.7	9717.9	3493.3	963.4	6377.8	7481.7	2711.1	1474.3
		± 360.3	± 5560.6	± 997.4	± 378.9	± 102.6	± 8463.8	± 1988.0	± 1003.0	± 2750.1	± 656.5	± 2199.8	± 1674.2
	40-50	962.1	10743.6	9746.1	691.0	1281.4	9044.8	7146.6	840.8	2707.1	6872.9	2899.5	1442.9
		± 621.5	± 7571.3	± 14674.9	± 637.4	± 252.7	± 4960.9	± 3614.6	± 484.7	± 1143.9	± 4255.0	± 2591.3	1271.6
	50-60	208.0	1980.3	554.1	123.5	96.0	2018.4	702.9	53.4	492.0	872.9	150.1	118.8
	± 208.0	± 1230.6	± 910.4	± 3.3	± 55.2	± 919.2	± 1052.2	± 31.8	± 581.7	± 1014.4	± 230.4	89.1	
	60-70	0.0	110.2	38.0	9.5	110.2	826.5	144.4	3.8	39.9	193.8	26.6	14.1
		± 0.0	± 133.6	± 3.3	± 8.7	± 162.2	± 1180.0	± 86.3	± 6.6	± 64.2	± 243.7	± 46.1	± 19.7
	>70	43.1	44.3	31.2	47.6	20.4	45.3	17.5	24.3	30.2	12.4	12.1	2.5 ± 2.3
		± 23.0	± 4.5	± 16.9	± 28.9	± 26.5	± 13.9	± 28.2	± 23.8	± 13.0	± 7.0	± 10.9	
Body Size ( $\text{ngC cell}^{-1}$ )	<0.1 ( $10^4$ )	12.7 ± 1.9	5.5 ± 2.6	4.4 ± 4.2	16.2 ± 14.2	14.6 ± 8.5	3.0 ± 0.8	1.7 ± 0.9	2.5 ± 2.4	15.4 ± 4.2	1.2 ± 1.0	1.0 ± 1.7	0.6 ± 0.6
	0.1-1	416.0	15097.9	2192.3	399.7	974.8	11454.2	6147.3	1104.0	7988.9	8365.0	3472.7	1678.7
		± 360.3	± 8258.5	± 3322.6	± 561.0	± 611.8	± 8610.1	± 3526.5	± 1200.4	± 1661.2	± 1406.7	± 1756.3	± 1775.2
	1-5	823.5	8784.2	8864.0	576.5	1045.8	8374.2	6565.7	861.7	2524.7	6414.1	2524.5	1323.3
		± 403.3	± 5577.8	± 13389.0	± 467.7	± 54.7	± 4850.9	± 3039.4	± 496.9	± 1272.0	± 3654.4	± 2768.0	± 1199.6
	5-10	138.7	152.9	279.3	72.2	7.6	1299.4	77.9	30.6	84.5	670.6	146.3	107.4
		± 240.7	± 74.3	± 439.3	± 6.6	± 13.2	± 821.6	± 120.4	± 43.6	± 126.7	± 1000.0	± 233.9	± 93.2
	10-50	208.0	68.4	38.0	19.0	116.9	911.0	142.5	1.9	43.7	85.5	0.0	11.6
	± 360.3	± 71.2	± 3.3	± 11.9	± 155.6	± 1306.5	± 89.6	± 3.3	± 70.8	± 70.0	± 0.0	± 20.0	
	50-100	43.1	89.9	31.2	47.6	20.4	51.0	19.4	28.1	30.2	130.2	38.7	16.6
		± 23.0	± 58.4	± 16.9	± 28.9	± 26.5	± 22.4	± 26.7	± 30.2	± 13.0	± 210.9	± 35.7	± 22.0
	>100	0.0 ± 0.0	0.0 ± 0.0	0.0 ± 0.0	0.0 ± 0.0	0.0 ± 0.0	5.7 ± 9.9	0.0 ± 0.0	0.0 ± 0.0	0.0 ± 0.0	0.0 ± 0.0	0.0 ± 0.0	0.0 ± 0.0

Table 3

Explanatory datasets	Mantel statistics	
	r	p
Taxons vs. Body Size	0.890	0.0001
Taxons vs. ESD	0.902	0.0001
Body Size vs. ESD	0.910	0.0001

Table 4.

Environmental conditions	Best parameters	Correlation coefficient r		
		<i>Taxon classification</i>	<i>ESD classification</i>	<i>Body Size classification</i>
Physical	Temperature and NO <sub>3</sub> concentration	0.451	0.436	0.430
Biological	Bacteria and Cyanobacteria biomasses			0.307
	Bacteria, Cyanobacteria and ANF biomasses	0.326	0.308	

For Peer Review




Figure 1  
172x201mm (72 x 72 DPI)






Figure 2  
163x119mm (72 x 72 DPI)




Figure 3  
215x279mm (72 x 72 DPI)




Figure 4  
268x90mm (72 x 72 DPI)




Figure 5  
296x210mm (72 x 72 DPI)

review




Figure 6  
296x210mm (72 x 72 DPI)

review




Figure 7  
173x254mm (72 x 72 DPI)




Figure 8  
192x127mm (72 x 72 DPI)






Figure 9  
289x231mm (300 x 300 DPI)