



**HAL**  
open science

# Community structure and grazing of the nano-microzooplankton on the continental shelf of the Bay of Biscay

Christine Dupuy, Agathe Talarmin, Hans J. Hartmann, Daniel Delmas, C  
Courties, Elise Marquis

## ► To cite this version:

Christine Dupuy, Agathe Talarmin, Hans J. Hartmann, Daniel Delmas, C Courties, et al.. Community structure and grazing of the nano-microzooplankton on the continental shelf of the Bay of Biscay. Estuarine, Coastal and Shelf Science, 2011, 10.1016/j.ecss.2011.05.002 . hal-01248044


**HAL Id: hal-01248044**

**<https://hal.science/hal-01248044>**

Submitted on 26 Dec 2016

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



## Community structure and grazing of the nano-microzooplankton on the continental shelf of the Bay of Biscay

Christine Dupuy<sup>a,\*</sup>, Agathe Talarmin<sup>b,c</sup>, Hans J. Hartmann<sup>a</sup>, Daniel Delmas<sup>d</sup>, C. Courties<sup>e</sup>, Elise Marquis<sup>f</sup>

<sup>a</sup> Laboratoire LIENSs, UMR 6250, Université de La Rochelle, Bâtiment ILE, 2 rue Olympe de Gouges, 17000 La Rochelle, France

<sup>b</sup> Université Aix-Marseille II- CNRS/INSU UMR 6117, LMGE, Centre d'Océanologie de Marseille, Case 901, Campus de Luminy, F-13288 Marseille, France

<sup>c</sup> UPMC Université Paris 06-CNRS, UMR 7621, LOMIC, Observatoire Océanologique, F-66651 Banyuls/mer, France

<sup>d</sup> IFREMER/Centre de Brest, DYNECO, Laboratoire Pelagos, BP 70, 29280 Plouzané, France

<sup>e</sup> Université Paris VI/ CNRS, Laboratoire Arago, BP 44, 66650 Banyuls-sur-mer, France

<sup>f</sup> Institute of Oceanography, National Taiwan University, No. 1, Section 4, Roosevelt, 8 Road, Taipei 10617, Taiwan

### ARTICLE INFO

#### Article history:

Received 16 February 2010

Accepted 2 May 2011

Available online 23 July 2011

#### Keywords:

heterotrophic protists  
size-fractionated phytoplankton  
bacteria  
mesozooplankton  
Bay of Biscay  
environmental factors  
grazing

### ABSTRACT

In order to investigate the parameters controlling the heterotrophic protists (nano-microzooplankton) on the continental shelf of the southern Bay of Biscay, plankton communities and their physico-chemical environment were studied 4 times in February, April, June and September–October 2004 at three stations in the euphotic zone in the Bay of Biscay. The abundance and carbon biomass of heterotrophic protists (ciliates, heterotrophic dinoflagellates and nanoflagellates) as well as all the others groups of plankton (picoplankton, nanophytoplankton, diatoms, autotrophic dinoflagellates, metazoan microzooplankton and mesozooplankton), the environmental parameters and the primary and bacteria production were evaluated at each sampling period. Microzooplankton grazing experiments were undertaken at the same time. Ciliates and heterotrophic dinoflagellates accounted for the main major component of nano- and microzooplankton communities in term of biomass. The total carbon biomass of heterotrophic protists was highest in spring and lowest at the end of summer. The development of heterotrophic protists started after a winter microphytoplankton bloom (principally large diatoms), the biomass was lower in June and was low in September (through inappropriate prey). The carbon requirement of microzooplankton ranged from 50 to more than 100% of daily primary, bacterial and nanoflagellate production. The heterotrophic protist community was predominantly constrained by bottom-up control in spring and at the end of summer via food availability and quality.

© 2011 Elsevier Ltd. All rights reserved.

### 1. Introduction

In pelagic environments, primary production is transferred to higher levels through two main pathways, according to the size of the main primary producers (Azam et al., 1983; Sherr et al., 1986; Sommaruga, 1995; Thingstad and Rassoulzadegan, 1999). Classically, the grazing food chain transfers energy directly from large diatoms to metazoans (herbivorous chain, Pomeroy, 1974). However, most of the primary production is due to phytoplankton less than 3 µm in size (Waterbury et al., 1979; Fenchel, 1988) not efficiently retained by mesozooplankton such as copepods (Sherr et al., 1986) but highly grazed by microzooplankton (Calbet and Landry, 2004; Landry and Calbet, 2004), with energy being transferred through the microbial food web (Azam et al., 1983). Ciliates (phylum Ciliophora) are a dominant component of microzooplankton in marine

waters (Pierce and Turner, 1992). They consume a wide range of prey, such as bacteria (Sherr and Sherr, 1987), pico- and nano-phytoplankton (Bernard and Rassoulzadegan, 1993), heterotrophic nanoflagellates (Jürgens et al., 1996), dinoflagellates (Stoecker et al., 1984), diatoms (Sime-Ngando et al., 1995) and also other ciliates (Dolan, 1991). Large heterotrophic dinoflagellates (>20 µm), another component of the nano-microzooplankton community, have been found to feed on mainly chain-forming diatoms, dinoflagellates, other flagellates and ciliates (Jacobson and Anderson, 1986; Gaines and Elbrachter, 1987; Hansen, 1991a,b; Jeong and Latz, 1994). Heterotrophic nanoflagellates (HNF) are assumed to feed on picoplankton (Fenchel, 1988) and maintain picoplankton populations at relatively stable concentrations in seawater (Pernthaler, 2005). Heterotrophic protists in turn are common prey for many consumers such as copepods (Vincent and Hartmann, 2001) and fish larvae (Fukami et al., 1999). Thus, nano-microzooplankton is recognized as a trophic intermediary in pelagic food webs, permitting the transfer of carbon from pico- and nanoplankton to the metazoans (microbial

\* Corresponding author.

E-mail address: [cdupuy@univ-lr.fr](mailto:cdupuy@univ-lr.fr) (C. Dupuy).

food web, Sherr et al., 1986; Pierce and Turner, 1992). Nano-microzooplankton occupied an important trophic node in food webs and their dynamics may be either controlled by predation or by resource availability as well as hydrography (e.g. Sanders, 1987; Cowlshaw, 2004).

The Bay of Biscay in the North East Atlantic Ocean has temperate climatic conditions where the hydrographic conditions of the water column on the continental shelf are seasonally governed by freshwater runoff from the Loire and the Gironde rivers. These discharges create a longshore flow which may dominate a wide part of the continental shelf at the end of winter (Lazure and Jegou, 1998; Labry et al., 2001). Haline stratification combined with an increase in solar radiation may induce late-winter phytoplankton blooms (Labry et al., 2001). Such blooms have been observed in both the Loire (March 2000, Gohin et al., 2003; Lunven et al., 2005; Guillaud et al., 2008; Maguer et al., 2009) and the Gironde estuarine plumes (March 1998, Labry et al., 2001). They considerably reduce the amount of nutrients available within the surface layer, in particular phosphorus. Phosphate limitation is enhanced by an excess of nitrate in both rivers (Labry et al., 2002; Loyer et al., 2006). As a result, small-sized cells tend to dominate in spring due to their high competitiveness in such an environment (Legendre and Rassoulzadegan, 1995). Consequently, spring represents a change from a diatom-dominated winter community (Herbland et al., 1998) to one of small phytoplankton cells that are mainly grazed by microzooplankton such as ciliates or heterotrophic dinoflagellates (Sautour et al., 2000; Marquis et al., 2007).

Components of the nano- and microzooplankton community were scarcely documented and especially in the Bay of Biscay (Verity, 1987; Kamiyama and Tsujino, 1996; Modigh, 2001; Urrutxurtu et al., 2003). It is important to consider all components of the nano- and microzooplankton when attempting to elucidate their function in planktonic food web processes. To our knowledge, this is the first investigation into all the major components of nano- and microzooplankton in the Bay of Biscay (temperate coastal waters). We hypothesize whether the heterotrophic protists community is controlled by hydrography, or/and resource availability. To answer this question, we investigated the structure and the grazing impact of heterotrophic protists (ciliates, heterotrophic nanoflagellates and heterotrophic dinoflagellates) as well as the environmental factors (abiotic and biotic factors) over the continental shelf of the Bay of Biscay at different months (February, April, June and September 2004) for three contrasting sites (river plume, coastal and offshore areas).

## 2. Material and methods

### 2.1. Study site and hydrography

The continental shelf of the Bay of Biscay (Fig. 1) is up to 200 km wide with a surface area of 223,000 km<sup>2</sup>. The hydrological structure over the shelf is principally influenced by the seasonal dynamics of the Loire and the Gironde river plumes (Lazure and Jegou, 1998). Moreover, the shelf ecology is highly variable in time related to temperate zone climatic fluctuations (Koutsikopoulos et al., 1998). Three stations of the Bay were studied in 2004 on board of R/V Thalia: 08 to 10 February, 23 to 25 April, 09 to 11 June, and 30 September to 02 October. They were located in front of the Gironde River estuary on a coast offshore triangle: “Gironde” (01°30W, 45°30N), “Coast” (01°30W, 45°01N) and “Offshore” (2°20W, 45°10N) (Fig. 1). Salinity and temperature profiles were measured using a CTD (Conductivity–Temperature–Density) probe (Sea-Bird SBE 9). Concurrently, irradiance and fluorescence profiles of the water column were measured in order to obtain the depths of the




Fig. 1. Map of the Bay of Biscay with the location of the three study sites. Dashed line is a schematic representation of the continental shelf limits.

photic zone and of maximum *in situ* fluorescence and to adapt the plankton sampling accordingly.

### 2.2. Plankton sampling and analysis

Water samples from subsurface (2 m), depth of maximum fluorescence and bottom of the photic zone were collected using 12-L Niskin bottles. Samples for dissolved inorganic nutrients [nitrate (NO<sub>3</sub>), nitrite (NO<sub>2</sub>), silicate (Si(OH)<sub>4</sub>) and phosphate (PO<sub>4</sub>)] were filtered immediately through Whatman GF/F filters (47 mm diameter) and the filtrate was stored at –20 °C until analysis with a Skalar autoanalyzer (Strickland and Parsons, 1972). Size-fractionated (<3 μm, 3–20 μm, >20 μm) primary production was assessed by *in situ* incubation of 300 ml seawater with <sup>14</sup>C-bicarbonate (10 μCi) from sunrise to sunset and filtered through different mesh and filter sizes, according to Labry et al. (2002). The incubations were processed on board under *in situ*-like irradiance conditions. Filters were placed in scintillation vials with 200 μl of 1 N HCl, dried overnight at 60 °C then recovered with 4 ml of scintillation cocktail before laboratory counting. Time zero filtration served as control for non-biological <sup>14</sup>C-adsorption onto the filters.

The bacterial production was estimated by the [<sup>3</sup>H] methylthymidine (TdR) incorporation method (Fuhrman and Azam, 1982). Control and triplicates (20 ml) were incubated for 60 min with 20 nM final concentration of [<sup>3</sup>H] TdR at *in situ* temperature. The incubations were stopped by addition of 0.75 ml de TCA à 50% and then stocked at 4 °C. In the laboratory, the precipitation of DNA was made into ice during 30 min. DNA was collected by filtration onto 0.2 μm polycarbonate membrane. Tubes were rinsed twice with cold TCA (5%), and samples were rinsed 4 times with cold TCA (5%), to eliminate all non-incorporation of [<sup>3</sup>H] methylthymidine. Filters

were placed in 5 mL of scintillation fluid and analyzed by scintillation counter (Perkin–Elmer). Rates of  $^3\text{H}$ -thymidine incorporation were transformed to cell production using a conversion factor of  $2.18 \times 10^{10}$  cells produced per mole of thymidine incorporated (Fuhrman and Azam, 1982). Because of on-board technical problems, bacterial and primary productions were not measured in February and in April at the Offshore station at two sampling depths (maximum of fluorescence and bottom of the photic zone).

Bacteria and picophytoplankton (cyanobacteria, picoeukaryotes) were fixed with formaldehyde (final concentration 2%), frozen in liquid  $\text{N}_2$ . Flow cytometry analyses (FCM) were run with a FACScan flow cytometer (BD-Biosciences) equipped with an air-cooled argon laser (488 nm, 15 mW). Picophytoplankton cells were detected according to their right-angle light scatter (SSC; related to cell size and structure) and their orange (FL2, 585/42 nm) and red (FL3 > 670 nm) fluorescence emissions, due to phycoerythrin and chlorophyll pigments, respectively. Beads of 1  $\mu\text{m}$  (Polysciences) were used as an internal standard and simultaneously analyzed with the sample in order to normalize and to compare the cytometric signatures defined as cell fluorescence emission and apparent cell size of each different discriminated cell type. During analysis, accurate volume analyzed (generally between 250 and 500  $\mu\text{l}$ ) and subsequent estimations of cell numbers were calculated by measuring the remaining volume and subtracting it from the initial subsample volume (1 ml). For bacterial counts, 1 mL subsamples were incubated with the nucleic acid stain SYBRGreen I (Molecular Probes) for 15 min at room temperature in the dark (Marie et al., 1997). Total bacterial cells were enumerated measuring green fluorescence of stained bacteria by SYBRGreen I collected at 530/30 nm and their side scatter properties.

Nanoflagellates were fixed with buffered paraformaldehyde (final concentration 1%) then stained with DAPI and counted on 0.8  $\mu\text{m}$  black polycarbonate filters by epifluorescence microscopy (Sherr et al., 1994). Heterotrophic nanoflagellates (HNF) were distinguished from pigmented (autotrophic) nanoflagellates (ANF) by the absence of chlorophyll fluorescence. Microphytoplankton (diatoms and dinoflagellates) was fixed with formaldehyde (final concentration 1%) plus alkaline lugol (final concentration 1%), enumerated and measured by inverse microscopy (Utermöhl, 1958). Heterotrophic and mixotrophic dinoflagellates (HDF) were determined from morphologic species recognition and relevant literature (e.g. Lessard and Swift, 1986). The cell sizes (length and width) were measured on at least 100 cells through a calibrated ocular micrometer. From cell size measurements, the mean cell volume of each taxon was calculated by equating the shape to standard geometric configurations. The cell volume was converted into carbon units (Table 1).

Ciliates were stained with alkaline lugol (final concentration 1%), counted and measured by inverse microscopy. The cell sizes (length and width) were measured for at least 100 cells through

a calibrated ocular micrometer. From cell size measurements, the mean cell volume of each taxon was calculated by equating the shape to standard geometric configurations. The cell volume was converted into carbon units (Table 1). Ciliate samples from the surface and bottom of the photic zone collected in February at the Gironde station were not analyzed due to poor preservation.

Samples of metazoan microplankton were obtained by gently filtering 10 L of collected seawater through a 63  $\mu\text{m}$  mesh. The retained organisms were then diluted in filtered (<63  $\mu\text{m}$ ) seawater and preserved in buffered formaldehyde (final concentration 2%). They were counted under a binocular microscope. The cell sizes (length and width) were measured for at least 100 cells through a calibrated ocular micrometer. From organism size measurements, the mean volume was calculated by equating the shape to standard geometric configurations. The cell volume was converted into carbon units (Table 1).

All the conversion factors and equations used to convert the abundance of pico-, nano- and microplankton into biomass were obtained from the literature (Table 1).

Mesozooplankton was collected from vertical tows through the entire photic zone using a 200  $\mu\text{m}$ -mesh WP2 net, preserved in buffered formaldehyde (final concentration 2%) and counted under a binocular microscope. A second replicate was used to measure the mesozooplankton dry weight. Mesozooplankton carbon biomass was determined by multiplying dry weights of each sample by a factor of 0.38 (Bode et al., 1998).

The planktonic community was classified by size: bacteria, autotrophic cells <3  $\mu\text{m}$  (cyanobacteria and picoeukaryotes), autotrophic cells between 3 and 20  $\mu\text{m}$  (ANF, and small autotrophic ciliates, *Myrionecta rubra*), autotrophic cells >20  $\mu\text{m}$  (diatoms and autotrophic dinoflagellates), heterotrophic cells <20  $\mu\text{m}$  (very small ciliates, HNF), heterotrophic cells between 20 and 50  $\mu\text{m}$  (small ciliates and heterotrophic dinoflagellates), heterotrophic cells between 50 and 100  $\mu\text{m}$  (ciliates and heterotrophic dinoflagellates) and heterotrophic organisms >100  $\mu\text{m}$  (nauplii, mesozooplankton).

### 2.3. Microzooplankton grazing experiments

Microzooplankton grazing impact was estimated using the dilution method (Landry and Hassett, 1982) in February, June and September 2004. Water samples from the « Maximum Fluorescence » depth of each station were collected using a 12-L Niskin bottle. Water was filtered through a 200  $\mu\text{m}$  mesh to isolate microzooplankton fraction from larger predators (e.g. copepods). Organisms were fractionated by gravity filtration and reverse flow through nylon screens (Ferrier-Pagès and Rassoulzadegan, 1994). The filtered water was therefore handled as carefully as possible to minimize production of bubbles and physical damage to fragile microplankton organisms. Particle-free water was obtained by

**Table 1**

Factors and formulas with their reference used to convert biovolume to carbon mass of each plankton organisms.

| Plankton organisms           | Conversion factor or formulas  | References   |
|------------------------------|--|--|
| Bacteria                     | 0.016 pgC cell <sup>-1</sup>   | Labry et al. (2002)  |
| Cyanobacteria                | 0.104 pgC cell <sup>-1</sup>   | Blanchot and Rodier (1996)   |
| Eucaryotic picophytoplankton | 0.22 pgC cell <sup>-1</sup>  | Shinada et al. (2005)  |
| Nanoflagellates              | 0.125 pgC $\mu\text{m}^{-3}$ = 3.14 pgC cell <sup>-1</sup><br>(with mean biovolume of 25.2 $\mu\text{m}^3$ )   | Pelegri et al. (1999) and our data of biovolumes                               |
| Dinoflagellates              | $\text{Log}_{10} \text{C (in pgC cell}^{-1}) = -0.353 + 0.864 \times \text{log}_{10} \text{V}$   | Menden-Deuer and Lessard (2000)  |
| Diatoms                      | $\text{Log}_{10} \text{C (in pgC cell}^{-1}) = -0.541 + 0.811 \times \text{log}_{10} \text{V}$<br>$\text{Log}_{10} \text{C (in pgC cell}^{-1}) = -0.933 + 0.881 \times \text{log}_{10} \text{V}$ | <3000 $\mu\text{m}^3$<br>>3000 $\mu\text{m}^3$ Menden-Deuer and Lessard (2000) |
| Naked Ciliates               | 0.19 pgC $\mu\text{m}^{-3}$  | Putt and Stoecker (1989)   |
| Ciliate Tintinnids           | $\text{C (in pgC cell}^{-1}) = 444.5 + 0.053 \times \text{LV}$   | Verity and Langdon (1984)  |
| Copepod Nauplii              | $\text{pgC ind}^{-3} = 0.08 \times \text{V}$   | Gowing et al. (2003)   |

V: biovolume (in  $\mu\text{m}^3$ ) and LV: Lorica volume (in  $\mu\text{m}^3$ ).




Fig. 2. Temperature (°C) and salinity (PSU) profiles of the water column at the three stations and the four sampling periods.

gently filtering a large volume of collected water, using a peristaltic pump, through a 0.2  $\mu\text{m}$  serial filtration unit. The filtered seawater was diluted with particle-free seawater to obtain the following proportions: 40, 60, 80 and 100% of filtered seawater. The different dilution treatments were distributed into 1.20-L polycarbonate bottles with three replicates for each treatment. All bottles were enriched with inorganic nutrients (100  $\mu\text{M}$   $\text{NaNO}_3$  and 3  $\mu\text{M}$   $\text{NaH}_2\text{PO}_4$ ) to minimize nutrient limitation during the incubations (Verity et al., 1996) (see Fig. 3 for *in situ* nutrient concentrations). The bottles were incubated during 24 h on board in a tank with a circulation of surface seawater to keep the temperature constant and the bottle free to move and covered with a screen to mimic *in situ* light conditions. Samples were collected for each experiment at the beginning ( $t_0$ , undiluted treatment) and at the end ( $t_{24}$ , each treatment) of the incubation to enumerate the pico-, nano- and microplankton composition and estimate the biomass of each group as described above. The prey was classified into five groups: autotrophic cells  $<3 \mu\text{m}$  (cyanobacteria and eukaryotic picophytoplankton), autotrophic cells between 3 and 20  $\mu\text{m}$  (ANF and small autotrophic ciliates, *M. rubra*), autotrophic cells  $>20 \mu\text{m}$  (diatoms and autotrophic dinoflagellates), bacteria (Bac) and heterotrophic nanoflagellates (HNF).

The grazing rates of the microzooplankton were calculated by the exponential growth model of Landry and Hassett (1982). The instantaneous rate coefficients of microzooplankton grazing ( $g$ ) and prey growth ( $k$ ) were estimated from the linear regression (95% of confidence limits) of the apparent growth rate:

$$P_t = P_0 e^{(k-g)t}$$

where  $P_t$  is prey carbon biomass at time  $t$  and  $P_0$  is the initial prey carbon biomass. Both  $g$  and  $k$  were used to calculate the grazing loss of potential production, while only  $g$  was employed to calculate the daily loss of initial standing stock.

#### 2.4. Statistical analysis

Spearman rank ( $r_s$ ) correlations were performed to investigate the relationships between heterotrophic biomasses and environmental parameters. All variables were logarithmically transformed to improve the linearity and homoscedasticity of residuals. The missing data were replaced by means calculated on the entire data set in order to minimize their role in the correlation. The following parameters were considered: salinity, temperature, concentrations of N and P, N/P ratio, and biomasses of bacteria, autotrophic cells  $<3 \mu\text{m}$  (picophytoplankton: cyanobacteria and picoeukaryotes), autotrophic cells between 3 and 20  $\mu\text{m}$  (nanophytoplankton: ANF, and small autotrophic ciliates, *Myrionecta rubra*), autotrophic cells

$>20 \mu\text{m}$  (microphytoplankton: diatoms and autotrophic dinoflagellates), heterotrophic cells  $<20 \mu\text{m}$  (very small ciliates, heterotrophic dinoflagellates, heterotrophic nanoflagellates (HNF)), between 20 and 50  $\mu\text{m}$  (small ciliates and heterotrophic dinoflagellates), heterotrophic cells between 50 and 100  $\mu\text{m}$  (ciliates and heterotrophic dinoflagellates) and heterotrophic organisms  $>100 \mu\text{m}$  (large ciliates and heterotrophic dinoflagellates). All statistical analyses were run using the statistical software Excel Stat Pro.

### 3. Results

#### 3.1. Environmental conditions of abiotic parameters, primary and bacterial productions

In winter (February, Fig. 2), the waters of the Gironde and Coast stations were halostratified with a halocline close to 10 m. The halocline was less developed in April, and was the strongest in June with surface salinities of 31.8 and 32.7 PSU respectively at the Gironde and Coast stations. The surface halostratification was absent in September. Except in June, the Offshore water column did not show significant surface halostratification. The water column temperature of the three stations showed progressive warming from February to September, contributing significantly to stratification in June (thermocline at 10–20 m) and September (thermocline at 30–60 m). Overall, at all three stations, the surface mixed layer was shallowest in February and June, followed by April, and deepest in September.

The concentration of total NO ( $\text{NO}_2$  and  $\text{NO}_3$ ) ranged from 0.2 to 12.1  $\mu\text{mol L}^{-1}$  on average in the photic zone of the three stations (Fig. 3), reaching the highest values in February (e.g. Gironde,  $12.1 \pm 1.6 \mu\text{mol L}^{-1}$ ). The average concentrations of  $\text{Si}(\text{OH})_4$  at the three stations were  $<6 \mu\text{mol L}^{-1}$  whereas the concentrations of  $\text{PO}_4$  were even lower ( $<1 \mu\text{mol L}^{-1}$ , Fig. 3). The lowest concentrations of the three studied nutrients were found in April at the Gironde and Coast stations and in June at the Offshore station. The N/P ratio was highest in April at the two coastal stations (77 in Gironde and 173 in Coast) and lowest in September (below the Redfield ratio, i.e. 10 and 11, respectively). The highest N/P ratio at the Offshore station was found in September but never exceeded 13. The highest N/Si ratios were found in April at the three stations (6 at Gironde and Coast, 156 Offshore).

Depth-integrated primary production varied from  $7.6 \pm 1$  to  $27.5 \pm 5.3 \times 10^2 \text{ mg C m}^{-2} \text{ d}^{-1}$  (Fig. 4a), being lowest in September, and peaking either in June (Coast, Gironde) or April (Offshore). In April, microphytoplankton ( $>20 \mu\text{m}$ ) was an important primary producer at all three stations, largely dominating at the Coast station (73% of the total integrated production, Fig. 4a). On the contrary, picophytoplankton was the main producer in September,





Fig. 3. Seasonal variations of nutrient concentration averages ( $\mu\text{M}$ ) over the photic zone of the three stations: Nitrites and Nitrates (N tot), Phosphates ( $\text{PO}_4$ ) and Silicates (Si).




**Fig. 4.** Seasonal variations of the primary production by size-classes (a) and the bacterial production (b) integrated over the photic zone at the three stations. PP<3 for primary production <3  $\mu\text{m}$ , PP 3–20 for primary production 3–20  $\mu\text{m}$ , PP > 20 for primary production >20  $\mu\text{m}$  and BP for bacterial production.

representing more than 56% of the total production at each station (Fig. 4a). Integrated bacterial production followed the pattern of primary production and was lowest in September at all three stations ( $66.7 \pm 8.3$  to  $84.7 \pm 7.3 \text{ mg C m}^{-2} \text{ d}^{-1}$ , Fig. 4b). Earlier in the year, it was considerably higher, with maxima in April ( $151.6 \pm 6.2 \text{ mg C m}^{-2} \text{ d}^{-1}$  Offshore) and June ( $235.6 \pm 14.8$  and  $206.1 \pm 34.5 \text{ mg C m}^{-2} \text{ d}^{-1}$  respectively at Gironde and Coast).

### 3.2. Variation in plankton abundance and biomass

#### 3.2.1. Autotrophic community abundance and biomass

Integrated phytoplankton biomass was maximal in February at Gironde ( $82.1 \times 10^2 \text{ mg C m}^{-2}$ , Fig. 5a) or in April at the Coast and Offshore (respectively  $33.1$  and  $59.5 \times 10^2 \text{ mg C m}^{-2}$ ), and low in June and September over the entire shelf ( $<5.8 \times 10^2 \text{ mg C m}^{-2}$  in



**Fig. 5.** Seasonal variations of integrated autotrophic biomass (a) and its relative composition (b) over the photic zone at the three stations. Diatoms <20 or >20  $\mu\text{m}$  for diatoms <20  $\mu\text{m}$  or >20  $\mu\text{m}$ , Auto. Dino. for autotrophic dinoflagellates, *M. Rubra* for *Myrionecta rubra*, ANF for autotrophic nanoflagellates, Euk. Picophyto. for eukaryotic picophytoplankton, Cyanobac. for cyanobacteria.

September, Fig. 5a). Large diatoms were the main component of the autotrophic community in February, April and June at the two coastal stations and in February and April at the Offshore station (Fig. 5b). In February, diatoms were dominated by large chain-forming cells (e.g. *Thalassiosira*), illustrated by the high biomass averages ( $371 \mu\text{g C L}^{-1}$ , Table 2) combined with relatively low abundance ( $4.8 \times 10^4 \text{ cells L}^{-1}$ , Table 2). In contrast, smaller diatoms (e.g. *Leptocylindrus minimus*) were very abundant in April at the Coast station (biomass:  $104.3 \mu\text{g C L}^{-1}$ ; abundance:  $50 \times 10^4 \text{ cells L}^{-1}$ , Table 2) and at the Offshore station (biomass:  $152.7 \mu\text{g C L}^{-1}$ ; abundance:  $46 \times 10^4 \text{ cells L}^{-1}$ , Table 2). The biomass of smaller autotrophs (ANF, cyanobacteria and picoeukaryotes) proportionally increased from April to September at all three stations and dominated in September (>60% of integrated biomass, Fig. 5b).

### 3.2.2. Heterotrophic community abundance, biomass and composition

At the three stations, integrated biomasses of heterotrophic plankton peaked in April ( $23.2$  and  $28.2$  and  $15.7 \times 10^2 \text{ mg C m}^{-2}$  at Gironde, Coast and Offshore, respectively, Fig. 6a) and were the lowest in February (less than  $9.2 \times 10^2 \text{ mg C m}^{-2}$ ) except at the Coast station (lowest biomass in September:  $17.2 \times 10^2 \text{ mg C m}^{-2}$ , Fig. 6a). The Offshore biomasses were consistently the lowest of the three stations.

The biomass of bacteria ranged from  $5$  to  $21.5 \mu\text{g C L}^{-1}$  (Table 2). The biomass was highest in June and September at Gironde and Coast. Finally, the concentrations were higher at the Gironde and Coast stations than at the Offshore station. Bacteria dominated in September (>57% of the integrated biomass, Fig. 6b).

Heterotrophic nanoflagellates (HNF) were  $<10 \mu\text{m}$  in size and typically ranged from  $2$  to  $5 \mu\text{m}$ . The biomass of HNF ranged from  $0.02$  to  $0.5 \mu\text{g C L}^{-1}$  (Table 2), and was relatively higher in February and September. The biomass of heterotrophic dinoflagellates (HDF) ranged from  $0.3$  to  $17 \mu\text{g C L}^{-1}$  (Fig. 6b; Table 2). The biomass of HDF was highest in April and June 2004, and showed a small peak in February and September. Finally, the concentrations were higher at the Gironde and Coast stations than at the Offshore station (Fig. 6b). HDF community was mainly composed of athecate dinoflagellates with the most abundant genus *Gymnodinium* spp.

Integrated ciliate biomass was highest in April at all three stations with a maximum of  $15.6 \times 10^2 \text{ mg C m}^{-2}$  at the Coast station. Ciliate abundances, averaging from  $0.2$  to  $9.4 \times 10^3 \text{ cells L}^{-1}$  over the sampling period, and the biomass followed the same pattern (Table 2). Finally, except for February, the concentrations were higher at the Gironde and Coast stations than at the Offshore station (Table 2). The ciliate community was mainly composed of naked ciliates (>82% of the total abundance). The most abundant were the genus *Strombidium*. The highest relative abundances of loricate ciliates (Tintinnina) were found in February at the Gironde and Coast stations (respectively 13 and 18% of the total ciliate abundance).


The biomass of copepod nauplii (microzooplankton metazoan) ranged from  $0.2$  to  $3.8 \mu\text{g C L}^{-1}$  (Table 2). The values were high in February and April, decreased in June, to enhance in September. Finally, the concentrations were higher at the Gironde and Coast stations than at the Offshore station.

The biomass of mesozooplankton varied from  $1.5$  to  $51.5 \mu\text{g C L}^{-1}$  (Table 2). The values were high in February at the Gironde and Coast stations and June at three stations and decreased in September. In fact, close to the coast, mesozooplankton dominated in winter (>65% of the total heterotrophic biomass at Gironde and Coast). In June, it dominated (>47%) at all three stations (Fig. 6b). At the coast, mesozooplankton was mainly composed of appendicularians in winter and of small copepods (e.g. *Oithona* sp.) in June. In Offshore, mesozooplankton was dominated by large copepods (e.g. *Calanus* sp.).

**Table 2** Seasonal abundances (bold font in cells  $\text{L}^{-1}$ ) and biomasses (in  $\mu\text{g C L}^{-1}$ ), with their standard deviations, of the different plankton components in average over the photic zone at the three stations.

| Station  | Month     | Ciliates ( $10^3 \text{ cells L}^{-1}$ ) | Bacteria ( $10^6 \text{ cells L}^{-1}$ ) | Cyanobacteria ( $10^6 \text{ cells L}^{-1}$ ) | Picoeukaryotes ( $10^6 \text{ cells L}^{-1}$ ) | HNF ( $10^4 \text{ cells L}^{-1}$ ) | ANF ( $10^4 \text{ cells L}^{-1}$ )  | Dinoflagellates ( $10^3 \text{ cells L}^{-1}$ ) | Diatoms ( $10^4 \text{ cells L}^{-1}$ ) | Metazoan microplankton (ind $\text{L}^{-1}$ ) | Mesozoo. (ind $\text{L}^{-1}$ )      |
|----------|-----------|--|--|---|--|-------------------------------------|--------------------------------------|---|---|---|--------------------------------------|
| Gironde  | February  | <b>0.18 ± 0.00</b><br>(0.2 ± 0.0)        | <b>6.55 ± 0.19</b><br>(10.5 ± 0.3)       | <b>6.38 ± 0.65</b><br>(0.7 ± 0.1)             | <b>1.48 ± 0.08</b><br>(0.3 ± 0.0)              | <b>12.75 ± 1.56</b><br>(0.4 ± 0.0)  | <b>21.19 ± 5.88</b><br>(0.7 ± 0.2)   | <b>2.01 ± 1.16</b><br>(4.9 ± 3.5)               | <b>4.82 ± 2.85</b><br>(370.9 ± 378.2)   | <b>66.12 ± 15.58</b><br>(3.4 ± 1.5)           | <b>4.55</b><br>(25.7)                |
|          | April     | <b>4.67 ± 1.47</b><br>(12.1 ± 7.2)       | <b>5.17 ± 0.92</b><br>(8.3 ± 1.5)        | <b>1.68 ± 0.18</b><br>(0.2 ± 0.0)             | <b>4.72 ± 5.73</b><br>(1.0 ± 1.3)              | <b>5.48 ± 2.12</b><br>(0.2 ± 0.1)   | <b>19.26 ± 2.43</b><br>(0.6 ± 0.1)   | <b>21.56 ± 10.56</b><br>(21.6 ± 4.6)            | <b>16.91 ± 1.59</b><br>(72.7 ± 48.6)    | <b>81.52 ± 13.11</b><br>(3.5 ± 0.3)           | <b>6.61</b><br>(33.6)                |
|          | June      | <b>3.83 ± 3.00</b><br>(10.9 ± 7.9)       | <b>11.75 ± 3.63</b><br>(18.8 ± 5.8)      | <b>16.06 ± 14.25</b><br>(1.7 ± 1.5)           | <b>3.53 ± 2.98</b><br>(0.3 ± 0.1)              | <b>2.66 ± 3.41</b><br>(0.1 ± 0.4)   | <b>57.64 ± 35.57</b><br>(1.8 ± 1.1)  | <b>10.43 ± 8.80</b><br>(16.4 ± 12.9)            | <b>17.27 ± 17.96</b><br>(0.45 ± 0.36)   | <b>52.37 ± 23.07</b><br>(2.5 ± 1.1)           | <b>29.88</b><br>(41.8)               |
|          | September | <b>0.64 ± 0.51</b><br>(1.8 ± 0.8)        | <b>13.42 ± 2.13</b><br>(21.5 ± 3.4)      | <b>74.34 ± 23.17</b><br>(7.7 ± 2.4)           | <b>2.59 ± 1.20</b><br>(0.3 ± 0.0)              | <b>16.24 ± 11.64</b><br>(0.5 ± 0.4) | <b>59.75 ± 32.65</b><br>(0.9 ± 0.4)  | <b>0.45 ± 0.36</b><br>(0.3 ± 0.2)               | <b>0.45 ± 0.36</b><br>(0.3 ± 0.2)       | <b>1.12 ± 0.75</b><br>(1.9 ± 1.4)             | <b>115.02 ± 57.10</b><br>(3.8 ± 1.9) |
| Coast    | February  | <b>1.80 ± 0.87</b><br>(5.4 ± 5.0)        | <b>6.65 ± 0.90</b><br>(10.6 ± 1.4)       | <b>5.00 ± 1.80</b><br>(0.5 ± 0.2)             | <b>0.90 ± 0.38</b><br>(0.2 ± 0.1)              | <b>14.64 ± 6.98</b><br>(0.5 ± 0.4)  | <b>25.42 ± 11.60</b><br>(0.8 ± 0.4)  | <b>0.52 ± 0.31</b><br>(0.7 ± 0.7)               | <b>2.58 ± 0.81</b><br>(33.3 ± 11.8)     | <b>38.63 ± 44.08</b><br>(1.6 ± 1.7)           | <b>4.00</b><br>(40.8)                |
|          | April     | <b>9.44 ± 5.48</b><br>(42.8 ± 37.5)      | <b>3.12 ± 0.58</b><br>(5.0 ± 0.9)        | <b>1.91 ± 0.51</b><br>(0.3 ± 0.1)             | <b>1.82 ± 1.66</b><br>(0.4 ± 0.4)              | <b>3.01 ± 0.69</b><br>(0.1 ± 0.0)   | <b>12.21 ± 1.99</b><br>(0.4 ± 0.1)   | <b>15.25 ± 8.15</b><br>(16.6 ± 9.7)             | <b>49.90 ± 4.21</b><br>(104.3 ± 19.5)   | <b>119.35 ± 57.93</b><br>(3.2 ± 0.4)          | <b>3.94</b><br>(29.0)                |
|          | June      | <b>4.26 ± 3.28</b><br>(9.5 ± 4.4)        | <b>11.40 ± 3.51</b><br>(18.2 ± 5.6)      | <b>12.55 ± 4.47</b><br>(1.3 ± 0.5)            | <b>2.31 ± 1.76</b><br>(1.2 ± 0.5)              | <b>1.55 ± 0.74</b><br>(0.0 ± 0.0)   | <b>125.17 ± 66.94</b><br>(3.9 ± 2.5) | <b>10.52 ± 5.96</b><br>(9.3 ± 8.1)              | <b>1.99 ± 1.10</b><br>(13.1 ± 12.7)     | <b>25.94 ± 32.73</b><br>(1.4 ± 1.8)           | <b>14.17</b><br>(51.5)               |
|          | September | <b>0.51 ± 0.55</b><br>(1.0 ± 1.1)        | <b>9.23 ± 2.76</b><br>(14.8 ± 4.4)       | <b>41.59 ± 43.94</b><br>(4.3 ± 4.6)           | <b>1.01 ± 0.09</b><br>(0.2 ± 0.1)              | <b>2.48 ± 1.98</b><br>(0.1 ± 0.1)   | <b>59.41 ± 12.19</b><br>(1.9 ± 0.4)  | <b>1.57 ± 0.85</b><br>(1.5 ± 1.2)               | <b>0.95 ± 0.59</b><br>(5.0 ± 2.9)       | <b>0.95 ± 0.59</b><br>(1.87 ± 1.09)           | <b>46.12 ± 15.77</b><br>(1.9 ± 1.5)  |
| Offshore | February  | <b>2.60 ± 1.39</b><br>(3.5 ± 2.4)        | <b>5.57 ± 0.37</b><br>(8.9 ± 0.6)        | <b>14.07 ± 4.43</b><br>(1.5 ± 0.5)            | <b>6.20 ± 1.41</b><br>(1.4 ± 0.3)              | <b>15.36 ± 3.43</b><br>(0.5 ± 0.1)  | <b>27.90 ± 1.72</b><br>(0.9 ± 0.1)   | <b>8.53 ± 1.78</b><br>(4.5 ± 2.0)               | <b>1.87 ± 1.09</b><br>(11.7 ± 6.3)      | <b>40.08 ± 12.92</b><br>(2.4 ± 0.8)           | <b>0.30</b><br>(1.5)                 |
|          | April     | <b>4.30 ± 1.74</b><br>(19.2 ± 14.0)      | <b>4.90 ± 0.93</b><br>(7.8 ± 1.5)        | <b>0.24 ± 0.10</b><br>(0.0 ± 0.0)             | <b>0.19 ± 0.11</b><br>(0.0 ± 0.0)              | <b>1.28 ± 0.86</b><br>(0.0 ± 0.0)   | <b>7.34 ± 1.94</b><br>(0.2 ± 0.1)    | <b>11.37 ± 3.72</b><br>(9.8 ± 5.5)              | <b>46.05 ± 13.57</b><br>(152.7 ± 34.2)  | <b>44.91 ± 25.38</b><br>(1.0 ± 0.4)           | <b>1.69</b><br>(5.1)                 |
|          | June      | <b>0.47 ± 0.10</b><br>(2.0 ± 2.2)        | <b>5.43 ± 2.41</b><br>(8.7 ± 3.9)        | <b>13.50 ± 10.27</b><br>(1.4 ± 1.1)           | <b>6.67 ± 6.44</b><br>(1.8 ± 0.1)              | <b>1.68 ± 1.15</b><br>(0.1 ± 0.0)   | <b>86.93 ± 3.55</b><br>(2.7 ± 0.1)   | <b>6.45 ± 3.57</b><br>(7.7 ± 4.5)               | <b>1.08 ± 0.94</b><br>(3.7 ± 3.6)       | <b>19.75 ± 14.55</b><br>(1.0 ± 0.7)           | <b>5.81</b><br>(32.9)                |
|          | September | <b>0.49 ± 0.32</b><br>(1.6 ± 1.7)        | <b>6.55 ± 2.83</b><br>(10.5 ± 4.5)       | <b>45.19 ± 25.22</b><br>(4.7 ± 2.6)           | <b>0.68 ± 0.54</b><br>(1.4 ± 0.3)              | <b>0.63 ± 0.49</b><br>(0.0 ± 0.0)   | <b>40.77 ± 17.75</b><br>(1.0 ± 0.5)  | <b>2.67 ± 2.35</b><br>(1.7 ± 1.3)               | <b>0.21 ± 0.25</b><br>(0.9 ± 1.2)       | <b>3.43 ± 2.82</b><br>(0.2 ± 0.1)             | <b>0.97</b><br>(4.9)                 |





**Fig. 6.** Seasonal variations of integrated heterotrophic biomass (a) and its relative composition (b) over the photic zone at the three stations. Mesozoo. for mesozooplankton, HNF for heterotrophic nanoflagellates, Hetero Dino. for heterotrophic dinoflagellates.

Ciliates and heterotrophic dinoflagellates dominated the integrated heterotrophic protist biomass in April particularly at the Coast and Offshore stations (Fig. 6b) with high percentage of ciliates. Heterotrophic nanoflagellates were consistently present, but never dominant (<2% of integrated biomasses, Fig. 6b).


### 3.2.3. Ratio heterotrophy/autotrophy

The heterotrophy/autotrophy ratio was calculated using the biomass data of every group (Fig. 7). In February, the heterotrophy/autotrophy ratio was close to 1 at Coast and Offshore sites which corresponded to an equilibrium between biomass of heterotrophic and autotrophic organisms. The ratio was very low at Gironde site, corresponding to the high bloom of diatoms in post-winter. In April, the ratio was lower at Coast and Offshore (0.5 and 0.3 respectively), corresponding to the large development of diatoms. In Gironde, the ratio was superior to 1 (1.5), in relation to a diminution of the biomass of diatom and a presence of ciliates and HDF. In June and September, the ratio was largely >1 at every site (higher in Gironde, 6.5 to 10), due to the bloom of heterotrophic organisms (bacteria, microzooplankton and mesozooplankton) and the bloom of small autotrophic cells. The HNF were abundant but the biomass was so low that it did not impact the heterotrophy/autotrophy ratio.

### 3.3. Correlation relationships between heterotrophic nano-microplankton, and size-fractionated phytoplankton and environmental factors

Correlations indicate potential relationships between organisms and give the direction of relationships. For the entire data set (all depths, stations and sampling periods combined), temperature showed no correlation with any heterotrophic nano-microplankton group (Table 3). Salinity was negatively correlated with HDF. NO<sub>3</sub> concentration was correlated with HNF biomass. PO<sub>4</sub> concentration was positively correlated with HNF and negatively correlated with ciliate biomass. Ciliates were significantly correlated with N/P ratio, microphytoplankton production and biomass and nanophytoplankton

biomass (Table 3). HDF were correlated to microphytoplankton production and bacterial production, and negatively correlated to picophytoplankton biomass. The correlations were also calculated with size-class of heterotrophic protists (HNF, HDF and ciliates). All the size-classes of heterotrophic protists were correlated with >20 μm primary production. The <20 μm size-fraction of heterotrophic protists was positively correlated with nitrites, N/P ratio and microphytoplankton biomass and negatively correlated with PO<sub>4</sub> concentration. The 20–50 μm size-fraction of heterotrophic protists, dominant in this study, was positively correlated with N/P ratio and bacterial production and negatively correlated with salinity, PO<sub>4</sub> and picophytoplankton biomass. The 50–100 μm size-fraction was correlated by all size of primary production and bacterial production and negatively correlated with salinity. The >100 μm size-fraction was correlated with N/P ratio and microphytoplankton biomass.



**Fig. 7.** Heterotrophic/autotrophic biomass ratio over the photic zone at the three stations.

**Table 3**

Simple (rs, Spearman rank) correlation coefficients between heterotrophic protists biomasses and environmental parameters (physical and chemical conditions and phytoplankton and bacteria biomass and production).

| A (Community structure)    |                        |          |           |         |
|----------------------------|------------------------|----------|-----------|---------|
|                            | HNF                    | Ciliates | HDF       |         |
| Temp                       |                        |          |           |         |
| Sal                        |                        |          | –0.555**  |         |
| Nitrites                   |                        |          |           |         |
| Nitrates                   | 0.466*                 |          |           |         |
| Phosphates                 | 0.447*                 | –0.417*  |           |         |
| Nitrogen/phosphorus        |                        | 0.562**  |           |         |
| Microphyto                 |                        | 0.604**  |           |         |
| Nanophyto                  |                        | 0.449*   |           |         |
| Picophyto                  |                        |          | –0.423*   |         |
| Bacteria                   | 0.426*                 |          |           |         |
| <3 µm primary production   |                        |          |           |         |
| 3–20 µm primary production |                        |          |           |         |
| >20 µm primary production  |                        | 0.684*** | 0.582**   |         |
| Bacterial production       |                        |          | 0.721***  |         |
| B (size structure)         |                        |          |           |         |
|                            | Heterotrophic protists |          |           |         |
|                            | <20 µm                 | 20–50 µm | 50–100 µm | >100 µm |
| Temp                       |                        |          |           |         |
| Sal                        |                        | –0.588*  | –0.764*** |         |
| Nitrites                   | 0.484*                 |          |           |         |
| Nitrates                   |                        |          |           |         |
| Phosphates                 | –0.474*                | –0.428*  |           |         |
| Nitrogen/Phosphorus        | 0.776***               | 0.512*   |           | 0.506*  |
| Microphyto                 | 0.605**                |          |           | 0.542** |
| Nanophyto                  |                        |          |           |         |
| Picophyto                  |                        | –0.465*  |           |         |
| Bacteria                   |                        |          |           |         |
| <3 µm primary production   |                        |          | 0.482*    |         |
| 3–20 µm primary production |                        |          | 0.421*    |         |
| >20 µm primary production  | 0.452*                 | 0.694*** | 0.628**   | 0.537** |
| Bacterial production       |                        | 0.702*** | 0.907***  |         |

Significant correlations were defined as \*\*\* $p < 0.001$ , \*\* $p < 0.01$ , \* $p < 0.05$ . Temp.: temperature; Sal: salinity; Microphyto: microphytoplankton; Nanophyto: nanophytoplankton; Picophyto: picophytoplankton.

### 3.4. Grazing experiment

Phytoplankton growth and grazing coefficients (Table 4) show that the relationship between the apparent growth rate and the dilution factor was linear ( $p < 0.05$ ). Phytoplankton growth rate ranged from 0.12 to 4.58  $d^{-1}$ , while microzooplankton grazing ranged from 0.10 to 3.25  $d^{-1}$ . This level of grazing corresponds to a daily loss varying between 50 and more than 100% of the daily potential primary production.

Bacteria and heterotrophic nanoflagellates growth and grazing coefficients are shown in Table 5. Bacteria growth rate ranged from 0.3 to 1.96  $d^{-1}$ , while microzooplankton grazing ranged from 0.73 to 1.98  $d^{-1}$ . This level of grazing corresponds to a daily loss varying between 62 and >100% of the daily potential bacteria production. Heterotrophic nanoflagellates growth rate ranged from 1.06 to 3.72  $d^{-1}$ , while microzooplankton grazing ranged from 0.62 to 4.95  $d^{-1}$ . This level of grazing corresponds to a daily loss varying between 48 and more than 100% of the daily potential heterotrophic nanoflagellates production.

## 4. Discussion

During our study period (i.e. 2004), the hydrography of the continental shelf of the Southern Bay of Biscay followed seasonal variations that appeared to be largely influenced by the Gironde plume, in a pattern similar to that described by Lazure and Jegou (1998). The seasonal runoff variation influenced both density

stratification and photic zone depth (through turbidity). Our data showed that, across the shelf, the photic zone deepened as the runoff signal decreased, between February and April (Figs. 2 and 8), and then was shallower again in June after the increasing runoff in May (Fig. 8), while finally showing typical offshore summer conditions (deep photic zone with deep thermal stratification) in September. Furthermore, the thermohaline profiles suggest that the photic zone depth exceeded the depth of the surface pycnoclines throughout the study period at all three stations.

### 4.1. Heterotrophic protists abundance, biomass and composition in the Bay of Biscay

The variations of the heterotrophy/autotrophy ratio were similar to typical patterns found in temperate ecosystems: a low ratio in winter and spring corresponding to the bloom of primary producers, and a high ratio in June corresponding to the bloom of grazers, such as mesozooplankton (Fig. 6) (Legendre and Rassoulzadegan, 1995; Lunven et al., 2005; Maguer et al., 2009).

Heterotrophic protist abundance and biomass showed strong variations at the three stations with highest values in mid-spring (April) and lowest in late summer (September, Fig. 6). This pattern is comparable to that observed in other temperate areas (e.g. Nielsen and Kiorboe, 1994; Johansson et al., 2004).

The range of measured ciliate abundances and biomass is similar to ciliate abundance commonly found in coastal and oceanic ecosystems (e.g. Montagnes et al., 1988; Dolan, 1991; Leakey et al., 1992; Chiang et al., 2003; Johansson et al., 2004) and encompasses concentrations observed previously in the Bay of Biscay (Vincent and Hartmann, 2001). The range of measured HDF abundances and biomass is a little higher than HDF abundance found in Northern Arabian Sea (Garisson et al., 1998). The range of measured HNF abundances and biomass is 10x higher all the year

**Table 4**

Summary of seasonal dilution experiment data for microphytoplankton (Micro, diatoms and autotrophic dinoflagellates), nanophytoplankton (Nano, autotrophic nanoflagellates and *Myrionecta rubra*) and picophytoplankton (Pico, eukaryotic picophytoplankton and cyanobacteria). The data include growth (Growth coef.), grazing rates (Grazing coef.), correlation coefficient ( $R^2$ ), percentage of primary production consumed per day (PP consumed).

|           |          |       | Growth coef ( $d^{-1}$ ) | Grazing coef ( $d^{-1}$ ) | $R^2$  | PP consumed (%) |
|-----------|----------|-------|--------------------------|---------------------------|--------|-----------------|
| February  | Gironde  | Micro | –                        | –                         | 0.58*  | –               |
|           |          | Nano  | 1.16                     | 1.23                      | 0.42*  | 105.4           |
|           |          | Pico  | 0.18                     | 0.13                      | 0.73*  | 74.6            |
|           | Coast    | Micro | 1.07                     | 0.97                      | 0.69*  | 90.4            |
|           |          | Nano  | 1.65                     | 1.63                      | 0.64*  | 99              |
|           |          | Pico  | 0.15                     | 0.11                      | 0.73*  | 69.1            |
|           | Offshore | Micro | 1.47                     | 1.49                      | 0.51*  | 101.2           |
|           |          | Nano  | –                        | –                         | 0; 47* | –               |
|           |          | Pico  | –                        | –                         | 0.45*  | –               |
| June      | Gironde  | Micro | 0.77                     | 0.82                      | 0.37   | 106.4           |
|           |          | Nano  | 1.26                     | 2.36                      | 0.91** | 187             |
|           |          | Pico  | 1.11                     | 1.58                      | 0.79** | 141.9           |
|           | Coast    | Micro | 1.77                     | 0.88                      | 0.45*  | 49.6            |
|           |          | Nano  | –                        | 1.18                      | 0.58*  | –               |
|           |          | Pico  | –                        | –                         | 0.38*  | –               |
|           | Offshore | Micro | 4.58                     | 3.25                      | 0.48*  | 70.9            |
|           |          | Nano  | –                        | –                         | 0.68*  | –               |
|           |          | Pico  | –                        | –                         | 0.5*   | –               |
| September | Gironde  | Micro | 2.97                     | 2.04                      | 0.45*  | 68.7            |
|           |          | Nano  | 1.09                     | 1.47                      | 0.65*  | 135.8           |
|           |          | Pico  | 0.23                     | 0.47                      | 0.75** | 204.6           |
|           | Coast    | Micro | –                        | –                         | 0.08   | –               |
|           |          | Nano  | 1.26                     | 1.12                      | 0.48*  | 89.4            |
|           |          | Pico  | –                        | –                         | 0.52*  | –               |
|           | Offshore | Micro | 1.97                     | 2.15                      | 0.32*  | 109.3           |
|           |          | Nano  | 0.51                     | 1.66                      | 0.33*  | 328.2           |
|           |          | Pico  | 0.12                     | 0.10                      | 0.62*  | 90.8            |

**Table 5**

Summary of seasonal dilution experiment data for bacteria (Bac) and heterotrophic nanoflagellates (HNF). The data include growth (Growth coef.), grazing rates (Grazing coef.), correlation coefficient ( $R^2$ ), percentage of production consumed per day.

|           |          | Growth coef. ( $d^{-1}$ ) | Grazing coef. ( $d^{-1}$ ) | $R^2$ | Production consumed (%) |       |
|-----------|----------|---------------------------|----------------------------|-------|-------------------------|-------|
| February  | Gironde  | Bac                       | 1.41                       | 1.3   | 0.73*                   | 92.6  |
|           |          | HNF                       | –                          | –     | 0.31                    | –     |
|           | Coast    | Bac                       | 0.97                       | 1.05  | 0.41*                   | 108.3 |
|           |          | HNF                       | 1.06                       | 2.07  | 0.69*                   | 196.1 |
|           | Offshore | Bac                       | 1.96                       | 1.33  | 0.46*                   | 67.6  |
|           |          | HNF                       | 2.97                       | 3.29  | 0.7*                    | 111   |
| June      | Gironde  | Bac                       | 0.87                       | –     | 0                       | –     |
|           |          | HNF                       | 3.72                       | 4.95  | 0.56*                   | 133.1 |
|           | Coast    | Bac                       | 0.81                       | 0.73  | 0.29                    | 90    |
|           |          | HNF                       | 2.41                       | 1.15  | 0.27                    | 47.9  |
|           | Offshore | Bac                       | 1.78                       | 1.11  | 0.35*                   | 62.5  |
|           |          | HNF                       | 1.28                       | 0.62  | 0.13                    | 48.4  |
| September | Gironde  | Bac                       | 0.3                        | –     | 0.42*                   | –     |
|           |          | HNF                       | 1.49                       | 1.5   | 0.84**                  | 101   |
|           | Coast    | Bac                       | 1.45                       | 1.98  | 0.87**                  | 137.2 |
|           |          | HNF                       | 1.52                       | 1.25  | 0.67*                   | 81.9  |
|           | Offshore | Bac                       | 0.77                       | –     | 0.34                    | –     |
|           |          | HNF                       | 3.64                       | 3.72  | 0; 91**                 | 102.3 |

The significance of the regression is showed with \*\* for  $p < 0.001$ , \* for  $p < 0.05$  and italic font for non-significant correlation. – stands negative values.

for the zones of Bay of Biscay compared to data of 6 oceans (Massana et al., 2006).

The composition of heterotrophic protists showed that naked oligotrichs dominated the ciliate abundances at the three stations over the sampling periods, as found in open waters (Leakey et al., 1992; Nielsen and Kiorboe, 1994; Fileman and Leakey, 2005). Tintinnids were not numerous in the Bay of Biscay ( $0\text{--}318\text{ cells L}^{-1}$ ) and are more concentrated in estuaries than in open waters (e.g. Urrutxurtu et al., 2003). HDF were dominated by the genus *Cyrodinium* which are common in temperate coastal waters (Yang et al., 2008).

#### 4.2. Heterotrophic protists community in relation to environmental parameters

In many previous studies, temperature, food availability and predator density seemed responsible in structuring the heterotrophic protist distribution, abundance and composition (e.g. Sanders, 1987; Nielsen and Kiorboe, 1994; Urrutxurtu et al., 2003). This leads to the question of which parameters are the most likely to structure the community of heterotrophic protists in the Bay of Biscay. Although the correlations do not necessary mean directly causal

relationships such as trophic links, they can give information on the biological parameters that influenced the heterotrophic protists. In addition, the microzooplankton grazing experiments, using the dilution method, allow determination of the grazing pressure on different size-classes of prey.

##### 4.2.1. Abiotic factors influence

Several authors have identified temperature as a key parameter influencing the rate of numerical response of heterotrophic protists to increases in phytoplankton due to temperature-enhancement growth and feeding (e.g. Sherr et al., 1988; Montagnes, 1996). However, in this study, we did not find significant correlations between heterotrophic protist biomass and temperature (Table 3). In June at Offshore, the water column was thermostratified (Fig. 2) which prevented subsurface-water nutrient injection and thus prohibited the development of resources for heterotrophic protists (low primary and bacterial productions). However, one may consider that the increase of temperature during spring (due to increasing thermostratification between April to June, Fig. 2) may have facilitated a rapid assimilation of primary production by heterotrophic protists especially ciliates (James and Hall, 1995). However, as shown previously, freshwater runoff, and thus salinity, is a major factor structuring the depth and development of the pyknocline over the shelf (Gironde and Coast). Sanders (1987) described a negative correlation with salinity and microzooplankton that may illustrate the higher abundances in waters where river runoff is significant. Our study showed that this is the case for HDF and 20–100  $\mu\text{m}$  size-class heterotrophic protists. However, ciliate biomasses and 2–50  $\mu\text{m}$  and >100  $\mu\text{m}$  size-class heterotrophic protists were not correlated to salinity but were negatively and significantly correlated to  $\text{PO}_4$  concentrations or positively correlated to N/P ratio, conditions of river runoff (Table 3). In this part of the Bay of Biscay (coast and continental shelf), Labry et al. (2002) showed that phosphate limitation is provoked by the development of a winter diatom bloom in the Gironde plume. Therefore, lower concentrations of phosphate or a high N/P ratio in winter-spring could indicate a recent development of microphytoplankton (mainly diatoms), as was observed in our study through high diatom biomasses in winter in Gironde (Figs. 3 and 5). In addition, ciliate biomass was correlated the nanomicrophytoplankton (Table 3). Our results thus suggest that the conditioning of the shelf water mass dynamics through freshwater runoff influenced the development of heterotrophic protist biomass together with other biological variables, with temperature being a secondary, non-significant factor.




Fig. 8. Gironde river discharge ( $\text{m}^3\text{ s}^{-1}$ ) over the year 2004. Red square: sampling date.

#### 4.2.2. Biological influences

**4.2.2.1. Resource availability.** Heterotrophic protists consume a wide range of prey items from bacteria (Sherr and Sherr, 1987), to nanophytoplankton (e.g. Burkill, 1982; Dolan and Coats, 1990) and diatoms (Sime-Ngando et al., 1995). In our study, the number of biomass doublings per day was lower than 3 in most of the cases for phytoplankton, bacteria and HNF. The very elevated numbers of biomass doublings per day found for the microphytoplankton in Offshore in June and in Gironde in September (6.6 and 4.3) were due to a very low initial biomass and high growth rates. Those situations may be more due to a bottle effect (e.g. addition of nutrients) than to *in situ* reality. In fact, phytoplankton growth is around  $0.7 \text{ d}^{-1}$  in temperate habitats, and higher ( $0.72 \text{ d}^{-1}$ ) in tropical/subtropical habitats (Calbet and Landry, 2004) but always lower in comparison to our data.

In our study, heterotrophic protists started to develop immediately after or during the peak of the microphytoplankton bloom, i.e. in April, when a mix of small and large sized-food began to develop (positive correlation with ciliates and nanomicrophytoplankton biomass, Table 3). The dilution experiment confirmed that in general there was a high predation pressure on nanophytoplankton and a moderate one on microphytoplankton (Table 4). In addition, the entire set of heterotrophic protist (except HNF) biomasses were positively correlated to microphytoplankton production. Verity (1987) and Lynn and Montagnes (1991) argued that the distribution of ciliates generally shows a close association with nanophytoplankton in temperate coastal ecosystems. Such a pattern was also described in the Northwestern Indian Ocean (Leakey et al., 1992), in Lake Erie (Hwang and Heath, 1997) and in Gyeonggi Bay, Yellow Sea (Yang et al., 2008).

In late summer (September), when the water column was deeply stratified over the shelf and primary and bacterial productions were minimal, heterotrophic protist biomasses and abundances were also the lowest. Despite the availability of picoplankton, especially cyanobacteria at this time (i.e. peak of cyanobacteria biomass and abundance, Fig. 5b), the picoplankton was little grazed, confirmed by dilution experiment (Table 4), the absence of correlation with picoplankton biomass and production, and a negative correlation with HDF and major fraction 20–50  $\mu\text{m}$  of heterotrophic protists (Table 3). We argue that this potential prey was not sufficiently grazed to maintain a high heterotrophic protist production in late summer on the shelf. In winter, heterotrophic protist biomass may have been low due too to the lack of suitable food (biomass dominance of large chain-forming diatoms and insufficient HDF to graze the bloom, Table 2). In many previous studies, heterotrophic protist abundance has been described as a function of food availability (e.g. Dolan and Coats, 1990).

Heterotrophic bacteria appear to be potentially important prey for heterotrophic protists. HNF are generally considered to be the main consumers of bacteria (e.g. Hagström et al., 1988; Shinada et al., 2003). Our data indicate this trophic link by a significant correlation between HNF and bacterial biomasses. However, ciliates may also be able to consume bacteria in high quantity (e.g. Sherr and Sherr, 1987). However, in Chesapeake Bay, the microphagous ciliate abundance (mostly bacterivorous species) was not related to bacterial biomass despite an obvious trophic link (Dolan, 1991). In such a situation, the heterotrophic protist biomass would be more related to bacterial productivity than to biomass. This is the case in our study: HDF and 20–100  $\mu\text{m}$  size-class heterotrophic protists were correlated with bacterial production (Table 3). The observed correlation between heterotrophic protists and bacterial production may then reveal a double link: first, a direct feeding link (driven by bacterivorous ciliate species) and secondly by a trophic link between ciliates and HNF. These conclusions were confirmed by our grazing experiment. The predation of

microzooplankton on HNF led to lower grazing rates on bacteria in the dilution experiments with microzooplankton. However, the growth rates of bacteria were always superior to the grazing rates (Table 5, Fonda Umani and Beran, 2003). That situation corresponded to a lower relationship between grazing activity of microzooplankton and growth of bacteria and heterotrophic nanoflagellate populations. These results may demonstrate a trophic cascade operating in the incubation bottles, with the grazing of bacteria by HNF.

**4.2.2.2. Predation.** Predation is an important potential structuring factor of the heterotrophic protists (Sanders, 1987). Although we did not study the predation pressure of metazoans on heterotrophic protists, we showed that in winter (February) along the Bay's inner shelf (Gironde plume and Coast), mesozooplankton density was relatively important (maxima up to  $66 \text{ ind L}^{-1}$ ) and ingestion rates ranged between 0.22 and  $0.33 \mu\text{g C}$  of ciliates  $\text{ind}^{-1} \text{ d}^{-1}$  (Vincent and Hartmann, 2001). The consumption of heterotrophic protists by mesozooplankton on heterotrophic protists can be high. In June, when heterotrophic protists biomass declined, mesozooplankton density peaked (maxima up to  $100 \text{ ind L}^{-1}$ ). As a consequence, we can predict the predation rates on heterotrophic protists could be high and greater than the winter values (Hartmann and Nejstgaard, pers. comm.) (Table 2). By contrast, in April, resource availability may have been the dominant structuring factor for the heterotrophic protists, since their biomass peaked while mesozooplankton biomass was less important (Table 2).

## 5. Conclusions

Overall, heterotrophic protists community did not differ from the typical pattern found in temperate areas where Gironde runoff influence is the highest (Gironde station) until the limits of the continental shelf. Among the three stations studied, heterotrophic protist followed a succession with highest biomass in spring, and lowest biomass under strong summer conditions. Heterotrophic protist community was predominantly constrained by bottom-up control in spring and in the end of summer (food availability and quality).

## Acknowledgments

This study was supported by the French "Programme National Environment Côtier" – Bay of Biscay working site – and Ifremer. The authors wish to thank the captains and crews of R/V Thalia for support activities, F. Mornet (IFREMER, UMS ELA, France) and V. Huet (LIENSs, France) for nutrient and chlorophyll measurements, P. Malterre and M-J Capdeville (students of University of La Rochelle, France) for metazoan counts.

## References

- Azam, F., Fenchel, T., Field, J.G., Gray, J.S., Meyer-Reil, L.A., Thingstad, F., 1983. The ecological role of water-column microbes in the sea. *Marine Ecology Progress Series* 10, 257–263.
- Bernard, C., Rassoulzadegan, F., 1993. The role of picoplankton (cyanobacteria and plastidic picoflagellates) in the diet of tintinnids. *Journal of Plankton Research* 15, 361–373.
- Blanchot, J., Rodier, M., 1996. Picophytoplankton abundance and biomass in the western tropical Pacific Ocean during the 1992 El Niño year: results from flow cytometry. *Deep Sea Research Part I* 43, 877–895.
- Bode, A., Alvarez-Ossorio, M., Gonzalez, N., 1998. Estimation of mesozooplankton biomass in a coastal upwelling area off NW Spain. *Journal of Plankton Research* 20, 1005–1014.
- Burkill, P.H., 1982. Ciliates and other microplankton components of a nearshore food-web: standing stocks and production processes. *Annales de l'Institut Océanographique* 58, 335–350.

- Calbet, A., Landry, M.R., 2004. Phytoplankton growth, microzooplankton grazing, and carbon cycling in marine systems. *Limnology and Oceanography* 49, 51–57.
- Chiang, K.-P., Lin, C.-Y., Lee, C.-H., Shiah, F.-K., Chang, J., 2003. The coupling of oligotrophic ciliate populations and hydrography in the East China Sea: spatial and temporal variations. *Deep Sea Research Part II* 50, 1279–1293.
- Cowlshaw, R.J., 2004. Seasonal coupling between ciliate and phytoplankton standing stocks in the South Slough of Coos Bay, Oregon. *Estuaries* 27, 539–550.
- Dolan, J.R., 1991. Guilds of ciliate microzooplankton in the Chesapeake Bay. *Estuarine, Coastal and Shelf Science* 33, 137–152.
- Dolan, J.R., Coats, D.W., 1990. Seasonal abundances of planktonic ciliates and microflagellates in mesohaline Chesapeake Bay Waters. *Estuarine, Coastal and Shelf Science* 31, 157–175.
- Fenchel, T., 1988. Marine plankton food chains. *Annual Review of Ecology, Evolution, and Systematics* 19, 19–38.
- Ferrier-Pagès, C., Rassoulzadegan, F., 1994. Seasonal impact of the microzooplankton on pico- and nanoplankton growth rates in the northwest Mediterranean Sea. *Marine Ecology Progress Series* 108, 283–294.
- Fileman, E.S., Leakey, R.J.G., 2005. Microzooplankton dynamics during the development of the spring bloom in the north-east Atlantic. *Journal of the Marine Biological Association of the United Kingdom* 85, 741–753.
- Fonda Umani, S., Beran, A., 2003. Seasonal variations in the dynamics of microbial plankton communities: first estimates from experiments in the Gulf of Trieste, Northern Adriatic Sea. *Marine Ecology Progress Series* 247, 1–16.
- Fuhrman, J., Azam, F., 1982. Thymidine incorporation as a measure of heterotrophic bacterioplankton production in marine surface waters: evaluation and field results. *Marine Biology* 66, 109–120.
- Fukami, K., Watanabe, A., Fujita, S., Yamaoka, K., Nishijima, T., 1999. Predation on naked protozoan microzooplankton by fish larvae. *Marine Ecology Progress Series* 185, 285–291.
- Gaines, G., Elbrächter, M., 1987. Heterotrophic nutrition. In: Taylor, F.J.R. (Ed.), *The Biology of Dinoflagellates*. Blackwell, Oxford, pp. 224–268.
- Garrison, D.L., Gowing, M.M., Hughes, M.P., 1998. Nano- and microplankton in the northern Arabian Sea during the Southwest Monsoon, August–September 1995. A US-JGOFS study. *Deep Sea Research II* 45, 2269–2299.
- Gohin, F., Lampert, L., Guillaud, J.-F., Herbland, A., Nezan, E., 2003. Satellite and in situ observations of a late winter phytoplankton bloom, in the northern Bay of Biscay. *Continental Shelf Research* 23, 1117–1141.
- Gowing, M.M., Garrison, D.L., Wishner, K.F., Gelfman, C., 2003. Mesopelagic microplankton of the Arabian Sea. *Deep Sea Research Part I: Oceanographic Research Papers* 50, 1205–1234.
- Guillaud, J.F., Aminot, A., Delmas, D., Gohin, F., Lunven, M., Labry, C., Herbland, A., 2008. Seasonal variation of riverine inputs in the northern Bay of Biscay (France) and patterns of marine phytoplankton response. *Journal of Marine Systems* 72, 309–319.
- Hagström, A., Azam, F., Andersson, A., Wikner, J., Rassoulzadegan, F., 1988. Microbial loop in an oligotrophic pelagic marine ecosystem: possible roles of cyanobacteria and nanoflagellates in the organic fluxes. *Marine Ecology Progress Series* 49, 171–178.
- Hansen, P.J., 1991a. Dinophysis – a planktonic dinoflagellate genus which act both as a prey and a predator of a ciliate. *Marine Ecology Progress Series* 69, 201–204.
- Hansen, P.J., 1991b. Quantitative importance and trophic role of heterotrophic dinoflagellates in a coastal pelagic food web. *Marine Ecology Progress Series* 73, 253–261.
- Herbland, A., Delmas, D., Laborde, P., Sautour, B., Artigas, F., 1998. Phytoplankton spring bloom of the Gironde plume waters in the Bay of Biscay: early phosphorus limitation and food-web consequences. *Oceanologica Acta* 21, 279–291.
- Hwang, S.-J., Heath, R.T., 1997. The distribution of protozoa across a trophic gradient, factors controlling their abundance and importance in the plankton food web. *Journal of Plankton Research* 19, 491–518.
- Jacobson, D.M., Anderson, D.M., 1986. Thecate heterotrophic dinoflagellates feeding behavior and mechanisms. *Journal of Phycology* 22, 249–258.
- James, M.R., Hall, J.A., 1995. Planktonic ciliated protozoa: their distribution and relationship to environmental variables in a marine coastal ecosystem. *Journal of Plankton Research* 1, 659–683.
- Jeong, H.J., Latz, M.I., 1994. Growth and grazing rates of the heterotrophic dinoflagellate *Protoperdinium* spp. on red tide dinoflagellates. *Marine Ecology Progress Series* 106, 173–185.
- Johansson, M., Gorokhova, E., Larsson, U., 2004. Annual variability in ciliate community structure, potential prey and predators in the open northern Baltic Sea proper. *Journal of Plankton Research* 26, 67–80.
- Jürgens, K., Wickham, S.A., Rothhaupt, K.O., Santer, B., 1996. Feeding rates of macro- and microzooplankton on heterotrophic nanoflagellates. *Limnology and Oceanography* 41, 1833–1839.
- Kamiyama, T., Tsujino, M., 1996. Seasonal variation in the species composition of tintinnid ciliates in Hiroshima Bay, the Seto Inland Sea of Japan. *Journal of Plankton Research* 18, 2312–2327.
- Koutsikopoulos, C., Beilouis, P., Leroy, C., Taillefer, F., 1998. Temporal trends and spatial structures of the sea surface temperature in the Bay of Biscay. *Oceanologica Acta* 21, 335–344.
- Labry, C., Herbland, A., Delmas, D., Laborde, P., Lazure, P., Froidefond, J.M., Jegou, A.M., Sautour, B., 2001. Initiation of winter phytoplankton blooms within the Gironde plume waters in the Bay of Biscay. *Marine Ecology Progress Series* 212, 117–130.
- Labry, C., Herbland, A., Delmas, D., 2002. The role of phosphorus on planktonic production of the Gironde plume waters in the Bay of Biscay. *Journal of Plankton Research* 24, 97–117.
- Landry, M.R., Calbet, A., 2004. Microzooplankton production in the oceans. *ICES Journal of Marine Science* 61, 501–507.
- Landry, M.R., Hassett, R.P., 1982. Estimating the grazing impact of marine microzooplankton. *Marine Biology* 67, 283–288.
- Lazure, P., Jegou, A.-M., 1998. 3D modelling of seasonal evolution of Loire and Gironde plumes on Biscay Bay continental shelf. *Oceanologica Acta* 21, 165–177.
- Leakey, R.J.G., Burkill, P.H., Sleight, M.A., 1992. Planktonic ciliates in Southampton water: abundance, biomass, production, and role in pelagic carbon flow. *Marine Biology* 114, 67–83.
- Legendre, L., Rassoulzadegan, F., 1995. Plankton and nutrients dynamics in marine waters. *Ophelia* 41, 153–172.
- Lessard, E.J., Swift, E., 1986. Dinoflagellates from the North Atlantic classified as phototrophic or heterotrophic by epifluorescence microscopy. *Journal of Plankton Research* 8, 1209–1215.
- Loyer, S., Lampert, L., Menesguen, A., Cann, P., Labasque, T., 2006. Seasonal evolution of the nutrient pattern on Biscay Bay continental shelf over the years 1999–2000. *Scientia Marina* 70, 31–46.
- Lunven, M., Guillaud, J.F., Youénu, A., Crassous, M.P., Berric, R., Le Gall, E., Kérouel, R., Labry, C., Aminot, A., 2005. Nutrient and phytoplankton distribution in the Loire River plume (Bay of Biscay, France) resolved by a new fine scale sampler. *Estuarine, Coastal and Shelf Science* 65, 94–108.
- Lynn, D.H., Montagnes, D.J.S., 1991. Global production of heterotrophic marine planktonic ciliates. In: Reid, P.C., Turley, C.M., Burkill, P.H. (Eds.), *Protozoa and Their Role in Marine Processes*. Springer, Berlin, Heidelberg, New York, pp. 281–307.
- Magueur, J.F., L'Helguen, S., Waeles, M., Morin, P., Riso, R., Caradee, J., 2009. Size-fractionated phytoplankton biomass and nitrogen uptake in response to high nutrient load in the North Biscay Bay in spring. *Continental Shelf Science* 29, 1103–1110.
- Marie, D., Partensky, F., Jacquet, S., Vault, D., 1997. Enumeration and cell cycle analysis of natural populations of marine picoplankton by flow cytometry using the nucleic acid stain SYBR Green I. *Applied and Environmental Microbiology* 63, 186–193.
- Marquis, E., Niquil, N., Delmas, D., Hartmann, H.J., Bonnet, D., Carlotti, F., Herbland, A., Labry, C., Sautour, B., Laborde, P., Vézina, A., Dupuy, C., 2007. Inverse analysis of the planktonic food web dynamics related to phytoplankton bloom development on the continental shelf of the Bay of Biscay, French coast. *Estuarine, Coastal and Shelf Science* 73, 223–235.
- Massana, R., Terrado, R., Forn, I., Lovejoy, C., Pedros-Alio, C., 2006. Distribution and abundance of uncultured heterotrophic flagellates in the world oceans. *Environmental Microbiology* 8, 1515–1522.
- Menden-Deuer, S., Lessard, E.J., 2000. Carbon to volume relationships for dinoflagellates, diatoms, and other protist plankton. *Limnology and Oceanography* 45, 569–579.
- Modigh, M., 2001. Seasonal variations of photosynthetic ciliates at a Mediterranean coastal site. *Aquatic Microbial Ecology* 23, 163–175.
- Montagnes, D.J.S., 1996. Growth responses of planktonic ciliates in the genera *Strobilidium* and *Strombidium*. *Marine Ecology Progress Series* 130, 241–254.
- Montagnes, D.J.S., Lynn, D.H., Roff, J.C., Taylor, W.D., 1988. The annual cycle of heterotrophic planktonic ciliates in the waters surrounding the Isles of Shoals, Gulf of Maine: an assessment of their trophic role. *Marine Biology* 99, 21–30.
- Nielsen, T.G., Kiorboe, T., 1994. Regulation of zooplankton biomass and production in a temperate, coastal ecosystem. 2. Ciliates. *Limnology and Oceanography* 39, 508–519.
- Pelegri, S.P., Dolan, J.R., Rassoulzadegan, F., 1999. Use of high temperature catalytic oxidation (HTCO) to measure carbon content of microorganisms. *Aquatic Microbial Ecology* 16, 273–280.
- Perthaler, J., 2005. Predation on prokaryotes in the water column and its ecological implications. *Nature Reviews Microbiology* 3, 537–546.
- Pierce, R.W., Turner, J.T., 1992. Ecology of planktonic ciliates in marine food webs. *Reviews in Aquatic Sciences* 6, 139–181.
- Pomeroy, L., 1974. The oceans food web, a changing paradigm. *International Journal of Biological Sciences* 24, 499–504.
- Putt, M., Stoecker, D.K., 1989. An experimentally determined carbon: volume ratio for marine "oligotrichous" ciliates from estuarine and coastal waters. *Limnology and Oceanography* 34, 1097–1103.
- Sanders, R.W., 1987. Tintinnids and other microzooplankton – seasonal distributions and relationships to resources and hydrography in a Maine estuary. *Journal of Plankton Research* 9, 65–77.
- Sautour, B., Artigas, L.F., Delmas, D., Herbland, A., Laborde, P., 2000. Grazing impact of micromesozooplankton during a spring situation in coastal waters off the Gironde estuary. *Journal of Plankton Research* 22, 531–552.
- Sherr, B.F., Sherr, E.B., Rassoulzadegan, F., 1988. Rates of digestion of bacteria by marine phagotrophic protozoa: temperature dependence. *Applied and Environmental Microbiology* 54, 1091–1095.
- Sherr, E., Sherr, B.F., Paffenhofer, G.-A., 1986. Phagotrophic protozoa as food for metazoans: a "missing" trophic link in marine pelagic food webs? *Marine Microbial Food Webs* 1, 61–80.
- Sherr, E.B., Caron, D.A., Sherr, B.F., 1994. Staining of heterotrophic protists for visualisation via epifluorescence microscopy. In: Kemp, P.F., Sherr, B.F., Sherr, E.B., Coll, J.J. (Eds.), *Handbook of Methods in Aquatic Microbial Ecology*. Lewis Publishers, Boca Raton, pp. 213–227.
- Sherr, E.B., Sherr, B.F., 1987. High rates of consumption of bacteria by pelagic ciliates. *Nature* 325, 710–711.

- Shinada, A., Ban, S., Ikeda, T., 2003. Seasonal changes in nano/micro-zooplankton herbivory and heterotrophic nano-flagellates bacterivory off Cape Esan, Southwestern Hokkaido, Japan. *Journal of Oceanography* 59, 609–618.
- Shinada, A., Ban, S., Yamada, Y., Ikeda, T., 2005. Seasonal variations of plankton food web structure in the coastal water off Usujiri Southwestern Hokkaido, Japan. *Journal of Oceanography* 61, 645–654.
- Sime-Ngando, T., Gosselin, M., Roy, S., Chanut, J.-P., 1995. Significance of planktonic ciliated protozoa in the lower St. Lawrence estuary: comparison with bacterial, phytoplankton, and particulate organic carbon. *Aquatic Microbial Ecology* 9, 243–258.
- Sommaruga, R., 1995. Microbial and classical food webs: a visit to a hypertrophic lake. *FEMS Microbiology Ecology* 17, 257–270.
- Stoecker, D.K., Davis, L.H., Anderson, D.M., 1984. Fine scale spatial correlations between planktonic ciliates and dinoflagellates. *Journal of Plankton Research* 6, 829–842.
- Strickland, J.D.H., Parsons, T.R., 1972. A practical handbook of seawater analysis. *Journal of the Fisheries Research Board of Canada* 167, 47–89.
- Thingstad, T.F., Rassoulzadegan, F., 1999. Conceptual models for the biogeochemical role of the photic zone microbial food web, with particular reference to the Mediterranean Sea. *Progress in Oceanography* 44, 271–286.
- Urrutxurtu, I., Orive, E., Sota, A., 2003. Seasonal dynamics of ciliated protozoa and their potential food in an eutrophic estuary (Bay of Biscay). *Estuarine, Coastal and Shelf Science* 57, 1169–1182.
- Utermöhl, H., 1958. Zur Vervollkommnung der quantitativen phytoplankton-methode. *Mitteilungen Internationale Vereinigung für Theoretische und Angewandte Limnologie* 9, 1–38.
- Verity, P.G., Langdon, C., 1984. Relationships between Lorica volume, carbon, nitrogen, and ATP content of tintinnids in Narragansett Bay. *Journal of Plankton Research* 6, 859–868.
- Verity, P.G., Paffenhofer, G.-A., Wallace, D., Sherr, E., Sherr, B., 1996. Composition and biomass of plankton in spring on the Cape Hatteras shelf, with implications for carbon flux. *Continental Shelf Research* 16, 1087–1116.
- Verity, P.G., 1987. Abundance, community composition, size distribution, and production rates of tintinnids in Narragansett Bay, Rhode Island. *Estuarine, Coastal and Shelf Science* 24, 671–690.
- Vincent, D., Hartmann, H.J., 2001. Contribution of ciliated Microprotozoans and Dinoflagellates to the diet of three copepod species in the Bay of Biscay. *Hydrobiologia* 443, 193–204.
- Waterbury, J.B., Watson, S.W., Guillard, R.R.L., Brand, L.E., 1979. Widespread occurrence of a unicellular, marine, planktonic, cyanobacterium. *Nature* 277, 293–294.
- Yang, E.J., Choi, J.K., Hyun, J.H., 2008. Seasonal variation in the community and size structure of nano- and microzooplankton in Gyeonggi Bay, Yellow Sea. *Estuarine, Coastal and Shelf Science* 77, 320–330.