



Gindikin-Karpelevich finiteness for Kac-Moody groups over local fields

Auguste Hébert

► To cite this version:

Auguste Hébert. Gindikin-Karpelevich finiteness for Kac-Moody groups over local fields. 2016. hal-01248091v2

HAL Id: hal-01248091

<https://hal.science/hal-01248091v2>

Preprint submitted on 9 Mar 2016 (v2), last revised 7 Nov 2017 (v4)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Gindikin-Karpelevich finiteness for Kac-Moody groups over local fields

Auguste HÉBERT

Univ Lyon, UJM-Saint-Etienne CNRS

UMR 5208 CNRS, F-42023, SAINT-ETIENNE, France

Abstract

In this paper, we prove some finiteness results about split Kac-Moody groups over local non-archimedean fields. Our results generalize those of "An affine Gindikin-Karpelevich formula" by Alexander Braverman, Howard Garland, David Kazhdan and Manish Patnaik. We do not require our groups to be affine. We use the hovel \mathcal{I} associated to this situation, which is the analogue of the Bruhat-Tits building for a reductive group.

1 Introduction

The classical Gindikin-Karpelevich formula was introduced in 1962 by Simon Gindikin and Fridrikh Karpelevich. It applies to real semi-simple Lie groups and it enables to compute certain Plancherel densities for semi-simple Lie groups. This formula was established in the non-archimedean case by Robert Langlands in [Lan71]. In 2014, in [BGKP14], Alexander Braverman, Howard Garland, David Kazhdan and Manish Patnaik obtained a generalisation of this formula in the affine Kac-Moody case.

Let \mathcal{F} be an ultrametric field. Let \mathcal{O} be its ring of integers, π be a generator of the maximal ideal of \mathcal{O} and q denote the cardinal of the residue field $\mathcal{O}/\pi\mathcal{O}$. Let \mathbf{G} be a split Kac-Moody group over \mathcal{F} . Let \mathbf{T} be a maximal split torus of \mathbf{G} . Choose a pair \mathbf{B}, \mathbf{B}^- of opposite Borel subgroups such that $\mathbf{B} \cap \mathbf{B}^- = \mathbf{T}$ and \mathbf{U}, \mathbf{U}^- be their unipotent radicals. We use boldface letters to denote schemes : $\mathbf{G}, \mathbf{T}, \dots$ and their sets of \mathcal{F} -points are denoted by G, T, \dots . Let Λ and Λ^\vee be the root lattice and the coroot lattice of \mathbf{T} . Let R be the set of roots \mathbf{G} and R^+ be its set of positive roots. Let $K = \mathbf{G}(\mathcal{O})$. For a coroot $\nu^\vee = \sum n_i \nu_i^\vee \in \Lambda^\vee$, where the ν_i^\vee are simple coroots, one sets $|\nu^\vee| = \sum n_i$. Let $\mathbb{C}[[\Lambda^\vee]]$ be the Looijenga's coweight algebra of \mathbf{G} , with generators e^{λ^\vee} , for $\lambda^\vee \in \Lambda^\vee$. Alexander Braverman, Howard Garland, David Kazhdan and Manish Patnaik proved that when \mathbf{G} is the affine Kac-Moody group associated to a simply connected semi-simple split group, we have the following formula:

$$\sum_{\mu^\vee \in \Lambda^\vee} |K \backslash K \pi^{\lambda^\vee} U^- \cap K \pi^{\lambda^\vee - \mu^\vee} U| e^{\lambda^\vee - \mu^\vee} q^{|\lambda^\vee - \mu^\vee|} = \frac{1}{H_0} \prod_{\alpha \in R_+} \left(\frac{1 - q^{-1} e^{-\alpha^\vee}}{1 - e^{-\alpha^\vee}} \right)^{m_\alpha} \quad (1)$$

where for $\alpha \in R_+$, m_α denotes the multiplicity of the coroot α^\vee in the Lie algebra \mathfrak{g} of \mathbf{G} , and H_0 is some term (there is a formula describing H_0 in [BGKP14]) depending on \mathfrak{g} . When \mathbf{G} , is a reductive group, $H_0 = 1$, the m_α are equals to 1 and this formula is equivalent to Gindikin-Karpelevich formula.

The aim of this article is to show the following four theorems:

Theorem 4.7: Let $\mu^\vee \in \Lambda^\vee$. Then if $\mu^\vee \notin \Lambda_-^\vee$, $K \backslash K\pi^{\lambda^\vee} U^- \cap K\pi^{\lambda^\vee + \mu^\vee} U$ is empty for all $\lambda^\vee \in \Lambda^\vee$. If $\mu^\vee \in \Lambda_+^\vee$, then for $\lambda^\vee \in \Lambda^\vee$ sufficiently dominant, $K \backslash K\pi^{\lambda^\vee} U^- \cap K\pi^{\lambda^\vee + \mu^\vee} U \subset K \backslash K\pi^{\lambda^\vee} K \cap K\pi^{\lambda^\vee + \mu^\vee} U$.

Corollary 5.3: Let $\mu^\vee \in \Lambda^\vee$. For all $\lambda \in \Lambda^\vee$, $K \backslash K\pi^{\lambda^\vee} K \cap K\pi^{\lambda^\vee + \mu^\vee} U$ is finite and is empty if $\mu^\vee \notin \Lambda_-^\vee$.

Theorem 5.6: Let $\mu^\vee \in \Lambda_-^\vee$ and $\lambda^\vee \in \Lambda^\vee$. Then $|K \backslash K\pi^{\lambda^\vee} U^- \cap K\pi^{\lambda^\vee + \mu^\vee} U| = |K \backslash K\pi^0 U^- \cap K\pi^{\mu^\vee} U|$ and these sets are finite.

Theorem 6.4: Let $\mu^\vee \in \Lambda^\vee$. Then for $\lambda^\vee \in \Lambda^\vee$ sufficiently dominant, $K \backslash K\pi^{\lambda^\vee} U^- \cap K\pi^{\lambda^\vee + \mu^\vee} U = K \backslash K\pi^{\lambda^\vee} K \cap K\pi^{\lambda^\vee + \mu^\vee} U$.

They correspond to Theorem 1.9 of [BGKP14], except Theorem 6.4, which seems weaker. Theorem 5.6 is named "Gindikin-Karpelevich finiteness" in [BKP14] and it enables to make sense to the left side of formula 1. Thus this could be a first step to a generalization of Gindikin-Karpelevich formula to general split Kac-Moody groups. However, we do not know yet how to express the term H_0 when \mathbf{G} is not affine.

The main tool that we use to prove these theorems is the hovel \mathcal{I} associated to \mathbf{G} , which was defined Stéphane Gaussent and Guy Rousseau in [GR14] and generalized by Guy Rousseau in [Rou12]. This hovel is a geometric object similar to a Bruhat-Tits building. It is a set covered by apartments isomorphic to a standard apartment \mathbb{A} on which \mathbf{G} acts. The standard apartment \mathbb{A} is an affine space with a structure obtained from a lattice Y containing Λ^\vee (which can be thought of as Λ^\vee in a first approximation) and from a set of hyperplanes called walls, defined by the roots of \mathbf{G} . In general, \mathcal{I} is not a building: there may be two points such that no apartment of \mathcal{I} contains these two points simultaneously. However some properties of euclidean buildings remain and, for example one can still define the retraction onto an apartment centred at a sector-germ. We can always define a fundamental vectorial chamber C_f^v in \mathbb{A} and its opposite, which gives rise to two retractions: $\rho_{+\infty}$ and $\rho_{-\infty}$. We also have a "vectorial distance" d^v defined on a subset of \mathcal{I}^2 . These applications enable to give other expressions to the sets defined above: for all $\lambda^\vee \in \Lambda^\vee$, $K \backslash K\pi^{\lambda^\vee} U^-$ corresponds to $\rho_{-\infty}^{-1}(\{\lambda^\vee\})$, $K \backslash K\pi^{\lambda^\vee} U$ corresponds to $\rho_{+\infty}^{-1}(\{\lambda^\vee\})$ and $K \backslash K\pi^{\lambda^\vee} K$ corresponds to $S^v(0, \lambda^\vee) = \{x \in \mathcal{I} \mid d^v(0, x) \text{ is defined and } d^v(0, x) = \lambda^\vee\}$. This enables to use properties of "Hecke paths", which are more or less the images of segments in the hovel by retractions, to prove these theorems. These paths were first defined by Misha Kapovich and John J. Millson in [KM08].

The framework of hovels is a bit more general than that of split Kac-Moody groups over local non-archimedean field. For example, we might also apply this to almost split Kac-Moody groups over local non-archimedean field (see [Rou12]).

In the sequel, the results are not stated as in this introduction. Their statements use retractions and vectorial distance. They are also a bit more general: they take into account the inessential part of the standard apartment \mathbb{A} , which will be defined later. Corollary 5.3, as it is stated in this introduction was proved in Section 5 of [GR14] and it is slightly generalized in the following. The lattice Λ^\vee will be denoted by Q^\vee .

In section 2 we set the general frameworks, we set the notation, we define hovels and we prove that $Y^{++} = Y \cap \overline{C_f^v}$ is a finitely generated monoid, which will be useful to prove Theorem 6.4.

In section 3 we first define two applications: $T_\nu : \mathcal{I} \rightarrow \mathbb{R}_+$ and $y_\nu : \mathcal{I} \rightarrow \mathbb{A}$, for a fixed $\nu \in C_f^v$. For $x \in \mathcal{I}$, $T_\nu(x)$ measures the distance between the point x and the apartment \mathbb{A}

along $\mathbb{R}_+\nu$ and $y_\nu(x)$ defines the projection of x on \mathbb{A} along $\mathbb{R}_+\nu$. We also determine the antecedents of some kinds of paths by $\rho_{-\infty}$.

In section 4, we show that under some conditions, T_ν is bounded and we deduce Theorem 4.7.

In section 5, we study some kinds of translations of \mathcal{I} , which enables us to show Corollary 5.3 and Theorem 5.6.

In section 6, we use the tools of the preceding sections to show Theorem 6.4.

Acknowledgements I thank Stéphane Gaussent for suggesting me this problem and for helping me improving the manuscript. I also thank Guy Rousseau for his remarks on a previous version of this paper.

2 General frameworks

2.1 Root generating system

A Kac-Moody matrix (or generalized Cartan matrix) is a square matrix $C = (c_{i,j})_{i,j \in I}$ with integers coefficients, indexed by a finite set I and satisfying:

1. $\forall i \in I, c_{i,i} = 2$
2. $\forall (i,j) \in I^2 | i \neq j, c_{i,j} \leq 0$
3. $\forall (i,j) \in I^2, c_{i,j} = 0 \Leftrightarrow c_{j,i} = 0$.

A root generating system is a 5-tuple $\mathcal{S} = (C, X, Y, (\alpha_i)_{i \in I}, (\alpha_i^\vee)_{i \in I})$ made of a Kac-Moody matrix C indexed by I , of two dual free \mathbb{Z} -modules X (of characters) and Y (of cocharacters) of finite rank $\text{rk}(X)$, a family $(\alpha_i)_{i \in I}$ (of simple roots) in X and a family $(\alpha_i^\vee)_{i \in I}$ (of simple coroots) in Y . They have to satisfy the following compatibility condition: $c_{i,j} = \alpha_j(\alpha_i^\vee)$ for all $i, j \in I$. We also suppose that the family $(\alpha_i)_{i \in I}$ is free in X and that the family $(\alpha_i^\vee)_{i \in I}$ is free in Y .

We now fix a Kac-Moody matrix C and a root generating system with matrix C .

Let $V = Y \otimes \mathbb{R}$. Every element of X induces a linear form on V . We will consider X as a subset of the dual V^* of V : the $\alpha_i, i \in I$ are viewed as linear form on V . For $i \in I$, we define an involution r_i of V by $r_i(v) = v - \alpha_i(v)\alpha_i^\vee$ for all $v \in V$. Its space of fixed points is $\ker \alpha_i$. The subgroup of $\text{GL}(V)$ generated by the α_i for $i \in I$ is denoted by W^v and is called the Weyl group of \mathcal{S} .

For $x \in V$ we let $\alpha(x) = (\alpha_i(x))_{i \in I} \in \mathbb{R}^I$.

Let $Q^\vee = \bigoplus_{i \in I} \mathbb{Z}\alpha_i^\vee$ and $P^\vee = \{v \in V | \alpha(v) \in \mathbb{Z}^I\}$. We call Q^\vee the *coroot-lattice* and P^\vee the *coweight-lattice* (but if $\bigcap_{i \in I} \ker \alpha_i \neq \{0\}$, this is not a lattice). Let $Q_+^\vee = \bigoplus_{i \in I} \mathbb{N}\alpha_i^\vee$, $Q_-^\vee = -Q_+^\vee$ and $Q_{\mathbb{R}}^\vee = \bigoplus_{i \in I} \mathbb{R}\alpha_i^\vee$. This enables us to define a preorder \leq_{Q^\vee} on V by the following way: for all $x, y \in V$, one writes $x \leq_{Q^\vee} y$ if $y - x \in Q_+^\vee$.

One defines an action of the group W^v on V^* by the following way: if $x \in V$, $w \in W^v$ and $\alpha \in V^*$ then $(w.\alpha)(x) = \alpha(w^{-1}.x)$. Let $\Phi = \{w.\alpha_i | (w, i) \in W^v \times I\}$, Φ is the set of *real roots*. Then $\Phi \subset Q$, where $Q = \bigoplus_{i \in I} \mathbb{Z}\alpha_i$. Let $W^a = Q^\vee \rtimes W^v \subset \text{GA}(V)$ the *affine Weyl group* of \mathcal{S} , where $\text{GA}(V)$ is the group of affine isomorphisms of V .

Let $V_{in} = \bigcap_{i \in I} \ker \alpha_i$. Then one has $Y + V_{in} \subset P^\vee$.

Remark 2.1. Let $C = (c_{i,j})_{i,j \in I}$ a Kac-Moody matrix. Then there exists a generating root system $(C, X, Y, (\alpha_i)_{i \in I}, (\alpha_i^\vee)_{i \in I})$ such that $Y + V_{in} = P^\vee$. For this, suppose $I = [1, k]$ for

some $k \in \mathbb{N}$. Let $n = 2k$ and let $(\alpha_i)_{i \in \llbracket 1, n \rrbracket}$ be the canonical basis of the dual of \mathbb{Z}^n . Let $(e_i)_{i \in \llbracket 1, n \rrbracket}$ be the canonical basis of \mathbb{Z}^n . For $i \in \llbracket 1, k \rrbracket$, let $\alpha_i^\vee = \sum_{j=1}^k c_{i,j} e_j + e_{i+k}$. Let $X = \bigoplus_{i \in I} \mathbb{Z} \alpha_i$, $X_{\mathbb{Q}} = \text{vect}_{\mathbb{Q}}(X)$, $Y_{\mathbb{Q}} = \bigoplus_{i \in I} \mathbb{Q} \alpha_i^\vee$ and $Y = \{y \in Y_{\mathbb{Q}} \mid x(y) \in \mathbb{Z} \forall i \in I\}$. Then $(C, X, Y, (\alpha_i)_{i \in I}, (\alpha_i^\vee)_{i \in I})$ is a generating root system.

Let $u \in P^\vee$. Then there exists $y \in Y$ such that $(\alpha_i(y))_{i \in I} = (\alpha_i(u))_{i \in I}$ and thus $u \in Y + V_{in}$. Therefore, $P^\vee = Y + V_{in}$.

2.2 Description of Y^{++}

In this subsection we show that $Y^{++} = Y \cap \overline{C_f^v}$ is a finitely generated monoid, which will be useful to prove Theorem 6.4.

Let $l \in \mathbb{N}^*$. Let us define a binary relation \prec on \mathbb{N}^l . Let $x, y \in \mathbb{N}^l$, and $x = (x_1, \dots, x_l)$, $y = (y_1, \dots, y_l)$, then one says $x \prec y$ if $x \neq y$ and for all $i \in \llbracket 1, l \rrbracket$, $x_i \leq y_i$.

Lemma 2.2. *Let $l \in \mathbb{N}^*$ and F be a subset of \mathbb{N}^l satisfying property $(\text{INC}(l))$:*

for all $x, y \in F$, x and y are not comparable for \prec .

Then F is finite.

Proof. This is clear for $l = 1$ because a set F satisfying $\text{INC}(1)$ is a singleton or \emptyset .

Suppose that $l > 1$ and that we have proven that all set satisfying $\text{INC}(l-1)$ is finite.

Let F be a set satisfying $\text{INC}(l)$ and suppose F infinite. Let $(\lambda_n)_{n \in \mathbb{N}}$ be an injective sequence of F . One writes $(\lambda_n) = (\lambda_n^1, \dots, \lambda_n^l)$. Let $(m_n) = (\min(\lambda_n))_{n \in \mathbb{N}}$ and $M = \max \lambda_0$. Then for all $n \in \mathbb{N}$, $m_n \leq M$ (if $m_n > M$, we would have $\lambda_0 \prec \lambda_n$). Maybe extracting a sequence of λ , one can suppose that $(m_n) = (\lambda_n^i)$ for some $i \in \llbracket 1, l \rrbracket$ and that (m_n) is constant and equal to $m_0 \in \llbracket 0, M \rrbracket$. For $x \in \mathbb{N}^l$, we define $\tilde{x} = (x_j)_{j \in \llbracket 1, l \rrbracket \setminus \{i\}} \in \mathbb{N}^{l-1}$.

Set $\tilde{F} = \{\tilde{\lambda}_n \mid n \in \mathbb{N}\}$. The set \tilde{F} satisfies $\text{INC}(l-1)$. By the induction hypothesis, \tilde{F} is finite and thus F is finite, which is absurd. Hence F is finite and the lemma is proven. \square

Lemma 2.3. *There exists a finite set $E \subset Y$ such that $Y^{++} = \sum_{e \in E} \mathbb{N}e$.*

Proof. The set $Y_{in} = Y \cap \mathbb{A}_{in}$ is a lattice in the vectorial space it spans. Consequently, it is a finitely generated \mathbb{Z} -module and thus a finitely generated monoid. Let E_1 be a finite set generating Y_{in} as a monoid.

Let $Y_{>0} = Y^{++} \setminus Y_{in}$. Let $\mathcal{P} = \{a \in Y_{>0} \mid a \neq b + c \forall b, c \in Y_{>0}\}$. Let $\alpha : Y^{++} \rightarrow \mathbb{N}^I$ such that $\alpha(x) = (\alpha_i(x))_{i \in I}$ for all $x \in Y^{++}$. Let $a, b \in \mathcal{P}$. If $\alpha(a) \prec \alpha(b)$, then $b = b - a + a$, with $a, b - a \in Y_{>0}$, which is absurd and by symmetry we deduce that $\alpha(a)$ and $\alpha(b)$ are not comparable for \prec . Therefore, by lemma 2.2, $\alpha(\mathcal{P})$ is finite. Let E_2 be a finite set of $Y_{>0}$ such that $\alpha(\mathcal{P}) = \{\alpha(x) \mid x \in E_2\}$. Then $Y^{++} = \sum_{e \in E_2} \mathbb{N}e + Y_{in} = \sum_{e \in E} \mathbb{N}e$, where $E = E_1 \cup E_2$. \square

2.3 Vectorial faces

Define $C_f^v = \{v \in V \mid \alpha_i(v) > 0, \forall i \in I\}$. We call it the *fundamental chamber*. For $J \subset I$, one sets $F^v(J) = \{v \in V \mid \alpha_i(v) = 0 \forall i \in J, \alpha_i(v) > 0 \forall i \in J \setminus I\}$. Then the closure $\overline{C_f^v}$ of C_f^v is the union of the $F^v(J)$ for $J \subset I$. The *positive* (resp. *negative*) *vectorial faces* are the sets $w.F^v(J)$ (resp. $-w.F^v(J)$) for $w \in W^v$ and $J \subset I$. A *vectorial face* is either a positive vectorial face or a negative vectorial face. We call *positive chamber* (resp. *negative*) every cone of the shape $w.C_f^v$ for some $w \in W^v$ (resp. $-w.C_f^v$). For all $x \in C_f^v$ and for all $w \in W^v$, $w.x = x$ implies that $w = 1$. In particular the action of w on the positive chambers is simply transitive. The *Tits cone* \mathcal{T} is defined by $\mathcal{T} = \bigcup_{w \in W^v} w.\overline{C_f^v}$. We also consider the negative cone $-\mathcal{T}$. We define a W^v invariant relation \leq on V by: $\forall (x, y) \in V^2$, $x \leq y \Leftrightarrow y - x \in \mathcal{T}$.

2.4 Filters

Définition 2.4. A filter in a set E is a nonempty set F of nonempty subsets of E such that, for all subsets S, S' of E , if $S, S' \in F$ then $S \cap S' \in F$ and, if $S' \subset S$, with $S' \in F$ then $S \in F$.

If F is a filter in a set E , and E' is a subset of E , one says that F contains E' if every element of F contains E' . If E' is nonempty, the set $F_{E'}$ of subsets of E containing E' is a filter. By abuse of language, we will sometimes say that E' is a filter by identifying $F_{E'}$ and E' . If F is a filter in E , its closure \bar{F} (resp. its convex envelope) is the filter of subsets of E containing the closure (resp. the convex envelope) of some element of F . A filter F is said to be contained in another filter F' : $F \subset F'$ (resp. in a subset Z in E : $F \subset Z$) if and only if any set in F' (resp. if Z) is in F .

If $x \in V$ and Ω is a subset of V containing x in its closure, then the *germ* of Ω in x is the filter $\text{germ}_x(\Omega)$ of subsets of V containing a neighbourhood in Ω of x .

A *sector* in V is a set of the shape $\mathfrak{s} = x + C^v$ with $C^v = \pm w.C_f^v$ for some $x \in \mathbb{A}$ and $w \in W^v$. The point x is its *base point* and C^v is its *direction*. The intersection of two sectors of the same direction contains a sector of the same direction.

A *sector-germ* of a sector $\mathfrak{s} = x + C^v$ is the filter \mathfrak{S} of subsets of V containing a V -translate of \mathfrak{s} . It only depends on the direction C^v . We denote by $+\infty$ (resp. $-\infty$) the sector-germ of C_f^v (resp. of $-C_f^v$).

A ray δ with base point x and containing $y \neq x$ (or the interval $]x, y] = [x, y] \setminus \{x\}$ or $[x, y]$) is called *preordered* if $x \leq y$ or $y \leq x$ and *generic* if $y - x \in \pm \overset{\circ}{\mathcal{T}}$, the interior of $\pm \mathcal{T}$.

In the next subsection, we define the notions of faces, enclosures and chimneys defined in [Rou11] 1.7 and 1.10 and in [GR14] 1.4. For a first reading, one can just know the following facts about these objects and skip this subsection:

1. To any filter F of V is associated its enclosure $\text{cl}_{\mathbb{A}}(F)$ which is a filter in \mathbb{A} containing the convex envelope of the closure of F .
2. A face or a chimney is a filter in V .
3. A sector is a chimney which is solid and splayed.
4. If a chimney is a sector, its germ as a chimney coincides with its germ as a sector.
5. Every $x \in V$ is in some face of V .
6. The group W^a permutes the sectors, the enclosures, the faces and the chimneys of V .

2.5 Definitions of enclosures, faces, chimneys and related notions

Let $\Delta = \Phi \cup \Delta_{im}^+ \cup \Delta_{im}^- \subset Q$ be the set of all roots (recall that $Q = \bigoplus_{i \in I} \mathbb{Z}\alpha_i$), defined in [Kac94]. The group W^v stabilizes Δ . For $\alpha \in \Delta$, and $k \in \mathbb{Z} \cup +\infty$, let $D(\alpha, k) = \{v \in V \mid \alpha(v) + k \geq 0\}$ (and $D(\alpha, +\infty) = V$ for all $\alpha \in \Delta$) and $D^\circ(\alpha, k) = \{v \in V \mid \alpha(v) + k > 0\}$ (for $\alpha \in \Delta$ and $k \in \mathbb{Z} \cup \{+\infty\}$).

Given a filter F of subsets of V , its *enclosure* $\text{cl}_V(F)$ is the filter made of the subsets of V containing an element of F of the shape $\bigcap_{\alpha \in \Delta} D(\alpha, k_\alpha)$ where $k_\alpha \in \mathbb{Z} \cup \{+\infty\}$ for all $\alpha \in \Delta$.

A *face* F in V is a filter associated to a point $x \in V$ and a vectorial face $F^v \subset V$. More precisely, a subset S of V is an element of the face $F = F(x, F^v)$ if and only if, it contains an intersection of half-spaces $D(\alpha, k_\alpha)$ or open half-spaces $D^\circ(\alpha, k_\alpha)$, with $k_\alpha \in \mathbb{Z}$ for all $\alpha \in \Delta$, that contains $\Omega \cap (x + F^v)$, where Ω is an open neighbourhood of x in V .

There is an order on the faces: if $F \subset \overline{F'}$ we say that " F is a face of F' " or " F' contains F ". The dimension of a face F is the smallest dimension of an affine space generated by some $S \in F$. Such an affine space is unique and is called its support. A face is said to be *spherical* if the direction of its support meets the open Tits cone $\overset{\circ}{\mathcal{T}}$ or its opposite $-\overset{\circ}{\mathcal{T}}$; then its pointwise stabilizer W_F in W^v is finite.

We have $W^a \subset P^\vee \rtimes W^v$. As $\alpha(P^\vee) \subset \mathbb{Z}$ for all α in $\bigoplus_{i \in I} \mathbb{Z}\alpha_i$, if τ is a translation of V of a vector $p \in P^\vee$, then for all $\alpha \in Q$, τ permutes the sets of the shape $D(\alpha, k)$ where k runs over \mathbb{Z} . As W^v stabilizes Δ , any element of W^v permutes the sets of the shape $D(\alpha, k)$ where α runs over Δ . Therefore, W^a permutes the sets $D(\alpha, k)$, where (α, k) runs over $\Delta \times \mathbb{Z}$ and thus W^a permutes the enclosures, faces, chimneys, ... of V .

A *chamber* (or alcove) is a maximal face, or equivalently, a face such that all its elements contains a nonempty open subset of V .

A *panel* is a spherical face maximal among faces that are not chambers or, equivalently, a spherical face of dimension $n - 1$.

A *chimney* in V is associated to a face $F = F(x, F_0^v)$ and to a vectorial face F^v ; it is the filter $\mathfrak{r}(F, F^v) = \text{cl}_{\mathbb{A}}(F + F^v)$. The face F is the basis of the chimney and the vectorial face F^v its direction. A chimney is *splayed* if F^v is spherical, and is *solid* if its support (as a filter, i.e., the smallest affine subspace of V containing \mathfrak{r}) has a finite pointwise stabilizer in W^v . A splayed chimney is therefore solid.

A *shortening* of a chimney $\mathfrak{r}(F, F^v)$, with $F = F(x, F_0^v)$ is a chimney of the shape $\mathfrak{r}((F(x + \xi), F_0^v), F^v)$ for some $\xi \in \overline{F^v}$. The *germ* of a chimney \mathfrak{r} is the filter of subsets of V containing a shortening of \mathfrak{r} (this definition of shortening is slightly different from the one of [Rou11] 1.12 but follows [Rou12] 3.6) and we obtain the same germs with these two definitions).

2.6 Hovel

We now denote by \mathbb{A} the affine space V equipped with its faces, chimneys, ...

An apartment of type \mathbb{A} is a set A with a nonempty set $\text{Isom}(\mathbb{A}, A)$ of bijections (called isomorphisms) such that if $f_0 \in \text{Isom}(\mathbb{A}, A)$ then $f \in \text{Isom}(\mathbb{A}, A)$ if and only if, there exists $w \in W^a$ satisfying $f = f_0 \circ w$. An isomorphism between two apartments $\phi : A \rightarrow A'$ is a bijection such that ($f \in \text{Isom}(\mathbb{A}, A)$ if, and only if, $\phi \circ f \in \text{Isom}(\mathbb{A}, A')$). We extend all the notions that are preserved by W^a to each apartment. By the fact 6 of the above subsection, sectors, enclosures, faces and chimneys are well defined in any apartment of type \mathbb{A} .

Définition 2.5. *An ordered affine hovel of type \mathbb{A} is a set \mathcal{I} endowed with a covering \mathcal{A} of subsets called apartments such that:*

(MA1) *Any $A \in \mathcal{A}$ admits a structure of an apartment of type \mathbb{A} .*

(MA2) *If F is a point, a germ of a preordered interval, a generic ray or a solid chimney in an apartment A and if A' is another apartment containing F , then $A \cap A'$ contains the enclosure $\text{cl}_A(F)$ of F and there exists an isomorphism from A onto A' fixing $\text{cl}_A(F)$.*

(MA3) *If \mathfrak{R} is the germ of a splayed chimney and if F is a face or a germ of a solid chimney, then there exists an apartment that contains \mathfrak{R} and F .*

(MA4) *If two apartments A, A' contain \mathfrak{R} and F as in (MA3), then there exists an isomorphism from A to A' fixing $\text{cl}_A(\mathfrak{R} \cup F)$.*

(MAO) *If x, y are two points contained in two apartments A and A' , and if $x \leq_A y$ then the two segments $[x, y]_A$ and $[x, y]_{A'}$ are equal.*

In this definition, we say that an apartment contains a germ of a filter if it contains at least one element of this germ. We say that an application fixes a germ if it fixes at least one element of this germ.

Remark 2.6. (consequence 2.2 3) of [Rou11]) By (MA2), the axioms (MA3) and (MA4) also apply in a hovel when F is a point, a germ of a preordered segment and when \mathfrak{R} or F is a germ of a generic ray or a germ of a spherical sector face.

Until the end of this article, \mathcal{I} will be an affine ordered hovel. We suppose that \mathcal{I} is thick of *finite thickness*: the number of chambers (=alcoves) containing a given panel has to be finite, greater or equal to 3. This assumption will be crucial to use some theorems of [GR14] but we will not use it directly.

We assume that \mathcal{I} has a strongly transitive group of automorphisms G , which means that all isomorphisms involved in the above axioms are induced by elements of G . We choose in \mathcal{I} a fundamental apartment, that we identify with \mathbb{A} . As G is strongly transitive, the apartments of \mathcal{I} are the sets $g.\mathbb{A}$ for $g \in G$. The stabilizer N of \mathbb{A} induces a group $\nu(N)$ of affine automorphisms of \mathbb{A} and we suppose that $\nu(N) = W^v \ltimes Y$.

An example of such a hovel \mathcal{I} is the hovel associated to a split Kac-Moody group over an ultrametric field constructed in [GR08] and in [Rou12].

Vectorial distance For $x \in \mathcal{I}$, we denote by x^{++} the unique element in $\overline{C_f^v}$ conjugated by W^v to x . Let $\mathcal{I} \times_{\leq} \mathcal{I} = \{(x, y) \in \mathcal{I}^2 | x \leq y\}$ be the set of increasing pairs in \mathcal{I} . Such a pair is always in the same apartment $g.\mathbb{A}$ for some $g \in G$. So $g^{-1}.y - g^{-1}.x \in \mathcal{T}$, and we define the *vectorial distance* $d^v(x, y) \in \overline{C_f^v}$ by $d^v(x, y) = (g^{-1}.y - g^{-1}.x)^{++}$. It does not depend on the choices we made. This "distance" is G -invariant.

For $x \in \mathcal{I}$ and $\lambda \in \overline{C_f^v}$, one defines $S^v(x, \lambda) = \{y \in \mathcal{I} | x \leq y \text{ and } d^v(x, y) = \lambda\}$.

Remark 2.7. a) If $a \in Y$ and $\lambda \in \overline{C_f^v}$, then $S^v(a, \lambda) = \{x \in \mathcal{I} | \exists g \in G | g.a = a \text{ and } g.x = a + \lambda\}$.

b) Let $x, y \in \mathcal{I}$ and suppose that for some $g \in G$, $g.y - g.x \in \overline{C_f^v}$. Then $x \leq y$ and $d^v(x, y) = g.y - g.x$.

2.7 Retractions and Hecke paths

Let \mathfrak{R} be the germ of a splayed chimney of an apartment A . Let $x \in \mathcal{I}$. By (MA3), for all $x \in \mathcal{I}$, there exists an apartment A_x of \mathcal{I} containing x and \mathfrak{R} . By (MA4), there exists an isomorphism of apartments $\phi : A_x \rightarrow A$ fixing \mathfrak{R} . By [Rou11] 2.6, $\phi(x)$ does not depend on the choices we made and thus we can let $\rho_{A, \mathfrak{R}}(x) = \phi(x)$.

The application $\rho_{A, \mathfrak{R}}$ is a retraction from \mathcal{I} onto A . It only depends on \mathfrak{R} and A and we call it the *retraction onto A centred at \mathfrak{R}* .

We denote by $\rho_{+\infty}$ (resp. $\rho_{-\infty}$) the retraction onto \mathbb{A} of center $+\infty$ (resp. $-\infty$).

We now define Hecke paths. They are more or less the images by $\rho_{-\infty}$ of preordered segments $[x, y]$ in \mathcal{I} . The definition is a bit technical but it expresses the fact that the image of such a path "goes nearer to $+\infty$ " when it crosses a wall. A consequence of that is Remark 2.9 and we will not use directly this definition in the following.

We consider piecewise linear continuous paths $\pi : [0, 1] \rightarrow \mathbb{A}$ such that the values of π' belong to some orbit $W^v.\lambda$ for some $\lambda \in \overline{C_f^v}$. Such a path is called a λ -path. It is increasing with respect to the preorder relation \leq on \mathbb{A} . For any $t \neq 0$ (resp. $t \neq 1$), we let $\pi'_-(t)$ (resp. $\pi'_+(t)$) denote the derivative of π at t from the left (resp. from the right).

Définition 2.8. A Hecke path of shape λ with respect to $-C_f^v$ is a λ -path such that $\pi'_+(t) \leq_{W_{\pi(t)}^v} \pi'_-(t)$ for all $t \in [0, 1] \setminus \{0, 1\}$, which means that there exists a $W_{\pi(t)}^v$ -chain from $\pi'_-(t)$ to $\pi'_+(t)$, i.e., a finite sequence $(\xi_0 = \pi'_-(t), \xi_1, \dots, \xi_s = \pi'_+(t))$ of vectors in V and $(\beta_1, \dots, \beta_s) \in \phi^s$ such that, for all $i \in \llbracket 1, s \rrbracket$,

1. $r_{\beta_i}(\xi_{i-1}) = \xi_i$.
2. $\beta_i(\xi_{i-1}) < 0$.
3. $r_{\beta_i} \in W_{\pi(t)}^v$; i.e., $\beta_i(\pi(t)) \in \mathbb{Z}$: $\pi(t)$ is in a wall of direction $\ker(\beta_i)$.
4. Each β_i is positive with respect to $-C_f^v$; i.e., $\beta_i(C_f^v) > 0$.

Remark 2.9. Let $\pi : [0, 1] \rightarrow \mathbb{A}$ be a Hecke path of shape $\lambda \in \overline{C_f^v}$ with respect to $-C_f^v$. Then if $t \in [0, 1]$ such that π is derivable in t and $\pi'(t) \in \overline{C_f^v}$, then for all $s \geq t$, π is derivable in s and $\pi'(s) = \lambda$.

3 Segments and Rays in \mathcal{I}

In this section we begin by defining for all $\nu \in C_f^v$ two applications $y_\nu : \mathcal{I} \rightarrow \mathbb{A}$ and $T_\nu : \mathcal{I} \rightarrow \mathbb{R}_+$, where for all $x \in \mathcal{I}$, $T_\nu(x)$ and $y_\nu(x)$ can be considered as the distance between x and \mathbb{A} along $\mathbb{R}_+\nu$ and the projection of x on \mathbb{A} along $\mathbb{R}_+\nu$.

We also show that the only antecedent of some paths for $\rho_{-\infty}$ are themselves (this is Lemma 3.4).

Recall that we want to prove Theorem 4.7: Let $\mu \in \mathbb{A}$. Then if $\mu \notin Q_{\mathbb{R}}^\vee$, $\rho_{+\infty}^{-1}(\{\lambda + \mu\}) \cap \rho_{-\infty}^{-1}(\{\lambda\})$ is empty for all $\lambda \in \mathbb{A}$. If $\mu \in Q_{\mathbb{R}}^\vee$, then for $\lambda \in \mathbb{A}$ sufficiently dominant, $\rho_{+\infty}^{-1}(\{\lambda + \mu\}) \cap \rho_{-\infty}^{-1}(\{\lambda\}) \subset S^v(0, \lambda) \cap \rho_{+\infty}^{-1}(\{\lambda + \mu\})$.

Let us sketch the proof of this theorem. We fix $\nu \in C_f^v$ and $\mu \in Q_{\mathbb{R}}^\vee$. Let $T^- = T_\nu^-$ and $y^- = y_\nu^-$ be the analogue of T_ν and y_ν with ν replaced by $-\nu$. We show that if $\lambda \in \mathbb{A}$, $T^-(\rho_{+\infty}^{-1}(\{\lambda + \mu\}) \cap \rho_{-\infty}^{-1}(\{\lambda\}))$ is bounded by some constant $h(\mu)$ (this is Corollary 4.2), which uses Lemma 3.4 and Lemma 4.1). Therefore, for λ sufficiently dominant, $y^-(\rho_{+\infty}^{-1}(\{\lambda + \mu\}) \cap \rho_{-\infty}^{-1}(\{\lambda\})) \subset C_f^v$ and Lemma 4.6 completes the proof.

Let us recall briefly the notion of parallelism in \mathcal{I} . This is done more completely in [Rou11] Section 3. Let δ and δ' be two generic rays in \mathcal{I} . Then there exists a splayed chimney R containing δ and a solid chimney F containing δ' . By (MA3) there exists an apartment A containing the germ \mathfrak{R} of R and F . Therefore A contains translates of δ and δ' and we say that δ and δ' are *parallel*, if these translates are parallel in A . Parallelism is an equivalence relation and its equivalence classes are called *directions*.

Lemma 3.1. *Let $x \in \mathcal{I}$ and δ be a generic ray. Then there exists a unique ray $x + \delta$ in \mathcal{I} with base point x and direction δ . In any apartment A containing x and a ray δ' parallel to δ , this ray is the translate in A of δ' having x as a base point.*

This lemma is analogous to Proposition 4.7 1) of [Rou11]. The difficult part of this lemma is the uniqueness of such a ray because second part of the lemma yields a way to construct a ray having direction δ and x as a base point. This uniqueness can be shown exactly in the same manner as the proof of Proposition 4.7.1) by replacing "spherical sector face" by "generic ray". This is possible by NB.a) of Proposition 2.7 and by 2.2 3) (or by Remark 2.6 of this paper) of [Rou11].

Definition of y_ν and T_ν (resp. y_ν^- and T_ν^-) Let $x \in \mathcal{I}$. Let $\nu \in C_f^v$ and $\delta = \mathbb{R}_+\nu$, which is a generic ray. According to axiom (MA3) applied to a face containing x and the splayed chimney C_f^v , there exists an apartment A containing x and $+\infty$. Then A contains $x + \delta$. The set $x + \delta \cap \mathbb{A}$ is nonempty. Let $z \in x + \delta \cap \mathbb{A}$. Then $A \cap \mathbb{A}$ contains z , $+\infty$ and by (MA4),

$A \cap \mathbb{A}$ contains $\text{cl}(z, +\infty)$. As $\text{cl}(z, +\infty) \supset z + \overline{C_f^v}$, $A \cap \mathbb{A} \supset z + \delta$ and thus $x + \delta \cap \mathbb{A} = y + \delta$ or $x + \delta \cap \mathbb{A} = y + \delta^\circ$ for some $y \in x + \delta$, where $\delta^\circ = \mathbb{R}_+^* \nu$.

Suppose $x + \delta \cap \mathbb{A} = y + \delta^\circ$. Let $z \in y + \delta^\circ$. Then by (MA2) applied to $\text{germ}_y([y, z] \setminus \{y\})$, $\mathbb{A} \cap A \supset \text{cl}(\text{germ}_y([y, z] \setminus \{y\})) \ni y$ because $\text{cl}(\text{germ}_y([y, z] \setminus \{y\}))$ contains the closure of $\text{germ}_y([y, z] \setminus \{y\})$. This is absurd and thus $\mathbb{A} \cap x + \delta = y + \delta$, with $y \in \mathbb{A}$. One sets $y_\nu(x) = y \in \mathbb{A}$ (actually, y_ν only depends on δ).

One has $\rho_{+\infty}(x + \delta) = \rho_{+\infty}(x) + \delta$ and $y \in \rho_{+\infty}(x) + \delta$. We define $T_\nu(x)$ as the unique element T of \mathbb{R}_+ such that $y = \rho_{+\infty}(x) + T\nu$.

Let $\delta^- = -\mathbb{R}_+ \nu$ and $x \in \mathcal{I}$. Similarly, one defines y_ν^- as the first point of $x + \delta^-$ meeting \mathbb{A} and $T_\nu^-(x)$ as the element T of \mathbb{R}_+ such that $\rho_{-\infty}(x) = y + T\nu$.

Remark 3.2. In the following, the choice of ν will not be very important. We will often need to choose $\nu \in Y \cap C_f^v$.

Let $\mathcal{I}_0 = G.0$.

Lemma 3.3. *Let $x \in \mathcal{I}$ and $\nu \in C_f^v$. Let $y = y_\nu(x)$ and $T = T_\nu(x)$.*

a) Then $x \leq y$ and $d^v(x, y) = T\nu$.

b) One has $\rho_{+\infty}(x) \in Y$ if and only if $\rho_{-\infty}(x) \in Y$ if and only if $x \in \mathcal{I}_0$. In this case, $\rho_{+\infty}(x) \leq_{Q^v} \rho_{-\infty}(x)$.

Proof. Let A be an apartment containing x and $+\infty$ and $g \in G$ fixing $+\infty$ such that $A = g^{-1}.\mathbb{A}$. Then $x + \delta$ is the translate of a shortening $\delta' \subset A$ of δ (which means $\delta' = z + \delta$, with $z \in \delta$). As for all $z' \in z + \delta$, $z \leq z'$, one has $x \leq y$. As $d^v(x, y) = d^v(g.x, g.y)$ and $g|_A = \rho_{+\infty}$ one gets a).

For $x \in \mathcal{I}$, there exists $g_-, g_+ \in G$ such that $\rho_{-\infty}(x) = g_- . x$ and $\rho_{+\infty}(x) = g_+ . x$, which shows the claimed equivalence because $Y = G.0 \cap \mathbb{A}$.

Suppose $x \in \mathcal{I}_0$. One chooses $\nu \in Y \cap C_f^v$. Let $S = \lfloor T \rfloor + 1$, where $\lfloor \cdot \rfloor$ is the floor function, and $z = \rho_{+\infty}(x) + S\nu \in x + \delta$. Then $d^v(x, z) = d^v(g_+ . x, g_+ . z) = d^v(\rho_{+\infty}(x), z) = S\nu \in Y \cap C_f^v$.

According to paragraph 2.3 of [GR14], the image π of $[x, z]$ by $\rho_{-\infty}$ is a Hecke path of shape $z - \rho_{+\infty}(x) = S\nu$ with respect to $-C_f^v$ (unless the contrary is specified, "Hecke path" will mean with respect to $-C_f^v$). By applying Lemma 2.4b) of [GR14] to π , one gets that $z - \rho_{-\infty}(x) \leq_{Q^v} d^v(x, z) = z - \rho_{+\infty}(x)$ and thus $\rho_{+\infty}(x) \leq_{Q^v} \rho_{-\infty}(x)$ and one has b). \square

Lemma 3.4. *Let $\tau : [0, 1] \rightarrow \mathcal{I}$ be a segment of \mathcal{I} such that $\tau(1) \in \mathbb{A}$ and $\rho_{-\infty} \circ \tau$ is a segment of \mathbb{A} satisfying $(\rho_{-\infty} \circ \tau)' = \nu \in C_f^v$. Then $\tau([0, 1]) \subset \mathbb{A}$ and thus $\rho_{-\infty} \circ \tau = \tau$.*

Proof. Suppose $\tau([0, 1]) \not\subset \mathbb{A}$. Let $u = \sup\{t \in [0, 1] \mid \tau(u) \notin \mathbb{A}\}$. Then by the same reasoning as in the proof that " $x + \delta \cap \mathbb{A} = y + \delta$ " in the paragraph "Definition of y_ν and T_ν ", $x = \tau(u) \in \mathbb{A}$. One has $\tau(0) \leq x \leq \tau(1)$ (by the same reasoning as in the proof of Lemma 3.3 a)).

By Remark 2.6, there exists an apartment $A = g^{-1}.\mathbb{A}$ with $g \in G$ containing $\mathfrak{R} = -\infty$ and $\text{germ}_x([x, \tau(0)])$. By axiom (MA4) (and Remark 2.6) applied to $\mathfrak{R} = -\infty$ and to x , we can suppose that g fixes $\text{cl}(x, -\infty) \supset x - C_f^v$. Let $x' \in [x, \tau(0)] \setminus \{x\}$ such that $[x, x'] \subset A$ and $x' \notin \mathbb{A}$. Then $g.x' = \rho_{-\infty}(x') \in x - \mathbb{R}_+ \nu \subset x - C_f^v$. Therefore, $x' = g.x' \in \mathbb{A}$, which is absurd. Hence $\tau([0, 1]) \subset \mathbb{A}$. \square

4 Bounding of T_ν and Proof of Theorem 4.7

One defines $h : Q_{\mathbb{R}}^\vee \rightarrow \mathbb{R}$

$$x = \sum x_i \alpha_i^\vee \mapsto \sum x_i.$$

Lemma 4.1. *Let $T \in \mathbb{R}_+$, $\mu \in \mathbb{A}$, $a \in \mathbb{A}$, $\nu \in Y^{++} = Y \cap \overline{C_f^v}$ and suppose there exists a Hecke path π from a to $a + T\nu - \mu$ of shape $T\nu$. Then*

- a) $\mu \in Q_{\mathbb{R}_+}^\vee = \bigoplus_{i \in I} \mathbb{R}_+ \alpha_i^\vee$. Consequently $h(\mu)$ is well defined.
- b) if $T > h(\mu)$, there exists t such that π is derivable on $(t, 1]$ and $\pi'_{|(t,1]} = \nu$. Furthermore, let t^* be the smallest $t \in [0, 1]$ having this property, then $t^* \leq \frac{h(\mu)}{T}$.

Proof. The main idea of b) is to use the fact that during the time when $\pi'(t) \neq T\nu$, $\pi'(t) = T\nu - T\lambda(t)$ with $\lambda(t) \in Q_+^\vee \setminus \{0\}$. Hence for T large, π decreases quickly for the Q^\vee order, but it cannot decrease too much because μ is fixed.

Let $t_0 = 0, t_1, \dots, t_n = 1$ be a subdivision of $[0, 1]$ such that for all $i \in \llbracket 0, n-1 \rrbracket$, $\pi_{|(t_i, t_{i+1})}$ is derivable and let $w_i \in W^v$ be such $\pi'_{|(t_i, t_{i+1})} = w_i \cdot T\nu$. If $w_i \cdot \nu = \nu$, one chooses $w_i = 1$.

For $i \in \llbracket 0, n-1 \rrbracket$, according to Lemma 2.4 a) of [GR08], $w_i \cdot \nu = \nu - \lambda_i$, with $\lambda_i \in Q_+^\vee$ and if $w_i \neq 1$, $\lambda_i \neq 0$. One has

$$\pi(1) - \pi(0) = T\nu - \sum_{i=0, w_i \neq 1}^{n-1} (t_{i+1} - t_i) T\lambda_i = T\nu - \mu$$

and one deduces a).

Suppose now $T > h(\mu)$. Let us show that there exists $i \in \llbracket 0, n-1 \rrbracket$ such that $w_i = 1$. Let $i \in \llbracket 0, n-1 \rrbracket$.

For all i such that $w_i \neq 1$, one has $h(\lambda_i) \geq 1$. Hence $T \sum_{i=0, w_i \neq 1}^{n-1} (t_{i+1} - t_i) \leq h(\mu)$, and $\sum_{i=0, w_i \neq 1}^{n-1} (t_{i+1} - t_i) < 1 = \sum_{i=0}^{n-1} (t_{i+1} - t_i)$. Thus there exists $i \in \llbracket 0, n-1 \rrbracket$ such that $w_i = 1$.

By Remark 2.9, if $w_i = 1$ for some i , then $w_j = 1$ for all $j \geq i$. This shows the existence of t^* . We also have $t^* \leq \sum_{i=0, w_i \neq 1}^{n-1} (t_{i+1} - t_i)$ and hence the claimed inequality follows. \square

From now on and until the end of this subsection, ν will be a fixed element of $C_f^v \cap Y$. We define $T_\nu : \mathcal{I} \rightarrow \mathbb{R}_+$ and $y_\nu : \mathcal{I} \rightarrow \mathbb{A}$ as in the paragraph "Definition of y_ν and T_ν ".

Corollary 4.2. *Let $\mu \in Q_{\mathbb{R}}^\vee$. Then for all $x \in \mathcal{I}$ such that $\rho_{-\infty}(x) - \rho_{+\infty}(x) = \mu$, $T_\nu(x) \leq h(\mu)$.*

Proof. Let $x \in \mathcal{I}$ such that $\rho_{-\infty}(x) - \rho_{+\infty}(x) = \mu$. Let $y = y_\nu(x)$. Let π be the image by $\rho_{-\infty}$ of $[x, y]$. This is a Hecke path from $\rho_{-\infty}(x)$ to $y = \rho_{+\infty}(x) + T\nu$, of shape $T\nu$, with $T = T_\nu(x)$. The minimality of T and Lemma 3.4 imply that $\pi'(t) \neq \nu$ for all $t \in [0, 1]$, where π is derivable. By applying Lemma 4.1, we deduce that $T \leq h(\mu)$. \square

Remark 4.3. With the same reasoning but by considering Hecke with respect to C_f^v we obtain an analogous bounding for T_ν^- .

Corollary 4.4. *Let $x \in \mathcal{I}$ such that $\rho_{+\infty}(x) = \rho_{-\infty}(x)$. Then $x \in \mathbb{A}$. Therefore, $\forall z \in \mathbb{A}$, $\rho_{+\infty}^{-1}(\{z\}) \cap \rho_{-\infty}^{-1}(\{z\}) = \{z\}$.*

Proof. Let $x \in \mathcal{I}$ such that $\rho_{+\infty}(x) = \rho_{-\infty}(x)$. Then by Corollary 4.2, $T_\nu(x) = 0$ and thus $x \in \mathbb{A}$. \square

Corollary 4.5. *Let $x \in \mathcal{I}$, then $\rho_{-\infty}(x) - \rho_{+\infty}(x) \in Q_{\mathbb{R}_+}^\vee$.*

Proof. Let $y = y_\nu(x)$ and $T = T_\nu(x)$. By Lemma 3.3, $x \leq y$ and $d^v(x, y) = T\nu$. Therefore, if $\tau : [0, 1] \rightarrow \mathcal{I}$ is the segment going from x to y , its image π by $\rho_{-\infty}$ is a Hecke path of shape $T\nu$ from $\rho_{-\infty}(x)$ to $y = \rho_{+\infty}(x) + T\nu$. Therefore, Lemma 4.1a) applied to $a = \rho_{-\infty}(x)$ and $\mu = \rho_{-\infty}(x) - \rho_{+\infty}(x)$ completes the proof. \square

Lemma 4.6. *Let $x \in \mathcal{I}$ such that $y_\nu^-(x) \in C_f^v$. Then $0 \leq x$ and $\rho_{-\infty}(x) = d^v(0, x)$.*

Proof. Let $y^- = y_\nu^-(x)$. Let A be an apartment containing x and $-\infty$. By (MA4) there exists $g \in G$ such that $A = g^{-1}.\mathbb{A}$ and g fixes $\text{cl}(y, -\infty) \supset y - \overline{C_f^v} \ni 0$. Then $g.x - g.y^- = \rho_{-\infty}(x) - y^- = T^-\nu \in C_f^v$ and $g.y^- - g.0 = y^-$. Thus $g.x - g.0 = \rho_{-\infty}(x) \in C_f^v$ and we can conclude by Remark 2.7. \square

Theorem 4.7. *Let $\mu \in \mathbb{A}$. Then if $\mu \notin Q_{\mathbb{R}}^\vee$, $\rho_{+\infty}^{-1}(\{\lambda + \mu\}) \cap \rho_{-\infty}^{-1}(\{\lambda\})$ is empty for all $\lambda \in \mathbb{A}$. If $\mu \in Q_{\mathbb{R}}^\vee$, then for $\lambda \in \mathbb{A}$ sufficiently dominant, $\rho_{+\infty}^{-1}(\{\lambda + \mu\}) \cap \rho_{-\infty}^{-1}(\{\lambda\}) \subset S^v(0, \lambda) \cap \rho_{+\infty}^{-1}(\{\lambda + \mu\})$.*

Proof. The condition on μ comes from Corollary 4.5.

Suppose $\mu \in Q_{\mathbb{R}}^\vee$. Let $\lambda \in \mathbb{A}$. Let $y^- = y_\nu^-$ and $T^- = T_\nu^-$. By Corollary 4.2 and remark 4.3, if $x \in \rho_{+\infty}^{-1}(\{\lambda + \mu\}) \cap \rho_{-\infty}^{-1}(\{\lambda\})$, then $y^-(x) \in [\lambda - T^-(x)\nu, \lambda] \subset \lambda - [0, h(\mu)]\nu$. For all $i \in I$, $\alpha_i([0, h(\mu)]\nu)$ is bounded. Consequently for λ sufficiently dominant, $\alpha_i(\lambda - [0, h(\mu)]\nu) \subset \mathbb{R}_+^*$ for all $i \in I$. For such a λ , $y^-(\rho_{+\infty}^{-1}(\{\lambda + \mu\}) \cap \rho_{-\infty}^{-1}(\{\lambda\})) \subset C_f^v$. We conclude the proof with Lemma 4.6. \square

5 Study of "Translations" of \mathcal{I} and Proof of Theorem 5.6

Let $\mathbb{A}_{in} = \bigcap_{i \in I} \ker \alpha_i$.

In this subsection we introduce some kinds of "translation" of \mathcal{I} of an inessential vector and show that they have very nice properties. It will be useful to "generalize theorem from Y to $Y + \mathbb{A}_{in}$ " by getting rid of the inessential part. First example of this technique will be Corollary 5.3 which generalizes a theorem of [GR14]. Then we study elements of G inducing a translation on \mathbb{A} . We show that they commute with $\rho_{+\infty}$ and $\rho_{-\infty}$. We then can see that for fixed $\mu \in Q^\vee$, the $\rho_{+\infty}^{-1}(\{\lambda + \mu\}) \cap \rho_{-\infty}^{-1}(\{\lambda\})$, for $\lambda \in Y + \mathbb{A}_{in}$, are some translates of $\rho_{+\infty}^{-1}(\{\mu\}) \cap \rho_{-\infty}^{-1}(\{0\})$, which enables to show theorem 5.6 by using Theorem 4.7 and Corollary 5.3.

Lemma 5.1. *Let $\nu \in \mathbb{A}_{in}$ and $a \in \mathcal{I}$. Then $|S^v(a, \nu)| = 1$. Moreover, if $a \in \mathbb{A}$, $S^v(a, \nu) = \{a + \nu\}$.*

Proof. Let $h \in G$ such that $h.a \in \mathbb{A}$. Then $S^v(a, \nu) = h^{-1}.S^v(h.a, \nu)$. Therefore one can assume $a \in \mathbb{A}$.

Let $x \in S^v(a, \nu)$. Let $g \in G$ such that $x, a \in g^{-1}.\mathbb{A}$ and $g.x - g.a = \nu$. Let $\tau : [0, 1] \rightarrow \mathcal{I}$ defined by $\tau(t) = g^{-1}.(g.a + (1 - t)\nu)$. Then $\pi = \rho_{-\infty} \circ \tau$ is a Hecke path of shape $-\nu$ and in particular, it is a $-\nu$ -path. For all t where π is derivable, there exists $w(t) \in W^v$ such that $\pi'(t) = -w(t).\nu = -\nu$. As W^v acts trivially on \mathbb{A}_{in} , $\pi'(t) = -\nu$ for all such t and thus π is derivable on $[0, 1]$ and $\pi' = -\nu$. As $\tau(1) = a \in \mathbb{A}$, one can apply Lemma 3.4 and we get that $\tau([0, 1]) \subset \mathbb{A}$. Therefore, $x \in \mathbb{A}$ and there exists $w \in W^v$ such that $x - a = w.\nu = \nu$. \square

This lemma enables us to define some kinds of translation of an inessential vector. Let $\nu \in \mathbb{A}_{in}$. Let $\tau_\nu : \mathcal{I} \rightarrow \mathcal{I}$ which associates to $x \in \mathcal{I}$ the unique element of $S^v(x, \nu)$. Then we have the following lemma:

Lemma 5.2. *Let $\nu \in \mathbb{A}_{in}$ and $\tau = \tau_\nu$. Then:*

1. *For all $x \in \mathbb{A}$, $\tau(x) = x + \nu$.*

2. For all $g \in G$, $g \circ \tau = \tau \circ g$. In particular, for all $x \in \mathcal{I}$, if x is in an apartment A , then $\tau(x) \in A$, and if $x = g.a$ with $g \in G$ and $a \in \mathbb{A}$, then $\tau(x) = g.(x + \nu)$.
3. The application τ is a bijection, its inverse being $\tau_{-\nu}$.
4. The application τ commutes with $\rho_{+\infty}$ and $\rho_{-\infty}$.
5. Let $x \in \mathcal{I}$ and $\lambda \in \overline{C_f^v}$, then $\tau(S^v(x, \lambda)) = S^v(\tau(x), \lambda) = S^v(x, \lambda + \nu)$.

Proof. part 1 is a part of Lemma 5.1.

Choose $x \in \mathcal{I}$ and $g \in G$. Then $d^v(x, \tau(x)) = \nu$ and thus $d^v(g.x, g.\tau(x)) = \nu$. Consequently $g.\tau(x) = \tau(g.x)$ by definition of τ . Let A be an apartment containing x , $A = h.\mathbb{A}$, with $h \in G$. Then $\tau(x) = \tau(h.a)$ with $a \in \mathbb{A}$, hence $\tau(x) = h.(x + \nu) \in A = h.\mathbb{A}$ and we have 2.

Let $x \in \mathcal{I}$, $x = g.a$ with $a \in \mathbb{A}$. By part 2 applied to τ and $\tau_{-\nu}$, one has $\tau_{-\nu}(\tau(g.a)) = g.\tau_{-\nu}(\tau(a)) = g.a$ and thus $\tau_{-\nu} \circ \tau = \text{Id}$. This is enough to show 3.

Let $x \in \mathcal{I}$ and $g \in G$ fixing $+\infty$ such that $g.x = \rho_{+\infty}(x)$. Then $\tau(x) \in g^{-1}.\mathbb{A}$, thus $g.\tau(x) = \rho_{+\infty}(\tau(x))$ and by part 2, $g.\tau(x) = \tau(g.x)$. Hence, τ and $\rho_{+\infty}$ commute and by the same reasoning, this is also true for τ and $\rho_{-\infty}$.

Let $x \in \mathcal{I}$ and $\lambda \in \overline{C_f^v}$. Let $u \in S^v(x, \lambda)$. There exists $g' \in G$ such that $x, u \in g'^{-1}.\mathbb{A}$ and $g'.u - g'.x = w.\lambda$, for some $w \in W^v$. Let $n \in N$ inducing w on \mathbb{A} and $g = n^{-1}g'$. Then $x, u \in g^{-1}.\mathbb{A}$ and $g.u - g.x = \lambda$. We have $g.\tau(u) - g.x = \tau(g.u) - g.x = \lambda + \nu$. Therefore, $\tau(S^v(x, \lambda)) \subset S^v(x, \lambda + \nu)$. Applying this result with $\tau_{-\nu}$ yields $\tau_{-\nu}(S^v(x, \lambda + \nu)) \subset S^v(x, \lambda)$ and thus $\tau(S^v(x, \lambda)) = S^v(x, \lambda + \nu)$.

One has $g.\tau(u) - g.\tau(x) = (g.u + \nu) - (g.x + \nu) = \lambda$ and thus $\tau(S^v(x, \lambda)) \subset S^v(\tau(x), \lambda)$. Again, by considering $\tau_{-\nu}$, we have that $\tau(S^v(x, \lambda)) = S^v(\tau(x), \lambda)$. \square

Corollary 5.3. *Let $\lambda \in \mathbb{A}$ and $\mu \in Y + \mathbb{A}_{in}$. Then for all $\lambda_{in} \in \mathbb{A}_{in}$, $\tau_{\lambda_{in}}(S^v(0, \lambda) \cap \rho_{+\infty}^{-1}(\{\mu\})) = S^v(0, \lambda + \lambda_{in}) \cap \rho_{+\infty}^{-1}(\{\mu + \lambda_{in}\})$. In particular, for all $\lambda \in Y + \mathbb{A}_{in}$, $S^v(0, \lambda) \cap \rho_{+\infty}^{-1}(\{\mu + \lambda\})$ is finite and is empty if $\mu \notin Q_-^v$.*

Proof. The first assertion is a consequence of Lemma 5.2 part 3, 4 and 5.

Let $\lambda \in Y + \mathbb{A}_{in}$ and $\lambda_{in} \in \mathbb{A}_{in}$ such that $\tau(\lambda) \in Y$, with $\tau = \tau_{\lambda_{in}}$. Then $\tau(S^v(0, \lambda) \cap \rho_{+\infty}^{-1}(\{\mu + \lambda\})) = S^v(0, \tau(\lambda)) \cap \rho_{+\infty}^{-1}(\{\tau(\mu + \lambda)\})$. Consequently, one can assume $\lambda \in Y$.

Suppose $S^v(0, \lambda) \cap \rho_{+\infty}^{-1}(\{\mu + \lambda\})$ is nonempty and let x be in this set. Then there exists $g, h \in G$ such that $g.x = \lambda$ and $h.x = \mu + \lambda$. Thus $\lambda + \mu = h.g^{-1}.\lambda \in \mathcal{I}_0 \cap \mathbb{A} = Y$ and therefore, $\mu \in Y$. We can now conclude because the finiteness and the condition on μ are shown in [GR14], Section 5: if $\lambda, \mu \in Y$ then $S^v(0, \lambda) \cap \rho_{+\infty}^{-1}(\{\mu\})$ is finite and it is empty if $\mu \notin Q_-^v$ (the cardinals of these sets correspond to the $n_\lambda(\nu)$ of loc. cit.). \square

We now show a lemma similar to Lemma 5.2 part 4 for translations of G :

Lemma 5.4. *Let $n \in G$ inducing a translation on \mathbb{A} . Then $n \circ \rho_{+\infty} = \rho_{+\infty} \circ n$ and $n \circ \rho_{-\infty} = \rho_{-\infty} \circ n$.*

Proof. Let $x \in \mathcal{I}$ and A be an apartment containing x and $+\infty$. Then $n.A$ is an apartment containing $+\infty$. Let $\phi : A \rightarrow \mathbb{A}$ an isomorphism fixing $+\infty$. We have $n.x \in n.\mathbb{A}$, and $n \circ \phi \circ n^{-1} : n.A \rightarrow \mathbb{A}$ fixes $+\infty$. Hence $\rho_{+\infty}(n.x) = n \circ \phi \circ n^{-1}(n.x) = n \circ \phi(x) = n \circ \rho_{+\infty}(x)$ and thus $n \circ \rho_{+\infty} = \rho_{+\infty} \circ n$. By the same reasoning applied to $\rho_{-\infty}$, we get the lemma. \square

Lemma 5.5. *Let $n \in G$ inducing a translation on \mathbb{A} . Let $\lambda_{in} \in \mathbb{A}_{in}$. Set $\tau = \tau_{\lambda_{in}} \circ n$. Let $\nu \in C_f^v$ and $y^- = y_\nu^-$. Then $\tau \circ y^- = y^- \circ \tau$.*

Proof. If $x \in \mathbb{A}$, then $y^-(x) = x$, $y^-(\tau(x)) = \tau(x)$ and there is nothing to prove.

Suppose $x \notin \mathbb{A}$. Then $[x, y^-(x)] \setminus \{y^-(x)\} \subset (x - \mathbb{R}_+\nu) \setminus \mathbb{A}$, thus $\tau([x, y^-(x)] \setminus \{y^-(x)\}) \subset (\tau(x) - \mathbb{R}_+\nu) \setminus \mathbb{A}$ and $\tau(y^-(x)) \in \mathbb{A}$. \square

Theorem 5.6. *Let $\mu \in \mathbb{A}$ and $\lambda \in Y + \mathbb{A}_{in}$. One writes $\lambda = \lambda_{in} + \Lambda$, with $\lambda_{in} \in \mathbb{A}_{in}$ and $\Lambda \in Y$. Let $n \in G$ inducing the translation of vector Λ and $\tau = \tau_{\lambda_{in}} \circ n$. Then $\rho_{+\infty}^{-1}(\{\lambda + \mu\}) \cap \rho_{-\infty}^{-1}(\{\lambda\}) = n(\rho_{+\infty}^{-1}(\{\mu\}) \cap \rho_{-\infty}^{-1}(\{0\}))$. Therefore these sets are finite and if $\mu \notin Q_-^\vee$ these sets are empty.*

Proof. First assertion is a consequence of Lemma 5.4 and Lemma 5.2 part 4. Then Lemma 3.3 b) shows that these sets are empty unless $\mu \in Q_+^\vee$.

By Theorem 4.7, for $\lambda' \in Y$ sufficiently dominant, $\rho_{+\infty}^{-1}(\{\lambda' + \mu\}) \cap \rho_{-\infty}^{-1}(\{\lambda'\}) \subset S^v(0, \lambda') \cap \rho_{+\infty}^{-1}(\{\lambda' + \mu\})$, which is a finite set by [GR14] Section 5 (or by Corollary 5.3). \square

6 Proof of Theorem 6.4

Recall that we want to prove Theorem 6.4:

Let $\mu \in Q_-^\vee$. Then for $\lambda \in Y^{++} + \mathbb{A}_{in}$ sufficiently dominant, $S^v(0, \lambda) \cap \rho_{+\infty}^{-1}(\{\lambda + \mu\}) = \rho_{-\infty}^{-1}(\{\lambda\}) \cap \rho_{+\infty}^{-1}(\{\lambda + \mu\})$.

The basic idea of the proof of this theorem, is that if $\mu \in Q_-^\vee$ there exists a finite set $F \subset Y^{++}$ such that for all $\lambda \in Y^{++} + \mathbb{A}_{in}$, $\rho_{+\infty}^{-1}(\{\lambda + \mu\}) \cap S^v(0, \lambda) \subset \bigcup_{f \in F} \rho_{+\infty}^{-1}(\{\lambda + \mu\}) \cap S^v(\lambda - f, f)$ (this is Corollary 6.3, which generalizes Lemma 6.1 and uses Section 2.2). Then we use Subsection 5 to show that $\rho_{+\infty}^{-1}(\{\lambda + \mu\}) \cap S^v(\lambda - f, f)$ is the image of $\rho_{+\infty}^{-1}(\mu + f) \cap S^v(0, f)$ by a "translation" $\tau_{\lambda-f}$ of G of vector $\lambda - f$ (which means that $\tau_{\lambda-f}$ induces the translation of vector $\lambda - f$ on \mathbb{A}). We fix $\nu \in C_f^v$ and set $y^- = y_\nu^-$. By Lemma 5.5,

$$y^-(\rho_{+\infty}^{-1}(\{\lambda + \mu\}) \cap S^v(0, \lambda)) \subset \bigcup_{f \in F} \tau_{\lambda-f} \circ y^-(S^v(0, f) \cap \rho_{+\infty}^{-1}(\{\mu + f\})).$$

According to Section 5 of [GR14], for all $f, \mu \in Y$, $S^v(0, f) \cap \rho_{+\infty}^{-1}(\{\mu + f\})$ is finite. Consequently, for λ sufficiently dominant, $\bigcup_{f \in F} \tau_{\lambda-f}(y^-(S^v(0, f) \cap \rho_{+\infty}^{-1}(\{\mu + f\}))) \subset C_f^v$ and one concludes with Lemma 4.6.

Lemma 6.1. *Let $\mu \in Q_-^\vee$ and $H = -h(\mu) + 1 \in \mathbb{N}$. Let $a \in Y$, $T \in [H, +\infty)$, $\nu \in Y^{++}$ and $x \in S^v(a, T\nu) \cap \rho_{+\infty}^{-1}(\{a + T\nu + \mu\})$. Let $g \in G$ such that $g.a = a$ and $g.x = T\nu + a$. Then g fixes $[a, a + (T - H)\nu]$ and in particular $x \in S^v(a + (T - H)\nu, H\nu)$.*

Proof. Let $\tau : [0, 1] \rightarrow \mathbb{A}$ defined by $\tau(t) = a + (1 - t)T\nu$. The main idea is to apply Lemma 4.1 to $\rho_{+\infty} \circ (g^{-1}.\tau)$ but we cannot do it directly because $\rho_{+\infty} \circ \tau$ is not a Hecke path with respect to $-C_f^v$. Let \mathbb{A}' be the vectorial space \mathbb{A} equipped with a structure of apartment of $-\mathbb{A}$: the fundamental chamber of \mathbb{A}' is $C_f'^v = -C_f^v$ etc ... Let \mathcal{I}' be the set \mathcal{I} , whose apartments are the $-A$ where A runs over the apartment of \mathcal{I} . Then \mathcal{I}' is a hovel of standard apartment \mathbb{A}' . We have $a \leq x$ in \mathcal{I} and so $x \leq' a$ in \mathcal{I}' .

Then the image π of $g^{-1}.\tau$ by $\rho_{+\infty}$ is a Hecke path of shape $-T\nu$ from $\rho_{+\infty}(x) = a + T\nu + \mu$ to a . By Lemma 4.1 (for \mathcal{I}'), for $t > -h(\mu)/T$, $\pi'(t) = -T\nu$, and thus Lemma 3.4 (for \mathcal{I}' and \mathbb{A}') implies $\rho_{+\infty}(g^{-1}.\tau(t)) = g^{-1}.\tau(t)$ for all $t > -h(\mu)/T$. Therefore, $g^{-1}.\tau|_{(-h(\mu)/T, 1]}$ is a segment in \mathbb{A} ending in a , with derivative $-T\nu$ and thus, for all $t \in (-h(\mu)/T, 1]$, $g^{-1}.\tau(t) = \tau(t)$. In particular, g fixes $[a, \tau(\frac{H}{T})] = [a, a + (T - H)\nu]$, and $d^v(a + (T - H)\nu, x) = d^v(g^{-1}.(a + (T - H)\nu), g^{-1}.x) = d^v(a + (T - H)\nu, a + T\nu) = H\nu$. \square

Let E be as in the Lemma 2.3.

Lemma 6.2. *Let $\mu \in Q_-^\vee$, $H = -h(\mu) + 1 \in \mathbb{N}$ and $a \in Y$. Let $\lambda \in Y^{++}$. One writes $\lambda = \sum_{e \in E} \lambda_e e$ with $\lambda_e \in \mathbb{N}$ for all $e \in E$. Let $e \in E$. Then if $\lambda_e \geq H$, $S^v(a, \lambda) \cap \rho_{+\infty}^{-1}(\{a + \lambda + \mu\}) \subset S^v(a + (\lambda_e - H)e, \lambda - (\lambda_e - H)e)$.*

Proof. Let $x \in S^v(a, \lambda) \cap \rho_{+\infty}^{-1}(\{a + \lambda + \mu\})$ and $g \in G$ fixing a such that $g.x = a + \lambda$. Let $z = g^{-1}(a + \lambda_e e)$.

Then one has $d^v(a, z) = \lambda_e e$ and $d^v(z, x) = \lambda - \lambda_e e$.

According to Lemma 2.4.b) of [GR14] (adapted because one considers Hecke paths with respect to C_f^v), one has:

$$\rho_{+\infty}(z) - a \leq_{Q^\vee} d^v(a, z) = \lambda_e e \text{ and } \rho_{+\infty}(x) - \rho_{+\infty}(z) \leq_{Q^\vee} d^v(z, x) = \lambda - \lambda_e e.$$

Therefore,

$$a + \lambda + \mu = \rho_{+\infty}(x) \leq_{Q^\vee} \rho_{+\infty}(z) + \lambda - \lambda_e e \leq_{Q^\vee} a + \lambda.$$

Hence, $\rho_{+\infty}(z) = a + \lambda_e e + \mu'$, with $\mu \leq_{Q^\vee} \mu' \leq_{Q^\vee} 0$. One has $-h(\mu') + 1 \leq H$. By Lemma 6.1, g fixes $[a, a + (\lambda_e - H)e]$ and thus g fixes $a + (\lambda_e - H)e$.

As $d^v(g^{-1}.(a + (\lambda_e - H)e), x) = \lambda - (\lambda_e - H)e$, $x \in S^v(a + (\lambda_e - H)e, \lambda - (\lambda_e - H)e)$. \square

Corollary 6.3. *Let $\mu \in Q_-^\vee$. Let $H = -h(\mu) + 1$. Let $\lambda \in Y^{++}$. We fix a writing $\lambda = \sum_{e \in E} \lambda_e e$, with $\lambda_e \in \mathbb{N}$ for all $e \in E$. Let $J = \{e \in E \mid \lambda_e \geq H\}$. Then $S^v(0, \lambda) \cap \rho_{+\infty}^{-1}(\{\lambda + \mu\}) \subset S^v(\lambda - H \sum_{e \in J} e - \sum_{e \notin J} \lambda_e e, H \sum_{e \in J} e + \sum_{e \notin J} \lambda_e e)$.*

Proof. this is a generalization by induction of Lemma 6.2. \square

Theorem 6.4. *Let $\mu \in Q^\vee$. Then for $\lambda \in Y^{++} + \mathbb{A}_{in}$ sufficiently dominant, $S^v(0, \lambda) \cap \rho_{+\infty}^{-1}(\{\lambda + \mu\}) = \rho_{-\infty}^{-1}(\{\lambda\}) \cap \rho_{+\infty}^{-1}(\{\lambda + \mu\})$.*

Proof. Theorem 4.7 yields one inclusion. It remains to show that $\rho_{+\infty}^{-1}(\{\lambda + \mu\}) \cap S^v(0, \lambda) \subset \rho_{+\infty}^{-1}(\{\lambda + \mu\}) \cap \rho_{-\infty}^{-1}(\{\lambda\})$ for λ sufficiently dominant.

Let $H = -h(\mu) + 1$ and $F = \{\sum_{e \in E} \nu_e e \mid (\nu_e) \in \llbracket 0, H - 1 \rrbracket^E\}$. This set is finite. Let $\lambda \in Y^{++} + \mathbb{A}_{in}$, $\lambda = \lambda_{in} + \Lambda$, with $\lambda_{in} \in \mathbb{A}_{in}$ and $\Lambda \in Y^{++}$.

Let $x \in S^v(0, \lambda) \cap \rho_{+\infty}^{-1}(\{\lambda + \mu\})$. Then by Lemma 6.3, there exists $f \in F$ such that $\lambda - f \in \overline{C_f^v}$ and $x \in S^v(\lambda - f, f)$. Let n an element of G inducing the translation of vector $\Lambda - f = \lambda - \lambda_{in} + f$ on \mathbb{A} and $\tau_{\lambda, f} = \tau_{\lambda_{in}} \circ n$. Then $x \in \tau_{\lambda, f}(B_f)$ where $B_f = S^v(0, f) \cap \rho_{+\infty}^{-1}(\{\mu + f\})$.

Let $B = \bigcup_{f \in F} B_f$. Then one has proven that

$$S^v(0, \lambda) \cap \rho_{+\infty}^{-1}(\{\lambda + \mu\}) \subset \bigcup_{f \in F} \tau_{\lambda, f}(B).$$

By Section 5 of [GR14] (or Corollary 5.3), B_f is finite for all $f \in F$ and thus $B = \bigcup_{f \in F} B_f$ is finite. Let $\nu \in C_f^v$ and $y^- = y_\nu^-$. Then $y^-(B)$ is finite and for λ sufficiently dominant, $\bigcup_{f \in F} \tau_{\lambda, f} \circ y^-(B) \subset C_f^v$. Moreover, according to Lemma 5.5, $\bigcup_{f \in F} \tau_{\lambda, f} \circ y^-(B) = \bigcup_{f \in F} y^- \circ \tau_{\lambda, f}(B)$. Hence

$$y^-(S^v(0, \lambda) \cap \rho_{+\infty}^{-1}(\{\lambda + \mu\})) \subset C_f^v$$

for λ sufficiently dominant. Eventually one concludes with Lemma 4.6. \square

References

- [BGKP14] Alexander Braverman, Howard Garland, David Kazhdan, and Manish Patnaik. An affine Gindikin-Karpelevich formula. *Perspectives in Representation Theory (Yale University, May 12–17, 2012)*(P. Etingof, M. Khovanov, and A. Savage, eds.), *Contemp. Math*, 610:43–64, 2014.
- [BKP14] Alexander Braverman, David Kazhdan, and Manish M Patnaik. Iwahori–hecke algebras for p-adic loop groups. *Inventiones mathematicae*, pages 1–96, 2014.
- [GR08] Stéphane Gaussent and Guy Rousseau. Kac-Moody groups, hovels and Littelmann paths. In *Annales de l’institut Fourier*, volume 58, pages 2605–2657, 2008.
- [GR14] Stéphane Gaussent and Guy Rousseau. Spherical Hecke algebras for Kac-Moody groups over local fields. *Annals of Mathematics*, 180(3):1051–1087, 2014.
- [Kac94] Victor G Kac. *Infinite-dimensional Lie algebras*, volume 44. Cambridge university press, 1994.
- [KM08] Michael Kapovich and John J. Millson. A path model for geodesics in Euclidean buildings and its applications to representation theory. *Groups Geom. Dyn.*, 2(3):405–480, 2008.
- [Lan71] Robert P Langlands. *Euler products*. Yale University Press New Haven, 1971.
- [Rou11] Guy Rousseau. Measures affines. *Pure and Applied Mathematics Quarterly*, 7(3):859–921, 2011.
- [Rou12] Guy Rousseau. Almost split Kac-Moody groups over ultrametric fields. *arXiv preprint arXiv:1202.6232*, 2012.