



HAL
open science

Common variation in PHACTR1 is associated with susceptibility to cervical artery dissection.

Stéphanie Debette, Yoichiro Kamatani, Tiina M Metso, Manja Kloss, Ganesh Chauhan, Stefan T Engelter, Alessandro Pezzini, Vincent Thijs, Hugh S Markus, Martin Dichgans, et al.

► **To cite this version:**

Stéphanie Debette, Yoichiro Kamatani, Tiina M Metso, Manja Kloss, Ganesh Chauhan, et al.. Common variation in PHACTR1 is associated with susceptibility to cervical artery dissection.. *Nature Genetics*, 2014, 47 (1), pp.78-83. 10.1038/ng.3154 . hal-01253336

HAL Id: hal-01253336

<https://hal.science/hal-01253336>

Submitted on 16 Feb 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Published in final edited form as:

Nat Genet. 2015 January ; 47(1): 78–83. doi:10.1038/ng.3154.

Common variation in *PHACTR1* is associated with susceptibility to cervical artery dissection

A full list of authors and affiliations appears at the end of the article.

Abstract

Cervical artery dissection (CeAD), a mural hematoma in a carotid or vertebral artery, is a major cause of ischemic stroke in young adults although relatively uncommon in the general population (incidence of 2.6/100,000 per year)¹. Minor cervical traumas, infection, migraine and hypertension are putative risk factors^{1–3}, and inverse associations with obesity and hypercholesterolemia are described^{3,4}. No confirmed genetic susceptibility factors have been identified using candidate gene approaches⁵. We performed genome-wide association studies (GWAS) in 1,393 CeAD cases and 14,416 controls. The rs9349379[G] allele (*PHACTR1*) was associated with lower CeAD risk (odds ratio (OR) = 0.75, 95% confidence interval (CI) = 0.69–0.82; $P = 4.46 \times 10^{-10}$), with confirmation in independent follow-up samples (659 CeAD cases and 2,648 controls; $P = 3.91 \times 10^{-3}$; combined $P = 1.00 \times 10^{-11}$). The rs9349379[G] allele was previously shown to be associated with lower risk of migraine and increased risk of myocardial infarction^{6–9}. Deciphering the mechanisms underlying this pleiotropy might provide important information on the biological underpinnings of these disabling conditions.

We organized an international initiative with the aim of collecting clinical data and DNA for the largest possible number of CeAD cases to identify genetic susceptibility loci. We obtained 942 CeAD cases from the Cervical Artery Dissections and Ischemic Stroke Patients (CADISP) study (CADISP-1: 170 Finns and 772 non-Finnish Europeans)¹⁰. An additional 451 CeAD cases of European origin were recruited specifically for the CADISP-genetics project at European and US centers (CADISP-2). These collections provided a total of 1,393

Reprints and permissions information is available online at <http://www.nature.com/reprints/index.html>.

Correspondence should be addressed to S.D. (stephdebette@wanadoo.fr).

⁶⁶A list of members and affiliations appears in the Supplementary Note.

⁶⁷These authors contributed equally to this work.

Note: Any Supplementary Information and Source Data files are available in the online version of the paper.

AUTHOR CONTRIBUTIONS

Study conception and design were performed by S.D., Y.K., M.L., D.L., P.A. and J.D. Acquisition of data was carried out by S.D., T.M.M., M.K., S.T.E., A. Pezzini, V.T., H.S.M., M.D., R.D., E.T., A.M.S., Y.S., S.A., Y.B., V.C., A.B., A.G., M.S., J.C., C. Lamy, E.M., S.B., L.H.B., A.J.G., P.M., J.J.M., P.S., E.B., B.G., E.G.v.d.H., I.F.-C., K.J., I.W., M.A.N., F.-E.D.L., C.J., Y.-C.C., A.J.M., C. Lichy, L.D., L.K., M.N., P.A.L., T. Brandt, G.B.B., H.-E.W., C.G., T. Böttcher, M.C., D.A., M.A.I., M.M.B.B., A. Padovani, J.F.M., G.K., A.R., B.B.W., E.-B.R., D.Z., T.T., M.L., D.L., P.A. and J.D. Statistical analysis and interpretation of the data were performed by S.D., Y.K., C.W., Y.-C.C., G.C., M.L., P.A. and J.D. The manuscript was drafted by S.D., Y.K., M.L., P.A. and J.D. Critical revision of the manuscript was performed by S.D., Y.K., T.M.M., S.T.E., C.W., M.L., P.A., J.D., A. Pezzini, V.T., H.S.M., E.T., A.M.S., J.C., J.J.M., P.S., I.F.-C., A.J.M., P.A.L., M.A.I., D.Z., T.T., M.L., D.L., P.A. and J.D. Annotation for expression quantitative trait loci was performed by A.D.J. Funding was obtained by S.D., S.T.E., A. Pezzini, V.T., H.S.M., M.D., S.B., A.J.G., P.M., J.J.M., P.S., B.G., F.-E.D.L., C.J., P.A.L., G.B.B., H.-E.W., M.C., D.A., M.M.B.B., J.F.M., A.R., B.B.W., E.-B.R., D.Z., T.T., M.L., D.L., P.A. and J.D.

COMPETING FINANCIAL INTERESTS

The authors declare no competing financial interests.

CeAD cases of European ancestry for the GWAS (discovery) phase. All CeAD cases were ascertained through departments of neurology specialized in stroke care (Supplementary Figs. 1–3 and Supplementary Note; [ClinicalTrials.gov](https://clinicaltrials.gov/ct2/show/study/NCT00657969) identifier NCT00657969)¹¹. DNA from 14,416 controls was available (287 Finns and 14,129 non-Finnish Europeans). We obtained 659 additional CeAD cases that could not be included in the GWAS because of the timing of their inclusion or the availability of only limited amounts of DNA. These cases were used along with 2,648 controls for follow-up of a small number of GWAS results. Finally, to examine the disease specificity of the genetic associations, we recruited 583 individuals with an ischemic stroke attributable to other causes (non-CeAD ischemic stroke: 162 Finns and 421 non-Finnish Europeans) who had similar age, sex and geographical origin characteristics as the CeAD cases. The clinical characteristics of the research subjects are shown in Table 1.

DNA samples were genotyped on an Illumina Human610-Quad or Human660W-Quad BeadChip. We performed imputation to the non-monomorphic SNPs described in the HapMap 2 and 1000 Genomes Project (August 2010 release) CEU (European-ancestry) panels. CADISP-1 and CADISP-2 data were analyzed separately because genotyping was carried out on different platforms (Supplementary Table 1). Moreover, because Finnish participants had a distinct ancestral origin (Supplementary Fig. 4), CADISP-1 was divided into CADISP-1 non-Finnish and CADISP-1 Finnish studies. We performed a fixed-effects inverse variance-weighted meta-analysis after applying a genomic control correction to each of the three GWAS results (CADISP-1 non-Finnish, CADISP-1 Finnish and CADISP-2; Supplementary Note). The quantile-quantile plot for the CeAD GWAS is shown in Supplementary Figure 5. We observed no overall inflation of P values or evidence for significant population stratification (genomic inflation factor $\lambda = 1.032$).

A Manhattan plot of the meta-analysis association results for genotyped SNPs is shown in Supplementary Figure 6 (and in Supplementary Fig. 7 by substudy). The evidence of association with CeAD reached genome-wide significance for two SNPs at two loci. These were rs9349379[G] in intron 2 of *PHACTR1* on chromosome 6p24.1-p23 (OR = 0.75, 95% CI = 0.69–0.82; $P = 4.46 \times 10^{-10}$) and rs11172113[C] in intron 1 of *LRPI* on chromosome 12q13.3 (OR = 0.78, 95% CI = 0.71–0.85; $P = 4.22 \times 10^{-8}$). Overall, we found 77 SNPs (at 51 loci) with association $P < 1 \times 10^{-4}$ (Supplementary Table 2) and 6 SNPs (at 5 loci) with association $P < 1 \times 10^{-5}$ (Table 2).

We restricted further analyses to loci with association $P < 1 \times 10^{-5}$. Two of these SNPs showed nominally significant heterogeneity in effect according to dissection site ($P = 6.49 \times 10^{-3}$ for rs1466535 (*LRPI*) and $P = 0.038$ for rs6820391 (*LNXI*)), with stronger associations for carotid than vertebral dissection (Table 3). None of these markers were associated with non-CeAD ischemic stroke (Supplementary Table 3).

We also identified two loci that were not included in the GWAS genotyping panel but reached genome-wide significance when imputed: rs2163474 in *CCDC102B* on chromosome 18q22.1 (OR = 1.78, 95% CI = 1.50–2.11; $P = 3.86 \times 10^{-11}$) and rs9915775 on chromosome 17q21.1 (OR = 3.39, 95% CI = 2.21–5.19; $P = 1.97 \times 10^{-8}$). Neither showed significant heterogeneity in effect by dissection site ($P = 0.72$ and 0.24, respectively), nor

were they associated with non-CeAD ischemic stroke (Supplementary Table 3). We removed rs9915775 from further consideration because of its low estimated minor allele frequency (MAF = 0.01) and relatively poor imputation quality ($R^2 = 0.59$).

We examined 6 SNP markers at the 5 most significantly associated loci from the genotyped panel ($P < 1 \times 10^{-5}$ in the GWAS) in the 659 CeAD case and 2,648 control samples available for follow-up (Supplementary Table 4). In addition, we selected two SNPs from the imputed locus at 18q22.1 for genotyping in follow-up samples (Supplementary Table 5). We also genotyped two proxies for two of the most significantly associated SNPs from the GWAS meta-analysis (rs12215208, $R^2 = 0.39$ with rs9349379; rs6761601, $R^2 = 0.67$ with rs6741522).

First, we calculated association statistics for the follow-up samples, using a threshold of $P = 5.00 \times 10^{-3}$ to assign significance (Bonferroni correction for ten SNPs; a conservative threshold because of the existence of positive linkage disequilibrium (LD) between some markers); irrespective of significance, we calculated combined association test statistics using meta-analysis and evaluated evidence of heterogeneity between the GWAS and follow-up data (Table 3 and Supplementary Table 5). Second, we applied a Bayesian approach to evaluate the significance of the associations using Wakefield's approximate Bayes factor (Online Methods)¹². Asymptotic Bayes factor (ABF) and Bayesian false discovery probability (BFDP) values are provided in Supplementary Tables 6 and 7.

rs9349379 (in *PHACTR1*), which provided the strongest evidence of association in the GWAS data, showed significant association with CeAD in the follow-up sample using the Bonferroni-corrected P -value threshold described above ($P = 5.00 \times 10^{-3}$), and the result from the meta-analysis of the discovery and follow-up studies maintained the finding of genome-wide significance (OR = 0.77, 95% CI = 0.72–0.83; $P = 1.00 \times 10^{-11}$) (Fig. 1 and Supplementary Fig. 8). Application of the Wakefield Bayesian approach yielded an ABF of 1.70×10^{-10} . Another SNP in *PHACTR1*, rs12215208, selected as a proxy ($P = 2.52 \times 10^{-5}$ in the GWAS), exhibited a nominally significant association in the follow-up samples ($P = 0.015$), but the combined GWAS and follow-up data did not show genome-wide significant association (Supplementary Table 5). In contrast, we found no evidence of association in follow-up samples at rs11172113 in *LRPI*, the second genotyped marker that had genome-wide significant association in the GWAS (Supplementary Fig. 8).

Of the other markers that we examined, one met the criteria for significance in the follow-up samples: rs6820391 in *LNX1* on chromosome 4q12 gave $P = 9.28 \times 10^{-4}$ in the follow-up samples and reached genome-wide significance in the combined meta-analysis (OR = 1.24, 95% CI = 1.15–1.34; $P = 2.36 \times 10^{-8}$) (Supplementary Fig. 8). However, as the Wakefield Bayesian approach yielded an ABF of 3.25×10^{-6} , we consider this association to be only suggestive, requiring further studies for confirmation. Data for the imputed 18q22.1 locus provided no evidence in favor of association in the follow-up collection ($P > 0.15$; Supplementary Table 5).

In sensitivity analyses, stratifying on sex, migraine status and recent cervical trauma status did not modify the genetic associations with CeAD for the top loci (Supplementary Tables

8–10). We observed a nominally significant association of the rs9349379 (*PHACTRI*) risk allele for CeAD with a younger age of onset (Supplementary Table 11). The effect of rs9349379 was more marked in individuals with CeAD without ischemia, whereas the effect of rs6820391 (*LNXI*) was more marked in individuals with CeAD with ischemia, although the confidence intervals largely overlapped and *P* values were similar (Supplementary Table 12). Secondary analyses of association performed separately for carotid and vertebral artery dissections did not yield any genome-wide significant association (Supplementary Table 13).

We did not find any association with CeAD for (i) SNPs reported to be associated with CeAD in candidate gene association studies^{13–15}; (ii) SNPs within a 100-kb window from the start and end of *COL3A1*, the gene harboring causal mutations for vascular Ehlers-Danlos syndrome, a rare etiology of CeAD¹⁶; and (iii) published genome-wide susceptibility SNPs for intracranial aneurysms^{17–19} and for thoracic aortic aneurysms and dissection²⁰ (Supplementary Tables 14–17).

Our most significant association with CeAD was for rs9349379 (*PHACTRI*), which has been associated with myocardial infarction and coronary calcifications in various ancestry groups^{6,7,21–26}, with effects in the opposite direction of that in CeAD. Two other susceptibility loci for myocardial infarction showed significant associations with CeAD in our data set with effects in the opposite direction of those for CeAD (rs2023938 (*HDAC9*)⁷ and rs9982601 (*SLC5A3*, *MRPS6* and *KCNE2*)⁶), and one susceptibility locus for myocardial infarction at 9p21 was associated with CeAD with the same direction of effect (rs3217992 (*CDKN2B-AS1*)⁷) (Supplementary Table 18). No significant association was found between rs9349379 and non-CeAD ischemic stroke in our data set, although such an association was recently described in a larger sample for the ischemic stroke subtype secondary to large artery atherosclerosis (LAA-IS), where the effect was in the opposite direction to the one we observed for CeAD²⁷. Another variant predisposing to LAA-IS (rs11984041 (*HDAC9*)^{28,29}; $R^2 = 1$ with rs2023938) also displayed an inverse association with CeAD (Supplementary Table 19). Opposite effects of the same genetic variant on different diseases have been described elsewhere³⁰ and suggest either that the same region harbors different causal variants or that the same causal variant has biological effects with opposite implications for each disease. The vascular risk factor profile^{3,4}, young age of occurrence¹ and heterogeneous echostructure of carotid arteries in individuals with CeAD³¹ all suggest that atherosclerosis is not a predisposing condition for CeAD, in contrast with aortic dissection³². With aging and arteriosclerosis, the increased synthesis and reduced degradation of extracellular matrix components, as well as increasing collagen and elastin cross-links, could be hypothesized to make the arterial walls of cervical arteries more resistant to tears^{33,34}, as could arterial wall calcifications, thus rendering the artery more stiff and resistant to the shear forces of lateral rotation and hyperextension that contribute to CeAD³⁵.

Migraine is more common in individuals with CeAD than in the community^{1,2}, and vascular mechanisms are thought to have a key role in the pathophysiology of this disease³⁶. Recently, rs9349379 (*PHACTRI*) and rs11172113 (*LRPI*) were identified as migraine susceptibility SNPs^{8,9,37}. Moreover, one additional migraine risk variant (rs13208321 in *FHL5*)⁹ was associated with CeAD at $P = 6.80 \times 10^{-4}$. All these SNPs showed effects in the

same direction as those for CeAD (Supplementary Table 20). Interestingly, associations with these shared variants were most significant for migraine without aura⁹, the migraine subtype most commonly associated with CeAD³⁸.

PHACTR1 is in a highly conserved genomic region, suggesting that it has a crucial involvement in biological processes³⁹, but its function is poorly understood. Experimental studies identified a pivotal role for the PHACTR1 protein in vascular tube formation and actin polymerization, suggesting that it possibly has a role in angiogenic processes^{40,41}. Upregulation of *PHACTR1* by transforming growth factor (TGF)- β has been described in breast cancer cell lines⁴², potentially pointing to a connection with the TGF- β signaling pathway, which is also implicated in genetic predisposition to migraine⁹ and has a key role in Marfan and Loeys-Dietz syndromes, two inherited connective tissue disorders causing aortic dissection^{43,44}. *In silico* functional annotation obtained from a wide array of published and unpublished expression quantitative trait locus (eQTL) data sets (Supplementary Table 21) provides some support for a functional effect of CeAD-associated SNPs in the 6p24.1-p23 locus on *PHACTR1* expression in whole blood and cerebellum cells (although the SNPs in the databases are in relatively weak LD with rs9349379)^{45,46}.

Although the follow-up analysis did not provide additional support for association with markers in *LRPI*, this locus remains of interest because of its association with migraine and abdominal aortic aneurysm, with effects in the same direction as in CeAD (Supplementary Table 17)⁴⁷.

CeAD is relatively uncommon (~2.6/100,000 cases per year)⁴⁸, and the current study is by far the largest genetic study of CeAD thus far (2,052 CeAD cases). Nevertheless, we might have lacked power to detect and replicate some associations, especially for loci showing heterogeneity in effect according to dissection site (Supplementary Table 22). We deliberately chose to focus on the discovery of genetic variants with overarching effects on CeAD risk, as we were underpowered to identify genetic variants underlying carotid or vertebral dissection exclusively. Future studies on larger samples are warranted to explore specific genetic susceptibility factors for carotid and vertebral artery dissections and to determine whether the 12q13.3 (*LRPI*), 4q12 (*LNX1*) and 18q22.1 (*CCDC102B*) loci are associated with CeAD.

In summary, we identified one previously unreported genome-wide significant risk locus for CeAD at *PHACTR1* and additional highly suggestive loci requiring confirmation in future independent studies. *PHACTR1* is also a major susceptibility locus for myocardial infarction and migraine. Understanding the mechanisms by which this locus appears to influence key vascular functions could have major applications for the treatment of these severe and disabling conditions.

URLs

CADISP, <http://www.cadisp.com/>; International Stroke Genetics Consortium, <http://www.strokegenetics.org/>.

METHODS

Methods and any associated references are available in the online version of the paper.

ONLINE METHODS

Study population

CeAD cases—We included 942 CeAD cases in the CADISP study in 2004–2009 (CADISP-1: 170 Finns and 772 non-Finnish Europeans)¹⁰. An additional 451 CeAD cases of European origin were recruited in 2008–2010, exclusively for the CADISP-genetics project, in some CADISP centers and additional European and US centers (CADISP-2). In total, 1,393 CeAD cases of European ancestry were available. All CeAD cases were ascertained through departments of neurology specialized in stroke care. Inclusion criteria, recruiting centers and participant selection are described in Supplementary Figures 1–3 and the Supplementary Note¹¹.

Individuals with non-CeAD ischemic stroke—As most individuals with CeAD sustained a cerebral ischemia, we planned to test whether genetic variants associated with CeAD were specific for CeAD, possibly through a predisposing vasculopathy, and did not confer generalized susceptibility to cerebral ischemia in young adults that could also predispose to other subtypes of ischemic stroke. We recruited 583 individuals with an ischemic stroke attributable to other causes (non-CeAD ischemic stroke: 162 Finns and 421 non-Finnish Europeans), frequency matched on the basis of age, sex and geographical origin to the CADISP-1 CeAD cases (Supplementary Fig. 3 and Supplementary Note).

Control populations—DNA for 14,416 controls was available. Most controls ($n = 14,203$: 74 Finns and 14,129 non-Finnish Europeans) were selected from an anonymized control genotype database at the CNG (Centre National de Génotypage) to match cases for ancestry, on the basis of the distribution of eigenvectors (Supplementary Fig. 4 and Supplementary Note)⁵⁰. In addition, 213 Finnish controls were recruited (Supplementary Note). Although controls were not screened for CeAD, given the low disease incidence in the community, a misclassification bias is unlikely.

Genotyping, quality control filters and imputation

DNA samples were genotyped on an Illumina Human610-Quad or Human660W-Quad BeadChip (Supplementary Table 1 and Supplementary Note). After quality control (Supplementary Note), we used 472,862 autosomal SNPs and 10,029 X-chromosomal SNPs for analyses. We performed genotype imputation to the non-monomorphic SNPs described in the HapMap 2 and 1000 Genomes Project (August 2010 release) CEU panels (Supplementary Note). Only SNPs with an imputation score of $R^2 > 0.3$ and a MAF of >0.01 were used for analysis.

Follow-up studies

We sought to replicate our strongest association signals in an independent sample comprising 659 CeAD cases, mostly drawn from existing databases of ischemic stroke

(recruited through neurology departments specialized in stroke care according to the same inclusion criteria as in the discovery GWAS), and 2,648 controls, all of European ancestry (Supplementary Table 4). Some of these (85 CeAD cases and 998 controls) had already been genotyped elsewhere on a genome-wide chip (Supplementary Table 1 and Supplementary Note). In addition, DNA samples from 238 CeAD cases were genotyped across the genome, and 1,584 controls were genotyped on a custom chip including the SNPs yielding the most significant associations in the GWAS (Supplementary Table 1 and Supplementary Note). Finally, DNA samples from 391 additional CeAD cases and 162 controls, recruited in the same centers as other individuals included in the discovery or follow-up analyses, were genotyped for the top 5 genotyped loci and the top imputed locus (1–2 SNPs per locus) using KASPAR technology. Of these 714 follow-up CeAD cases, 55 were excluded because of the unavailability of information on the dissection site, leaving us with 659 CeAD cases for analysis. To avoid population stratification bias, the association of the top SNPs with CeAD was analyzed separately by inclusion region (see Supplementary Table 4 for the matching algorithm), and meta-analysis (fixed effects, inverse variance weighted) was performed on the results.

Statistics

Genome-wide association analysis comparing CeAD cases to healthy controls

—Analyses were based on an additive genetic model. We used logistic regression to estimate OR values with corresponding 95% CIs, adjusting for sex. The CADISP-1 and CADISP-2 cohorts were analyzed separately because genotyping was carried out on different platforms (Supplementary Table 1). Moreover, because Finnish participants had a distinct ancestral origin relative to other populations (Supplementary Fig. 4), CADISP-1 was divided into CADISP-1 non-Finnish and CADISP-1 Finnish cohorts. Hence, sample sizes (CeAD-cases/controls) were as follows: 772/8,972 for CADISP-1 non-Finnish, 170/287 for CADISP-1 Finnish and 451/5,157 for CADISP-2. The first ten principal component scores were used as covariates for the CADISP-1 non-Finnish and CADISP-2 cohorts, and the first principal component score was used for the CADISP-1 Finnish cohort (Supplementary Note). Genomic control was applied to each of the three GWAS results (CADISP-1 non-Finnish, CADISP-1 Finnish and CADISP-2). Thereafter, a combined analysis was performed by inverse variance-weighted meta-analysis with a fixed-effects model (Supplementary Note). $P < 5 \times 10^{-8}$ was considered genome-wide significant.

Follow-up study—Because of the limits on DNA availability, six loci (ten SNPs) were chosen for follow-up genotyping in the phase 3 samples uniquely on the basis of statistical rankings in the GWAS meta-analysis of phase 1 and 2 samples (CADISP-1 and CADISP-2). We examined 6 SNP markers at the 5 most significantly associated loci from the genotyped panel ($P < 1 \times 10^{-5}$ in the GWAS) in the 659 CeAD and 2,648 control samples available for follow-up. In addition, two SNPs from the imputed locus on chromosome 18q22.1 were selected for genotyping in follow-up samples (Supplementary Table 5). We also genotyped two proxies for two of the most significantly associated SNPs ($P < 1 \times 10^{-5}$) from the GWAS meta-analysis (rs12215208, $R^2 = 0.39$ with rs9349379; rs6761601, $R^2 = 0.67$ with rs6741522).

First, we calculated the association statistics for the follow-up samples, using a threshold of $P = 5.00 \times 10^{-3}$ to assign significance (with Bonferroni correction for ten SNPs, which is conservative because of positive LD between some markers); irrespective of significance, we calculated combined association test statistics using meta-analysis and evaluated evidence of heterogeneity between GWAS and follow-up data. Second, we applied a Bayesian approach to evaluate the significance of associations using Wakefield's approximate Bayes factor¹². ABF and BFDP values are provided in Supplementary Tables 6 and 7.

Bayesian approach—We followed the approach developed by Wakefield¹² to consider the posterior odds on the null hypothesis (no association of a SNP with CeAD) for sequential studies from a Bayesian perspective. In summary, we obtained estimates of the log relative risk $\hat{\theta}$, standard error \sqrt{V} and Wald statistic $z^2 = \hat{\theta}^2 / V$ for a SNP from the logistic regression model (based on an additive genetic model), which was performed as our primary analysis to generate frequentist P values (Table 3 and Supplementary Table 5). Thereafter, we calculated Wakefield's ABF as follows:

$$\text{ABF} = \sqrt{\frac{W}{RV_1V_2}} \exp \left\{ -\frac{1}{2} \left(z_1^2 RV_2 + 2z_1 z_2 R \sqrt{V_1 V_2} + z_2^2 RV_1 \right) \right\}$$

where W is the variance on the prior for the effect size, $R = W / (V_1 W + V_2 W + V_1 V_2)$, V_1 and V_2 are squared standard errors, and z_1 and z_2 are Wald statistics, each from the discovery GWAS (stages 1 and 2 combined) and the follow-up study (stage 3). A smaller ABF value corresponds to more evidence of the alternative hypothesis, i.e., the association of a SNP with CeAD.

Wakefield proposed three different specifications of priors for the log relative risk. Because our aim in this study was to find common susceptibility variants for CeAD, we followed an effect-MAF independence prior as $W = \left\{ \log \text{RR}_u / \Phi^{-1}(1 - q) \right\}^2$, with $\text{RR}_u = 1.5$. RR_u is an upper value above which we believe that relative risks will occur with low probability; q is the prior probability of a relative risk above RR_u ; $\Phi(\cdot)$ is the distribution function of a standard normal random variable.

Considering $\text{PO} = \pi_0 / (1 - \pi_0)$ to be the prior odds on the null hypothesis, where π_0 is the prior probability of the null ($\text{Pr}[H_0]$), the posterior probability of H_0 is given by the BFDP, with $\text{BFDP} = \text{ABF} \times \text{PO} / (1 + \text{ABF} \times \text{PO})$.

Specificity analysis—We tested whether the SNPs yielding the most significant associations with CeAD were also associated with non-CeAD ischemic stroke. We ran a logistic regression adjusted for sex and principal components, as described above, for non-Finnish Europeans (421 cases and 8,972 controls) and Finns (162 cases and 287 controls), followed by a fixed-effects inverse variance-weighted meta-analysis. Statistical significance was defined by false discovery rate (FDR)-corrected $P < 0.05$.

Sensitivity analyses—For the most significant associations, we stratified on and tested for interaction with sex, and we examined whether the most significant SNPs predicted an earlier age of onset of CeAD. We then evaluated the stability of the associations according to dissection site (carotid or vertebral) and the presence of cerebral ischemia (ischemic stroke or transient ischemic attack). Finally, we performed analyses stratified on the presence of two CeAD risk factors, migraine and cervical trauma in cases; case-only association analyses of the top CeAD SNPs with dissection site, the presence of cerebral ischemia, migraine and trauma were used as surrogate tests of heterogeneity or interaction, with $P < 0.05$ being considered significant⁵¹.

Candidate gene approach—We examined the associations of CeAD with published CeAD susceptibility SNPs and with SNPs in *COL3A1*, the gene harboring causal mutations for vascular Ehlers-Danlos syndrome, a rare etiology of CeAD¹⁶. We also tested for pleiotropic effects of susceptibility SNPs for intracranial aneurysms, thoracic aortic aneurysms and dissections, other subtypes of ischemic stroke and migraine (Supplementary Note). An FDR-corrected P value of <0.05 was considered to be statistically significant. We performed a sign test to evaluate whether associations of CeAD with alleles previously associated with migraine and ischemic stroke subtypes involved the same risk alleles more often than predicted by chance. A right-tailed P value of 0.05 was considered to be statistically significant.

Ethics

The study protocol was approved by local authorities in all participating centers and conducted according to national rules concerning ethics committee approval and informed consent.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Authors

Stéphanie Debette^{1,2,3,4,5,6}, Yoichiro Kamatani^{7,8,9}, Tiina M Metso¹⁰, Manja Kloss¹¹, Ganesh Chauhan⁶, Stefan T Engelter¹², Alessandro Pezzini¹³, Vincent Thijs^{14,15}, Hugh S Markus¹⁶, Martin Dichgans^{17,18}, Christiane Wolf^{6,19}, Ralf Dittrich²⁰, Emmanuel Touzé^{21,22}, Andrew M Southerland²³, Yves Samson²⁴, Shérine Abboud²⁵, Yannick Béjot²⁶, Valeria Caso^{27,28}, Anna Bersano²⁹, Andreas Gschwendtner^{17,18}, Maria Sessa³⁰, John Cole³¹, Chantal Lamy³², Elisabeth Medeiros³³, Simone Beretta³⁴, Leo H Bonati¹², Armin J Grau³⁵, Patrik Michel³⁶, Jennifer J Majersik³⁷, Pankaj Sharma^{38,39}, Ludmila Kalashnikova⁴⁰, Maria Nazarova^{40,41}, Larisa Dobrynina⁴⁰, Eva Bartels⁴², Benoit Guillon⁴³, Evita G van den Herik⁴⁴, Israel Fernandez-Cadenas^{45,46}, Katarina Jood⁴⁷, Michael A Nalls⁴⁸, Frank-Erik De Leeuw⁴⁹, Christina Jern⁴⁹, Yu-Ching Cheng³¹, Inge Werner¹¹, Antti J Metso¹⁰, Christoph Lichy¹¹, Philippe A Lyrer¹², Tobias Brandt⁵⁰, Giorgio B Boncoraglio²⁹, Heinz-Erich Wichmann⁵¹, Christian Gieger⁵², Andrew D Johnson⁵³, Thomas Böttcher⁵⁴, Maurizio Castellano⁵⁵, Dominique Arveiler⁵⁶, M Arfan

Ikram^{44,57,58,59}, Monique M B Breteler^{57,60}, Alessandro Padovani¹³, James F Meschia⁶¹, Gregor Kuhlenbäumer⁶², Arndt Rolfs⁵⁴, Bradford B Worrall^{23,63}, International Stroke Genetics Consortium⁶⁴, Erich-Bernd Ringelstein²⁰, Diana Zelenika⁷, Turgut Tatlisumak¹⁰, Mark Lathrop^{7,8,65}, Didier Leys^{3,4,5,6,67}, Philippe Amouyel^{1,2,3,4,67}, Jean Dallongeville^{1,2,3,67}, and for the CADISP group⁶⁶

Affiliations

¹INSERM U744, Lille, France ²Institut Pasteur de Lille, Lille, France ³Université Lille Nord de France, Lille, France ⁴Lille University Hospital, Lille, France ⁵Department of Neurology, Equipe d'accueil 1046, Lille, France ⁶INSERM U897, University of Bordeaux, Bordeaux, France ⁷Centre National de Génotypage, Evry, France ⁸Fondation Jean Dausset, Centre d'Etude du Polymorphisme Humain (CEPH), Paris, France ⁹Center for Integrative Medical Sciences, RIKEN, Yokohama, Japan ¹⁰Department of Neurology, Helsinki University Central Hospital, Helsinki, Finland ¹¹Department of Neurology, Heidelberg University Hospital, Heidelberg, Germany ¹²Department of Neurology, Basel University Hospital, Basel, Switzerland ¹³Department of Clinical and Experimental Sciences, Neurology Clinic, Brescia University Hospital, Brescia, Italy ¹⁴Department of Neurology, Leuven University Hospital, Leuven, Belgium ¹⁵Vesalius Research Center, VIB, Leuven, Belgium ¹⁶Department of Clinical Neurosciences, University of Cambridge, Cambridge, UK ¹⁷Institute for Stroke and Dementia Research, Klinikum der Universität München, Ludwig Maximilians Universität, Munich, Germany ¹⁸Munich Cluster for Systems Neurology (SyNergy), Munich, Germany ¹⁹Department of Statistical Genetics, Max Planck Institute for Psychiatry, Munich, Germany ²⁰Department of Neurology, University Hospital of Münster, Münster, Germany ²¹Department of Neurology, University Hospital of Sainte-Anne, Paris, France ²²Department of Neurology, University Hospital of Caen, Caen, France ²³Department of Neurology, University of Virginia, Charlottesville, Virginia, USA ²⁴Urgences Cérébro-Vasculaire, Assistance Publique-Hôpitaux de Paris Salpêtrière Urgences Cérébro-Vasculaires and Université Pierre et Marie Curie Paris Paris VI Université, Paris, France ²⁵Laboratory of Experimental Neurology, Université Libre de Bruxelles, Brussels, Belgium ²⁶Department of Neurology, Dijon University Hospital, Dijon, France ²⁷Stroke Unit, University of Perugia Santa Maria della Misericordia Hospital, Sant'Andrea delle Fratte, Perugia, Italy ²⁸Division of Cardiovascular Medicine, University of Perugia Santa Maria della Misericordia Hospital, Sant'Andrea delle Fratte, Perugia, Italy ²⁹Department of Cerebrovascular Diseases, Fondazione Istituto di Ricovero e Cura a Carattere Scientifico Istituto Neurologico Carlo Besta, Milan, Italy ³⁰Department of Neurology, Milan, San Raffaele University Hospital, Milan, Italy ³¹Maryland Stroke Center, Department of Neurology, University of Maryland School of Medicine, Baltimore, Maryland, USA ³²Department of Neurology, Amiens University Hospital, Amiens, France ³³Department of Neurology, Besançon University Hospital, Besançon, France ³⁴Department of Neurology, University of Milano Bicocca, San Gerardo Hospital, Monza, Italy ³⁵Department of Neurology, Klinikum Ludwigshafen, Ludwigshafen, Germany ³⁶Department of Neurology, Lausanne University Hospital, Lausanne, Switzerland ³⁷Division of Vascular Neurology, Department of Neurology,

University of Utah, Salt Lake City, Utah, USA ³⁸Institute of Cardiovascular Research Royal Holloway University of London (ICR2UL), London, UK ³⁹Ashford and St Peter's Hospitals, London, UK ⁴⁰Research Center of Neurology, Russian Academy of Medical Sciences, Moscow, Russia ⁴¹Centre for Cognition and Decision Making, Higher School of Economics, Moscow, Russia ⁴²Center for Neurological Vascular Diagnostics, Munich, Germany ⁴³Department of Neurology, Nantes University Hospital, Nantes, France ⁴⁴Department of Neurology, Erasmus MC University Medical Center, Rotterdam, the Netherlands ⁴⁵Stroke Pharmacogenomics and Genetics, Fundació Docència i Recerca MutuaTerrassa, Terrassa, Spain ⁴⁶Laboratorio Neurovascular, Institut de Recerca, Hospital Vall d'Hebron, Barcelona, Spain ⁴⁷Institute of Neuroscience and Physiology, Sahlgrenska Academy at the University of Gothenburg, Gothenburg, Sweden ⁴⁸Molecular Genetics Section, Laboratory of Neurogenetics, National Institute on Aging, US National Institutes of Health, Bethesda, Maryland, USA ⁴⁹Donders Institute for Brain, Cognition and Behaviour, Centre for Neuroscience, Department of Neurology, Radboud University Nijmegen Medical Centre, Nijmegen, the Netherlands ⁵⁰Department of Rehabilitation, Schmieder Klinik, Heidelberg, Germany ⁵¹Institute of Epidemiology, Helmholtz Center Munich, Neuherberg, Germany ⁵²Institute of Genetic Epidemiology, Helmholtz Center Munich, Neuherberg, Germany ⁵³National Heart, Lung, and Blood Institute's Framingham Heart Study Cardiovascular Epidemiology and Human Genomics Branch, Framingham, Massachusetts, USA ⁵⁴Department of Neurology, Rostock University Hospital, Rostock, Germany ⁵⁵Department of Clinical and Experimental Sciences, Clinica Medica, Brescia University Hospital, Brescia, Italy ⁵⁶Department of Epidemiology and Public Health, EA 3430, Strasbourg University, Strasbourg, France ⁵⁷Department of Epidemiology, Erasmus MC University Medical Center, Rotterdam, the Netherlands ⁵⁸Department of Radiology, Erasmus MC University Medical Center, Rotterdam, the Netherlands ⁵⁹Netherlands Consortium for Healthy Aging, Leiden, the Netherlands ⁶⁰Population Health Sciences, German Center for Neurodegenerative Diseases (DZNE), Bonn, Germany ⁶¹Department of Neurology, Mayo Clinic Florida, Jacksonville, Florida, USA ⁶²Institute of Experimental Medicine, University of Kiel, Kiel, Germany ⁶³Department of Public Health Sciences, University of Virginia, Charlottesville, Virginia, USA ⁶⁴A list of contributing members appears in the Supplementary Note ⁶⁵Génome Québec, McGill University, Montreal, Quebec, Canada

Acknowledgments

The authors thank the staff and participants of all CADISP centers for their important contributions. The CADISP study has been supported by INSERM, Lille 2 University, Institut Pasteur de Lille and Lille University Hospital and received funding from the European Regional Development Fund (FEDER funds) and Région Nord-Pas-de-Calais in the framework of Contrat de Projets Etat-Region 2007–2013 Région Nord-Pas-de-Calais (grant 09120030), Centre National de Génotypage, the Emil Aaltonen Foundation, the Paavo Ilmari Ahvenainen Foundation, the Helsinki University Central Hospital Research Fund, the Helsinki University Medical Foundation, the Päivikki and Sakari Sohlberg Foundation, the Aarne Koskelo Foundation, the Maire Taponen Foundation, the Aarne and Aili Turunen Foundation, the Lilly Foundation, the Alfred Kordelin Foundation, the Finnish Medical Foundation, the Orion Farnos Research Foundation, the Maud Kuistila Foundation, the Finnish Brain Foundation, the Biomedicum Helsinki Foundation, Projet Hospitalier de Recherche Clinique Régional, Fondation de France, Génomopôle de Lille,

Adrinord, the Basel Stroke Funds, the Käthe-Zingg-Schwichtenberg-Fonds of the Swiss Academy of Medical Sciences and the Swiss Heart Foundation.

L.H.B., S.T.E. and P.A.L. were supported, in part, by a grant from the Swiss National Science Foundation (33CM30-124119). S.D. is supported by a Chair of Excellence from the French National Research Agency (ANR). S.D. and M.D. are supported by a grant from the Leducq Foundation. M.D. is supported by the Vascular Dementia Research Foundation. I.F.-C. is supported by the Miguel Servet programme (CP12/03298) from the Spanish Ministry of Health (Instituto de Salud Carlos III). G.K. is a member of the Deutsche Forschungsgemeinschaft Cluster of Excellence 'Inflammation at Interfaces'. P.S. is supported by a Department of Health (UK) senior fellowship. A.M.S. is supported by the American Heart Association/American Stroke Association National Clinical Research Program (AHA 3CRP14140001). V.T. is supported by Fonds Wetenschappelijk Onderzoek Flanders.

More detailed acknowledgments can be found in the Supplementary Note.

References

1. Debette S, Leys D. Cervical-artery dissections: predisposing factors, diagnosis, and outcome. *Lancet Neurol.* 2009; 8:668–678. [PubMed: 19539238]
2. Rubinstein SM, Peerdeman SM, van Tulder MW, Riphagen I, Haldeman S. A systematic review of the risk factors for cervical artery dissection. *Stroke.* 2005; 36:1575–1580. [PubMed: 15933263]
3. Debette S, et al. Association of vascular risk factors with cervical artery dissection and ischemic stroke in young adults. *Circulation.* 2011; 123:1537–1544. [PubMed: 21444882]
4. Arnold M, et al. Vascular risk factors and morphometric data in cervical artery dissection: a case-control study. *J Neurol Neurosurg Psychiatry.* 2009; 80:232–234. [PubMed: 19151021]
5. Debette S, Markus HS. The genetics of cervical artery dissection: a systematic review. *Stroke.* 2009; 40:e459–e466. [PubMed: 19390073]
6. Kathiresan S, et al. Genome-wide association of early-onset myocardial infarction with single nucleotide polymorphisms and copy number variants. *Nat Genet.* 2009; 41:334–341. [PubMed: 19198609]
7. Deloukas P, et al. Large-scale association analysis identifies new risk loci for coronary artery disease. *Nat Genet.* 2013; 45:25–33. [PubMed: 23202125]
8. Freilinger T, et al. Genome-wide association analysis identifies susceptibility loci for migraine without aura. *Nat Genet.* 2012; 44:777–782. [PubMed: 22683712]
9. Anttila V, et al. Genome-wide meta-analysis identifies new susceptibility loci for migraine. *Nat Genet.* 2013; 45:912–917. [PubMed: 23793025]
10. Debette S, et al. Differential features of carotid and vertebral artery dissections: the CADISP study. *Neurology.* 2011; 77:1174–1181. [PubMed: 21900632]
11. Debette S, et al. CADISP-genetics: an international project searching for genetic risk factors of cervical artery dissections. *Int J Stroke.* 2009; 4:224–230. [PubMed: 19659826]
12. Wakefield J. Bayes factors for genome-wide association studies: comparison with P-values. *Genet Epidemiol.* 2009; 33:79–86. [PubMed: 18642345]
13. Pezzini A, et al. Plasma homocysteine concentration, C677T MTHFR genotype, and 844ins68bp CBS genotype in young adults with spontaneous cervical artery dissection and atherothrombotic stroke. *Stroke.* 2002; 33:664–669. [PubMed: 11872884]
14. Pezzini A, et al. Migraine mediates the influence of C677T MTHFR genotypes on ischemic stroke risk with a stroke-subtype effect. *Stroke.* 2007; 38:3145–3151. [PubMed: 17962595]
15. Longoni M, et al. The ICAM-1 E469K gene polymorphism is a risk factor for spontaneous cervical artery dissection. *Neurology.* 2006; 66:1273–1275. [PubMed: 16636253]
16. Pepin M, Schwarze U, Superti-Furga A, Byers PH. Clinical and genetic features of Ehlers-Danlos syndrome type IV, the vascular type. *N Engl J Med.* 2000; 342:673–680. [PubMed: 10706896]
17. Low SK, et al. Genome-wide association study for intracranial aneurysm in the Japanese population identifies three candidate susceptible loci and a functional genetic variant at *EDNRA*. *Hum Mol Genet.* 2012; 21:2102–2110. [PubMed: 22286173]
18. Bilguvar K, et al. Susceptibility loci for intracranial aneurysm in European and Japanese populations. *Nat Genet.* 2008; 40:1472–1477. [PubMed: 18997786]

19. Yasuno K, et al. Genome-wide association study of intracranial aneurysm identifies three new risk loci. *Nat Genet.* 2010; 42:420–425. [PubMed: 20364137]
20. Lemaire SA, et al. Genome-wide association study identifies a susceptibility locus for thoracic aortic aneurysms and aortic dissections spanning FBN1 at 15q21.1. *Nat Genet.* 2011; 43:996–1000. [PubMed: 21909107]
21. O'Donnell CJ, et al. Genome-wide association study for coronary artery calcification with follow-up in myocardial infarction. *Circulation.* 2011; 124:2855–2864. [PubMed: 22144573]
22. Lu X, et al. Genome-wide association study in Han Chinese identifies four new susceptibility loci for coronary artery disease. *Nat Genet.* 2012; 44:890–894. [PubMed: 22751097]
23. Hager J, et al. Genome-wide association study in a Lebanese cohort confirms PHACTR1 as a major determinant of coronary artery stenosis. *PLoS ONE.* 2012; 7:e38663. [PubMed: 22745674]
24. Mehta NN. A genome-wide association study in Europeans and South Asians identifies 5 new loci for coronary artery disease. *Circ Cardiovasc Genet.* 2011; 4:465–466. [PubMed: 21846871]
25. Schunkert H, et al. Large-scale association analysis identifies 13 new susceptibility loci for coronary artery disease. *Nat Genet.* 2011; 43:333–338. [PubMed: 21378990]
26. Coronary Artery Disease (C4D) Genetics Consortium. A genome-wide association study in Europeans and South Asians identifies five new loci for coronary artery disease. *Nat Genet.* 2011; 43:339–344. [PubMed: 21378988]
27. Bevan S, et al. Genetic heritability of ischemic stroke and the contribution of previously reported candidate gene and genomewide associations. *Stroke.* 2012; 43:3161–3167. [PubMed: 23042660]
28. Bellenguez C, et al. Genome-wide association study identifies a variant in HDAC9 associated with large vessel ischemic stroke. *Nat Genet.* 2012; 44:328–333. [PubMed: 22306652]
29. Traylor M, et al. Genetic risk factors for ischaemic stroke and its subtypes (the METASTROKE Collaboration): a meta-analysis of genome-wide association studies. *Lancet Neurol.* 2012; 11:951–962. [PubMed: 23041239]
30. Smyth DJ, et al. Shared and distinct genetic variants in type 1 diabetes and celiac disease. *N Engl J Med.* 2008; 359:2767–2777. [PubMed: 19073967]
31. Calvet D, et al. Increased stiffness of the carotid wall material in patients with spontaneous cervical artery dissection. *Stroke.* 2004; 35:2078–2082. [PubMed: 15243148]
32. Golledge J, Eagle KA. Acute aortic dissection. *Lancet.* 2008; 372:55–66. [PubMed: 18603160]
33. Norman PE, Davis TM, Le MT, Golledge J. Matrix biology of abdominal aortic aneurysms in diabetes: mechanisms underlying the negative association. *Connect Tissue Res.* 2007; 48:125–131. [PubMed: 17522995]
34. Shantikumar S, Ajjan R, Porter KE, Scott DJ. Diabetes and the abdominal aortic aneurysm. *Eur J Vasc Endovasc Surg.* 2010; 39:200–207. [PubMed: 19948418]
35. Callaghan FM, et al. Wall stress of the cervical carotid artery in patients with carotid dissection: a case-control study. *Am J Physiol Heart Circ Physiol.* 2011; 300:H1451–H1458. [PubMed: 21297030]
36. Kurth T, et al. Headache, migraine, and structural brain lesions and function: population based Epidemiology of Vascular Ageing-MRI study. *Br Med J.* 2011; 342:c7357. [PubMed: 21245119]
37. Chasman DI, et al. Genome-wide association study reveals three susceptibility loci for common migraine in the general population. *Nat Genet.* 2011; 43:695–698. [PubMed: 21666692]
38. Metso TM, et al. Migraine in cervical artery dissection and ischemic stroke patients. *Neurology.* 2012; 78:1221–1228. [PubMed: 22491867]
39. Allen PB, Greenfield AT, Svenningsson P, Haspeslagh DC, Greengard P. Phactrs 1–4: a family of protein phosphatase 1 and actin regulatory proteins. *Proc Natl Acad Sci USA.* 2004; 101:7187–7192. [PubMed: 15107502]
40. Allain B, et al. Neuropilin-1 regulates a new VEGF-induced gene, Phactr-1, which controls tubulogenesis and modulates lamellipodial dynamics in human endothelial cells. *Cell Signal.* 2012; 24:214–223. [PubMed: 21939755]
41. Jarray R, et al. Depletion of the novel protein PHACTR-1 from human endothelial cells abolishes tube formation and induces cell death receptor apoptosis. *Biochimie.* 2011; 93:1668–1675. [PubMed: 21798305]

42. Fils-Aimé N, et al. MicroRNA-584 and the protein phosphatase and actin regulator 1 (PHACTR1), a new signaling route through which transforming growth factor- β mediates the migration and actin dynamics of breast cancer cells. *J Biol Chem*. 2013; 288:11807–11823. [PubMed: 23479725]
43. Goumans MJ, Liu Z, ten Dijke P. TGF- β signaling in vascular biology and dysfunction. *Cell Res*. 2009; 19:116–127. [PubMed: 19114994]
44. Loeys BL, et al. A syndrome of altered cardiovascular, craniofacial, neurocognitive and skeletal development caused by mutations in TGFBR1 or TGFBR2. *Nat Genet*. 2005; 37:275–281. [PubMed: 15731757]
45. Zhang X, et al. Genetic associations with expression for genes implicated in GWAS studies for atherosclerotic cardiovascular disease and blood phenotypes. *Hum Mol Genet*. 2014; 23:782–795. [PubMed: 24057673]
46. Westra HJ, et al. Systematic identification of trans eQTLs as putative drivers of known disease associations. *Nat Genet*. 2013; 45:1238–1243. [PubMed: 24013639]
47. Bown MJ, et al. Abdominal aortic aneurysm is associated with a variant in low-density lipoprotein receptor-related protein 1. *Am J Hum Genet*. 2011; 89:619–627. [PubMed: 22055160]
48. Lee VH, Brown RD Jr, Mandrekar JN, Mokri B. Incidence and outcome of cervical artery dissection: a population-based study. *Neurology*. 2006; 67:1809–1812. [PubMed: 17130413]
49. Pruim RJ, et al. LocusZoom: regional visualization of genome-wide association scan results. *Bioinformatics*. 2010; 26:2336–2337. [PubMed: 20634204]
50. Price AL, et al. Principal components analysis corrects for stratification in genome-wide association studies. *Nat Genet*. 2006; 38:904–909. [PubMed: 16862161]
51. Thomas D. Gene—environment-wide association studies: emerging approaches. *Nat Rev Genet*. 2010; 11:259–272. [PubMed: 20212493]




Figure 1. Regional association plot centered on rs9349379 (*PHACTR1*). Regional plot for associations in the region centered on rs9349379, drawn using LocusZoom software⁴⁹. All SNPs, on the basis of 1000 Genomes Project imputed results (dots), are plotted with their GWAS meta-analysis association P values against their genomic position. The color of the dots represents the LD between SNPs. The purple line represents the estimated recombination rates. Genes and exons are shown as dark blue arrows and vertical lines, respectively.

Table 1

Baseline clinical characteristics of the study subjects

	CeAD cases				
	GWAS		Follow-up studies		Non-CeAD ischemic stroke
	CADISP-1	CADISP-2	CADISP-1	CADISP-2	
<i>n</i>	942		659		583
Mean age (s.d.)	44.2 (10.0)	44.6 (10.5)	43.6 (9.4)		44.6 (10.5)
Men, <i>n</i> (%)	544 (57.7)	264 (58.5)	387 (58.7)		257 (61.2)
Location					
Carotid ^a , <i>n</i> (%)	603 (64.0)	253 (60.7) ^b	375 (56.9)		–
Vertebral, <i>n</i> (%)	305 (32.4)	147 (35.3) ^b	256 (38.8)		–
Both, <i>n</i> (%)	33 (3.5)	17 (4.0) ^b	26 (3.9)		–
Clinical presentation					
Cerebral ischemia ^c , <i>n</i> (%)	729 (77.4)	333 (79.5) ^b	544 (82.5)		–
Ischemic stroke, <i>n</i> (%)	606 (64.3)	281 (67.1) ^b	508 (77.1)		–

CeAD, cervical artery dissection.

^aInternal carotid artery dissection (one additional case had a common carotid artery dissection in CADISP-1).

^bInformation on dissection site and associated ischemia was missing for 34 and 32 cases, respectively, in CADISP-2.

^cCerebral ischemia corresponds to ischemic stroke or transient ischemic attack (including transient monocular blindness).

Table 2

SNPs yielding the most significant associations with CeAD in the GWAS

SNP ^a	Chr.	Position (bp)	EA	Nearest gene ^b	Stage	EAF	OR (95% CI)	P
rs12402265	1	59,463,190	A	<i>FGGY</i>	CADISP-1 non-Finnish	0.26	1.21 (1.07–1.36)	2.70×10^{-3}
					CADISP-1 Finnish	0.28	1.16 (0.87–1.54)	0.315
					CADISP-2	0.28	1.28 (1.10–1.49)	1.30×10^{-3}
					Meta-analysis		1.23 (1.13–1.35)	6.12×10^{-6}
rs6741522	2	185,544,143	A	<i>ZNF804A</i>	CADISP-1 non-Finnish	0.14	1.33 (1.15–1.54)	1.99×10^{-4}
					CADISP-1 Finnish	0.10	1.32 (0.85–2.05)	0.230
					CADISP-2	0.13	1.38 (1.14–1.66)	1.21×10^{-3}
					Meta-analysis		1.34 (1.19–1.50)	5.65×10^{-7}
rs6820391	4	54,109,453	A	<i>LNX1</i>	CADISP-1 non-Finnish	0.29	1.21 (1.07–1.36)	2.33×10^{-3}
					CADISP-1 Finnish	0.29	1.15 (0.86–1.54)	0.345
					CADISP-2	0.28	1.26 (1.09–1.47)	2.69×10^{-3}
					Meta-analysis		1.23 (1.12–1.35)	6.35×10^{-6}
rs9349379	6	13,011,943	G	<i>PHACTR1</i>	CADISP-1 non-Finnish	0.40	0.76 (0.68–0.86)	8.90×10^{-6}
					CADISP-1 Finnish	0.45	0.84 (0.63–1.10)	0.216
					CADISP-2	0.40	0.73 (0.62–0.84)	3.39×10^{-5}
					Meta-analysis		0.75 (0.69–0.82)	4.46×10^{-10}
rs11172113	12	55,813,550	C	<i>LRPI</i>	CADISP-1 non-Finnish	0.38	0.77 (0.68–0.86)	1.66×10^{-5}
					CADISP-1 Finnish	0.39	0.90 (0.69–1.18)	0.456
					CADISP-2	0.40	0.77 (0.67–0.89)	7.61×10^{-4}
					Meta-analysis		0.78 (0.71–0.85)	4.22×10^{-8}
rs1466535 ^c	12	55,820,737	A	<i>LRPI</i>	CADISP-1 non-Finnish	0.32	0.75 (0.66–0.85)	1.06×10^{-5}
					CADISP-1 Finnish	0.28	1.11 (0.83–1.49)	0.473
					CADISP-2	0.34	0.80 (0.68–0.93)	4.04×10^{-3}
					Meta-analysis		0.80 (0.73–0.88)	2.07×10^{-6}

Only results for genotyped SNPs with $P < 1 \times 10^{-5}$ are presented here (for all associations with $P < 1 \times 10^{-4}$, see Supplementary Table 2). Chr., chromosome; EA, effect allele (minor allele); EAF, effect allele frequency in controls. Alleles and chromosomal positions were identified on the basis of the plus strand of the NCBI Build 36 reference genome.

^a rs9349379, rs11172113, rs1466535 and rs6820391 are intronic; rs6741522 and rs12402265 are intergenic.

^bThe names for the nearest genes are shown according to the Human Gene Organization (HUGO) Gene Nomenclature System.

^crs1466535 is in LD with rs11172113 ($r^2 = 0.72$).

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 3

SNPs yielding the most significant associations with CeAD and their proxies in GWAS and follow-up studies

SNP	Chr.	EA	EAF	Gene	CeAD site	P_{het}	CeAD site ^b	GWAS meta-analysis (<i>n</i> = 1,393/14,416)			Follow-up meta-analysis (<i>n</i> = 659/2,648)			Overall meta-analysis (<i>n</i> = 2,052/17,064)		
								OR (95% CI)	<i>P</i>	OR (95% CI)	<i>P</i>	OR (95% CI)	<i>P</i>	OR (95% CI)	<i>P</i>	P_{het}
rs12402265 ^a	1	A	0.27	<i>FGGY</i>	All	0.70		1.23 (1.13–1.35)	6.12×10^{-6}	1.21 (1.04–1.40)	0.012	1.23 (1.14–1.33)	2.30×10^{-7}	0.83		
					Carotid			1.23 (1.10–1.38)	2.95×10^{-4}	1.40 (1.17–1.68)	2.52×10^{-4}	1.28 (1.16–1.41)	5.46×10^{-7}	0.24		
					Vertebral			1.27 (1.10–1.47)	1.36×10^{-3}	1.05 (0.85–1.31)	0.64	1.20 (1.06–1.35)	3.47×10^{-3}	0.16		
rs6741522	2	A	0.13	<i>ZNF804A</i>	All	0.64		1.34 (1.19–1.50)	5.65×10^{-7}	1.09 (0.91–1.31)	0.36	1.26 (1.15–1.39)	2.29×10^{-6}	0.06		
					Carotid			1.37 (1.19–1.57)	1.17×10^{-5}	1.05 (0.83–1.32)	0.68	1.27 (1.13–1.44)	7.33×10^{-5}	0.06		
					Vertebral			1.32 (1.10–1.60)	3.53×10^{-3}	1.24 (0.96–1.61)	0.10	1.29 (1.11–1.51)	8.92×10^{-4}	0.69		
rs6820391	4	A	0.29	<i>LNX1</i>	All	0.038		1.23 (1.12–1.35)	6.35×10^{-6}	1.28 (1.10–1.47)	9.28×10^{-4}	1.24 (1.15–1.34)	2.36×10^{-8}	0.68		
					Carotid			1.30 (1.16–1.45)	3.88×10^{-6}	1.29 (1.08–1.55)	4.84×10^{-3}	1.30 (1.18–1.43)	6.41×10^{-8}	0.97		
					Vertebral			1.09 (0.94–1.26)	0.28	1.23 (1.00–1.52)	0.049	1.13 (1.00–1.28)	0.042	0.33		
rs9349379	6	G	0.40	<i>PHACTR1</i>	All	0.52		0.75 (0.69–0.82)	4.46×10^{-10}	0.81 (0.71–0.94)	3.91×10^{-3}	0.77 (0.72–0.83)	1.00×10^{-11}	0.36		
					Carotid			0.76 (0.68–0.85)	1.27×10^{-6}	0.76 (0.64–0.90)	1.79×10^{-3}	0.76 (0.69–0.83)	8.24×10^{-9}	0.96		
					Vertebral			0.72 (0.62–0.84)	1.38×10^{-5}	0.92 (0.75–1.13)	0.42	0.79 (0.70–0.88)	6.32×10^{-5}	0.06		
rs11172113	12	C	0.39	<i>LRP1</i>	All	0.061		0.78 (0.71–0.85)	4.22×10^{-8}	0.93 (0.81–1.07)	0.34	0.82 (0.76–0.89)	3.03×10^{-7}	0.03		
					Carotid			0.74 (0.66–0.83)	1.14×10^{-7}	0.88 (0.75–1.05)	0.16	0.78 (0.71–0.86)	1.90×10^{-7}	0.08		
					Vertebral			0.89 (0.77–1.03)	0.11	0.99 (0.81–1.21)	0.93	0.92 (0.82–1.04)	0.17	0.39		
rs1466535 ^a	12	A	0.32	<i>LRP1</i>	All	6.49×10^{-3}		0.80 (0.73–0.88)	2.07×10^{-6}	0.92 (0.80–1.07)	0.30	0.83 (0.77–0.90)	4.94×10^{-6}	0.10		
					Carotid			0.73 (0.65–0.82)	2.50×10^{-7}	0.85 (0.71–1.01)	0.07	0.76 (0.69–0.84)	1.17×10^{-7}	0.18		

SNP	Chr.	EA	EAF	Gene	CeAD site	P_{het}	CeAD site ^b	GWAS meta-analysis (<i>n</i> = 1,393/14,416)			Follow-up meta-analysis (<i>n</i> = 659/2,648)			Overall meta-analysis (<i>n</i> = 2,052/17,064)		
								OR (95% CI)	<i>P</i>	P_{het}	OR (95% CI)	<i>P</i>	P_{het}	OR (95% CI)	<i>P</i>	P_{het}
					Vertebral			0.97 (0.84–1.12)	0.68		0.99 (0.80–1.22)	0.92		0.98 (0.87–1.10)	0.69	0.87

Chr., chromosome; EA, effect allele (minor allele); EAF, effect allele frequency (in controls from the GWAS); OR, odds ratio; CI, confidence interval; P_{het} , *P* value for heterogeneity between the discovery GWAS and the follow-up sample.

^aThese SNPs were imputed (HapMap 2) in the Maryland follow-up sample with $R^2 = 0.93$ (rs1466535) and 1 (rs12402265).

^bThe *P* value of association with the dissection site in a case-only analysis.