



HAL
open science

Multivariate Multi-scaled Student Distributions : brain tumor characterization from multiparametric MRI

Alexis Arnaud, Florence Forbes, Benjamin Lemasson, Emmanuel Barbier

► To cite this version:

Alexis Arnaud, Florence Forbes, Benjamin Lemasson, Emmanuel Barbier. Multivariate Multi-scaled Student Distributions : brain tumor characterization from multiparametric MRI. *Statistique Mathématique et Applications* 2015, Aug 2015, Fréjus, France. hal-01253595

HAL Id: hal-01253595

<https://hal.science/hal-01253595v1>

Submitted on 11 Jan 2016

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

MULTIVARIATE MULTI-SCALED STUDENT DISTRIBUTIONS : BRAIN TUMOR CHARACTERIZATION FROM MULTIPARAMETRIC MRI.

Alexis Arnaud¹ & Florence Forbes¹
Benjamin Lemasson² & Emmanuel Barbier²

¹ *Inria Grenoble - Rhône-Alpes, 655 avenue de l'Europe, 38330 Montbonnot-Saint-Martin,
France, firstName.lastName@inria.fr*

² *GIN, 31 Chemin Fortuné Ferrini, 38706 La Tronche, France,
firstName.lastName@ujf-grenoble.fr*

Keywords.

Multivariate multi-scaled Student distribution, mixture model, EM algorithm, clustering, multiparametric MRI, tumor dictionary.

Abstract.

Brain tumor characterization is very useful for patients treatment, but it can be time-consuming for medical experts. Furthermore, the reference method to characterize tissues is biopsy which is a local and invasive technique. Because of this, there is a huge interest for automatic and non-invasive approaches in order to characterize tumor.

In this study we use a statistical model-based method to classify multiparametric MRI of brain rat tumors, which allows data quality control with atypical observations detection, and may provide a dictionary of tumor signatures.

A previous study, [1], used a Gaussian mixture model to characterize pixels inside tumors. With this model, the observations are gathered into classes resulting from Gaussian distributions. However, this model is sensitive to outliers which degrade the relevance of the obtained groups. And inside a tumor, there could be a huge variability and so a lot of outliers.

To account for this biological variability, we propose to use generalized Student distributions : the multivariate multi-scaled Student distributions (MMSD, [2]). The MMSD distribution extends the standard multivariate Student distribution by using the Gaussian scale mixture representation of Student distributions. This representation allows us to introduce multi-dimensional weights, which control different tail thickness of the distribution for each dimension, and provide a way to detect outlier data. In this way, we obtain a finer regulation of the influence of atypical data on the groups shapes, and so a greater flexibility of the clustering model.

We use an Expectation-Maximization algorithm (EM) to adjust a MMSD mixture on brain tumor MRI. The number of classes inside the mixture is selected by minimizing the Bayesian information criterion (BIC).

Our sample consists of healthy rats (n=8) and 4 groups of rats bearing a brain tumor model (n=8 per group), and 5 quantitative MRI parameter maps for each rat. We adjust a MMSD mixture on the healthy sample to detect tumor area in the tumor sample through the multi-dimensional weights. Then we characterize the tumor areas with another MMSD mixture and build a tumor dictionary which discriminates the 4 tumor.

Bibliography

[1] N. Coquery, O. Francois, B. Lemasson, C. Debacker, R. Farion, C. Rémy, and E. Barbier (2014), *Microvascular MRI and unsupervised clustering yields histology-resembling images in two rat models of glioma*, Journal of Cerebral Blood Flow & Metabolism, 34(8), 1354–62.

[2] F. Forbes, and D. Wraith (2014), *A new family of multivariate heavy-tailed distributions with variable marginal amounts of tailweights : Application to robust clustering*, Statistics and Computing, 24(6), 971–984.