



HAL
open science

Jet schemes and minimal toric embedded resolutions of rational double point singularities

Hussein Mourtada, Camille Plenat

► **To cite this version:**

Hussein Mourtada, Camille Plenat. Jet schemes and minimal toric embedded resolutions of rational double point singularities. 2015. hal-01255233

HAL Id: hal-01255233

<https://hal.science/hal-01255233>

Preprint submitted on 29 Nov 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Jet schemes and minimal toric embedded resolutions of rational double point singularities

Hussein Mourtada, Camille Plénat

July 16, 2018

Abstract

Using the structure of the jet schemes of rational double point singularities, we construct "minimal embedded toric resolutions" of these singularities. We also establish, for these singularities, a correspondence between a natural class of irreducible components of the jet schemes centered at the singular locus and the set of divisors which appear on every "minimal embedded toric resolution". We prove that this correspondence is bijective except for the E_8 singularity. This can be thought as an embedded Nash correspondence for rational double point singularities.

1 Introduction

In this article, we construct embedded resolutions of surfaces having rational double point singularities; also called simple singularities. The word simple refers to the fact that they have no moduli (i.e., a hypersurface singularity with the same topological type of a simple singularity is analytically isomorphic to the simple singularity [LeT]). But at the same time, they are simple from a resolution of singularities point of view, i.e., easy to resolve. The traditional approach to resolve singularities is to iterate blowing ups at smooth centers in order to make an invariant drop. This invariant takes values in a discrete ordered set with a smallest element (which detects smoothness). It should not only detect smoothness, but also be easy to compute so that its behavior can be followed when iterating the blowing ups. In this article, we adopt a different strategy to resolve a simple singularity $X \subset \mathbf{C}^3$: we construct an embedded toric resolution of $X \subset \mathbf{C}^3$. This construction is based on a deep invariant, the set \mathbf{EC} (essential components) of irreducible components of the jet schemes satisfying some natural properties. Recall that for $m \geq 0$, the m -th jet scheme of a variety X parametrizes morphisms $\text{Spec } \mathbf{C}[t]/(t^{m+1}) \rightarrow X$ (see section 2 for details). These are finite dimensional approximations of the space of arcs which parametrize germs of curves drawn on X . Actually the geometry of the m -th jet scheme is intimately related with the geometry of the set of arcs in a smooth ambient space containing X and which have "contact" with X larger than m . This explains in parts why while the arc

2010 Mathematics Subject Classification. 14E15, 14E18, 14M25.

Keywords Embedded Nash problem, Resolution of singularities, Toric Geometry.

This research was partially supported by the ANR-12-JS01-0002-01 SUSI. The second author is supported by Labex Archimède (ANR-11-LABX-0033) and the A*MIDEX project (ANR-11-IDEX-0001-02), funded by the "Investissements d'Avenir" French Government programme managed by the French National Research Agency (ANR)", and by the CLVA.

space of X detect information about abstract resolution of singularities (see [N]), jet schemes detect information about embedded resolution of singularities (see [Mo5], [LMR], [ELM]).

The other subject which this article considers is the minimality of embedded resolutions of singularities. In contrast with the abstract resolution case, there is no "universal" minimal embedded resolution for surface singularities. Therefore we need to make this notion more precise. Actually, the embedded resolution of $X \subset \mathbf{C}^3$ that we construct is toric and is obtained from a particular regular subdivision of the Newton dual fan Γ associated with X (see section 3 for the definition of Γ). Since simple singularities are Newton non-degenerate (see section 3), regular subdivisions of Γ give toric embedded resolutions of $X \subset \mathbf{C}^3$ (see [Ho],[Va],[LJ]). This is equivalent to say that an abstract resolution of singularities of the toric variety Z_Γ defined by the fan Γ gives an embedded toric resolution of $X \subset \mathbf{C}^3$. But the toric variety Z_Γ is of dimension 3 and hence thanks to a theorem by Bouvier and Gonzalez-Sprinberg [BGS], we know that there exists a toric resolution of singularities of Z_Γ where all the irreducible divisors of the exceptional locus are essential, i.e., the centers of the divisorial valuations associated with these divisors give irreducible components of the exceptional locus of any other toric resolution of singularities of Z_Γ . We call such a resolution of singularities a minimal toric resolution; in [BGS], it is called G -desingularization; we call a toric embedded resolution of $X \subset \mathbf{C}^3$ minimal if it corresponds to a minimal toric resolution of Z_Γ . Note that in general, a minimal toric embedded resolution of singularities is not unique, but the divisorial valuations associated with the irreducible divisors of its exceptional locus are the same for all minimal toric embedded resolution of singularities. We say that these last divisorial valuations are *embedded essential*.

We will prove that the toric embedded resolution of $X \subset \mathbf{C}^3$ which we construct from jet schemes is minimal for all simple singularities except for E_8 (Note that the E_8 singularity behaves exceptionally also from many other points of view [LeT2]). In particular we describe a bijection between the set of essential components of the jet schemes of simple singularities (except for E_8) and the set of embedded essential divisorial valuations. This can be thought as a solution of an embedded version of the Nash problem.

The choice of this class of singularities is related to the following facts:

First, for m big enough the number of irreducible components of their m -th jet scheme is constant; this simplifies the classification of the irreducible components we are interested in, when m varies. This is not the case in general (see [Mo2], [Mo3]).

Second, these singularities are Newton non-degenerate with respect to their Newton polyhedron, and therefore they have a toric embedded resolution; see [AGS],[O1],[O2]. Defining the class of irreducible components of jet schemes, mentioned above, is more subtle for Newton degenerate singularities [LMR]. This is related to a conjecture of Teissier on embedding any singularity in such way that it can be resolved by a toric morphism (see [T1], [T2], [Mo5]).

We would like to thank Shihoko Ishii, Monique Lejeune-Jalabert, Patrick Popescu-Pampu and Bernard Teissier for several discussions about this article.

2 Jet schemes

In this section, we begin by giving some preliminaries on jet schemes; we then recall from [Mo1] the structure of jet schemes of rational double point singularities and we extract from this structure some information about particular irreducible components of these jet schemes. These information will be used in the next section to obtain the canonical resolution of these

singularities. We will all the cases of simple singularities (see their defining equations below); the case of E_6 (see below the defining equation) has been treated in [Mo1], but we consider it briefly here for the convenience of the reader.

Let k be an algebraically closed field of arbitrary characteristic and X be a k -algebraic variety. For $m \in \mathbb{N}$, the functor

$$F_m : k\text{-Schemes} \longrightarrow \text{Sets}$$

$$\text{Spec}(A) \longrightarrow \text{Hom}_k(\text{Spec}A[t]/(t^{m+1}), X)$$

where A is a k -algebra, is representable by a k -scheme X_m [Is]. X_m is the m -th jet scheme of X , and F_m is isomorphic to its functor of points. In particular the closed points of X_m are in bijection with the $k[t]/(t^{m+1})$ points of X .

For $m, p \in \mathbb{N}, m > p$, the truncation homomorphism $A[t]/(t^{m+1}) \longrightarrow A[t]/(t^{p+1})$ induces a canonical projection $\pi_{m,p} : X_m \longrightarrow X_p$. These morphisms verify $\pi_{m,p} \circ \pi_{q,m} = \pi_{q,p}$ for $p < m < q$, and they are affine morphisms, they define a projective system whose limit is a scheme that we denote by X_∞ . This is the arc space of X .

Note that $X_0 = X$. We denote the canonical projection $\pi_{m,0} : X_m \longrightarrow X_0$ by π_m , and denote by Ψ_m the canonical morphisms $X_\infty \longrightarrow X_m$.

We now assume that $X \subset \mathbf{C}^3$ is defined by one of the following equations:

$$A_n, n \in \mathbb{N} : xy - z^{n+1} = 0.$$

$$D_n, n \in \mathbb{N}, n \geq 4 : z^2 - x(y^2 + x^{n-2}) = 0.$$

$$E_6 : z^2 + y^3 + x^4 = 0$$

$$E_7 : x^2 + y^3 + yz^3 = 0.$$

$$E_8 : z^2 + y^3 + x^5 = 0.$$

We denote by $X_m^0 := \pi_{m,0}^{-1}(O)$, where O is the origin of \mathbf{C}^3 ; it is the singular locus of X .

We will now associate a divisorial valuation over \mathbf{C}^3 with the irreducible component of $\mathcal{C}_m \subset X_m^0$ satisfying the property (\star) that for every irreducible component $\mathcal{C}_{m+1} \subset X_{m+1}^0$,

$$\text{if } \pi_{m+1,m}(\mathcal{C}_{m+1}) \subset \mathcal{C}_m \text{ then } \text{codim}(\mathcal{C}_{m+1}) > \text{codim}(\mathcal{C}_m). \quad (\star)$$

For $m \in \mathbb{N}$, let $\psi_m^a : \mathbf{C}_\infty^3 \longrightarrow \mathbf{C}_m^3$ be the canonical morphism, here the exponent a stands for "ambient". For $p \in \mathbb{N}$, we consider the following cylinder in the arc space

$$\text{Cont}^p(f) = \{\gamma \in \mathbf{C}_\infty^3; \text{ord}_t f \circ \gamma = p\}.$$

Since ψ_m^a is a trivial fibration, for every irreducible component $\mathcal{C}_m \subset X_m^0$, we have that

$$\psi_m^{a-1}(\mathcal{C}_m) \cap \text{Cont}^{m+1}(f)$$

is an irreducible component of $\text{Cont}^{m+1}(f)$, whenever this last intersection is non-empty. But from the property (\star) we have $\text{codim}(\mathcal{C}_{m+1}) > \text{codim}(\mathcal{C}_m)$, this implies

$$\psi_m^{a-1}(\mathcal{C}_m) \cap \text{Cont}^{m+1}(f) \neq \emptyset.$$

Let γ be the generic point of $\psi_m^{a-1}(\mathcal{C}_m) \cap \text{Cont}^{m+1}(f)$, then for every $h \in \mathbf{C}[x, y, z]$, we define the map $\nu_{\mathcal{C}_m} : \mathbf{C}[x, y, z] \rightarrow \mathbb{N}$, as follows

$$\nu_{\mathcal{C}_m}(h) = \text{ord}_t h \circ \gamma. \quad (1)$$

It follows from corollary 2.6 in [ELM], that \mathcal{C}_m is a divisorial valuation (see also [dFEI], [Re], prop. 3.7 (vii) applied to $\psi_m^{a-1}(\mathcal{C}_m)$).

Given $m \geq 1$, with an irreducible component \mathcal{C}_m of X_m^0 , we associate the following vector that we call the weight vector:

$$v(\mathcal{C}_m) = (\nu_{\mathcal{C}_m}(x), \nu_{\mathcal{C}_m}(y), \nu_{\mathcal{C}_m}(z)) \in \mathbb{N}^3.$$

In the following definition, we consider the irreducible components of X_m^0 that will be meaningful for the problem we are considering.

Definition 2.1. *The following set of particular irreducible components of X_m^0 :*

$$EC(X) := \{\mathcal{C}_m \subset X_m^0, m \geq 1 \text{ is an irreducible component satisfying } (\star) \text{ and}$$

$$v(\mathcal{C}_m) \neq v(\mathcal{C}_{m-1}) \text{ for any component } \mathcal{C}_{m-1} \text{ verifying } \pi_{m,m-1}(\mathcal{C}_m) \subset \mathcal{C}_{m-1}\}, \quad (2)$$

is called the set of essential components of X .

We also consider the following set of associated valuations :

$$EE(X) := \{\nu_{\mathcal{C}_m}, \mathcal{C}_m \in EC(X)\},$$

Where EE stands for Embedded-Essential, to say that these valuations are supposed to appear in every embedded toric resolution.

Note that the definitions of the sets $EC(X)$ and $EE(X)$ are ad-hoc to simple singularities. Indeed, the fact that these singularities are non-degenerate with respect to their Newton polyhedron implies that the meaningful valuations are monomial; this motivates the definition above of the set $EC(X)$. From another point of view, if we check carefully the equations, we will figure out that the components which does not belong to the set $EC(X)$, are at their generic points trivial fibrations above components which belong to $EC(X)$; that is why they are not useful and this phenomenon reflects the fact that the singularities we consider are Newton non-degenerate.

Note also that on one hand we are interested in the embeddings of the singularities in \mathbf{C}^3 that are defined by the equations we gave at the beginning of this section; this includes that we consider the variables x, y, z as given and hence the embedding of the torus in \mathbf{C}^3 is also given. On the other hand, since we are interested in toric resolutions, the exceptional locus of such a resolution will be defined as the inverse image of the complement of the torus, i.e., of the coordinate hyperplanes. So the fact that the x axis (defined by $y = z = 0$ in \mathbf{C}^3) and the y axis (defined by $x = z = 0$ in \mathbf{C}^3) belong to the singularities A_n , the y axis belongs to the singularities D_n and the z axis belongs to the singularity E_7 , suggests that if we want to find a toric resolution, we need also to consider the jets which go through respectively these coordinate axis. But the family of m -th jets whose center is a generic point on one of these axis is irreducible because our surfaces have isolated singularities at the origin. So the components

which come from those families are easy to determine; again since our singularities are Newton non-degenerate, we are only interested in the vectors associated with these components, so we only consider such components for small m , whenever the associated vector changes. For m large enough, the associated vector stabilizes.

A careful reading of [Mo4] (and of [Mo2]) produces a determination of the set EC for a rational double point $(X, 0)$. We will denote it by $EC(X)$. Let $f(x, y, z) \in \mathbf{C}[x, y, z]$ be the defining equation of a rational double point singularity X . We write

$$f\left(\sum_{i=0}^m x_i t^i, \sum_{i=0}^m y_i t^i, \sum_{i=0}^m z_i t^i\right) = \sum_{i=0}^{i=m} F_i t^i \pmod{t^{m+1}}, \quad (\diamond)$$

then the m -th jet scheme X_m of X is naturally embedded in

$$\mathbf{C}^{3(m+1)} = (\mathbf{C}^3)_m = \text{Spec} \mathbf{C}[x_i, y_i, z_i, i = 0, \dots, m]$$

and is defined by the ideal $I_m = (F_0, F_1, \dots, F_m)$. From Section 3 in [Mo4] and section 3.2 in [Mo1], we are able to determine the set $EC(X)$. This is the subject of the following lemma. We do not treat the E_8 singularity (which will be in some sense an exception to part of our formulation) because it takes too much space; we only give the weight vectors associated with the components in $EC(E_8)$ in the corollary that follows the lemma.

Lemma 2.2. *For an A_n singularity, $EC(A_n)$ is given by the components centered at the origin*

$$\{V(x_0, \dots, x_{l-1}, y_0, \dots, y_{m-l}, z_0), l = 1, \dots, m \text{ and } m = 1, \dots, n\},$$

and the components centered at the x respectively y axis are

$$\{V(y_0, \dots, y_l, z_0), l = 0, \dots, n\},$$

respectively

$$\{V(x_0, \dots, x_l, z_0), l = 0, \dots, n\}.$$

For a D_{2n} singularity, $EC(D_{2n})$ is given by the components centered at the origin

$$\{X_1^0 = V(x_0, y_0, z_1),$$

$$X_2^0 = V(x_0, y_0, z_0, z_1),$$

$$H_{2k} = V(x_0, y_0, z_0, z_1, y_1, y_2, \dots, y_{k-1}, z_2, z_3, \dots, z_k), k = 2, \dots, n-1,$$

$$H_{2k+1} = V(x_0, y_0, z_0, z_1, y_1, y_2, \dots, y_k, z_2, z_3, \dots, z_k), k = 1, \dots, n-2,$$

$$L_{2k+1} = V(x_0, x_1, y_0, z_0, z_1, y_1, y_2, \dots, y_{k-1}, z_2, z_3, \dots, z_k), k = 1, \dots, 2n-1\},$$

and the components centered at the y -axis are

$$V(x_0, z_0), \text{ and } V(x_0, x_1, z_0).$$

For an E_6 singularity we have

$$EC(E_6) = \{V(x_0, y_0, z_0), V(x_0, y_0, z_0, z_1), V(x_0, y_0, y_1, z_0, z_1),$$

$$V(x_0, x_1, y_0, y_1, z_0, z_1, z_2), V(x_0, x_1, y_0, y_1, y_2, z_0, z_1, z_2, z_3), \\ V(x_0, x_1, x_2, y_0, y_1, y_2, y_3, z_0, z_1, z_2, z_3, z_4, z_5)\}.$$

For an E_7 singularity, $EC(E_7)$ is given by the components centered at the origin

$$\{V(x_0, y_0, z_0), V(x_0, x_1, y_0, z_0), V(x_0, x_1, y_0, y_1, z_0), V(x_0, x_1, x_2, y_0, y_1, z_0), \\ V(x_0, x_1, x_2, y_0, y_1, y_2, z_0), V(x_0, x_1, x_2, y_0, y_1, z_0, z_1), V(x_0, x_1, x_2, x_3, y_0, y_1, y_2, z_0, z_1), \\ V(x_0, x_1, x_2, x_3, x_4, y_0, y_1, y_2, z_0, z_1), V(x_0, x_1, x_2, x_3, x_4, y_0, y_1, y_2, y_3, z_0, z_1), \\ V(x_0, x_1, x_2, x_3, x_4, x_5, y_0, y_1, y_2, y_3, z_0, z_1, z_2), V(x_0, x_1, x_2, x_3, x_4, x_5, x_6, y_0, y_1, y_2, y_3, y_4, z_0, z_1, z_2), \\ V(x_0, x_1, x_2, x_3, x_4, x_5, x_6, x_8, y_0, y_1, y_2, y_3, y_4, z_0, z_1, z_2, z_3)\},$$

and those centered above the z axis

$$\{V(x_0, y_0), V(x_0, y_0, y_1)\}.$$

From lemma 2.2, we deduce directly the set EE . First recall that a monomial valuation, defined on the ring $\mathbf{C}[x, y, z]$ and associated with a vector $a = (a_1, a_2, a_3) \in \mathbb{N}^3$, is defined by: for $h = \sum_{i=(i_1, i_2, i_3) \in \mathbb{N}^3} c_i x^{i_1} y^{i_2} z^{i_3} \in \mathbf{C}[x, y, z]$ then

$$\nu_a(h) = \min_{i \in \mathbb{N}^3; c_i \neq 0} a_1 i_1 + a_2 i_2 + a_3 i_3. \quad (3)$$

From lemma 2.2 we deduce:

Proposition 2.3. *The valuations that belong to $EE(A_n)$ are monomial. They are associated with the vectors $v(\mathcal{C}_m)$ where $\mathcal{C}_m \in EC(A_n)$. These vectors are*

$$(l, m - l + 1, 1), l = 1, \dots, m \text{ and } m = 1, \dots, n \\ (l, 0, 1), l = 1, \dots, n + 1, \\ (0, l, 1), l = 1, \dots, n + 1.$$

The valuations that belong to $EE(D_{2n})$ are monomial. They are associated with the vectors $v(\mathcal{C}_m)$ where $\mathcal{C}_m \in EC(D_{2n})$. These vectors are

$$U_1 = (1, 1, 1), U_2 = (1, 2, 2), \dots, U_{n-1} = (1, n-1, n-1), \\ W_1 = (1, 1, 2), W_2 = (1, 2, 3), \dots, W_{n-1} = (1, n-1, n), \\ V_1 = (2, 1, 2), V_2 = (2, 2, 3), \dots, V_{2n-2} = (2, 2n-2, 2n-1), \\ W_0 = (1, 0, 1), V_0 = (2, 0, 1).$$

The valuations that belong to $EE(E_6)$ are monomial. They are associated with the vectors $v(\mathcal{C}_m)$ where $\mathcal{C}_m \in EC(E_6)$:

$$(1, 1, 1), (1, 1, 2), (1, 2, 2), (2, 2, 3), (2, 3, 4), (3, 4, 6)$$

The valuations that belong to $EE(E_7)$ are monomial. They are associated with the vectors $v(\mathcal{C}_m)$ where $\mathcal{C}_m \in EC(E_7)$:

$$(1, 1, 0), (1, 2, 0), (1, 1, 1), (2, 1, 1), (2, 2, 1), (3, 2, 1), (3, 3, 1), (3, 2, 2),$$

$$(4, 3, 2), (5, 3, 2), (5, 4, 2), (6, 4, 3), (7, 5, 3), (9, 6, 4)$$

The valuations that belong to $EE(E_8)$ are monomial. They are associated with the vectors $v(\mathcal{C}_m)$ where $\mathcal{C}_m \in EC(E_8)$:

$$(1, 1, 1), (1, 1, 2), (1, 2, 2), (1, 2, 3), (2, 2, 3), (2, 3, 4), (2, 3, 5), (2, 4, 5), (3, 4, 6), \\ (3, 5, 7), (3, 5, 8), (4, 6, 8), (4, 6, 9), (4, 7, 10), (5, 7, 11), (5, 8, 11), (5, 8, 12), (5, 9, 13) \\ (5, 9, 14), (6, 10, 14), (6, 10, 15)$$

Proof. Let $a = (a_1, a_2, a_3) \in \mathbb{N}^3$. Assume that

$$\mathcal{C}_m = V(x_0, \dots, x_{a_1-1}, y_0, \dots, y_{a_2-1}, z_0, \dots, z_{a_3-1}) \subset \mathbf{C}_m^3$$

is an irreducible component of X_m , which belongs to $EC(X)$. Then, a generic point of $\Psi_m^{a-1}(\mathcal{C}_m) \cap \text{Cont}^{m+1}(f)$ is of the shape $(x_{a_1}t^{a_1} + \dots, y_{a_2}t^{a_2} + \dots, z_{a_3}t^{a_3} + \dots)$ where $(x_{a_1}, y_{a_2}, z_{a_3})$ is generic. For $h \in \mathbf{C}[x, y, z]$, it follows from the definition of $\nu_{\mathcal{C}_m}(h)$ (see equation (1)) and from the definition of $\nu_a(h)$ (see equation (3)) that $\nu_{\mathcal{C}_m}(h) = \nu_a(h)$. The proposition follows. \square

3 Toric minimal embedded resolutions

In this section, we define minimal toric embedded resolutions, and we give the motivations for this definition.

Let $N \simeq \mathbb{Z}^3$ be a lattice and let $\sigma \subset N_{\mathbb{R}^3} = N \otimes_{\mathbb{Z}} \mathbb{R}$ be the strictly convex cone generated by the vectors v_1, \dots, v_r , i.e.

$$\sigma = \langle v_1, \dots, v_r \rangle = \left\{ \sum_{i=1}^r \lambda_i v_i, \lambda_i \in \mathbb{R}_+ \right\}.$$

Let $f = \sum_{i=(i_1, i_2, i_3) \in \mathbb{N}^3} c_i x^{i_1} y^{i_2} z^{i_3}$ be a polynomial in $\mathbf{C}[x, y, z]$ with $f(0) = 0$. We denote by $NP^+(f)$ its Newton polyhedron at the origin: this is the convex hull in \mathbb{R}_+^3 of the set

$$\{(i_1, i_2, i_3) + \mathbb{R}_+^3, c_i \neq 0\},$$

where we have supposed that f is given by its Taylor expansion at the origin and let $NP(f)$ be the union of the compact boundaries of $NP^+(f)$, called Newton boundary. This polyhedron and its boundary depend on the choice of coordinates. Let $\Gamma(f)$ be the dual fan of $NP(f)$; we will call covectors vectors which lie in $\Gamma(f)$. A covector defines a linear map on \mathbb{R}^3 . For a positive covector L , i.e. with positive coefficients, we define the distance $d(L, f)$ as the minimal value of the linear map L at $x \in NP^+(f)$; let $\Delta(L, f) := \{x \in NP^+(f) : L(x) = d(L, f)\}$ be the dual face to L . We can then define $f_L(x)$ as the restriction of f to the dual face of L , $\Delta(L, f)$; that is

$$f_L(x) = \sum_{i=(i_1, i_2, i_3) \in \Delta(L, f)} c_i x^{i_1} y^{i_2} z^{i_3}.$$

We can find more details about Newton polyhedra, their dual fans and Newton non-degeneracy in [O2](see also [AGS]).

Definition 3.1. Let $f \in k[x, y, z]$ be such that $f(0) = 0$. We say that the hypersurface $\{f = 0\}$ is non-degenerate with respect to a covector L if the variety

$$F^*(L) = \{u \in k^{*3} / f_L(u) = 0\}$$

is a reduced smooth hypersurface in the torus k^{*3} . We say that f is non degenerate at the origin 0 with respect to its Newton Polyhedron $NP(f)$ if it is non degenerate for each covector.

Recall that a fan Σ is said to be regular if every cone $\sigma \in \Sigma$ is regular, i.e. the vectors generating σ are part of a basis of \mathbb{Z}^3 as a \mathbb{Z} -module, or equivalently (for σ a cone maximal dimension), if $\sigma = \langle v_1, v_2, v_3 \rangle$ then the absolute value $|\det(v_1, v_2, v_3)| = 1$. A very useful result about toric embedded resolutions of this type of singularities goes as follows.

Theorem 3.2. (*[Va],[O1],[AGS]*) We consider a pair $X \subset \mathbb{C}^3$ where $X = \{f = 0\}$ is a Newton non-degenerate singularity, then the following properties are equivalent:

- 1) A subdivision Σ of $\Gamma(f)$ is regular,
- 2) The proper birational morphism $\mu_\Sigma : Z_\Sigma \rightarrow \mathbb{C}^3 = \text{Spec } k[x, y, z]$ induced by Σ is an embedded resolution of singularities of the pair. Here Z_Σ is the toric variety associated with the fan Σ .

Note that the morphism μ_Σ is explicit in term of Σ ; more precisely, a regular cone $\sigma = \langle v_1, v_2, v_3 \rangle \in \Sigma$ of maximal dimension determines an affine chart of Z_Σ which is isomorphic to the affine space $\mathbb{C}^3 = \text{Spec } k[q, r, s]$ and the restriction of μ_Σ to this chart is given by

$$\begin{aligned} x &= q^{v_{1,1}} r^{v_{2,1}} s^{v_{3,1}} \\ y &= q^{v_{1,2}} r^{v_{2,2}} s^{v_{3,2}} \\ z &= q^{v_{1,3}} r^{v_{2,3}} s^{v_{3,3}}, \end{aligned}$$

where $v_i = (v_{i,1}, v_{i,2}, v_{i,3})$. Moreover, an edge of Σ (a cone of dimension 1) determines an orbit of Z_Σ whose Zariski closure defines a divisor on Z_Σ which is an irreducible component of the exceptional divisor of μ_Σ ($[O]$).

And the strict transform of $X = \{f = 0\}$ is defined as follows:

Definition 3.3. Let Σ be a regular subdivision of $\Gamma(f)$. The strict transform of $\{f = 0\}$ by μ_Σ is the Zariski closure of $(\mu_\Sigma)^{-1}(\mathbb{C}^{*3} \cap \{f = 0\})$.

We now recall the notion of “essential divisor”, which is one of the motivations of this work. For a singular variety, we can find infinitely many resolutions of singularities; the notion of essential divisor searches for intrinsic data in all resolutions of singularities of a given singular variety.

Definition 3.4. (*Essential divisors*)

Let X be a singular variety and $\pi : (\tilde{X}, E) \rightarrow (X, \text{sing}(X))$ be a resolution of singularities of $(X, \text{Sing}(X))$. Let $E_i \in E$ be a divisor which is an irreducible component of the exceptional locus E . We say that E_i is an essential divisor if for any other resolution of singularities $\pi' : (X', E') \rightarrow (X, \text{Sing } X)$, the center on X' of the divisorial valuation determined by E_i is an irreducible component of E' . A divisor E_i is said inessential if it is not essential.

It should be noted that the notion of essential divisor given above is associated with abstract resolutions of singularity and not with embedded resolutions of singularities. For surface singularities, a divisor is essential if it appears on the minimal resolution of singularities. Note

that a minimal resolution does not exist in general for a singular variety of dimension larger than 2.

For normal toric singularities, it is possible to determine essential divisors. For that we need the following definition of a minimal system of generators of a cone.

Let $N \cong \mathbf{Z}^n$ be a lattice and let $N_{\mathbf{R}}$ be the real vector space $N \otimes_{\mathbf{Z}} \mathbf{R}$.

Definition 3.5. (*[BGS], see also [AM]*) Let $\sigma \subset N_{\mathbf{R}}$ be a strongly convex rational polyhedral cone. The minimal system of generators of σ is the set:

$$G_{\sigma} = \{x \in \sigma \cap N \setminus 0 \mid \forall n_1, n_2 \in \sigma \cap N, x = n_1 + n_2 \Rightarrow n_1 = 0 \text{ or } n_2 = 0\}$$

An element in G_{σ} is also called irreducible. It is primitive by definition.

It follows from proposition 1.3 in [BGS], that the elements of G_{σ} appear on every regular fan which subdivides σ as an extremal vector, i.e. a primitive generator of a 1-dimensional cone of the fan. Note that such a subdivision determines a (toric) resolution of singularities of the toric variety defined by σ , and dimension 1 cones determine the irreducible components of the exceptional divisors. This gives a feeling for the following theorem which characterizes essential divisors on toric varieties.

Theorem 3.6. (*[BGS], [IK]*)

Let $\sigma \subset N_{\mathbf{R}}$ be a strongly convex rational polyhedral cone. The minimal system of generators G_{σ} is in bijection with the set of essential divisors of the toric variety V_{σ} . Therefore a primitive vector is in G_{σ} if and only if it is an extremal vector of any regular subdivision of the cone σ .

Actually, the fact that the divisors which correspond to elements of G_{σ} are toric essential (i.e. essential for toric resolutions of singularities) follows from [BGS] and the fact that they are essential for all resolutions of singularities follows from [IK]. It also follows from [BGS] that for a three dimensional toric variety defined by a cone σ there exists a toric resolution of singularities which is associated with a subdivision of σ all of whose 1-dimensional cones are rays associated to elements of G_{σ} . Such a resolution is called a G_{σ} -resolution. We think of such a resolution as a minimal toric resolution. Note that such a resolution does not exist in general for normal toric varieties of dimension larger than 3. Together with theorems 3.2 and 3.6, this discussion motivates the following two definitions:

Definition 3.7. (*Toric embedded-essential divisors*)

Let $X \subset \mathbf{C}^3$ where $X = \{f = 0\}$ is a Newton non-degenerate singularity. A toric embedded essential divisor E is a divisor which appears on every toric resolution of singularities of X (such a resolution exists thanks to theorem 3.2) as an irreducible component of the exceptional divisor. This is equivalent to say that E corresponds to an element of G_{σ} for some cone $\sigma \in \Gamma(f)$ of maximal dimension.

Definition 3.8. (*Minimal toric embedded resolution*) Let $X = \{f = 0\} \subset \mathbf{C}^3$ be a Newton non-degenerate singularity at the origin. A toric embedded resolution of $X \subset \mathbf{C}^3$ which is associated with a subdivision Σ of $\Gamma(f)$ is minimal if the only 1-dimensional cones which appear in Σ are determined by elements of G_{σ} for some cone $\sigma \in \Gamma(f)$ of maximal dimension and the 1-dimensional cones of $\Gamma(f)$. This is equivalent to say that all the irreducible components of the exceptional divisor of a minimal toric embedded resolution are toric embedded essential divisors.

Note that these definitions make sense also for non-isolated singularities as in [ACT] for instance.

4 Toric minimal embedded resolutions for simple singularities


In this section, we construct toric embedded resolutions from the data of the jet schemes of simple singularities that we gave in section 2. We also prove that for all singularities of this type, except for the E_8 singularity, the essential components of the jet schemes (introduced in section 2) correspond to toric embedded-essential divisors. We will treat each type of singularity separately: first we construct a toric resolution of singularities using the weight vectors associated in section 2 with essential components, then we construct a regular subdivision of the dual fan of the singularity embedded in \mathbf{C}^3 , and finally by using a theorem from [BGS] (see also [AM]) which characterizes toric essential divisor for toric varieties, we will prove for simple singularities (except for the singularity of type E_8) the minimality of our toric embedded resolution of singularities.

Remark 4.1. *Instead of drawing the dual fan of each polyhedron in \mathbb{R}^3 (with coordinates x, y, z) we use the trace of the dual fan on the hyperplane defined by the equation $x + y + z = 1$. Vectors are then represented by points. We call this trace the **Newton face** or simply, with some abuse of notation **dual fan**.*

4.1 The A_n singularities

Theorem 4.2. *The weight vectors of A_n give a toric minimal embedded resolution of the singularity. In particular the set of essential components $EC(A_n)$ is in bijection with the divisors which appear on every toric minimal embedded resolution of these singularities.*

The Newton polyhedron and its dual fan are the following:



The dual fan is not simplicial; a resolution of this singularity is given by a regular and simplicial subdivision of the Newton dual fan.

The set $EE(A_n)$ of valuations, found after a careful reading of [Mo1], is

$$(l, m - l + 1, 1), l = 1, \dots, m \quad \text{and} \quad m = 1, \dots, n$$

$$(l, 0, 1), l = 1, \dots, n + 1,$$

$$(0, l, 1), l = 1, \dots, n + 1.$$

(see lemma 2.2 and proposition 2.3). We reorder this set in the following way in order to study them more easily (see the remarks below):

$$\begin{aligned}
EE(A_n) := & \{(n, 0, 1), \dots, (1, 0, 1), (0, n, 1), \dots, (0, 1, 1), (1, 1, 1), (2, 1, 1), \dots, (n, 1, 1), \\
& (1, 2, 1) \dots (1, n, 1), (2, 2, 1), (3, 2, 1) \dots (n-1, 2, 1), \\
& (2, 3, 1) \dots (2, n-1, 1) \dots (\lceil \frac{n+1}{2} \rceil, \lceil \frac{n+1}{2} \rceil, 1)\}
\end{aligned}$$

to which we add , if n is even, $(\lceil \frac{n+1}{2} \rceil + 1, \lceil \frac{n+1}{2} \rceil, 1)$ et $(\lceil \frac{n+1}{2} \rceil, \lceil \frac{n+1}{2} \rceil + 1, 1)$.

The weight vectors are positioned on the following figure :




Figure 1 : Positions of $EE(A_4)$ and $EE(A_5)$

CLAIM: these vectors are irreducible, i.e. they represent essential components in the resolution.

Before proving this claim, we study the position of the vectors in the Newton face of dual fan, and show that they provide a toric embedded resolution.

Some remarks on the vectors of $EE(A_n)$:

(with abuse of notation, the vectors are considered as points on the Newton face of the cone $[(e_3, (n+1, 0, 1), (0, n+1, 1))]$).

1. We first prove that the vectors are rational sums of $e_1, e_2, e_3, v_1 = (n+1, 0, 1), v_2 = (0, n+1, 1)$. By the symmetry of the singularity, we only have to look at half of the vectors:

we have :

$$(k, l, 1) = \frac{k}{n+1}v_1 + \frac{l}{n+1}v_2 + \frac{n-k-l+1}{n+1}e_3$$

for all $1 \leq k \leq n$ and $1 \leq l \leq n-k+1$.

Moreover $(k, 0, 1) = \frac{k}{n+1}v_1 + \frac{n}{n+1}e_3$.

This implies that they all belong to the cone $(e_3, (n+1, 0, 1), (0, n+1, 1))$.

2. One can remark the symmetry we obtain. It will follow that we have to do only half of the computations;
3. The vectors $(1, 1, 1) \dots (n, 1, 1)$ are on the line $((1, 1, 1), (1, 0, 0))$ as $(k, 1, 1) = (1, 1, 1) + (k-1)(1, 0, 0)$. The same holds for $(k, k, 1) \dots (n+1-k, k, 1)$ for $k \leq \lceil \frac{n+1}{2} \rceil$;

4. Homogeneous vectors (α, β, γ) such that $\alpha + \beta + \gamma = k$, for $3 \leq k \leq n + 2$ are also on the same line.

All these remarks show us a way to obtain an almost symmetric resolution from the vectors. Many algorithms work, we propose one of them to the reader.

ALGORITHM

First place $(1, 1, 1)$ and make the triangle $((0, 0, 1), (n + 1, 0, 1), (0, n + 1, 1))$ simplicial. Then place successively the vectors $(k, 0, 1), \dots, (1, 0, 1)$, joining each new vector to $(1, 1, 1)$, to make the resolution simplicial. Do the same symmetrically with $(0, k, 1), \dots, (0, 1, 1)$.

One has

$$(k, 0, 1) = \frac{k}{n+1}(n+1, 0, 1) + \frac{n+1-k}{n+1}(0, 0, 1)$$

Thus these vectors are primitive vectors.

For A_{2n} :

Place the vectors $(n, 1, 1)$ and $(1, n, 1)$ and the corresponding edges. One can remark that the vectors $(k, 1, 1)$ lie on the line $((n, 1, 1), (1, 1, 1))$. It is at this point that the resolution is not symmetrical: if one chooses $(n, 1, 1)$, then one makes the trapezium simplicial by adding the edges $((n, 1, 1), (0, 1, 0))$ and $((n, 1, 1), (1, 0, 0))$. So in the toric resolution, one will have the edges joining the vectors on $((n + 1, 0, 1), (0, n + 1, 1))$ to $(0, 1, 0)$. Conversely, if one chooses $(1, n, 1)$, one will have the edges joining the vectors on $((n + 1, 0, 1), (0, n + 1, 1))$ to $(1, 0, 0)$.

For A_{2n+1} , one had $[(\frac{n+1}{2}, \frac{n+1}{2}, 1)]$ first, the trapezium thus becomes divided into three parts and the resolution become symmetrical.

Then add alternatively $(2, 2, 1), (n - 1, 2, 1), (2, n - 1, 1), (3, 3, 1) \dots ([\frac{n+1}{2}], [\frac{n+1}{2}], 1)$, and if n is even $([\frac{n+1}{2}] + 1, [\frac{n+1}{2}], 1), ([\frac{n+1}{2}], [\frac{n+1}{2}] + 1, 1)$. Then add the remaining vectors in the natural order, with the corresponding edges.




Figure 2 : Resolution of A_4 step by step

CLAIM : This is a toric resolution for A_{2n} .

Proof. • Let us look at triangles $[(1, 1, 1), (k, 0, 1), (k + 1, 0, 1)]$ and $[(1, 1, 1), (0, k, 1), (0, k + 1, 1)]$ for $1 \leq k \leq n$,
we have:

$$\begin{vmatrix} 1 & 0 & 0 \\ 1 & k & k + 1 \\ 1 & 1 & 1 \end{vmatrix} = -1 \quad \text{and} \quad \begin{vmatrix} 1 & k & k + 1 \\ 1 & 0 & 0 \\ 1 & 1 & 1 \end{vmatrix} = 1$$

• We also have the triangles $[(1, 0, 1), (1, 1, 1), (0, 0, 1)]$ et $(0, 1, 1), (1, 1, 1), (0, 0, 1)]$ of determinant 1.

• Now, we look at $[(k, n - k, 1), (k - 1, n - (k - 1), 1), (0, 1, 0)]$.

We have

$$\begin{vmatrix} k & k - 1 & 0 \\ n - k & n - (k - 1) & 1 \\ 1 & 1 & 0 \end{vmatrix} = 1$$

If n is even, we have the triangles $t [(\lceil \frac{n+1}{2} \rceil - 1, \lceil \frac{n+1}{2} \rceil, 1), (\lceil \frac{n+1}{2} \rceil, \lceil \frac{n+1}{2} \rceil + 1, 1), (0, 1, 0)]$ and if n is odd, we have the triangles $[(\lceil \frac{n+1}{2} \rceil + 1, \frac{n+1}{2}, 1), (\frac{n+1}{2}, \frac{n+1}{2}, 1), (0, 1, 0)]$ et $[(\frac{n+1}{2}, \frac{n+1}{2} - 1, 1), (\frac{n+1}{2}, \frac{n+1}{2}, 1), (0, 1, 0)]$ with determinant 1.

• The cones $[e_1, (n + 1, 0, 1), (n, 1, 1)]$ and $[e_1, e_2, (n, 1, 1)]$ are non singular.

• Finally we have to look at the triangles $[(k + l, k, 1), (k + l - 1, k, 1), (k + 1, k + 1, 1)]$ and $[(k + l, k, 1), (k + l - 1, k, 1), (n + 2 - k, k - 1, 1)]$ for all $1 \leq k \leq \lceil \frac{n+1}{2} \rceil$ and $k \leq l \leq n - 1$ (and to the symmetrical one too).

One has

$$\begin{vmatrix} k + l & k + l - 1 & k + 1 \\ k & k & 1 \\ 1 & 1 & 1 \end{vmatrix} = 1 \quad \text{and} \quad \begin{vmatrix} k + l & k + l - 1 & n + 2 - k \\ k & k & k - 1 \\ 1 & 1 & 1 \end{vmatrix} = 1$$

□

Now we have to prove that in fact this resolution is minimal or a G -resolution (according to C. Bouvier and G. Gonzalez-Sprinberg).

First of all, the weight vectors $(l, m - l + 1, 1), l = 1, \dots, m$ and $m = 1, \dots, n$ are clearly primitive as their third component is 1.

Consider the simplicial cone generated by $e_3, (n + 1, 0, 1), (0, n + 1, 1)$; all the weight vectors belong to this cone.

Let us give the same names to the points in \mathbb{R}^3 which are the end of the vectors starting from 0 and the vectors themselves.

Lemma 4.3. *The points in $\mathbb{R}^3 \{e_3, (n+1, 0, 1), (0, n+1, 1), (l, m-l+1, 1), l = 1, \dots, m$ and $m = 1, \dots, n\}$ belong to the face $(e_3, (n+1, 0, 1), (0, n+1, 1))$ of the tetrahedron $[0, e_3, (n+1, 0, 1), (0, n+1, 1)]$. This implies that the set $\{e_3, (n+1, 0, 1), (0, n+1, 1), (l, m-l+1, 1), l = 1, \dots, m$ and $m = 1, \dots, n\}$ is free over \mathbb{Z} , i.e. none of the vectors is reducible into other vectors of this set.*

Suppose that the lemma is proved. Then the only primitive vectors that could generate the characteristic vectors should be vectors inside the tetrahedron $[0, e_3, (n+1, 0, 1), (0, n+1, 1)]$. Suppose that there is such a vector. By th. 1.10 of [BGS], it corresponds to an essential divisor in the toric resolution of the cone $[e_3, (n+1, 0, 1), (0, n+1, 1)]$, i.e., it appears in every regular subdivision as an extremal vector. But the vectors $(l, m-l+1, 1), l = 1, \dots, m$ and $m = 1, \dots, n$, we get such a regular subdivision, so there is no primitive vector in the tetrahedron $[0, e_3, (n+1, 0, 1), (0, n+1, 1)]$.

Let us prove the lemma:

Proof. The equation of the face $(e_3, (n+1, 0, 1), (0, n+1, 1))$ is $z-1=0$. Clearly all the points $\{e_3, (n+1, 0, 1), (0, n+1, 1), (k, n-k, 1), (k-1, n-(k-1), 1), (0, 1, 0), (l, m-l+1, 1), l = 1, \dots, m$ and $m = 1, \dots, n\}$ belongs to it. \square

4.2 The D_n singularities

Theorem 4.4. *The weight vectors of D_n give a "canonical" toric minimal embedded resolution of the singularity. In particular the set of essential components $EC(D_n)$ is in bijection with the divisors which appear on every toric minimal embedded resolution of these singularities.*

Proof. Let us do it for D_{2n} .

Let us recall what are the Newton polyhedron for D_{2n} and the associated dual cone:




Figure 3 : Newton polyhedron of D_{2n} and its dual cone

First we make the dual polyhedron simplicial by adding the edge $[e_1, (2, 2n-2, 2n-1)]$. Let us call \mathcal{C}_1 be the cone $[e_1, (2, 0, 1), (2, 2n-2, 2n-1)]$, \mathcal{C}_2 be the cone $[e_1, e_2, (2, 2n-2, 2n-1)]$, \mathcal{C}_3 be the cone $[e_3, (2, 0, 1), (2, 2n-2, 2n-1)]$, \mathcal{C}_4 be the cone $[e_2, e_3, (2, 2n-2, 2n-1)]$.

First of all, to obtain a toric resolution of singularities of D_{2n} with equation $z^2 + x(y^2 + x^{2n-2}) = 0$, as it is not commode, one has to blow up the y axes, i.e. to add the vector $(1, 0, 1)$. Let us now consider the weight vectors and show some of their properties:

- The vectors $U_i = (1, i, i)$ ($1 \leq i \leq n-1$) are primitive and belong to the cone \mathcal{C}_2 . More precisely they lie onto the line $(e_1, (0, 1, 1))$:

In fact we have the following equalities:

$$U_i = \frac{2n-2i-1}{2n-1}e_1 + \frac{i}{2n-1}e_2 + \frac{i}{2n-1}(2, 2n-2, 2n-1)$$

and if we note $U_0 = (1, 0, 0)$, then for $1 \leq i \leq n-1$,

$$U_i = U_{i-1} + (0, 1, 1)$$

- The vectors $W_i = (1, i, i + 1)$ ($1 \leq i \leq n - 1$) are primitive and belong to the cone \mathcal{C}_3 . More precisely they lie onto the line $((1, 0, 1), (0, 1, 1))$:
In fact we have the following equalities:

$$W_i = \frac{n-1}{2n-2}e_3 + \frac{n-i-1}{2n-2}(2, 0, 1) + \frac{i}{2n-2}(2, 2n-2, 2n-1)$$

and if we note $W_0 = (1, 0, 1)$, then for $1 \leq i \leq n - 1$, $W_i = W_{i-1} + (0, 1, 1)$

- The vectors $V_i = (2, i, i + 1)$ ($1 \leq i \leq 2n - 2$) are primitive and belong to the cones \mathcal{C}_1 and \mathcal{C}_3 . More precisely they lie onto the line $((2, 0, 1), (0, 1, 1))$:
In fact we have the following equalities:

$$V_i = \frac{2n-2-i}{2n-2}(2, 0, 1) + \frac{i}{2n-2}(2, 2n-2, 2n-1)$$

and, if we note $V_0 = (2, 0, 1)$, then for $1 \leq i < 2n - 2$, $V_i = V_{i-1} + (0, 1, 1)$.

CLAIM: the simplicial decomposition of the dual Newton cone of D_{2n} obtained by adding, the edge $[e_1, (2, 2n-2, 2n-1)]$, the vector $(1, 0, 1)$, then the vectors U_1 to U_{n-1} in the natural order, the vectors W_1 to W_{n-1} in the natural order and the vectors V_1 to V_{2n-2} in the natural order too, give a toric minimal embedded resolution of D_{2n} .

The resolved dual cone is of the form:




Figure 4 : Resolution of D_{2n}

Proof of the claim: by the previous observations, the vectors U_i, V_i, W_i above are primitive; thus it remains to prove that the cones obtained are non singular, i.e. they have determinant $+1$ or -1 and using the same proof as for A_n , we prove then that the vectors are irreducible.

We add first the vector $(1, 0, 1)$, which corresponds to the blowing up of the y -axes. Then, the cone \mathcal{C}_3 is decomposed into two cones : $\mathcal{C}_{31} = ((1, 0, 1), (2, 0, 1), (2, 2n - 2, 2n - 1))$ and $\mathcal{C}_{32} = ((1, 0, 1), e_3, (2, 2n - 2, 2n - 1))$. By the above observation, we have now that each group of characteristic vectors are in different cones.

- Decomposition of the cone \mathcal{C}_1 :

Let us note $U_0 = e_1$.

We add the vectors from U_1 to U_{n-1} and show by induction that we obtain at the end regular cones. At each step the cone where lie the new A_i is decomposed into three cones:




Figure 5 : Resolution of D_4 step by step

(U_i, U_{i-1}, e_2) , $((U_i, U_{i-1}, (2, 2n - 2, 2n - 1))$ and $(U_i, e_2, (2, 2n - 2, 2n - 1))$. That last one contains U_{i+1} .

$$\begin{vmatrix} 1 & 1 & 0 \\ i & i-1 & 1 \\ i & i-1 & 0 \end{vmatrix} = 1, \quad \begin{vmatrix} 1 & 1 & 2 \\ i & i-1 & 2n-2 \\ i & i-1 & 2n-1 \end{vmatrix} = -1, \quad \begin{vmatrix} 1 & 2 & 0 \\ i & 2n-2 & 1 \\ i & 2n-1 & 0 \end{vmatrix} = n-1-2i$$

That last determinant is not equal to $+1$ or -1 , so we add U_{i+1} and so on.
For $i = n - 1$ the last determinant is equal to one.

- Decomposition of the cone \mathcal{C}_{31} :

Let us note $W_0 = (1, 0, 1)$.

We add the vectors from W_1 to W_{n-1} and show by induction that we obtain at the end regular cones. At each step the cone where lie the new W_i is decomposed into three cones: (W_i, W_{i-1}, e_3) , $((W_i, W_{i-1}, (2, 2n - 2, 2n - 1))$ and $(W_i, e_3, (2, 2n - 2, 2n - 1))$. That last one contains W_{i+k} for $k \geq 1$.

We have :

$$\begin{vmatrix} 1 & 1 & 0 \\ i & i-1 & 0 \\ i+1 & i & 1 \end{vmatrix} = -1, \quad \begin{vmatrix} 1 & 1 & 2 \\ i & i-1 & 2n-2 \\ i+1 & i & 2n-1 \end{vmatrix} = 1, \quad \begin{vmatrix} 1 & 2 & 0 \\ i & 2n-2 & 0 \\ i+1 & 2n-1 & 1 \end{vmatrix} = 2n-2-2i$$

That last determinant is not equal to $+1$ or -1 , so we add A_{i+1} and so on.
For $i = n - 1$ the last determinant is equal to one.

- Decomposition of the cones \mathcal{C}_{32} and \mathcal{C}_2 :

Let us note $V_0 = (2, 0, 1)$.

We add the vectors from V_1 to V_{2n-2} and show by induction that we obtain at the end regular cones. At each step the two cones where lie the new C_i are decomposed into two cones themselves: (V_i, V_{i-1}, e_1) , $(V_i, V_{i-1}, (1, 0, 1))$ and $(V_i, e_3, (2, 2n - 2, 2n - 1))$, $(V_i, (1, 0, 1), (2, 2n - 2, 2n - 1))$. Those last ones contain V_{i+1} and are not regular.

$$\begin{vmatrix} 2 & 2 & 1 \\ i & i-1 & 0 \\ i+1 & i & 0 \end{vmatrix} = 1$$

$$\begin{vmatrix} 2 & 2 & 1 \\ i & i-1 & 0 \\ i+1 & i & 1 \end{vmatrix} = \begin{vmatrix} 2 & 2 & 1 \\ i & 2n-2 & 0 \\ i+1 & 2n-1 & 0 \end{vmatrix} = 2ni - i - 2ni + 2i - 2n + 2 = i - 2n + 2$$

and

$$\begin{vmatrix} 2 & 2 & 1 \\ i & 2n-2 & 0 \\ i+1 & 2n-1 & 1 \end{vmatrix} = 2n - 2 - i$$

For $i = 2n - 3$ the two last determinants are equal to $+1$ or -1 .

Moreover we have obtained a G -resolution:

As for A_n singularity, one can show that W_i (resp. U_i and V_i) belongs to the base of the cone $(e_3, (2, 0, 1), (2, 2n-2, 2n-1))$ (resp. $(e_1, (2, 0, 1), (2, 2n-2, 2n-1)) \cap (e_3, (2, 0, 1), (2, 2n-2, 2n-1))$, and $(e_1, e_2, (2, 2n-2, 2n-1))$, whose equation is $y-z+1=0$ (resp. $y-z+1=0; x+y-z-1=0$ and $x+y-z-1=0$). With these vectors we obtain a regular subdivision of each cone, which implies that there are no other primitive vectors inside each cone and that the characteristic vectors are irreducible. So we also obtain a G -resolution for D_{2n} . \square

Remark: the fact that each vector "belongs" to the base of the cone is equivalent to saying that the volume of the subdivisions stay the same during the process of the resolution.

4.3 The E_6 singularity, according to [Mo4]

Now consider the singularity E_6 given by the equation $z^2 + y^3 + x^4 = 0$. A simplicial dual Newton polyhedron associated to it is given on figure 6.

Theorem 4.5. *The set of weight vectors $EE(E_6)$ is*

$$(1, 1, 1), (1, 1, 2), (1, 2, 2), (2, 2, 3), (2, 3, 4), (3, 4, 6)$$

Call them in the lexicographic order.

The weight vectors of E_6 give a "canonical" toric minimal embedded resolution of the singularity. In particular the set of essential components $EC(E_6)$ is in bijection with the divisors which appear on every toric minimal embedded resolution of these singularities.




Figure 6 : Newton dual polyhedron of E_6 and its resolution

Proof. The vectors are in different subcones of the dual fan. To make the fan regular, we divide each cone using the vectors. So we have to look first at the positions of each vector. One has :

$$\{(1, 1, 1), (1, 2, 2), (2, 2, 3), (2, 3, 4)\} \subset C_1\}$$

$$\{(2, 3, 4), (1, 2, 2)\} \subset C_2\}$$

$$\{(1, 1, 2), (2, 2, 3)\} \subset C_3\}$$

As for the previous singularities, one can prove that adding the weight vectors in each cone (in the alphabetic order, see fig. 6), gives a minimal resolution for each of them. See fig. 6 for the picture. □

4.4 The E_7 singularity

Now consider the singularity E_7 given by the equation $x^2 + y^3 + yz^3 = 0$. A simplicial dual Newton polyhedron associated to it is given on figure 7.

Theorem 4.6. *The set of weight vectors is*

$$(1, 1, 0), (1, 2, 0) = O_2, (1, 1, 1), (2, 1, 1), (2, 2, 1), (3, 2, 1), (3, 3, 1),$$

$$(3, 2, 2), (4, 3, 2), (5, 3, 2), (5, 4, 2), (6, 4, 3), (7, 5, 3), (9, 6, 4) = O_1$$

Call them in the lexicographic order (except O_1 and O_2).

The weight vectors of E_7 give a "canonical" toric minimal embedded resolution of the singularity. In particular the set of essential components $EC(E_7)$ is in bijection with the divisors which appear on every toric minimal embedded resolution of these singularities.




Figure 7 : Newton dual polyhedron of E_7 and its resolution

Proof. The vectors are in different subcones of the dual fan (see fig.7 for denomination of the cones). To make the fan regular, we divide each cone thanks to the vectors. So first we have to look at the positions of each vector. One has :

$$\{(1, 1, 1), (2, 2, 1), (3, 2, 2), (4, 3, 2), (6, 4, 3)\} \subset C_1\}$$

$$\{(2, 1, 1), (3, 2, 2), (5, 3, 2), (6, 4, 3)\} \subset C_2\}$$

$$\{(1, 1, 0), (3, 2, 1), (3, 3, 1), (5, 3, 2), (5, 4, 2), (7, 5, 3)\} \subset C_3\}$$

$$\{(3, 3, 1), (5, 4, 2), (7, 5, 3)\} \subset C_4\}$$

As for the previous singularities, one can prove that adding the previous vectors in each cone (in the lexicographic order), gives a minimal resolution for each of them. See fig.7 for the picture.

□

4.5 The E_8 singularity

Now consider the singularity E_8 given by the equation $z^2 + y^3 + x^5 = 0$. A simplicial dual Newton polyhedron associated to it is given on figure 8.

Proposition 4.7. *The weight vector gives an embedded resolution of E_8 but this resolution is not minimal.*

Proof. Recall that the set of weight vectors are in this case :

$$\begin{aligned} &(1, 1, 1), (1, 1, 2), (1, 2, 2), (1, 2, 3), (2, 2, 3), (2, 3, 4), (2, 3, 5), (2, 4, 5), (3, 4, 6), \\ &(3, 5, 7), (3, 5, 8), (4, 6, 8), (4, 6, 9), (4, 7, 10), (5, 7, 11), (5, 8, 11), (5, 8, 12), (5, 9, 13) \\ &(5, 9, 14), (6, 10, 14), (6, 10, 15) = O \end{aligned}$$




Figure 8 : Newton dual polyhedron of E_8 and its resolution

One can show that we get a regular subdivision of each cone thanks to the following characteristic vectors

$$\begin{aligned} &(1, 1, 1), (1, 1, 2), (1, 2, 2), (1, 2, 3), (2, 2, 3), \\ &(2, 3, 4), (2, 3, 5), (2, 4, 5), (3, 4, 6), (3, 5, 7), (3, 5, 8), (4, 6, 9), (4, 7, 10), (5, 8, 12), (6, 10, 15) = O \end{aligned}$$

(We call them in the lexicographical order on fig.8)

Moreover they are irreducible (one can show it in the same way as above), so they give subdivision which induces a minimal embedded toric resolution for E_8 .

In fact we have

- $(4, 6, 8) = (2, 3, 4) + (2, 3, 4)$
- $(5, 7, 11) = (4, 6, 9) + (1, 1, 2)$
- $(5, 8, 11) = (4, 7, 10) + (1, 1, 1)$
- $(5, 9, 13) = (3, 5, 8) + (2, 4, 5)$
- $(6, 9, 13) = (3, 4, 6) + (3, 5, 7)$

- $(5, 9, 14) = (3, 4, 6) + (3, 5, 8)$
- $(6, 10, 14) = (3, 5, 7) + (3, 5, 7)$

□

4.6 A remark on abstract resolution of singularities of simple singularities

For a simple singularity, the restriction of a minimal embedded resolution to the strict transform of the singularity gives its minimal abstract resolution; note that not all the divisors that appear on the minimal embedded resolutions meet the strict transform, only those which corresponds to weight vectors which belong to the two dimensional cones of the Newton dual fan. Moreover, these last divisors may meet the strict transform along more than one irreducible component.

References

- [AM] S. Aguzzoli, D. Mundici, An algorithmic desingularization of 3-dimensional toric varieties. *Tohoku Math. J. (2)* 46 (1994), no. 4, 557-572.
- [ACT] A. Altıntaş Sharland, G. Cevik, M. Tosun, Nonisolated forms of rational triple point singularities of surfaces and their resolutions. *Rocky Mountain J. Math.* 46 (2016), no. 2, 357-388.
- [AGS] F. Aroca, Fuensanta, M. Gomez-Morales, K. Shabbir, Torical modification of Newton non-degenerate ideals. *Rev. R. Acad. Cienc. Exactas Fs. Nat. Ser. A Math. RACSAM* 107 (2013), no. 1, 221-239.
- [B] C. Bouvier, Diviseurs essentiels, composantes essentielles des variétés toriques singulières, *Duke Math. J.* **91** (1998), 609-620.
- [BGS] C. Bouvier, G. Gonzalez-Sprinberg, Système générateur minimal, diviseurs essentiels et G-désingularisations des variétés toriques, *Tohoku Math. J. (2)* **47** (1995), 125-149.
- [CP] V. Cossart, O. Piltant, Olivier Resolution of singularities of threefolds in positive characteristic. II. *J. Algebra* 321 (2009), no. 7, 1836-1976.
- [ELM] L. Ein, R. Lazarsfeld, M. Mustata, Contact loci in arc spaces. *Compos. Math.* 140 (2004), no. 5, 1229-1244.
- [dFEI] T. de Fernex, L. Ein, S. Ishii, Divisorial valuations via arcs. *Publ. Res. Inst. Math. Sci.* 44 (2008), no. 2, 425-448.
- [GGS] G. Gonzalez-Sprinberg, M. Lejeune-Jalabert Modèles canoniques plongés. I. *Kodai Math. J.* 14 (1991), no. 2, 194-209. doi:10.2996/kmj/1138039394.
- [Hi] H. Hironaka, Resolution of singularities of an algebraic variety over a field of characteristic zero. I, II. *Ann. of Math. (2)* 79 (1964), 109-203; *ibid.* (2) 79 1964 205-326.
- [Ho] A.G. Hovanskii, Newton polyhedra, and toroidal varieties. (Russian) *Funkcional. Anal. i Priložen.* 11 (1977), no. 4, 56-64, 96.
- [Is] S. Ishii, Jet schemes, arc spaces and the Nash problem, *C.R.Math. Rep. Acad. Canada*, 29 (2007) 1-21.

- [IK] S. Ishii, J. Kollár, The Nash problem on arc families of singularities. *Duke Math. J.* 120 (2003), no. 3, 601-620.
- [LeT] D.T. Lê, M. Tosun, Simple singularities and simple Lie algebras. *TWMS J. Pure Appl. Math.* 2 (2011), no. 1, 97-111.
- [LeT2] D.T. Lê, M. Tosun, Combinatorics of rational singularities. *Comment. Math. Helv.* 79 (2004), no. 3, 582-604.
- [LJ] M. Lejeune–Jalabert, Désingularisation explicite des surfaces quasi-homogènes dans \mathbf{C}^3 , *Nova Acta Leopoldina*, **NF 52**, Nr 240 (1981), 139–160.
- [LMR] M. Lejeune-Jalabert, H. Mourtada, A. Reguera Jet schemes and minimal embedded desingularization of plane branches. *Revista de la Real Academia de Ciencias Exactas, Fisicasy Naturales. Serie A. Matematicas: Volume 107, Issue 1 (2013)*, Page 145-157.
- [Mo1] H. Mourtada, Jet schemes of rational double point singularities, *Valuation Theory in Interaction*, EMS Ser. Congr. Rep., Eur. Math. Soc., Sept. 2014, pp: 373-388, DOI: 10.4171/149-1/18.
- [Mo2] H. Mourtada, Jet schemes of complex branches and equisingularity, *Ann. Inst. Fourier* 61, No. 6, 2313-2336 (2011).
- [Mo3] H. Mourtada, Jet schemes of toric surfaces, *C. R., Math., Acad. Sci. Paris* 349, No. 9-10, 563-566 (2011).
- [Mo4] H. Mourtada, Jet schemes of normal toric surfaces, to appear in *Bulletin de la SMF*.
- [Mo5] H. Mourtada, Jet schemes and minimal generating sequences of divisorial valuations in dimension 2, *Michigan Math. J.* Volume 66, Issue 1 (2017), 155-174.
- [N] J. F. Nash, Arc structure of singularities, *Duke Math. J.* 81 (1995), no. 1, 31-38 (1996).
- [O] T. Oda, Convex bodies and algebraic geometry. An introduction to the theory of toric varieties. *Ergebnisse der Mathematik und ihrer Grenzgebiete (3) [Results in Mathematics and Related Areas (3)]*, 15. Springer-Verlag, Berlin, 1988.
- [O1] M. Oka, On the resolution of the hypersurface singularities. *Complex analytic singularities*, 405-436, *Adv. Stud. Pure Math.*, 8, North-Holland, Amsterdam, 1987.
- [O2] M. Oka, Non-degenerate complete intersection singularity. *Actualités Mathématiques*. Hermann, Paris, 1997.
- [Re] A. Reguera, Towards the singular locus of the space of arcs. *Amer. J. Math.* 131 (2009), no. 2, 313-350.
- [T1] B. Teissier, Valuations, deformations, and toric geometry. *Valuation theory and its applications*, Vol. II (Saskatoon, SK, 1999), 361-459, *Fields Inst. Commun.*, 33, Amer. Math. Soc., Providence, RI, 2003.
- [T2] B. Teissier, Overweight deformations of affine toric varieties and local uniformization, to appear in *Proceedings of the second international conference on Valuation theory*.
- [Va] A.N. Varchenko, Zeta-function of monodromy and Newton’s diagram. *Invent. Math.* 37 (1976), no. 3, 253-262.

Hussein Mourtada,
Equipe Géométrie et Dynamique,
Institut Mathématique de Jussieu-Paris Rive Gauche,
Université Paris 7,
Bâtiment Sophie Germain, case 7012,
75205 Paris Cedex 13, France.
Email: hussein.mourtada@imj-prg.fr

Camille Plénat
I2M, Groupe AGT, Aix Marseille Université
CMI, Technopôle Château-Gombert
39, rue F. Joliot Curie, 13453 Marseille Cedex 13
Email: camille.plenat at univ-amu.fr