



HAL
open science

Linear rigidity of stationary stochastic processes

Alexander I. Bufetov, Yoann Dabrowski, Yanqi Qiu

► **To cite this version:**

Alexander I. Bufetov, Yoann Dabrowski, Yanqi Qiu. Linear rigidity of stationary stochastic processes. 2016. hal-01256215

HAL Id: hal-01256215

<https://hal.science/hal-01256215>

Preprint submitted on 14 Jan 2016

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Linear rigidity of stationary stochastic processes

Alexander I. Bufetov ^{*} Yoann Dabrowski [†] Yanqi Qiu [‡]

Abstract

We consider stationary stochastic processes X_n , $n \in \mathbb{Z}$ such that X_0 lies in the closed linear span of X_n , $n \neq 0$; following Ghosh and Peres, we call such processes linearly rigid. Using a criterion of Kolmogorov, we show that it suffices, for a stationary stochastic process to be rigid, that the spectral density vanish at zero and belong to the Zygmund class $\Lambda_*(1)$. We next give sufficient condition for stationary determinantal point processes on \mathbb{Z} and on \mathbb{R} to be rigid. Finally, we show that the determinantal point process on \mathbb{R}^2 induced by a tensor square of Dyson sine-kernels is *not* linearly rigid.

1 Introduction

This paper is devoted to rigidity of stationary determinantal point processes.

Recall that stationary determinantal point processes are strongly chaotic: they have the Kolmogorov property (Lyons [9]) and the Bernoulli property (Lyons and Steif [10]); and they satisfy the Central Limit Theorem (Costin and Lebowitz [2], Soshnikov [13]). On the other hand, Ghosh [5] and Ghosh-Peres [6] proved, for the determinantal point processes such as Dyson sine process and Ginibre point process, that number of particles in a finite window is measurable with respect to the

^{*}bufetov@mi.ras.ru; Aix-Marseille Université, Centrale Marseille, CNRS, I2M, UMR7373, 39 Rue F. Joliot Curie 13453, Marseille Cedex; Steklov Institute of Mathematics, Moscow; Institute for Information Transmission Problems, Moscow; National Research University Higher School of Economics, Moscow.

[†]dabrowski@math.univ-lyon1.fr, Université de Lyon, Université Lyon 1, Institut Camille Jordan, 43 blvd. du 11 novembre 1918, F-69622 Villeurbanne cedex, France.

[‡]yqi.qiu@gmail.com, Aix-Marseille Université, Centrale Marseille, CNRS, I2M, UMR7373, 39 Rue F. Joliot Curie 13453, Marseille cedex, France.

completion of the sigma-algebra describing the configurations outside that finite window. Their argument is spectral: they construct, for any small ε , a compactly supported smooth function φ_ε , such that φ_ε equals 1 in a fixed finite window and the linear statistic corresponding to φ_ε has variance smaller than ε .

In the same spirit, we consider general stationary stochastic processes (in broad sense) $X_n, n \in \mathbb{Z}$ such that X_0 lies in the closed linear span of $X_n, n \neq 0$; following Ghosh and Peres, we call such processes linearly rigid. Using a criterion of Kolmogorov, we show that it suffices, for a stationary stochastic process to be rigid, that the spectral density vanish at zero and belong to the Zygmund class $\Lambda_*(1)$. We next give sufficient condition for stationary determinantal point processes on \mathbb{Z} and on \mathbb{R} to be rigid. Finally, we show that the determinantal point process on \mathbb{R}^2 induced by a tensor square of Dyson sine-kernels is *not* linearly rigid.

We now turn to more precise statements. Let $X = \{X_n : n \in \mathbb{Z}^d\}$ be a multi-dimensional time stationary stochastic process of real-valued random variables defined on a probability space (Ω, \mathbb{P}) . Let $H(X) \subset L^2(\Omega, \mathbb{P})$ denote the closed subspace linearly spanned by $\{X_n : n \in \mathbb{Z}^d\}$ and let $\check{H}_0(X)$ denote the one linearly spanned by $\{X_n : n \in \mathbb{Z}^d \setminus \{0\}\}$.

Definition 1.1. The stochastic process X is said to be linearly rigid if

$$X_0 \in \check{H}_0(X). \quad (1)$$

Let $\text{Conf}(\mathbb{R}^d)$ be the set of configurations on \mathbb{R}^d . For a bounded Borel subset $B \subset \mathbb{R}^d$, we denote $N_B : \text{Conf}(\mathbb{R}^d) \rightarrow \mathbb{N} \cup \{0\}$ the function defined by

$$N_B(\mathcal{X}) := \text{the cardinality of } B \cap \mathcal{X}.$$

The space $\text{Conf}(\mathbb{R}^d)$ is equipped with the Borel σ -algebra which is the smallest σ -algebra making all N_B 's measurable. Recall that a point process with phase space \mathbb{R}^d is, by definition, a Borel probability measure on the space $\text{Conf}(\mathbb{R}^d)$. For the background on point process, the reader is referred to Daley and Vere-Jones' book [3].

Given a stationary point process on \mathbb{R}^d and $\lambda > 0$, we introduce the stationary stochastic process $N^{(\lambda)} = (N_n^{(\lambda)})_{n \in \mathbb{Z}^d}$ by the formula

$$N_n^{(\lambda)}(\mathcal{X}) := \text{the cardinality of } \mathcal{X} \cap (n\lambda + [-\lambda/2, \lambda/2]^d). \quad (2)$$

Definition 1.2. A stationary point process \mathbb{P} on \mathbb{R}^d is called **linearly rigid**, if for any $\lambda > 0$, the stationary stochastic process $N^{(\lambda)} = (N_n^{(\lambda)})_{n \in \mathbb{Z}^d}$ is linearly rigid, i.e.,

$$N_0^{(\lambda)} \in \check{H}_0(N^{(\lambda)}).$$

The above definition is motivated by the definition due to Ghosh and Peres of rigidity of point processes on \mathbb{R}^d , see [5] and [6]. Given a Borel subset $C \subset \mathbb{R}^d$, we will denote

$$\mathcal{F}_C = \sigma(\{N_B : B \subset C, B \text{ bounded Borel}\})$$

the σ -algebra generated by all random variables of the form N_B where $B \subset C$ ranges over all bounded Borel subsets of C . Let \mathbb{P} be a point process on \mathbb{R}^d , i.e., \mathbb{P} is a Borel probability on $\text{Conf}(\mathbb{R}^d)$, and denote $\mathcal{F}_C^{\mathbb{P}}$ for the \mathbb{P} -completion of \mathcal{F}_C .

Definition 1.3 (Ghosh [5], Ghosh-Peres [6]). A point process \mathbb{P} on \mathbb{R}^d is called **rigid**, if for any bounded Borel set $B \subset \mathbb{R}^d$ with Lebesgue-negligible boundary ∂B , the random variable N_B is $\mathcal{F}_{\mathbb{R}^d \setminus B}^{\mathbb{P}}$ -measurable.

Remark 1.1. Of course, in the above definition, it suffices to take Borel sets B of the form $[-\gamma, \gamma]^d$ for $\gamma > 0$, cf. [6].

A linear rigid stationary point process on \mathbb{R}^d is of course rigid in the sense of Ghosh and Peres. Observe that proofs for rigidity in [5], [6] and [1] in fact establish linear rigidity. We would like also to mention a notion of insertion-deletion tolerance studied by Holroyd and Soo in [7], which is in contrast to the notion of rigidity property.

2 The Kolmogorov criterion for linear rigidity

In this note, the Fourier transform of a function $f : \mathbb{R}^d \rightarrow \mathbb{C}$ is defined as

$$\hat{f}(\xi) = \int_{\mathbb{R}^d} f(x) e^{-i2\pi x \cdot \xi} dx.$$

Denote by $\mathbb{T}^d = \mathbb{R}^d / \mathbb{Z}^d$ the d -dimensional torus. In what follows, we identify \mathbb{T}^d with $[-1/2, 1/2]^d$. The Fourier coefficients of a measure μ on \mathbb{T}^d are given, for any $k \in \mathbb{Z}^d$, by the formula

$$\hat{\mu}(k) = \int_{\mathbb{T}^d} e^{-i2\pi k \cdot \theta} d\mu_X(\theta), \text{ where } k \cdot \theta := k_1\theta_1 + \dots + k_d\theta_d.$$

Denote by μ_X the spectral measure of X , i.e.,

$$\forall k \in \mathbb{Z}^d, \quad \mathbb{E}(X_0 X_k) = \mathbb{E}(X_n X_{n+k}) = \int_{\mathbb{T}^d} e^{-i2\pi k \cdot \theta} d\mu_X(\theta) = \hat{\mu}_X(k). \quad (3)$$

Recall that we have the following natural isometric isomorphism

$$H(X) \simeq L^2(\mathbb{T}^d, \mu_X), \quad (4)$$

by assigning to $X_n \in H(X)$ the function $\theta \mapsto e^{i2\pi n \cdot \theta} \in L^2(\mathbb{T}^d, \mu_X)$.

Let $\mu_X = \mu_a + \mu_s$ be the Lebesgue decomposition of μ_X with respect to the normalized Lebesgue measure $m(d\theta) = d\theta_1 \cdots d\theta_d$ on \mathbb{T}^d , i.e., μ_a is absolutely continuous with respect to m and μ_s is singular to m . Set

$$\omega_X(\theta) := \frac{d\mu_a}{dm}(\theta).$$

Lemma 2.1 (The Kolmogorov Criterion). *We have*

$$\text{dist}(X_0, \check{H}_0(X)) = \left(\int_{\mathbb{T}^d} \omega_X^{-1} dm \right)^{-1/2}.$$

The right side is to be interpreted as zero if $\int_{\mathbb{T}^d} \omega_X^{-1} dm = \infty$.

When the measure μ is assumed to be absolutely continuous with respect to m , Lemma 2.1 is a result of Kolmogorov, see Remark 5.17 in Lyons-Steif [10].

Corollary 2.2. *The stationary stochastic process $X = (X_n)_{n \in \mathbb{Z}^d}$ is linearly rigid if and only if*

$$\int_{\mathbb{T}^d} \omega_X^{-1} dm = \infty.$$

Proof of Lemma 2.1. We follow the argument of Lyons-Steif [10]. By the Lebesgue decomposition of μ , we may take a subset $A \subset \mathbb{T}^d$ of full Lebesgue measure $m(A) = 1$, such that $\mu_a(A) = 1$ and $\mu_s(A) = 0$.

Denote

$$L_0 = \overline{\text{span}}^{L^2(\mathbb{T}^d, \mu_X)}[e^{i2\pi n \cdot \theta} : n \neq 0].$$

By the isometric isomorphism (4), it suffices to show that

$$\text{dist}(1, L_0) = \left(\int_{\mathbb{T}^d} \omega_X^{-1} dm \right)^{-1/2}, \quad (5)$$

where 1 is the constant function taking value 1. Write

$$1 = p + h, \text{ such that } p \perp L_0, h \in L_0.$$

Modifying, if necessary, the values of p and h on a μ -negligible subset, we may assume that

$$1 = p(\theta) + h(\theta) \text{ for all } \theta \in \mathbb{T}^d.$$

Since $p \perp L_0$, we have

$$0 = \langle p, e^{i2\pi n \cdot \theta} \rangle_{L^2(d\mu)} = \int_{\mathbb{T}^d} p(\theta) e^{-i2\pi n \cdot \theta} d\mu(\theta), \text{ for any } n \in \mathbb{Z}^d \setminus 0. \quad (6)$$

Therefore, the complex measure $p \cdot d\mu$ is a multiple of Lebesgue measure, i.e., there exists $\xi \in \mathbb{C}$, such that

$$p \cdot d\mu = \xi dm.$$

It follows that p must vanish almost everywhere with respect to the singular component μ_s of μ , and $p(\theta)\omega_X(\theta) = \xi$ for m -almost every $\theta \in \mathbb{T}^d$. Thus we have

$$\|p\|_{L^2(d\mu)} = \|p\|_{L^2(d\mu_a)}, \quad (7)$$

and

$$h(\theta) = 1 - \xi\omega_X(\theta)^{-1} \text{ for } m\text{-almost every } \theta \in \mathbb{T}^d. \quad (8)$$

Case 1: $\int_{\mathbb{T}^d} \omega_X^{-1} dm < \infty$.

Define a function $f : \mathbb{T}^d \rightarrow \mathbb{C}$ by $f = \omega_X^{-1} \chi_A$. Then $f \in L^2(d\mu) \ominus L_0$. Indeed,

$$\|f\|_{L^2(d\mu)}^2 = \int_{\mathbb{T}^d} \omega_X^{-2} \chi_A d\mu = \int_{\mathbb{T}^d} \omega_X^{-2} d\mu_a = \int_{\mathbb{T}^d} \omega_X^{-1} dm < \infty.$$

And, for all $n \in \mathbb{Z}^d \setminus 0$,

$$\langle f, e^{i2\pi n \cdot \theta} \rangle_{L^2(d\mu)} = \int_{\mathbb{T}^d} \omega_X(\theta)^{-1} \chi_A(\theta) e^{-i2\pi n \cdot \theta} d\mu(\theta) = \int_{\mathbb{T}^d} e^{-i2\pi n \cdot \theta} dm(\theta) = 0.$$

It follows that $f \perp h$, i.e.,

$$0 = \langle h, f \rangle_{L^2(d\mu)} = \int_{\mathbb{T}^d} h \omega_X^{-1} \chi_A d\mu = \int_{\mathbb{T}^d} h dm.$$

By (8), we get

$$\int_{\mathbb{T}^d} (1 - \xi \omega_X^{-1}) dm = 0,$$

and hence

$$\xi = \left(\int_{\mathbb{T}^d} \omega_X^{-1} dm \right)^{-1}.$$

It follows that

$$\text{dist}(1, L_0)^2 = \|p\|_{L^2(d\mu)}^2 = \|p\|_{L^2(d\mu_a)}^2 = \xi^2 \int_{\mathbb{T}^d} \omega_X^{-2} \omega_X dm = \xi.$$

This shows the desired equality (5).

Case 2: $\int_{\mathbb{T}^d} \omega_X^{-1} dm = \infty$.

We claim that $\xi = 0$. If the claim were verified, then we would get the desired identity in this case

$$\text{dist}(1, L_0) = 0.$$

So let us turn to the proof of the claim. We argue by contradiction. If $\xi \neq 0$, then $p \neq 0$ and

$$\|p\|_{L^2(d\mu)}^2 = \|p\|_{L^2(d\mu_a)}^2 = \xi^2 \|\omega_X^{-1}\|_{L^2(d\mu_a)}^2 = \xi^2 \int_{\mathbb{T}^d} \omega_X^{-1} dm = \infty.$$

This contradicts the fact that $p \in L^2(d\mu)$. □

Remark 2.1. The same argument shows that, in the case of one-dimensional time, the following assertions are equivalent:

- $\sum_{k=-n}^n X_k \in \overline{\text{span}}\{X_j : |j| \geq n + 1\}$;
- for any $\alpha_1, \dots, \alpha_n \in (-1/2, 1/2) \setminus \{0\}$, we have

$$\int_{\mathbb{T}} \frac{\prod_{j=1}^n |e^{i2\pi\theta} - e^{i2\pi\alpha_j}|^2 |e^{i2\pi\theta} - e^{-i2\pi\alpha_j}|^2}{\omega_X(\theta)} dm(\theta) = \infty.$$

It would be interesting to find a similar characterization for multi-dimensional time as well.

Denote by $\text{Cov}(U, V)$ the covariance between two random variables U and V :
 $\text{Cov}(U, V) = \mathbb{E}(UV) - \mathbb{E}(U)\mathbb{E}(V)$.

If $X = (X_n)_{n \in \mathbb{Z}^d}$ is a stochastic process such that

$$\sum_{n \in \mathbb{Z}^d} |\text{Cov}(X_0, X_n)| < \infty, \quad (9)$$

then we may define a continuous function on \mathbb{T}^d by the formula

$$\omega_X(\theta) := \sum_{n \in \mathbb{Z}^d} \text{Cov}(X_0, X_n) e^{i2\pi n \cdot \theta}. \quad (10)$$

Lemma 2.3. *Let $X = (X_n)_{n \in \mathbb{Z}^d}$ be a stationary stochastic process satisfying condition (9). Then we have the following explicit Lebesgue decomposition of μ_X :*

$$\mu_X = (\mathbb{E}X_0)^2 \cdot \delta_0 + \omega_X \cdot m, \quad (11)$$

where δ_0 is the Dirac measure on the point $0 \in \mathbb{T}^d$ and ω_X is the function on \mathbb{T}^d defined by (10).

Proof. Note that, under the assumption (9), the function $\omega_X(\theta)$ is well-defined and continuous on \mathbb{T}^d . For proving the decomposition (11), it suffices to show that the Fourier coefficients of μ_X coincide with those of $\nu_X := (\mathbb{E}X_0)^2 \cdot \delta_0 + \omega_X \cdot m$. But if $n \in \mathbb{Z}^d$, then

$$\hat{\nu}_X(n) = (\mathbb{E}X_0)^2 + \text{Cov}(X_0, X_n) = \mathbb{E}(X_0 X_n) = \hat{\mu}_X(n).$$

The lemma is completely proved. \square

3 A sufficient condition for linear rigidity

Theorem 3.1. *Let $X = (X_n)_{n \in \mathbb{Z}}$ be a stationary stochastic process. If*

$$\sup_{N \geq 1} \left(N \sum_{|n| \geq N} |\text{Cov}(X_0, X_n)| \right) < \infty, \quad (12)$$

and

$$\sum_{n \in \mathbb{Z}} \text{Cov}(X_0, X_n) = 0. \quad (13)$$

Then X is linearly rigid.

Remark 3.1. The condition (12) is a sufficient condition such that the spectral density ω_X is a function in the Zygmund class $\Lambda_*(1)$, see below for definition. The condition (13) implies in particular that ω_X vanishes at the point $0 \in \mathbb{T}$.

We shall apply a result of F. Móricz [12, Thm. 3] on absolutely convergent Fourier series and Zygmund class functions. Recall that a continuous 1-periodic function φ defined on \mathbb{R} is said to be in the *Zygmund class* $\Lambda_*(1)$, if there exists a constant C such that

$$|\varphi(x+h) - 2\varphi(x) + \varphi(x-h)| \leq Ch \quad (14)$$

for all $x \in \mathbb{R}$ and for all $h > 0$.

Theorem 3.2 (Móricz, [12]). *If $\{c_n\}_{n \in \mathbb{Z}} \in \mathbb{C}$ is such that*

$$\sup_{N \geq 1} \left(N \sum_{|n| \geq N} |c_n| \right) < \infty, \quad (15)$$

then the function $\varphi(\theta) = \sum_{n \in \mathbb{Z}} c_n e^{i2\pi n\theta}$ is in the Zygmund class $\Lambda_(1)$.*

Proof of Theorem 3.1. First, in view of (10), our assumption (13) implies

$$\omega_X(0) = 0.$$

Next, by Theorem 3.2, under the assumption (12), we have

$$\omega_X \in \Lambda_*(1).$$

Since all Fourier coefficients of ω_X are real, we have

$$\omega_X(\theta) = \omega_X(-\theta).$$

Consequently, there exists $C > 0$, such that

$$\omega_X(\theta) = \frac{\omega_X(\theta) + \omega_X(-\theta)}{2} = \frac{\omega_X(\theta) + \omega_X(-\theta) - 2\omega_X(0)}{2} \leq C|\theta|,$$

whence

$$\int_{\mathbb{T}} \omega_X^{-1} dm = \infty,$$

and the stochastic process $X = (X_n)_{n \in \mathbb{Z}}$ is linearly rigid by the Kolmogorov criterion. \square

4 Applications to stationary determinantal point processes

In this section, we first give a sufficient condition for linear rigidity of stationary determinantal point processes on \mathbb{R} and then give an example of a very simple stationary, but not linearly rigid, determinantal point process on \mathbb{R}^2 . We briefly recall the main definitions. Let $B \subset \mathbb{R}^d$ be a bounded Borel subset. Let $K_B : L^2(\mathbb{R}^d) \rightarrow L^2(\mathbb{R}^d)$ be the operator of convolution with the Fourier transform $\widehat{\chi_B}$ of the indicator function χ_B . In other words, the kernel of K_B is

$$K_B(x, y) = \widehat{\chi_B}(x - y). \quad (16)$$

In particular, if $d = 1$ and $B = (-1/2, 1/2)$, then we find the well-known Dyson sine kernel

$$K_{\text{sine}}(x, y) = \frac{\sin(\pi(x - y))}{\pi(x - y)}.$$

Note that we always have $K_B(x, x) = K_B(0, 0)$.

Denote by \mathbb{P}_{K_B} the determinantal point process induced by K_B . For the background on the determinantal point processes, the reader is referred to [8], [9], [11], [13].

Proposition 4.1. *Let \mathbb{P}_{K_B} be the stationary determinantal point process on \mathbb{R}^d induced by the kernel K_B in (16). For any $\lambda > 0$, denote by $N^{(\lambda)} = (N_n^{(\lambda)})_{n \in \mathbb{Z}^d}$ the stationary stochastic process associated to \mathbb{P}_{K_B} as in (2). Then*

$$\sum_{n \in \mathbb{Z}^d} |\text{Cov}(N_0^{(\lambda)}, N_n^{(\lambda)})| < \infty \quad (17)$$

and

$$\sum_{n \in \mathbb{Z}^d} \text{Cov}(N_0^{(\lambda)}, N_n^{(\lambda)}) = 0. \quad (18)$$

Proof. Fix a number $\lambda > 0$, for simplifying the notation, let us denote $N_n^{(\lambda)}$ by N_n . Denote for any $n \in \mathbb{Z}^d$,

$$Q_n = n\lambda + [-\lambda/2, \lambda/2]^d.$$

By definition of a determinantal point process, we have

$$\mathbb{E}(N_n) = \mathbb{E}(N_0) = \int_{Q_0} K_B(x, x) dx = \lambda^d K_B(0, 0).$$

If $n \neq 0$, we have

$$\begin{aligned}\mathbb{E}(N_0 N_n) &= \iint \chi_{Q_0}(x) \chi_{Q_n}(y) \begin{vmatrix} K_B(x, x) & K_B(x, y) \\ K_B(y, x) & K_B(y, y) \end{vmatrix} dx dy \\ &= \lambda^{2d} K_B(0, 0)^2 - \iint_{Q_0 \times Q_n} |K_B(x, y)|^2 dx dy,\end{aligned}$$

whence

$$\text{Cov}(N_0, N_n) = - \iint_{Q_0 \times Q_n} |K_B(x, y)|^2 dx dy. \quad (19)$$

We also have

$$\begin{aligned}\mathbb{E}(N_0^2) &= \mathbb{E} \left[\sum_{x, y \in \mathcal{X}} \chi_{Q_0}(x) \chi_{Q_0}(y) \right] \\ &= \mathbb{E} \left[\sum_{x \in \mathcal{X}} \chi_{Q_0}(x) \right] + \mathbb{E} \left[\sum_{x, y \in \mathcal{X}, x \neq y} \chi_{Q_0}(x) \chi_{Q_0}(y) \right] \\ &= \int_{Q_0} K_B(x, x) dx + \iint \chi_{Q_0}(x) \chi_{Q_0}(y) \begin{vmatrix} K_B(x, x) & K_B(x, y) \\ K_B(y, x) & K_B(y, y) \end{vmatrix} dx dy \\ &= \lambda^d K_B(0, 0) + \lambda^{2d} K_B(0, 0)^2 - \iint_{Q_0 \times Q_0} |K_B(x, y)|^2 dx dy,\end{aligned}$$

whence

$$\text{Cov}(N_0, N_0) = \text{Var}(N_0) = \lambda^d K_B(0, 0) - \iint_{Q_0 \times Q_0} |K_B(x, y)|^2 dx dy. \quad (20)$$

Now recall that K_B is an orthogonal projection. Thus we have

$$K_B(0, 0) = K_B(x, x) = \int |K_B(x, y)|^2 dy = \sum_{n \in \mathbb{Z}^d} \int_{Q_n} |K_B(x, y)|^2 dy. \quad (21)$$

The identities (19), (20) and (21) imply that

$$\begin{aligned}\sum_{n \in \mathbb{Z}^d} \text{Cov}(N_0, N_n) &= \lambda^d K_B(0, 0) - \int_{Q_0} dx \sum_{n \in \mathbb{Z}^d} \int_{Q_n} |K_B(x, y)|^2 dy \\ &= \lambda^d K_B(0, 0) - \lambda^d K_B(0, 0) = 0.\end{aligned}$$

Moreover, the above series converge absolutely. Proposition 4.1 is completely proved. \square

Remark 4.1. By Lemma 2.3 and Proposition 4.1, we see that for any stationary determinantal point process induced by a projection, the spectral density of the associated stochastic process $N^{(\lambda)}$ always vanishes at 0.

4.1 Stationary determinantal point processes on \mathbb{R}

Theorem 4.2. *Assume that $B \subset \mathbb{R}$ satisfies*

$$\sup_{R>0} \left(R \int_{|\xi| \geq R} |\widehat{\chi}_B(\xi)|^2 d\xi \right) < \infty. \quad (22)$$

Then the stationary determinantal point process \mathbb{P}_{K_B} is linearly rigid.

Proof. By definition of linear rigidity, we need to show that for any $\lambda > 0$, the stochastic process $N^{(\lambda)} = (N_n^{(\lambda)})_{n \in \mathbb{Z}}$ is linearly rigid. As in the proof of Proposition 4.1, we denote $N_n^{(\lambda)}$ by N_n . By Theorem 3.1, it suffices to show that

$$\sup_{N \geq 1} \left(N \sum_{|n| \geq N} |\text{Cov}(N_0, N_n)| \right) < \infty, \quad (23)$$

and

$$\sum_{n \in \mathbb{Z}} \text{Cov}(N_0, N_n) = 0. \quad (24)$$

By Proposition 4.1, the identity (24) holds in general case. It remains to prove (23). By (19), we have

$$\begin{aligned} \sup_{N \geq 1} \left(N \sum_{|n| \geq N} |\text{Cov}(N_0, N_n)| \right) &= \sup_{N \geq 1} N \iint_{\bigcup_{|n| \geq N} Q_n} |\widehat{\chi}_B(x-y)|^2 dx dy \\ &\leq \sup_{N \geq 1} \lambda N \int_{|\xi| \geq (N-1)\lambda} |\widehat{\chi}_B(\xi)|^2 d\xi < \infty \end{aligned}$$

where in the last inequality, we used our assumption (22). Theorem 4.2 is proved completely. \square

Remark 4.2. When B is a finite union of finite intervals on the real line, the rigidity of the stationary determinantal point process \mathbb{P}_{K_B} is due to Ghosh [5].

4.2 Tensor product of sine kernels

In higher dimension, the situation becomes quite different. Let

$$S = I \times I = (-1/2, 1/2) \times (-1/2, 1/2) \subset \mathbb{R}^2.$$

Then the associate kernel K_S has a tensor form: $K_S = K_{\text{sine}} \otimes K_{\text{sine}}$, that is, for $x = (x_1, x_2)$ and $y = (y_1, y_2)$ in \mathbb{R}^2 , we have

$$K_S(x, y) = K_{\text{sine}}(x_1, y_1)K_{\text{sine}}(x_2, y_2) = \frac{\sin(\pi(x_1 - y_1))}{\pi(x_1 - y_1)} \frac{\sin(\pi(x_2 - y_2))}{\pi(x_2 - y_2)}.$$

Proposition 4.3. *The determinantal point process \mathbb{P}_{K_S} is not linearly rigid. More precisely, let $N^{(1)} = (N_n^{(1)})_{n \in \mathbb{Z}^2}$ be the stationary stochastic process given as in Definition 1.2, then*

$$N_0^{(1)} \notin \check{H}_0(N^{(1)}).$$

To prove the above result, we need to introduce some extra notation. First, we define the multiple Zygmund class Λ_* as follows. A continuous function $\varphi(x, y)$ periodic in each variable with period 1 is said to be in the multiple Zygmund class $\Lambda_*(1, 1)$ if for the double difference difference operator $\Delta_{2,2}$ of second order in each variable, applied to φ , there exists a constant $C > 0$, such that for all $x = (x_1, x_2) \in (-1/2, 1/2) \times (-1/2, 1/2)$ and $h_1, h_2 > 0$, we have

$$|\Delta_{2,2}\varphi(x_1, x_2; h_1, h_2)| \leq Ch_1h_2, \quad (25)$$

where

$$\begin{aligned} \Delta_{2,2}\varphi(x_1, x_2; h_1, h_2) &:= \varphi(x_1 + h_1, x_2 + h_2) + \varphi(x_1 - h_1, x_2 + h_2) \\ &\quad + \varphi(x_1 + h_1, x_2 - h_2) + \varphi(x_1 - h_1, x_2 - h_2) - 2\varphi(x_1 + h_1, x_2) \\ &\quad - 2\varphi(x_1 - h_1, x_2) - 2\varphi(x_1, x_2 + h_2) - 2\varphi(x_1, x_2 - h_2) + 4\varphi(x_1, x_2). \end{aligned}$$

The following result is due to Fülöp and Móricz [4, Thm 2.1 and Rem. 2.3]

Theorem 4.4 (Fülöp-Móricz). *If $\{c_{jk}\}_{j,k \in \mathbb{Z}} \in \mathbb{C}$ is such that*

$$\sup_{N \geq 1, M \geq 1} \left(MN \sum_{|j| \geq N, |k| \geq M} |c_{jk}| \right) < \infty, \quad (26)$$

then the function

$$\varphi(\theta_1, \theta_2) = \sum_{j,k \in \mathbb{Z}} c_{jk} e^{i2\pi(j\theta_1 + k\theta_2)}$$

is in the Zygmund class $\Lambda_*(1, 1)$.

Let us turn to the study of the density function $\omega_{N^{(1)}}$.

Lemma 4.5. *There exists $c > 0$, such that for any $\theta_1, \theta_2 \in (-1/2, 1/2)$, we have*

$$\omega_{N^{(1)}}(\theta_1, \theta_2) \geq c(|\theta_1| + |\theta_2|).$$

Proof. To make notation lighter, in this proof we simply write ω for $\omega_{N^{(1)}}$.

Denote $S_n = S \times (n + S)$ where $n + S := (-1/2 + n_1, 1/2 + n_1) \times (-1/2 + n_2, 1/2 + n_2)$. By the same argument as in the proof of Theorem 4.2, we obtain that for any $n = (n_1, n_2) \in \mathbb{Z}^2 \setminus 0$,

$$\widehat{\omega}(n) = - \int_{S_n} |K_S(x, y)|^2 dx dy,$$

and

$$\widehat{\omega}(0) = K_S(0, 0) - \int_{S_0} |K_S(x, y)|^2 dx dy.$$

The following properties can be easily checked.

- $\sum_{n \in \mathbb{Z}^2} \widehat{\omega}(n) = 0$.
- $\widehat{\omega}(\varepsilon_1 n_1, \varepsilon_2 n_2) = \widehat{\omega}(n_1, n_2)$, where $\varepsilon_1, \varepsilon_2 \in \{\pm 1\}$.
- there exist $c, C > 0$, such that

$$\frac{c}{(1 + n_1^2)(1 + n_2^2)} \leq |\widehat{\omega}(n_1, n_2)| \leq \frac{C}{(1 + n_1^2)(1 + n_2^2)}.$$

For instance, $\sum_{n \in \mathbb{Z}^2} \widehat{\omega}(n) = 0$ follows from Proposition 4.1. These properties combined with Theorem 4.4 yield that

- $\omega(0, 0) = 0$.
- $\omega(\varepsilon_1 \theta_1, \varepsilon_2 \theta_2) = \omega(\theta_1, \theta_2)$ for any $\varepsilon_1, \varepsilon_2 \in \{\pm 1\}$ and $\theta_1, \theta_2 \in (-1/2, 1/2)$.
- the function $\omega(\theta_1, \theta_2)$ is in the multiple Zygmund class $\Lambda_*(1, 1)$.

Hence there exists $C > 0$, such that

$$|\omega(\theta_1, \theta_2) - \omega(\theta_1, 0) - \omega(0, \theta_2)| \leq C|\theta_1 \theta_2|. \quad (27)$$

Lemma 4.6. *There exists $c > 0$, such that*

$$\omega(\theta_1, 0) \geq c|\theta_1| \text{ and } \omega(0, \theta_2) \geq c|\theta_2|. \quad (28)$$

Let us postpone the proof of Lemma 4.6 and proceed to the proof of Lemma 4.5. The inequalities (27) and (28) imply that

$$\omega(\theta_1, \theta_2) \geq c(|\theta_1| + |\theta_2|) - C|\theta_1\theta_2|.$$

To prove the lower bound of type as in the lemma, it suffices to prove it when $|\theta_1|$ and $|\theta_2|$ are small enough, for instance, $2C|\theta_1| \leq c$, then we have

$$\omega(\theta_1, \theta_2) \geq \frac{c}{2}(|\theta_1| + |\theta_2|).$$

□

Now let us turn to the

Proof of Lemma 4.6. By symmetry, it suffices to prove that there exists $c > 0$, such that $\omega(\theta_1, 0) \geq |\theta_1|$. To this end, let us denote $\omega_1(\theta_1) := \omega(\theta_1, 0)$. Then $\omega_1(0) = 0$ and there exists $c > 0$ such that if $k \neq 0$, then

$$\widehat{\omega}_1(k) < 0 \text{ and } |\widehat{\omega}_1(k)| \geq c/(1 + k^2),$$

Indeed, we have

$$\omega_1(\theta_1) = \sum_{k \in \mathbb{Z}} \sum_{n_2 \in \mathbb{Z}} \widehat{\omega}(k, n_2) e^{i2\pi k \theta_1},$$

if $k \neq 0$, then $\widehat{\omega}(k, n_2) < 0$ and hence

$$|\widehat{\omega}_1(k)| = \sum_{n_2 \in \mathbb{Z}} |\widehat{\omega}(k, n_2)| \geq \sum_{n_2 \in \mathbb{Z}} \frac{c}{(1 + n_2^2)(1 + k^2)} \geq \frac{c'}{1 + k^2}.$$

Note also that $\omega_1(0) = \omega(0, 0) = 0$, hence

$$\sum_{k \in \mathbb{Z}} \widehat{\omega}_1(k) = 0.$$

It follows that

$$\begin{aligned}
\omega_1(\theta_1) &= \sum_{k \in \mathbb{Z}} \widehat{\omega}_1(k) e^{i2\pi k \theta_1} = \sum_{k \in \mathbb{Z}} \widehat{\omega}_1(k) \left(\frac{e^{i2\pi k \theta_1} + e^{-i2\pi k \theta_1}}{2} - 1 \right) \\
&= \sum_{k \in \mathbb{Z}, k \neq 0} -\widehat{\omega}_1(k) (1 - \cos(2\pi k \theta_1)) = \sum_{k \in \mathbb{Z}, k \neq 0} |\widehat{\omega}_1(k)| (1 - \cos(2\pi k \theta_1)) \\
&\geq c'' \sum_{j=1}^{\infty} \frac{1}{(2j-1)^2} (1 - \cos(2\pi(2j-1)\theta_1)).
\end{aligned}$$

Combining with the classical formulae

$$\begin{aligned}
\sum_{j=1}^{\infty} \frac{1}{(2j-1)^2} &= \frac{\pi^2}{8}, \\
|\alpha| &= \frac{1}{4} - \frac{2}{\pi^2} \sum_{j=1}^{\infty} \frac{\cos(2(2j-1)\pi\alpha)}{(2j-1)^2}, \text{ for } \alpha \in (-1/2, 1/2);
\end{aligned}$$

we obtain that

$$\omega_1(\theta_1) \geq c'' \frac{\pi^2}{2} |\theta_1|.$$

□

Proof of Proposition 4.3. By Lemma 2.1, it suffices to show that

$$\int_{\mathbb{T}^2} \omega_{N(1)}^{-1} dm < \infty. \quad (29)$$

By Lemma 4.5, the inequality (29) follows from the following elementary inequality

$$\int_{|\theta_1| < 1/2, |\theta_2| < 1/2} \frac{1}{|\theta_1| + |\theta_2|} d\theta_1 d\theta_2 < \infty.$$

□

Acknowledgements

The authors would like to thank Guillaume Aubrun for many valuable discussions. A.B. and Y.Q. are supported by A*MIDEX project (No. ANR-11-IDEX-0001-02), financed by Programme “Investissements d’Avenir” of the Government of the French Republic managed by the French National Research Agency (ANR). A. B. is also supported in part by the Grant MD-2859.2014.1 of the President of the Russian Federation, by the Programme “Dynamical systems and mathematical control theory” of the Presidium of the Russian Academy of Sciences, by the ANR under the project “VALET” of the Programme JCJC SIMI 1, and by the RFBR grants 11-01-00654, 12-01-31284, 12-01-33020, 13-01-12449.

References

- [1] Alexander I. Bufetov. Rigidity of Determinantal Point Processes with the Airy, the Bessel and the Gamma Kernel. *arXiv:1506.07581*, Jun 2015.
- [2] Ovidiu Costin and Joel L. Lebowitz. Gaussian fluctuation in random matrices. *Phys. Rev. Lett.*, 75(1):69–72, 1995.
- [3] Daryl J. Daley and David Vere-Jones. *An introduction to the theory of point processes. vol. I. , Elementary theory and methods*. Probability and its applications. Springer, New York, Berlin, Paris, 2003.
- [4] Vanda Fülöp and Ferenc Móricz. Absolutely convergent multiple Fourier series and multiplicative Zygmund classes of functions. *Analysis*, 28(3/2008):345–354, 2008.
- [5] Subhroshekhar Ghosh. Determinantal processes and completeness of random exponentials: the critical case. *Probability Theory and Related Fields*, pages 1–23, 2014.
- [6] Subhroshekhar Ghosh and Yuval Peres. Rigidity and tolerance in point processes: Gaussian zeros and Ginibre eigenvalues,. *arXiv:1211.3506*.
- [7] Alexander E. Holroyd and Terry Soo. Insertion and deletion tolerance of point processes. *Electron. J. Probab.*, 18:no. 74, 24, 2013.
- [8] J. Ben Hough, Manjunath Krishnapur, Yuval Peres, and Bálint Virág. Determinantal processes and independence. *Probab. Surv.*, 3:206–229, 2006.

- [9] Russell Lyons. Determinantal probability measures. *Publ. Math. Inst. Hautes Études Sci.*, (98):167–212, 2003.
- [10] Russell Lyons and Jeffrey E. Steif. Stationary determinantal processes: phase multiplicity, Bernoullicity, entropy, and domination. *Duke Math. J.*, 120(3):515–575, 2003.
- [11] Odile Macchi. The coincidence approach to stochastic point processes. *Advances in Appl. Probability*, 7:83–122, 1975.
- [12] Ferenc Móricz. Absolutely convergent Fourier series and function classes. *J. Math. Anal. Appl.*, 324(2):1168–1177, 2006.
- [13] Alexander Soshnikov. Determinantal random point fields. *Uspekhi Mat. Nauk*, 55(5(335)):107–160, 2000.