

Geomorphic impacts, age and significance of two giant landslide dams in the Nepal Himalayas: Ringmo-Phoksundo (Dolpo District) and Dhampu-Chhoya (Mustang District).

Monique Fort, Regis Braucher, Didier Bourles, Valery Guillou, Lila Nath

Rimal, Natasha Gribenski, Etienne Cossart

▶ To cite this version:

Monique Fort, Regis Braucher, Didier Bourles, Valery Guillou, Lila Nath Rimal, et al.. Geomorphic impacts, age and significance of two giant landslide dams in the Nepal Himalayas: Ringmo-Phoksundo (Dolpo District) and Dhampu-Chhoya (Mustang District).. EGU General Assembly 2014, Apr 2014, Vienne, Austria. 1 p. hal-01262118

HAL Id: hal-01262118 https://hal.science/hal-01262118

Submitted on 26 Jan 2016 $\,$

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Large catastrophic slope failures have recently

retained much attention in the northern dry

Himalayas (Hewitt 2009). They play a prominent

role in the denudation history of active orogens

at a wide range of spatial and time scales

(Korup & Clague, 2009), and they impact

durably landforms and process evolution in

upstream catchments. Their occurrence mostly

results from three different potential triggers:

earthquakes, post-glacial debuttressing, and

permafrost melting. We focus on two examples

of giant rock slope failures that occurred across

and north of the Higher Himalaya of Nepal and

assess their respective influence on the

rockslides bodies derived from different parts of

From Fort, 2011

3. RINGMO-PHOKSUNDO SITE

regional, geomorphic evolution.

1. INTRODUCTION: MAIN ISSUES

Geomorphic impacts, age and significance of two giant landslide dams in the Nepal Himalayas: Ringmo-Phoksundo (Dolpo District) and Dhampu-Chhova (Mustang District). Monique FORT ⁽¹⁾, Régis BRAUCHER ⁽²⁾, Didier BOURLES ⁽²⁾, Valéry GUILLOU ⁽²⁾, Lila Nath RIMAL ⁽³⁾, Natasha GRIBENSKI ⁽⁴⁾, Etienne COSSART⁽⁵⁾

rigger: sism

ectonic event + glacial

debuttressi

YE KANNER

3 massive landslides (volume > 1 km^3)

reported in Nepal

Right: Manang (# 4)

(Fort 2011)

(1) Dent of Geography LIMR 8586 PRODIG CC 7001 Liniv Paris-Didero SPC E - 75 251 PARIS Cedex 05 France (fort@univ-paris-diderot fr)* (2) Aix-Marseille Université, CEREGE CNRS UMR 6635, 13545 Aix en Provence, France; (3) Department of Mines and Geology, Lainchaur, Kathmandu; (4) Department of Physical Geography and Quaternary Geology, Stockholm University, SE-106 91, Stockholm, Sweden: (5) UMR 8586 PRODIG, Univ. Paris1-Sorbonne, 2 rue Valette 75005 Paris, Erance



2. DATING METHODS: COSMIC RAY EXPOSURE DATING

AMS measurements of the ³⁶Cl and ¹⁰Be concentrations were performed at the 5 MV nation AMS facility ASTER located at CEREGE, Aix en Provence, France, All ³⁶Cl concentrations were normalized to the KNSTD1600 calibration material $({}^{36}Cl/{}^{35}Cl = 1.6*10^{-12}$ provided by Nishiizumi) and all ¹⁰Be concentrations were normalized to the SRM 4325 NIST reference material with an assigned ¹⁰Be/⁹Be value of (2,79±0.03)*10⁻¹¹. The ³⁶Cl decay constant of (2,303 ± 0.016)*10-6 a-1 corresponds to a half-life (T1/2) of (3.014±0.021)*105 years and the 10Be deca constant of (4.987±0.036)*10⁻⁷ a⁻¹ to a half-life of (1.39±0.01)*10⁶ years. Analytica uncertainties include the counting statistics, machine stability (~0.5%) and blank correction the associated ³⁶Cl/³⁵Cl blank ratio being ~10⁻¹⁴ and the ¹⁰Be/⁹Be blank ratio ~2*10⁻¹⁵. Cosmi ray exposure ages derived from these cosmogenic nuclide concentrations were calculated usin production rates corrected for topographic shielding following Dunne et al. (1999) and scaled to the sites position (latitude, altitude) using the Stone scheme (Stone, 2000) from a sea leve and high latitude production rate of 4.5±0.3 atoms of ¹⁰Be g(SIO2)-1.yr-1 for ¹⁰Be and integrating all production pathways (spallation, including a spallation production rate of 42.0 \pm 2.0 atom of ³⁶Cl g⁻¹_{Ca}⁻¹ yr⁻¹ and at sea level and high latitude, thermal neutron, muon captures and radiogenic production) and the rock's chemical composition for ³⁶Cl.

the second	Two consistent ³⁶ Cl ages of 20,885 ±1675 argue for a single massive event of paradacial origin								
	PHOK COS 3	72,015	54,59%	314,8	10,42	13,82	19,5	20888	1661
	PHOK COS 2	69,08	54,70%	382,0	24,08	4,58	19,8	20882	1687
	PHOK COS 1	67,29	55,18%	133,8	35,64	2,79	6,8	6987	566
	Sample ID	dissolved [g]		(10 ⁻¹⁴)		(ppm)	[10° at/g]	(uneroded surface)	Uncertainties
		Sample weight	% CaO	"CI/"CI	"CL/"CI	Cl	"CI	Exposure Age of surface	Description

Field assumptions are confirmed by cosmic ray exposure dating: the landslide dam was formed in response to the collapse of the limestone cliff in one single, massive event. The ³⁶Cl dates confirm the paraglacial/post-glacial origin of the failure, an event that fits well with the last chronologies available on the Last Glacial Maximum in the Nepal Himalaya (Owen & Dortch 2014). IRSL dating of silts suggests a progressive trend to dryness of these Northern Himalayan regions from Late Pleistocene to Holocene.

Note on Infrared Stimulated Luminescence (IRSL) dating

The IRSL samples, PHOK 2 and PHOK 1, were collected respectively at the top and the bottom of the silty deposit, below the surface soil. IRSL measu were performed at Montreal laboratory by S. Balescu, S. Huot and M. Lamothe. The corrected IRSL ages of PHOK 2 and PHOK 1, respectively 4.7 ± 0.3 ka and 12.6 ± 0.7 ka, are stratigraphically consistent (*Fort et al.*, 2013).

Letter and Letters 1,000 1 8 - m 2 10 - a 12 - 2 - 10 / 10 / 10 / 10 / 10

Annapurna, 8078 m

landslide dar lake, volum 160 M m³

Linear, guily



Volume (10⁶ m³) Age Geological Type of Nature of Geomorphic context failure material impact References

HIC
Cozartanic Egencidados
Denner of Ferrito (1997), 1981. (1994)
Denner of Ferrito (1994)
Denner of Ferrito (1997), 1981. (1994)
Denner of Ferrit

recemented blocks Sledge-like sleding Pott glacial Pott glacial gravity instability pseudotachylites)

biotitic conisses)

and incrusive orianite)

(gneisses) Post glacial gravity

The Phoksumdo lake (3600 m asl.; area of 4.5-to-5 km²) is the second largest lake of Nepal: it owes its origin to the damming of the Suli Gad River by the large (4.5 km3) collapse of a mountain wall (Dhaulagiri limestones) culminating at 5148 m, SE of the lake (Fort & Rimal 2010), According to Yaai (1997), the collapse may have occurred 30 to 40 ka, an interpretation revised by this study.



The landslide morphology

Sampling site of Ti09-02

The Pokhara gravels fill (#3): result of a gigantic rock-

avalanche 500-years old. Mechanism? Catastrophic collapse of the Annapurna IV (thrust) front, possibly

seismically triggered in 1505 (Fort 1987; 2010)

presents a series of mounds (metres to tens of metres high) and depressions of varying size. Its mass is composed of large blocks of massive dolomites found as far as west of Ringmo Village and down to the southern rim of the landslide dam

Most of the Suli scarp is uphold by limestones outcrops, but along the central part, the former Phoksumdo gorge is filled in by dark grey, fine sediments (till material) and overlain by orange conglomerates (including dolomites boulders), then by yellowish, unsorted conglomeratic material blocks and calcareous silty matrix). The presence of glacial till at the very base of the sequence suggests the rockslide event is post-glacial.

peaks

The Dhampu-Chhova rock avalanche (10⁹

m³, area extent 10 km²) was derived from

the northward failure of the Kaiku ridge uphold by north-dipping, upper crystal

lines of the Higher Himalava (Fort 1974;

2000). It dammed the Kali Gandaki River,

with complex interactions with the Late

Pleistocene ice tongues derived from the



5. COMPARISONS and CONCLUDING REMARKS

Phoksundo-Ringmo

- Efficient drainage blockage, but no backwash sedimentation (predominant carbonates + low discharge)

- Prominant knick point: karst + boulder pavement reducing incision => prominent hanging valley

- Triggering factor: glacial retreat («paraglacial collapse»), cf. ³⁶Cl dates Dhampu-Choova

- Efficient drainage blockage, and extensive lacustrine sedimentation (glacial sediment reworking + Spiti shales + glacial- and snow-melt discharges) Knick points and associated epigenetic gorges + braided Kali Gandaki valley: delayed incision => relatively hanging valley - No climate forcing, cf. ¹⁰Be dates => Triggering factor related to the North Himalavan Detachment Fault + foliation dip of Upper Himalayan Gneisses

Both examples confirm that giant landslides play a significant role in the: Destruction of Himalayan topography (Fort 1988), Bedrock protection from river incision (Korup et al. 2010), Delay of sediment transport outward from the mountain zone (cf. sediment storages & budgets; Fort & Cossart 2013; Blöthe J. & Korup O., 2013) - Local controls exerted by topography and climate (eg. magnitude of glaciation + snow <--> discharge) + lithology (eg. sediment supply <--> storage)

- Complex interplay between glaciers and mountain wall collapses (Fort 2000)

the rocky face bounding the lake in its SE part. 4. DHAMPU-CHOOYA SITE

caused



e of Quaternary elaciation in the Himalayan - Tibetan duction / Geophys Res 105 23753-23759 - Landslide dams of Tal, Latamrang, Ghatta khola, Ringr hazards. J. Nepol Geol. Soc., 22, 371-380. Wepal ministry of the Photosundo Tal, Dolpa district, western Nepal. Jo. 4, 1977, Origin of the Photosundo Tal, Dolpa district, western Nepal. Jo. A, Zech M, Kubik P.W., Kharki K., Zech W. (2009). Deglaciation and land A and Annual Control of the Annual Control of Control Nepal. Jour. Nepal Geol. Soc., 15:.1-7.

MCT

Both the rock avalanche and

and level of the "Marpha

Kagbeni) (Fort 1980).

glaciers controlled the existence

Lake" (lacustrine deposits up to



Dhaulagiri (8167 m) and Nilgiris (7061 m) Be-10 cosmic ray exposure ages (CEREGE, Gribenski 2010) i09-01 : 29 674 ± 1 022 ka Ti09-02 · 24 844 + 628 ka Ti09-03 : 29 694 + 1 011 ka The discrepancy in the dating of both lacustrine and failure events raise the issue of the nature of the lacustrine dam: (i) rockavalanche as initially thought (Fort 1974: 1980: Baade 2000) or (ii) possibly a glacial tongue controlling the lake

level (Fort and Cossart 2013); (iii) repeated failures cannot

by Paris-Diderot Univ. (UFRs GHSS and STEP) and by LN2C (Aix-Marseille Univ.). ASTER AMS national facility (CEREGE, Aix-en-Provence) is supported by the INSU/CNRS, the French ion IRD and CEA



Acknowleaments: ermission for geological and

geomorphological investigations was given by the Dpt of National Parks, Ministry of Forests, and free entrance in the Shey-Phoksundo National Park was delivered by Home Ministry, Government of Nepal. Our very deep appreciation goes to Dr. Som Sapitota (Dpt of Mines and Geology) for his bein