

The influence of tree species mixture on ecosystem-level carbon accumulation and water use in a mixed boreal plantation

Charlotte Grossiord, André Granier, Arthur Gessler, Martina Pollastrini,

Damien Bonal

▶ To cite this version:

Charlotte Grossiord, André Granier, Arthur Gessler, Martina Pollastrini, Damien Bonal. The influence of tree species mixture on ecosystem-level carbon accumulation and water use in a mixed boreal plantation. Forest Ecology and Management, 2013, 298, pp.82-92. 10.1016/j.foreco.2013.03.001 . hal-01268099

HAL Id: hal-01268099 https://hal.science/hal-01268099

Submitted on 29 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

"Does Drought Influence the Relationship between Biodiversity and Ecosystem Functioning in Boreal Forests?"

- 3
- 4 Charlotte Grossiord,^{1,2} André Granier,^{1,2} Arthur Gessler,^{3,4} Tommaso Jucker,⁵ Damien Bonal^{1,2*}
- 5

6	¹ INRA, UMR	1137 Ecologie et Ec	cophysiologie Fo	restières, 54280 (Champenoux, France
-) -				F ,

- 7 ²Université de Lorraine, UMR 1137 Ecologie et Ecophysiologie Forestières, 54500
- 8 Vandœuvre-lès-Nancy, France
- 9 ³Leibniz Centre for Agricultural Landscape Research (ZALF), Institute for Landscape
- 10 Biogeochemistry, Eberswalderstrasse 84 15374 Müncheberg, Germany
- ⁴Berlin-Brandenburg Institute of Advanced Biodiversity Research (BBIB), 14195 Berlin,
 Germany
- ⁵Forest Ecology and Conservation Group, Department of Plant Sciences, University of
 Cambridge, Downing Street, Cambridge CB2 3EA, United Kingdom
- 15
- 16 Corresponding Author: Damien Bonal (bonal@nancy.inra.fr)
- 17 Tel + 33 3 83 39 73 43 Fax +33 3 83 39 40 22
- 18
- 19 Author contribution:
- 20 DB, AG¹ and AG² designed the experimental study. DB and CG conducted the field work. TJ
- 21 provided the productivity data. CG, DB, TJ, AG¹ and AG² analyzed the results. CG and DB
- 22 wrote the first draft of this manuscript and all authors substantially contributed to revisions.
 - 1

23	
24	
25	Total word count:
26	Abstract: 269
27	Manuscript (without abstract): 6236
28	Figures: 3
29	Tables: 2
30	
31	Shortened title: Biodiversity-Ecosystem Functioning Relationship

33 Abstract

34 In mixed forests, interactions among species influence ecosystem functioning but 35 environmental conditions also play an important role in shaping relationships between biodiversity and ecosystem functioning. In the context of climate change, the carbon and 36 37 water balance in pure vs. mixed forest stands may be differentially influenced by changing soil water availability. To test this hypothesis, we compared the influence of biodiversity on 38 39 water use efficiency (WUE_s) in boreal forests between wet and dry years. We assessed the 40 carbon isotope composition (δ^{13} C) of tree rings in *Betula pendula*, *Pinus sylvestris* and *Picea* 41 *abies* growing in pure vs. mixed stands. In addition, we tested whether differences in WUEs affected patterns of basal area increment (BAIs). No biodiversity effect was found for stand 42 δ^{13} C (δ^{13} C_s) during the wet year. However, there was a significant increase in δ^{13} C_s between 43 the wet and the dry year and a significant effect of biodiversity on $\delta^{13}C_s$ in the dry year. The 44 increase in $\delta^{13}C_s$ in mixed stands was associated with both selection and complementarity 45 46 effects. Although BAI_s decreased significantly in the dry year, changes in $\delta^{13}C_s$ did not 47 translate into variations in BAI_s along the biodiversity gradient. Our results confirmed that the 48 physiological response of boreal forest ecosystems to changing soil water conditions is 49 influenced by species interactions and that during dry growing seasons, species interactions in 50 mixed stands can lead to lower soil moisture availability. This illustrates that biodiversity 51 effects can also be negative in mixed stands in the sense that soil resources can be more 52 intensively exhausted. Overall, our results confirm that in boreal forests, the biodiversity-53 ecosystem functioning relationship depends on local environmental conditions.

54 **Key-words**: biodiversity, boreal forest, drought, δ^{13} C, mixed forest, water use efficiency

55 INTRODUCTION

The biodiversity and ecosystem functioning (B-EF) relationship has received growing interest 56 57 from ecologists throughout the last two decades due to the rapid loss of diversity observed 58 during the last century (Symstad and others 2003). Several mechanisms have been put 59 forward to explain how community composition and richness can either negatively or 60 positively influence ecosystem functions. A negative influence on biodiversity can result from 61 direct competition for resources among species in a given ecosystem. Negative interactions 62 commonly arise when ecological niches overlap and/or when species share the same 63 functional characteristics and are thus functionally redundant in the ecosystem (Naeem 2008; 64 Vilà and Sardans 1999). In contrast, positive B-EF relationships are commonly attributed to 65 two other mechanisms: complementarity and selection. Complementary use of resources 66 among species refers both to ecological niche partitioning and facilitation (Loreau and Hector 67 2001) and implies species coexistence without major interspecific competition for resources. 68 The selection effect recognises that the probability of occurrence of high-performing species 69 is greater in highly diverse ecosystems (Loreau and Hector 2001). Studies investigating 70 complementarity and selection effects mainly focused on ecosystem productivity (e.g. 71 Bradford 2011; Fargione and others 2007; Isbell and others 2009; Morin and others 2011; 72 Zhang and others 2012), while only a few were interested in other ecosystem functional traits 73 (e.g. Forrester and others 2010; Kunert and others 2012; Meinen and others 2009). 74 In addition to competition, complementarity and selection effects, local environmental 75 conditions also play an important role in shaping B-EF relationships (Belote and others 2011; 76 Hooper and Dukes, 2004). Under the assumptions of the "stress-gradient" hypothesis, which predicts that the net outcome of biotic interactions (competition and facilitation) shifts from 77 78 negative to positive along gradients of limiting physical conditions (Bertness and Callaway 79 1994), positive biodiversity effects are expected to be more common in severely resource80 limited conditions while negative effects should prevail in richer and milder environments. 81 This general conceptual model has recently been refined (Maestre and others 2009) and is 82 widely supported in the literature (Herbert and others 2004; Jucker and Coomes 2012; Steudel 83 and others 2012; He and others 2013; Wang and others 2013). In the context of climatic 84 change, most regions around the world are expected to encounter more extreme 85 environmental conditions (IPCC, 2007). The "stress-gradient" hypothesis is therefore of great 86 interest since B-EF relationships are likely to change in the future. Whether or not more 87 diverse ecosystems might be better adapted and/or more resilient to these changes is an 88 important issue to investigate.

89 In the boreal climate zone, simulations predict a general shift from short, cool summers 90 towards longer, warmer summers (IPCC 2007; Jylhä and others 2010). Thus, boreal forest 91 ecosystems are expected to encounter more frequent and intense reduced soil water 92 availability in summer. Contrasted responses of carbon and water fluxes in forest ecosystems 93 to warmer and drier climate have already been observed (reviewed in Boisvenue and Running 94 2006). Few studies so far were conducted on the response of boreal forests to these 95 conditions. Nevertheless, Dulamsuren and others (2010) showed that increasing summer 96 temperatures accompanied by decreasing precipitation lead to reduced productivity in taiga 97 forests in Mongolia.

98 Plants adapt to reduced soil water conditions through numerous physiological and/or 99 morphological processes (review in Kozlowski and Pallardy 2002). At the leaf level, under 100 drought conditions, they must manage the trade-off between optimum carbon gain for growth 101 and loss of water through transpiration (Farquhar and others 1982) which usually results in an 102 increase in intrinsic water use efficiency (WUE_{int}), defined as the ratio between CO₂ 103 assimilation during photosynthesis and stomatal conductance for water vapour (e.g. Zhang 104 and Marshall 1994). At ecosystem level, carbon and water fluxes are influenced by species-

specific functional responses to environmental conditions and by intra- and inter-specific
interactions both below- and above-ground. Both positive and negative interactions among
species may occur in mixed stands and this may lead to differing spatial and temporal
resource availability and physiological and morphological adaptations within species. These
complex interactions mean that general predictions on stand-level water use efficiency
(WUE_s) under dry conditions cannot solely be based on individual species responses to these
conditions.

112 In this context, little information is available concerning the relationship between tree 113 biodiversity and the regulation of carbon and water fluxes in boreal forest ecosystems. 114 Gamfeldt and others (2013) found positive relationships between tree species richness and 115 multiple ecosystem services (tree biomass, soil carbon storage, berry production and game production potential) in production forests in Sweden and explained the observed positive B-116 117 EF relationship by facilitation processes among tree species. In contrast, Grossiord and others 118 (2013a) found no complementary effect for biomass production and WUE_s in a young boreal 119 plantation, though they did find a weak selection effect. These studies, however, did not 120 compare the B-EF relationship in contrasted environmental conditions.

121 In this study, we tested the following assumptions: i) in boreal forest ecosystems, the stand-122 level carbon and water balance under non-limiting soil water conditions (wet year) should not 123 necessarily depend on biodiversity effects, and ii) under limiting soil water conditions (dry year), species identity, species combinations and richness levels should influence ecosystem 124 125 functioning. We analyzed the influence of species richness on time-integrated WUEs estimated from the carbon isotope composition (δ^{13} C) (Farquhar and others 1982) of ring 126 whole wood measured in tree rings from two years with contrasting soil water conditions (wet 127 128 year in 2004 and dry year in 2006). Samples were taken from 26 mature stands in Eastern 129 Finland, which were either pure or mixed with varying percentages of *Betula pendula*, *Pinus*

- *sylvestris* and *Picea abies*. We also tested whether changes in WUE_s would influence stand
- 131 basal area increment (BAI_s).

134 MATERIALS AND METHODS

135 Site description

The study was conducted in August 2012 in 26 stands of boreal forest that are distributed over a 2000 km² area around Joensuu, Finland (62.60°N, 29.76°E). The stands were 30 m x 30 m and included pure, 2-species mixture, or 3-species mixture of Scots pine (*Pinus sylvestris*), Norway spruce (*Picea abies*) and Silver birch (*Betula pendula*). Four replicates of each combination were selected except for pure birch stands that were sampled only twice. Mean tree age within the stands was 33.9 ± 5.8 years. Tree species characteristics are given in Table 142 1.

143 The stands were selected according to the following specifications: no change in management 144 had occurred in the last 5 years; confounding factors like altitude, topography or soil type were kept to a minimum between selected stands; in a given forest patch, the stands were 145 146 more than 500 m apart to avoid spatial autocorrelation; stands were mostly even-aged and 147 single-layered. More detailed information on the selection procedure of the study stands can 148 be found at http://www.fundiveurope.eu. The selected stands were situated between 80 and 149 200 m a.s.l. on Podzols soils (FAO classification) above mica schist bedrock. Mean annual 150 rainfall in the region is around 700 mm and mean annual temperature is 2.1°C.

151

152 Selection of the target years

The daily water balance model "BILJOU" (Granier and others 1999) was implemented in order to quantify soil water availability at a daily time-scale in the study region and then to select the driest and the wettest year over the past fifteen years. We used above-canopy measurements of daily meteorological conditions (rainfall, global radiation, air temperature and humidity, wind speed) from the Hyytiälä forestry field station (61.84°N, 24.28°E, 153 m

a.s.l.) to run the model. Although this meteorological station is situated 300 km south-west of 158 159 Joensuu, it is the nearest source from which data could be obtained to estimate daily relative extractable soil water (REW) over the 1997-2011 period (Fig. 1). The model was initialized 160 assuming that soil was at field capacity on 1st January. We used a leaf area index of 5.0 m² m⁻² 161 162 and a soil holding capacity of 125 mm as input data for model simulations. We discarded the vears 1998 and 2010 because of too many missing data. Soil water deficit was assumed to 163 164 occur when the REW dropped below 0.4, a threshold value that has been shown to induce 165 stomatal closure and interrupt radial growth in forest trees (Granier and others 1999). For each 166 remaining year from 1997 to 2011, we characterized annual water stress using the starting day 167 and the duration of the period when REW dropped below the threshold limit coupled with a 168 water stress index which cumulates the difference between simulated REW and the 0.4 threshold (Granier and others 1999). These calculations revealed that in 2006 water stress was 169 170 the most severe and in 2004 it was the least severe (Fig. 1). Drought started on June 16 in 171 2006 while there was no drought in 2004. Total precipitation during the growing season (May 172 to September) was 157 and 303 mm in 2006 and 2004, respectively.

173

174 Wood carbon isotope composition

175 Six trees per species in each pure stand and three trees per species in mixed stands were 176 randomly selected among the ten trees with the largest diameter at breast height. For each 177 selected tree we extracted one 5-mm-diameter wood core at breast height (1.3 m) from the 178 south side of the trunk. The fresh samples were shipped to INRA Nancy and oven-dried at 179 50°C for 72 hours to reach constant weight before being filed with a scalpel for tree-ring dating. For both target years, we carefully extracted the late wood with a scalpel. The wood 180 181 fragments from a given species in a given stand were then pooled and ground into a fine 182 powder. We combined the wood fragments from individuals of the same species in the same 183 stand to obtain enough material for carbon isotope analyses, to ensure adequate sample 184 homogeneity, and to reduce experimentation costs. Although we sampled only a few trees per 185 species in each stand, the level of replication was adequate given the imprint of environmental 186 signals on tree-ring carbon isotope composition (e.g. Peñuelas and other 2008). From 0.4 to 187 1.5 mg of the ground wood samples were weighed on a high-precision scale (MX5, Mettler 188 Toledo, Viroflay, FR) and placed into tin capsules (Elemental Microanalysis Limited, Devon, 189 UK) for δ^{13} C analyses. Samples were analyzed with a PDZ Europa 20-20 isotope ratio mass 190 spectrometer (Sercon Ltd., Cheshire, UK) at the Stable Isotope Facility of UC Davis, CA, 191 USA. All isotopic measurements are reported in the standard delta notation (δ , ∞):

192

[1]

193 where R_{sample} and $R_{standard}$ are the ${}^{13}C/{}^{12}C$ ratios of the samples and the Vienna Pee Dee 194 Belemnite (VPDB) standard.

195

196 Wood basal area increment

197 To quantify productivity, we randomly selected five trees per species from each pure stand, and three trees per species in mixed stands, among the twelve (pure stands) or eight (mixed 198 199 stands) trees with the largest diameter at breast height. For each tree, one 5-mm-diameter 200 wood core was extracted at breast height from the south side of the trunk. These cores were different from the ones taken for δ^{13} C analysis. Once air dried, the core samples were mounted 201 202 on wooden boards and then sanded with paper of progressively finer grit sizes. A high 203 resolution (2400 dpi) flatbed scanner was then used to image the cores, following which 204 diameter growth measurements and crossdating were performed with the CDendro software 205 suite (Cybis Elektronik & Data, Saltsjöbaden, Sweden). For each of the studied years, diameter growth measurements of the individual trees were converted to basal area increment, 206

then standardized basal area increment (BAI_i, cm² cm⁻² yr⁻¹) was obtained by dividing basal
area increment by the basal area of each tree.

209

210 Stand-level carbon isotope composition and basal area increment

Based on species-specific δ^{13} C values (δ^{13} C_i) for each stand, we calculated stand-level carbon isotope composition (δ^{13} C_s). The contribution of a given species to δ^{13} C_s depends on its species-specific contribution to stand CO₂ assimilation rates (Lloyd and Farquhar 1994). Consequently, when δ^{13} C_i values are scaled up from species level to stand level, δ^{13} C_i values should be weighted by species assimilation rates. However, since species-specific CO₂ assimilation rates were not available in this study, we used species total basal area (BA_i, cm²) in each stand as a proxy, following Bonal and others (2000). δ^{13} C_s was thus calculated as:

219 where *N* is the number of species in the mixture.

220 δ^{13} C in C₃ plants constitutes a record of the intercellular / atmospheric CO₂ concentration ratio 221 during the period in which the carbon was fixed and is strongly positively correlated to leaf 222 intrinsic water use efficiency (WUE_{int}) (Farquhar and others 1982) as shown in the following 223 equation:

where C_a is the CO₂ concentration in the atmosphere, δ_a and $\delta^{13}C$ are respectively the carbon isotope compositions of the atmosphere and of the sample, *a* represents the fractionation occurring due to diffusion in air (4.4 ‰), and *b* is the fractionation caused by carboxylation (27 ‰). We thus used Equation 3 to calculate stand-level water use efficiency (WUE_s) based on $\delta^{13}C_s$. To estimate stand basal area increment (BAI_s, cm² cm⁻² yr⁻¹) of the two studied years (2004 and 2006) and of the year following the driest year (2007), we considered the relative importance of each species in terms of contribution to total basal area. Thus, BAI_s was calculated as:

235

236 Data analysis

237 All analyses were performed using the statistical software R 2.14.1 (R Development Core Team, 2011). We tested three effects that could impact $\delta^{13}C_i$, $\delta^{13}C_s$, BAI_i and BAI_s. Firstly, tree 238 239 biodiversity, which we characterised for each stand with the Shannon biodiversity index. This 240 index expresses the uncertainty of predicting the proportion of stand basal area a given tree 241 species represents in binary digits (ranging from zero for pure stands to the logarithm in base two of the total basal area of the species in the stand). Secondly, competition intensity in each 242 243 stand, which we expressed as the total basal area of the stand. Thirdly, year. We applied a 244 series of mixed-effect models to analyze the impact of the three effects and their interactions on $\delta^{13}C_i$, $\delta^{13}C_s$, BAI_i and BAI_s with "stand" as a random effect. To test for any biodiversity 245 effect during the recovery phase to drought (year 2007), we applied the same mixed-effect 246 247 model on BAIs. Contrasts were used to test for relevant differences between fixed effects when the overall model was significant. We calculated the difference in $\delta^{13}C_S$ and BAI_S 248 between the dry and wet year ($\Delta \delta^{13}C_s$ and ΔBAI_s , respectively) and used Spearman 249 regressions to test the relationship between the Shannon biodiversity index and $\Delta \delta^{^{13}}C_s$ or 250 251 ΔBAI_s .

252 When a significant effect of biodiversity on $\delta^{13}C_s$ or BAI_s was observed, we evaluated the net 253 biodiversity effect, the complementarity effect, and the selection effect on $\delta^{13}C_s$ or BAI_s with

the method developed by Grossiord and others (2013b). We used one-way ANOVA and posthoc Tukey tests to check for differences in these biodiversity effects among species richness
levels. We used t-tests to check all three biodiversity effects for differences from 0 for each
mixture level and for each mixture type (identity of the mixture).

258 **RESULTS**

259 Wood carbon isotope composition

We observed considerable variability in $\delta^{13}C_i$ across species, species mixtures and years, 260 261 ranging from -28.7 ‰ for birch in a 2-species mixture with spruce to -26.2 ‰ for spruce in a 2-species mixture with pine during the wet year, and from -27.4 ‰ for birch in a mixture with 262 263 spruce to -24.2 ‰ for pine in a 3-species mixture during the dry year. Mixed-effect models revealed a significant species effect (P < 0.001), year effect (P < 0.001) and an interaction 264 between Shannon index and year (P = 0.019), but no effect of the Shannon index alone, the 265 266 interactions of the Shannon index and the species, or the interactions of the year and the species (P > 0.050). The significant effect of the interaction between Shannon index and year 267 268 was only observed for pine and spruce (P = 0.012 and P = 0.020, respectively), indicating for these species a differential response of $\delta^{13}C_i$ to the mixture in the two years. 269

For each species, whatever the richness level, we found significantly higher values of $\delta^{13}C_i$ during the dry year than during the wet year (P = 0.003 for birch, P = 0.017 for pine and P = 0.037 for spruce). For the wet year, we did not observe any effect of the Shannon index or of competition intensity on $\delta^{13}C_i$ for any species (P > 0.050). In contrast, during the dry year, pine and spruce showed a significant positive Shannon index effect (P = 0.013 and P = 0.027, respectively) and a competition intensity effect (P < 0.001 and P = 0.010, respectively), whereas birch did not display any significant patterns (P > 0.050).

At plot level, $\delta^{13}C_s$ values ranged from -28.1 ‰ for a pine-spruce mixture to -26.6 ‰ for a pure stand of spruce during the wet year and from -26.9 ‰ in a pure stand of pine to -24.6 ‰ for a 3-species mixture during the dry year (Fig. 2). We observed a significant effect of year on $\delta^{13}C_s$ (P < 0.001), with higher values of $\delta^{13}C_s$ during the dry year, as well as a significant interaction between Shannon index and year (P = 0.012). Specifically, while there was no significant relationship between the Shannon index or the competition index and $\delta^{13}C_s$ during the wet year (P > 0.050), a highly significant and positive Shannon index effect was apparent (P = 0.008) during the dry year, along with a marginal influence of competition intensity (P =0.067). Differences in $\delta^{13}C_s$ between the dry and the wet year ($\Delta\delta^{13}C_s$) for each stand were positively correlated with the Shannon index (P = 0.038).

287

288 Wood basal area increment

The basal area increment (BAI_i) of individual species ranged from 0.3×10⁻³ cm² cm⁻² yr⁻¹ for 289 pine mixed with birch to 52.9×10^{-3} cm⁻² yr⁻¹ for spruce mixed with pine during the wet 290 year, and from 0.7×10^{-3} cm⁻² yr⁻¹ for pine mixed with birch to 41.7×10^{-3} cm⁻² yr⁻¹ for 291 292 birch mixed with spruce during the dry year. Mixed-effect models revealed a significant species effect (P < 0.001), year effect (P < 0.001) and an interaction between the species and 293 the year (P < 0.001), but no effect of the Shannon index alone, the interactions between the 294 Shannon index and species, or the interactions between the Shannon index and the year (P >295 296 0.050). For spruce, we found significantly lower values of BAI_i during the dry year than during the wet year (P < 0.001), while for birch and pine, no significant changes in BAI_i were 297 found. For the wet year, we did not observe any effect of the Shannon index or competition 298 intensity on BAI_i for any species (P > 0.05). For the dry year, pine showed a significant 299 300 positive Shannon index effect (P = 0.035) and no competition intensity effect (P = 0.813). Birch and spruce did not display any significant Shannon or competition effects during the 301 dry year (*P* > 0.050). 302

BAI_s values ranged from 14.9×10⁻³ cm² cm⁻² yr⁻¹ in a birch-pine mixture to 61.3×10⁻³ cm² cm⁻² yr⁻¹ in a pure stand of spruce for the wet year and from 16.9×10⁻³ cm² cm⁻² yr⁻¹ in a birch-pine mixture to 59.8×10⁻³ cm² cm⁻² yr⁻¹ in a birch-spruce mixture for the dry year (Fig. 2). In 2007,

the year following the driest year, BAI_s values ranged from 16.8×10^{-3} cm⁻² yr⁻¹ in a pure 306 pine stand to 59.4×10⁻³ cm⁻² cm⁻² vr⁻¹ in a pine-spruce mixture (Fig. 2). The interaction between 307 the Shannon index and the year was not significant (P = 0.976). However, we observed a 308 significant effect of the year on BAI_{S} (*P* = 0.001), with higher values during the wet year in 309 comparison to the dry and the recovery years (Fig. 2). We did not detect any significant effect 310 of the Shannon index or the competition index on BAI_s in either year (P > 0.050). We found 311 312 that differences in BAI_s between the dry and the wet year (Δ BAI_s) were not correlated with 313 the Shannon index (P = 0.143).

314

315 Biodiversity effects on stand carbon isotope composition

316 Overall, there was no significant change in either net, complementarity or selection effect on $\delta^{13}C_s$ with increasing species richness during the dry year (P > 0.050) (Fig. 3). However, 317 when each mixture level was considered independently, complementarity, selection and net 318 effects were significantly higher than zero in the 2-species mixtures (P < 0.050), and mainly 319 positive - although not significantly different from 0 - in the 3-species mixtures (P > 0.050). 320 321 When we tested each different species mixture for significance of the biodiversity effects, we 322 found that none of the three effects were significantly different from zero (P > 0.050) in the pine-spruce mixtures. However, for birch-pine and birch-spruce mixtures, net and 323 324 complementarity effects were significantly higher than zero (P < 0.050), while no selection 325 effects emerged (P > 0.050).

326 **DISCUSSION**

327 **1. Climate effect**

328 Our stable isotope analysis revealed a strong physiological response in boreal trees to 329 changing soil water conditions and helped clarify the mechanisms responsible for observed 330 changes both at the leaf and stand level. The increase in both individual species' $\delta^{13}C_i$ and stand level $\delta^{13}C_s$ from wet to dry years (Fig. 2) reflects an active regulation of carbon and 331 332 water fluxes in this boreal ecosystem and the strong sensitivity of the three species to limiting 333 soil water conditions. Assuming that the difference in δ_a between the two selected years was 334 negligible, the magnitude of the increase in $\delta^{13}C_s$ reflected a mean increase in WUE_s of 17.0 ± 1.8 μ mol mol⁻¹ (i.e. +19.2 ± 1.5 %). This increase is consistent with the extensive body of 335 336 literature documenting the functional response of boreal trees to drought conditions (e.g. 337 Vaganov and others 2009; Zhang and Marshall 1994; Saurer and others 2004). This pattern 338 indicates that in the context of climate change, when more frequent periods of water shortage 339 in north-European regions can be expected (IPCC 2007), the three locally-adapted species 340 studied here will be more efficient in using available water resources for carbon acquisition. 341 However, carbon sequestration and storage rates in wood may well decline; indeed, we found 342 that the increase in stand $\delta^{13}C_s$ during the dry year was associated with a decrease in BAI_s. 343 This is unsurprising as numerous studies conducted in boreal forests have shown lower 344 productivity under water limiting conditions (e.g. Kljun and others 2007).

345

346 2. Biodiversity effects under wet conditions

Our results confirmed our first hypothesis that under non-limiting soil water conditions there would be little or no effect of biodiversity on $\delta^{13}C_s$ and BAI_s (Fig. 2). Interactions among species under non-limiting soil water conditions did not lead to any clear B-EF relationships in our boreal study forests, which is consistent with the expectations of the "stress-gradient"

hypothesis (Bertness and Callaway 1994). This result is also consistent with a recent study we 351 352 conducted over a wet summer on the influence of biodiversity on species- and stand-level 353 carbon and water balance in a young boreal plantation in Finland (Grossiord and others 354 2013a). Therein, we argued that in addition to non-limiting soil water conditions, the young 355 ontogenic stage of the plantation likely caused the absence of any biodiversity effect. 356 However, the present study does not support this assumption. Similar observations made in 357 other ecosystem types also concluded on the absence and/or the decrease of biodiversity 358 effects under non-limiting environmental conditions (e.g. Hughes and Stachowicz 2004; 359 Steudel and others 2012).

360

361 **3. Contrasting biodiversity effects under wet and dry conditions**

The major result of this study was the positive relationship between biodiversity and the 362 difference in $\delta^{13}C_s$ between dry and wet years ($\Delta\delta^{13}C_s$) (Fig. 2). Drier conditions globally 363 enhanced $\delta^{13}C_s$, and thus WUE_s, but this influence depended on species interactions. Even 364 365 though the relationship was rather dispersed, stands with greater diversity showed the greatest increase in $\delta^{13}C_s$, pointing to a greater compromise between carbon accumulation and water 366 consumption under limiting soil water conditions. However, those changes in ecosystem-scale 367 368 physiological acclimatization to drought failed to minimize the decrease in BAIs during the 369 dry year and did not influence the productivity of these plots the following year (Fig. 2). The 370 literature actually mentions this apparent inconsistency (e.g. Vaganov and others 2009). The 371 compromise between carbon acquisition and water loss at leaf or plant level usually has a 372 moderate effect on how much assimilated carbon is invested in secondary growth (Flanagan 373 and Johnsen 1995). Carbon allocation to above- or below-ground components in response to 374 soil water conditions strongly varies among species (e.g. Axelsson and Axelsson 1986) and 375 could partially explain the absence of enhanced BAIs in our study.

377 4. Origin of the biodiversity effects

Partitioning the influence of biodiversity on $\delta^{13}C_s$ into selection and complementarity effects 378 379 revealed that during the dry year the proportionally greater increase in WUE_s observed in the 2-species mixtures was a result of both mechanisms (Fig. 3). Selection effects refer to a higher 380 381 performance of one or several species in mixtures on the cost of others. Therefore, the pattern of increasing $\delta^{13}C_s$ we observed was most likely driven by the two conifers rather than by 382 birch, since both conifers showed higher $\delta^{13}C_i$ in 2-species mixtures while birch did not (Table 383 1). Higher $\delta^{13}C_s$ values in the 2-species mixtures with birch were also partially explained by a 384 complementarity effect, probably as a result of different rooting strategies among the species. 385 386 Birch is known to have deeper rooting systems that explore larger volumes than those of pine 387 and spruce, thus enabling birch to access both shallow and deep soil layers (Kalliokoski and others 2008). The fact that birch $\delta^{13}C_i$ values did not respond to increasing diversity during the 388 389 dry year supports this assumption and suggests that the effects of intra- and inter-specific 390 interactions for water acquisition for this species are similar. In contrast, the complementarity effect for $\delta^{13}C_s$ was not significant in stands where the two conifers cohabited. *Pinus* 391 sylvestris and Picea abies are known to present similar rooting system development when 392 393 cohabiting in mixed stands (Kalliokoski and others 2008). It therefore appears that the two 394 conifers were in part functionally redundant in terms of carbon and water acquisition and use 395 in these mixed stands.

Although in the 3-species mixtures none of the three biodiversity effects were significantly different from zero, overall they were generally consistent with results for the 2-species mixed stands (Fig. 3). Numerous B-EF studies have revealed such saturating or even hump-shaped relationships (e.g. Gamfeld and others 2013; Vilà and others 2013). One explanation for this type of pattern is increasing functional redundancy at higher levels of species richness (as 401 observed for the two conifers here), which leads to faster and more complete exploitation of402 available resources in stands with greater richness.

403

404 5. Other potential origins of the biodiversity effects

Variations in δ^{13} C in tree rings reflect physiological processes of acclimatization to 405 environmental conditions and physical archives of short- or long-term changes in soil 406 407 moisture and/or air humidity in forests (Saurer and others 2004; Andreu and others 2008; 408 Mölder and others 2011). Therefore, the significantly higher $\delta^{13}C_s$ values we observed along the diversity gradient in the dry year could be viewed not only as a consequence of species 409 410 interactions, but also as a result of lower soil water availability in mixed stands as compared 411 to pure ones. High soil moisture exhaustion in mixed stands during dry years was already reported in other hydrological investigations in mixed species forests (Gebauer and others 412 413 2012). Such a feedback effect could arise because of the particularly high competitiveness of birch for water resources and its high water consumption (Kalliokoski and others 2008). This 414 415 would lead to a higher drought exposure for the conifers in the mixed stands and thus reduced 416 transpiration through stomatal regulation. This increased drought exposure may have been 417 also the cause for the absence of any increase in BAIs in the mixed stands. Nevertheless, 418 differences in canopy structure inducing different atmospheric humidity and light conditions 419 between pure and mixed stands (Kelty 2006) as well as below-ground competition among these species for nutrients (nitrogen, phosphorus...) may also have contributed to the mixture 420 421 effects we detected; however, this is beyond the scope of the present study.

422

424 CONCLUSION

425 In our boreal study region, we demonstrated a clear interaction between the 426 biodiversity/ecosystem carbon and water flux relationship and environmental conditions, consistent with the "stress gradient" hypothesis. The increase in $\delta^{13}C_s$, and thus in ecosystem-427 428 level water use efficiency, with drier soil conditions was much higher in mixed stands than in 429 pure ones, but this did not prevent a decrease in above-ground growth rates with drought. At 430 the species level, functional similarities between the two conifers lead to a redundant 431 contribution of these two species to ecosystem-level carbon and water balance. In contrast, the 432 presence of birch in mixed stands strongly impacted the water use efficiency of the conifers 433 under limiting water conditions. Our results suggest that species interactions in mixed stands 434 during dry growing seasons can lead to lower soil moisture availability as compared to pure 435 ones. This illustrates that biodiversity effects can also be negative in mixed stands in the sense 436 that soil resources can be more intensively exhausted. Our study highlights the need to take 437 into account not only species growth characteristics but also species functional traits when 438 managing boreal mixed forest in order to bolster mixed-forests in a context of climate change.

440 Acknowledgments

We thank Leena Finér and Timo Domisch for the selection of the stands and for their help 441 442 when preparing this study, and Timo Vesala for sharing the climatic data. We would like to 443 acknowledge the two anonymous reviewers who greatly contributed to the improvement of a previous version of this manuscript. The research leading to these results received funding 444 from the European Union Seventh Framework Programme (FP7/2007-2013) under grant 445 agreement n° 265171. CG was supported by a grant from the INRA Nancy in the framework 446 of the FunDivEUROPE project. This work was conducted in the framework of the Laboratory 447 448 of Excellence ARBRE (ANR-12- LABXARBRE-01) supported the French National Research 449 Agency.

451 **References**

- 452 Andreu L, Planells O, Gutiérrez E, Helle G, Schleser GH. 2008. Climatic significance of tree-453 ring width and δ^{13} C in a Spanish pine forest network. Tellus B 60:771-781.
- 454 Axelsson E, Axelsson B. 1986. Changes in carbon allocation patterns in spruce and pine trees
 455 following irrigation and fertilization. Tree Physiology 2:189-204.
- 456 Belote RT, Prisley S, Jones RH, Fitzpatrick M, de Beurs K. 2011. Forest productivity and tree
- 457 diversity relationships depend on ecological context within mid-Atlantic and Appalachian
- 458 forests (USA). Forest Ecology and Management 261: 1315–1324.
- 459 Bertness M, Callaway R.M. 1994. Positive interactions in communities. Trends in Ecology460 and Evolution 9:191-193.
- 461 Boisvenue C, Running SW. 2006. Impacts of climate change on natural forest productivity–
 462 evidence since the middle of the 20th century. Global Change Biology 12:862-882.
- 463 Bonal D, Sabatier D, Montpied P, Tremeaux D, Guehl JM. 2000. Interspecific variability of
- 464 δ¹³C among trees in rainforests of French Guiana: functional groups and canopy integration.
 465 Oecologia 124:454-468.
- Bradford JB. 2011. Divergence in forest-type response to climate and weather: evidence for
 regional links between forest-type evenness and net primary productivity. Ecosystems 14:
 975-986.
- 469 Dulamsuren C, Hauck M, Khishigjargal M, Leuschner H, Leuschner C. 2010. Diverging
 470 climate trends in Mongolian taiga forests influence growth and regeneration of *Larix sibirica*.
- 471 Oecologia 163:1091-1102.
- 472 Fargione J, Tilman D, Dybzinski R, Lambers JHR, Clark C, Harpole WS, Knops JMH, Reich
- 473 PB, Loreau M. 2007. From selection to complementarity: shifts in the causes of biodiversity-
- 474 productivity relationships in a long-term biodiversity experiment. Proceedings of the Royal
- 475 Society B: Biological Sciences 274: 871-876.

- 476 Farquhar GD, O'Leary MH, Berry JA. 1982. On the relationship between carbon isotope
 477 discrimination and the intercellular carbon dioxide concentration in leaves. Australian Journal
 478 of Plant Physiology 9:121-137.
- Flanagan LB, Johnsen KH. 1995. Genetic variation in carbon isotope discrimination and its
 relationship to growth under field conditions in full-sib families of Picea mariana. Canadian
 Journal of Forest Research 25:39-47.
- 482 Forrester DI, Theiveyanathan S, Collopy JJ, Marcar NE. 2010. Enhanced water use efficiency
 483 in a mixed *Eucalyptus globules* and *Acacia mearnsii* plantation. Forest Ecology and
 484 Management 259:1761-1770.
- 485 Gamfeldt L, Snäll T, Bagchi R, Jonsson M, Gustafsson L, Kjellander P, Ruiz-Jaen MC,
- 486 Fröberg M, Stendahl J, Philipson CD, Mikusinski G, Andersson A, Westerlund B, Andren H,
- 487 Moberg F, Moen J, Bengtsson J. 2013. Higher levels of multiple ecosystem services are found488 in forests with more tree species. Nature Communications 4:1340.
- 489 Gebauer T, Horna V, Leuschner C. 2012. Canopy transpiration of pure and mixed forest490 stands with variable abundance of European beech. Journal of Hydrology 442-443:2-14.
- 491 Granier A, Bréda N, Biron P, Villette S. 1999. A lumped water balance model to evaluate
- 492 duration and intensity of drought constraints in forest stands. Ecological Modelling 116:269-493 283.
- Grossiord C, Granier A, Gessler A, Pollastrini M, Bonal D. 2013a. The influence of tree
 species mixture on ecosystem-level carbon accumulation and water use in a mixed boreal
 plantation. Forest Ecology and Management 298:82-92.
- 497 Grossiord C, Granier A, Gessler A, Scherer-Lorenzen M, Pollastrini M, Bonal D. 2013b.
- 498 Application of Loreau & Hector's (2001) partitioning method to complex functional traits.
- 499 Methods in Ecology and Evolution. In press

- He Q, Bertness MD, Altieri AH. 2013. Global shifts towards positive species interactions with
 increasing environmental stress. Ecology Letters 16:695-706.
- Herbert DA, Rastetter EB, Gough L, Shaver GR. 2004. Species diversity across nutrient
 gradients: An analysis of resource competition in model ecosystems. Ecosystems 7:296-310.
- Hooper DU, Dukes JS. 2004. Overyielding among plant functional groups in a long-term
 experiment. Ecology Letters 7:95-105.
- Hughes AR, Stachowicz JJ. 2004. Genetic diversity enhances the resistance of a seagrass
 ecosystem to disturbance. Proceedings of the National Academy of Sciences of the United
 States of America 101:8998-9002.
- 509 IPCC. 2007. Climate change 2007: Synthesis report. Pachauri RK, Reisinger A, editors.
- 510 Contribution of Working Groups I, II and III to the Fourth Assessment Report of the
- 511 Intergovernmental Panel on Climate Change. Cambridge University Press, Cambridge, p104.
- 512 Isbell FI, Polley HW, Wilsey BJ. 2009. Biodiversity, productivity and the temporal stability of
- 513 productivity: patterns and processes. Ecology Letters 12:443–451.
- 514 Jylhä K, Tuomenvirta H, Ruosteenoja K, Niemi-Hugaerts H, Keisu K, Karhu JA. 2010.
- 515 Observed and projected future shifts of climatic zones in Europe and their use to visualize
- 516 climate change information. Weather Climate and Society 2:148-167.
- 517 Jucker T, Coomes DA. 2013. Comment on "Plant Species Richness and Ecosystem
 518 Multifunctionality in Global Drylands". Science 337:155-155.
- 519 Kalliokoski T, Nygren P, Sievänen R. 2008. Coarse root architecture of three boreal tree
 520 species growing in mixed stands. Silva Fennica 42:189–210.
- 521 Kelty MJ. 2006. The role of species mixtures in plantation forestry. Forest Ecology and522 Management 233: 195-204.

- Kljun N, Black TA, Griffis TJ, Barr AG, Gaumont-Guay D, Morgenstern K, McCaughey JH,
 Nesic Z. 2007. Response of net ecosystem productivity of three boreal forest stands to
 drought. Ecosystems 10:1039-1055.
- Kozlowski TT, Pallardy SG. 2002. Acclimation and adaptive responses of woody plants toenvironmental stresses. The Botanical Review 68:270-334.
- 528 Kunert N, Schwendenmann L, Potvin C, Hölscher D. 2012. Tree diversity enhances tree 529 transpiration in a Panamanian forest plantation. Journal of Applied Ecology 49:135-144.
- 530 Lloyd J, Farquhar GD. 1994. ¹³C discrimination during CO₂ assimilation by the terrestrial
 531 biosphere. Oecologia 99:201-215.
- 532 Loreau M, Hector A. 2001. Partitioning selection and complementarity in biodiversity533 experiments. Nature 412:72-76.
- Maestre FT, Callaway RM, Valladares F, Lortie CJ. 2009. Refining the stress-gradient
 hypothesis for competition and facilitation in plant communities. Journal of Ecology 97:199205.
- 537 Meinen C, Hertel D, Leuschner C. 2009. Root growth and recovery in temperate broad-leaved
 538 forest stands differing in tree species diversity. Ecosystems 12:1103-1116.
- 539 Mölder I, Leuschner C, Leuschner HH. 2011. δ^{13} C signature of tree rings and radial increment
- 540 of *Fagus sylvatica* as dependent on tree neighbourhood and climate. Trees 25:215-229.
- 541 Morin X, Fahse L, Scherer-Lorenzen M, Bugmann H. 2011. Tree species richness promotes
- 542 productivity in temperate forests through strong complementary between species. Ecology
- 543 Letters 14:1211-1219.
- Naeem S. 2008. Species redundancy and ecosystem reliability. Conservation Biology 12:39-45.

- 546 Peñuelas J, Hunt JM, Ogaya R, Jump AS. 2008. Twentieth century changes of tree-ring δ^{13} C at
- 547 the southern range-edge of *Fagus sylvatica*: increasing water-use efficiency does not avoid the
- 548 growth decline induced by warming at low altitudes. Global Change Biology 14:1076–1088.
- 549 R Development Core Team. 2011. R: a language and environment for statistical computing.
- 550 Vienna, Austria: R Foundation for Statistical Computing.
- 551 Saurer M, Siegwolf RT, Schweingruber FH. 2004. Carbon isotope discrimination indicates
- improving water-use efficiency of trees in northern Eurasia over the last 100 years. GlobalChange Biology 10:2109-2120.
- 554 Steudel B, Hector A, Friedl T, Lofke C, Lorenz M, Wesche M, Kessler M. 2012. Biodiversity
- 555 effects on ecosystem functioning change along environmental stress gradients. Ecology
- 556 Letters 15:1397–1405.
- 557 Symstad AJ, Chapin FS, Wall DH, Gross KL, Huenneke LF, Mittelbach GG, Peters DPC,
- 558 Tilman D. 2003. Long-term and large-scale perspectives on the relationship between559 biodiversity and ecosystem functioning. BioScience 53:89-98.
- 560 Vaganov EA, Schulze ED, Skomarkova MV, Knohl A, Brand WA, Roscher C. 2009. Intra-
- annual variability of anatomical structure and δ^{13} C values within tree rings of spruce and pine
- in alpine, temperate and boreal Europe. Oecologia 161:729-745.
- Vilà M, Sardans J. 1999. Plant competition in mediterranean-type vegetation. Journal of
 Vegetal Sciences 10:281-294.
- 565 Vilà M, Carrillo-Gavilán A, Vayreda J, Bugmann H, Fridman J, Grodzki W, Haase J, Kunstler
- 566 G, Schelhaas M, Trasobares A. 2013. Disentangling biodiversity and climatic determinants of
- 567 wood production. PLoS ONE 8: e53530.
- 568 Wang J, Zhang CB, Chen T, Li WH. 2013. From selection to complementarity: the shift along
- the abiotic stress gradient in a controlled biodiversity experiment. Oecologia 171:227-235.

- 570 Zhang J, Marshall JD. 1994. Population differences in water-use efficiency of well-watered
- and water-stressed western larch seedlings. Canadian Journal of Forest Research 24:92-99.

Zhang Y, Chen HYH, Reich PB. 2012. Forest productivity increases with evenness, species richness and trait variation: a global meta-analysis. Journal of Ecology 100:742-749.

Figure 1











579 Figure legends:

Figure 1: Time-course of relative extractable water (REW) for the years 1997 to 2012
simulated with the "BILJOU" model. The two selected years are highlighted in grey (wet for
2004 and dry for 2006).

583

Figure 2: Stand carbon isotope composition ($\delta^{13}C_s$, ‰), stand size-standardized basal area increment (BAI_s, cm² cm⁻² yr⁻¹) for the wet, the dry and the recovery year (DRY+1, white circles), and the differences in $\delta^{13}C_s$ and BAI_s between the wet and the dry year ($\Delta\delta^{13}C_s$ and Δ BAI_s, respectively) as a function of the Shannon biodiversity index. The black lines represent the fitted regression model when the model was significant (R² = 0.22 and 0.23 for $\delta^{13}C_s$ and $\Delta\delta^{13}C_s$, respectively).

590

Figure 3: Net, complementarity, and selection effects on stand carbon isotope composition ($\delta^{13}C_s$, ‰) for the dry year as a function of species richness. Black circles indicate birch and pine mixtures, white squares indicate birch and spruce mixtures, black diamonds indicate pine and spruce mixtures, and black triangles indicate 3-species mixtures. Asterisks denote significant differences from zero in the biodiversity effects for each species richness level (ttest, * *P* < 0.05; ** *P* < 0.01).

597

Table 1: Overall mean species diameter at breast height (cm) and mean height (m) with600 standard errors for *Betula pendula*, *Pinus sylvestris* and *Picea abies*.

Species	Mean diameter at breast height (cm)	Mean height (m)	
Betula pendula	16.4 ± 0.2	17.9 ± 0.2	
Pinus sylvestris	19.2 ± 0.1	17.6 ± 0.1	
Picea abies	17.7 ± 0.2	16.1 ± 0.1	

604 **Table 2:** Mean species-specific Shannon biodiversity index, wood carbon isotope 605 composition ($\delta^{13}C_i$, ∞), and size-standardized basal area increment (BAI_i, cm² cm⁻² yr⁻¹) for 606 the dry and wet years, with standard errors for *Betula pendula*, *Pinus sylvestris* and *Picea* 607 *abies*, and for each mixture level.

608

Species	Mixture	Shannon	δ ¹³ C _i Wet (‰)	δ ¹³ C _i Dry (‰)	BAI _i Wet (cm ² cm ⁻² yr ⁻¹)	BAI _i Dry (cm² cm²² yr¹)
	1	0.2 ± 0.1	-27.1 ± 0.3	-25.6 ± 0.1	$17.1 \times 10^{-3} \pm 0.8 \times 10^{-3}$	$18.2 \times 10^{-3} \pm 0.9 \times 10^{-3}$
Betula pendula	2	0.9 ± 0.1	-27.8 ± 0.2	-25.8 ± 0.3	$19.2 \times 10^{-3} \pm 0.9 \times 10^{-3}$	$19.2 \times 10^{-3} \pm 0.9 \times 10^{-5}$
1	3	1.4 ± 0.1	-27.8 ± 0.2	-25.9 ± 0.5	$21.3 \times 10^{-3} \pm 1.5 \times 10^{-3}$	$19.8 \times 10^{-3} \pm 1.4 \times 10^{-3}$
	1	0.1 ± 0.1	-27.3 ± 0.2	-26.2 ± 0.2	$11.9 \times 10^{-3} \pm 0.7 \times 10^{-3}$	$11.4 \times 10^{-3} \pm 0.7 \times 10^{-3}$
Pinus sylvestris	2	0.9 ± 0.0	-27.2 ± 0.1	-25.7 ± 0.3	$16.4 \times 10^{-3} \pm 0.9 \times 10^{-3}$	$15.1 \times 10^{-3} \pm 0.7 \times 10^{-3}$
5	3	1.4 ± 0.1	-26.9 ± 0.2	-25.4 ± 0.6	$20.6 \times 10^{-3} \pm 1.2 \times 10^{-3}$	$16.5 \times 10^{-3} \pm 0.8 \times 10^{-3}$
	1	0.1 ± 0.1	-26.9 ± 0.1	-26.2 ± 0.2	22.9×10 ⁻³ ± 1.0×10 ⁻³	$17.9 \times 10^{-3} \pm 0.8 \times 10^{-3}$
Picea abies	2	0.9 ± 0.0	-27.2 ± 0.2	-25.4 ± 0.3	$21.3 \times 10^{-3} \pm 1.3 \times 10^{-3}$	$18.4 \times 10^{-3} \pm 1.1 \times 10^{-3}$
	3	1.4 ± 0.1	-27.2 ± 0.2	-25.3 ± 0.2	$26.6 \times 10^{-3} \pm 1.5 \times 10^{-3}$	$21.9 \times 10^{-3} \pm 1.3 \times 10^{-3}$