



HAL
open science

Uptake of water via branches helps timberline conifers refill embolized xylem in late winter

Stefan Mayr, Peter Schmid, Joan Laur, Sabine Rosner, Katline
Charra-Vaskou, Birgit Daemon, Uwe G. Hacke

► **To cite this version:**

Stefan Mayr, Peter Schmid, Joan Laur, Sabine Rosner, Katline Charra-Vaskou, et al.. Uptake of water via branches helps timberline conifers refill embolized xylem in late winter. *Plant Physiology*, 2014, 164 (4), pp.1731-1740. 10.1104/pp.114.236646 . hal-01268574

HAL Id: hal-01268574

<https://hal.science/hal-01268574>

Submitted on 27 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.




Fig. S1. Embolism and water potential in different sections of branches during late winter. Percent loss of conductivity (PLC) in stem xylem and water potential (Ψ) of basal, and distal stem sections of branches. Mean \pm SE.