

# Tree diversity does not always improve resistance of forest ecosystems to drought

Damien Bonal, Charlotte Grossiord, André Granier, Arthur Gessler

#### ▶ To cite this version:

Damien Bonal, Charlotte Grossiord, André Granier, Arthur Gessler. Tree diversity does not always improve resistance of forest ecosystems to drought. Reunion annuelle du Labex ARBRE, Oct 2014, Nancy, France. 15 p. hal-01268786

HAL Id: hal-01268786

https://hal.science/hal-01268786

Submitted on 5 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.





# Tree diversity does not always improve resistance of forest ecosystems to drought

Damien BONAL Charlotte GROSSIORD Arthur GESSLER André GRANIER





















Ecotourism

Increasing species diversity can lead to important benefits for ecosystems



#### www.cedarcreek.umn.edu


#### Biomass Production



Loreau et al. 2001

Higher biomass production in more diverse ecosystems


Increasing species diversity can lead to important benefits for ecosystems



www.worldofstock.com





www.bbc.co.uk



Lower foliar pathogen load in more diverse ecosystems

Are more diverse ecosystems more resistant to drought?





Increasing drought resistance in more diverse communities

## Objectives of the study

Test whether higher tree species diversity in forest ecosystems across Europe is associated with less drought exposure



Grossiord C.




www.nature.com

# **Study Sites**




6 regions across North - South gradient in Europe to cover the major forest types and climatic conditions


THERMOPHILOUS DECIDUOUS (Italy)





### Material and Methods

Comparison of ecosystem functioning for different levels of species mixture (1-5)










### Material and Methods

# Comparison of ecosystem functioning for different levels of species mixture (1-5)

- Selection of 25 to 42 stands (30×30m) varying in species richness in each studied region
- Extraction of wood cores from a selection of dominant and codominant trees of each species in each stand
- Selection of years (tree rings) with contrasting climatic conditions (dry vs. wet)
- Analyses of carbon isotope composition ( $\delta^{13}$ C) of latewood of each ring
- Calculation of plot-level  $\delta^{13}$ C

### Material and Methods

• The difference in plot-level  $\delta^{13}C$  between the DRY and the WET year can be used as a proxy of drought exposure



If difference in  $\delta^{13}$ C is high, low resistance

If difference in  $\delta^{13}$ C is low, high resistance



# Tree diversity does not always improve resistance of forest ecosystems to drought

Charlotte Grossiord<sup>a</sup>, André Granier<sup>a</sup>, Sophia Ratcliffe<sup>b</sup>, Olivier Bouriaud<sup>c</sup>, Helge Bruelheide<sup>d,e</sup>, Ewa Chećko<sup>f</sup>, David Ian Forrester<sup>g</sup>, Seid Muhie Dawud<sup>h</sup>, Leena Finér<sup>i</sup>, Martina Pollastrini<sup>j</sup>, Michael Scherer-Lorenzen<sup>k</sup>, Fernando Valladares<sup>l</sup>, Damien Bonal<sup>a, 1, 2</sup>, and Arthur Gessler<sup>m,n, 2</sup>

<sup>a</sup>Unité Mixte de Recherche 1137, Ecologie et Ecophysiologie Forestiéres, Institut National de la Recherche Agronomique, 54280 Champenoux, France; <sup>b</sup>Institut für Spezielle Botanik und Funktionelle Biodiversität, University Leipzig, 04103 Leipzig, Germany; <sup>c</sup>Forestry Faculty, University of Suceava, 720229 Suceava, Romania; <sup>d</sup>Institute of Biology/Geobotany and Botanical Garden, Martin Luther University Halle Wittenberg, 06108 Halle, Germany; <sup>e</sup>German Centre for Integrative Biodiversity Research, 04103 Leipzig, Germany; <sup>f</sup>Faculty of Biology, Bialowieza Geobotanical Station, University of Warsaw, 17-230 Bialowieza, Poland; <sup>g</sup>Faculty of Environment and Natural Resources, Freiburg University, 79108 Freiburg, Germany; <sup>h</sup>Department of Geosciences and Natural Resource Management, University of Copenhagen, 1958 Copenhagen, Denmark; <sup>i</sup>Finnish Forest Research Institute, 80101 Joensuu, Finland; <sup>j</sup>Department of Agri-Food and Environmental Sciences, Section of Soil and Plant Science, University of Florence, 50144 Florence, Italy; <sup>k</sup>Faculty of Biology/Geobotany, University Freiburg, 79104 Freiburg, Germany; <sup>i</sup>Museo Nacional de Ciencias Naturales, Consejo Superior de Investigaciones Cientificas, E-28006 Madrid, Spain; <sup>m</sup>Long-Term Forest Ecosystem Research, Swiss Federal Institute for Forest, Snow and Landscape Research, 8903 Birmensdorf, Switzerland; and <sup>n</sup>Berlin-Brandenburg Institute of Advanced Biodiversity Research, 14195 Berlin, Germany

#### Results No effect of species Lower drought resistance diversity in species-rich forests Boreal = 0.69x + 1.04 Hemi-boreal P = 0.159Increasing drought $R^2 = 0.23$ P = 0.038Increase in stand-level $\delta^{13} C$ between a wet and a dry year (%) Increasing drought y = -0.40x + 1.44 $R^2 = 0.25$ P= 0.297 Temperate beech Mountainous beech 3 P = 0.003exposure 2 Thermophilous deciduous y = -0.50x + 1.70Mediterranean P = 0.716Increasing drought $R^2 = 0.20$ P = 0.007exposure

1.5

2.0

2.5 0.0

ъиannon diversity inaex

0.5

1.0

1.5

**Increasing diversity** 

2.0


2.5

Higher drought resistance in species-rich forests

0.0

#### Results

Tree species diversity effects seem partially dependent on **climatic conditions** 



Frequent events and high intensity of drought

Positive effects

### **Conclusions**

- Tree species diversity influences the functioning of forest ecosystems in terms of water and carbon acquisition and use
- More diverse forests may be more resistant to soil drought than pure ones, but not necessarily
- Favoring mixed forest ecosystems could be preferred in certain regions, not only to preserve biodiversity or sustain productivity in the context of climate changes, but also to delay the negative effect of drought

### Thanks for your attention







