



HAL
open science

Linking litter quality to microbial community structure and functioning in an agricultural soil

Marie Sauvadet, Mathieu Chauvat, Daniel Cluzeau, Pierre-Alain Maron,
Isabelle Bertrand

► To cite this version:

Marie Sauvadet, Mathieu Chauvat, Daniel Cluzeau, Pierre-Alain Maron, Isabelle Bertrand. Linking litter quality to microbial community structure and functioning in an agricultural soil. 1. Global Soil Biodiversity Initiative Conference (GSBI), Institut National de la Recherche Agronomique (INRA). FRA., Dec 2014, Dijon, France. 703 p. hal-01269063

HAL Id: hal-01269063

<https://hal.science/hal-01269063v1>

Submitted on 2 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

THE FIRST GLOBAL
SOIL BIODIVERSITY CONFERENCE

Assessing soil biodiversity and role in ecosystem services



GLOBAL
SOIL BIODIVERSITY
INITIATIVE

Palais des Congrès, Dijon, France

2-5 DECEMBER 2014

THE FIRST GLOBAL
SOIL BIODIVERSITY
CONFERENCE

December 2-5, 2014, Dijon, France

PROGRAM AND ABSTRACTS

THE FIRST GLOBAL **SOIL BIODIVERSITY** **CONFERENCE**



GLOBAL
SOIL BIODIVERSITY
INITIATIVE

December 2-5, 2014

Dijon – France

[P1.155]

Linking litter quality to microbial community structure and functioning in an agricultural soil

M. Sauvadet*, M. Chauvat, D. Cluzeau, P-A. Maron, I. Bertrand

¹*Ecodiv EA 1293 Université de Rouen, France*, ²*UMR 6553 Ecobio, CNRS, Rennes, France*,

³*UMR 1347 Agroécologie, INRA, Dijon, France*, ⁴*UMR 614 FARE, INra, Reims, France*

Anthropogenic pressures on agricultural soils are known to alter their biodiversity, which may affect the capacity of ecosystems to deliver functions and services according to the insurance hypothesis on diversity – function relationship. Cropping systems is fuelling belowground biota through the quality of litter supplied, which varies in quality from year to year. Therefore we hypothesized that soil biota structure and functioning strongly respond to crop residues quality. Aerial and underground parts of maize plants were used as labile and recalcitrant litter, respectively. A dynamic experiment was performed using soil columns filled with a silty loam agricultural soil (Estrées-Mons, Northern France) in which we incorporated in the 0-5cm layer either maize leaves or maize roots (and a control treatment without litter). We measured regularly soil respiration, litter quality, enzyme activities, microbial biomass, bacterial and fungal community structure (using 16S and 18S pyrosequencing) during the incubation period.

Litter quality strongly influenced degradation rates of litter soluble fraction and polysaccharides, leading to faster C mineralization of maize leaves compared to roots. Enzyme efficiency was higher in the recalcitrant litter demonstrating different microbial strategies. After 15 days of decomposition, leaves stimulated the relative proportion of Proteobacteria and Bacteroidetes (classified as copiotrophic bacteria by Fierer et al. 2007). Similarly in the fungal community, Tremellomycetes and Sordariomycetes exhibited the most copiotrophic behavior. General functions such as soluble compounds mineralization rates were better correlated with microbial biomass ($R^2=0.65$) than with community structure. Nevertheless, more specific functions such as complex C forms degradation were related to bacterial community structure. We concluded that the decomposition of complex C forms, having a high energetic cost for microorganisms, strongly impacts the community structure. Thus, ecosystem functioning depends on biota community structure linked to the resources provided to them.

Keywords: Litter quality, Microbial communities, Enzymatic activities, pyrosequencing