



HAL
open science

Genetic variability for susceptibility to the invasive ascomycete *Hymenoscyphus fraxineus* in a common Ash (*Fraxinus excelsior*) provenance and progeny trial indicates significant potential for breeding

Arnaud Dowkiw, Facundo Munoz, Benoit Marçais, Fabrice Bonne, Vincent Rousselet, Jean Dufour

► **To cite this version:**

Arnaud Dowkiw, Facundo Munoz, Benoit Marçais, Fabrice Bonne, Vincent Rousselet, et al.. Genetic variability for susceptibility to the invasive ascomycete *Hymenoscyphus fraxineus* in a common Ash (*Fraxinus excelsior*) provenance and progeny trial indicates significant potential for breeding. 5. International Workshop on the Genetics of Tree-Parasite Interactions, Aug 2015, Orléans, France. , 2015, 5th International Workshop on the Genetics of Tree-Parasite Interactions. Book of abstracts. hal-01269238

HAL Id: hal-01269238

<https://hal.science/hal-01269238>

Submitted on 3 Jun 2020


HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

5th International Workshop on the
Genetics of Tree-Parasite Interactions
Orléans, France, 23-28 August 2015



Book of Abstracts



An event sponsored by :



Genetic variability for susceptibility to the invasive ascomycete *Hymenoscyphus fraxineus* in a common Ash (*Fraxinus excelsior*) provenance and progeny trial indicates significant potential for breeding

Arnaud DOWKIW¹, MUNOZ F¹, MARÇAIS B², BONNE F³, ROUSSELET V³, DUFOUR J¹

¹ Institut National de la Recherche Agronomique, UR AGPF 0588, 2163 avenue de la Pomme de Pin, ARDON CS4001, France.

² INRA, Université de Lorraine, UMR 1136 Interactions Arbres – Microorganismes (IAM), F–54280 Champenoux, France

³ Institut National de la Recherche Agronomique, UE UEFL 1261, 54280 CHAMPENOUX, France

Severe dieback of European common ash (*Fraxinus excelsior*) was first reported in Poland and Lithuania in the early 1990s. It was fourteen years after the first reports that an ascomycete, now referred to as *Hymenoscyphus fraxineus* (Queloz et al. 2011), was finally identified as the primary causal agent (Kowalski 2006). Several findings pledge for the species being of Asian origin and acting as an invasive pathogen in Europe (Husson et al. 2011, McKinney et al. 2012, Bengtsson et al. 2012, Zhao et al. 2013). Detailed quantifications are scarce, but consequences of the disease can be severe. In Lithuania, 10 years after the first report, over 30,000 ha of common ash stands were reported to be affected and mortality was estimated to be 60% state wide while in some parts of the country only about 2% of the trees remained visually healthy (Juodvakis and Vasiliauskas 2002). First detection of the disease in France occurred in 2008, in the North-Eastern part of the country. Since then, the disease has been progressing southwestwards at a pace of approximately 50 km per year. The fungus now occupies half of the country. Eventhough millions of *Fraxinus excelsior* trees died from the disease, recent results from clonal or progeny trials raised hopes that resistant/tolerant trees could be selected. Significant heritability estimates and provenance and/or family effects seem to indicate that breeding programs could be implemented against the disease (Pliura et al. 2011, Kjaer et al. 2012). The present study is only third to analyze the genetic variability of common ash for resistance/tolerance to *H. fraxineus* using open-pollinated progenies. It is based on a 20-year old field trial with 23 half-sib families from 3 French provenances. The trial is located very close to the place where the disease was first detected in France. Because the stand has been monitored every year since 2010 for disease incidence, spatio-temporal components of disease spread could be taken into account to avoid confounding disease escape and resistance. Contrary to previous studies, a Bayesian approach was used instead of the classical frequentist analysis. Bayesian methods accommodate spatio-temporal structures in a more straightforward way. They are also more suitable for modelling data which deviates from normality as commonly observed during early stages of disease spread. This study is also the first to report on genetic parameters for collar lesions and on the genetic correlation with crown defoliation. Collar lesions due to *H. fraxineus* have been described in France only recently (Husson et al. 2012) and this symptom has certainly been overlooked in countries with a longer infection history.

References

- Queloz, V. et al. (2011). Cryptic speciation in *Hymenoscyphus albidus*. *Forest Pathology*41(2): 133-142.
- Husson, C., et al. (2011). *Chalara fraxinea* is an invasive pathogen in France. *European Journal of Plant Pathology* 130(3): 311-324.
- McKinney, L. V., et al. (2012). Rapid invasion by an aggressive pathogenic fungus (*Hymenoscyphus pseudoalbidus*) replaces a native decomposer (*Hymenoscyphus albidus*): a case of local cryptic extinction? *Fungal Ecology*5(6): 663-669.
- Bengtsson, S. B. K., et al. (2012). Population structure of *Hymenoscyphus pseudoalbidus* and its genetic relationship to *Hymenoscyphus albidus*. *Fungal Ecology*5(2): 147-153.
- Zhao, Y. J., et al. (2013). *Hymenoscyphus pseudoalbidus*, the correct name for *Lambertella albida* reported from Japan. *Mycotaxon*122(1): 25-41.
- Juodvakis, A. and A. Vasiliauskas (2002). Drying extent of Lithuanian ash-tree woods and factors predetermining it. *Vagos* 56: 17-22.
- Pliura, A., et al. (2011). Performance of Twenty Four European *Fraxinus excelsior* Populations in Three Lithuanian Progeny Trials with a Special Emphasis on Resistance to *Chalara Fraxinea*. *Baltic Forestry*17(1): 17-34.
- Kjaer, E. D., et al. (2012). Adaptive potential of ash (*Fraxinus excelsior*) populations against the novel emerging pathogen *Hymenoscyphus pseudoalbidus*. *Evolutionary Applications*5(3): 219-228.
- Husson, C., et al. (2012). Occurrence of *Hymenoscyphus pseudoalbidus* on infected ash logs. *Plant Pathology* 61(5): 889-895.

Keywords

Fraxinus excelsior, European Ash dieback, resistance, tolerance, breeding, progeny trial, genetic variability.