



HAL
open science

Metabolic genome-wide-association unravels trait architecture and natural variation in tomato primary and secondary metabolisms

Guillaume Bauchet, Stéphane Grenier, Nicolas Samson, Vincent Segura, Aniko Kende, Jules Beekwilder, Katarina Cankar, Arnaud G. Bovy, Jean-Luc Gallois, Julien Bonnet, et al.

► To cite this version:

Guillaume Bauchet, Stéphane Grenier, Nicolas Samson, Vincent Segura, Aniko Kende, et al.. Metabolic genome-wide-association unravels trait architecture and natural variation in tomato primary and secondary metabolisms. 12. Solanaceae Conference, Oct 2015, Bordeaux, France. hal-01269376

HAL Id: hal-01269376


<https://hal.science/hal-01269376>

Submitted on 5 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

SOL2015



The 12th Solanaceae Conference

October 25 -29, 2015
ENSEIRB Building, Talence

Bordeaux, France



Metabolic genome-wide-association unravels trait architecture and natural variation in tomato primary and secondary metabolisms

Guillaume Bauchet^{1,2, *}, Stéphane Grenier², Nicolas Samson², Vincent Segura³, Aniko Kende⁴, Jules Beekwilder⁵, Katarina Cankar⁵, Arnaud Bovy⁵, Jean-Luc Gallois¹, Julien Bonnet², Laurent Grivet², Elek Bolygo³, Charles Baxter³ and Mathilde Causse¹

1 INRA –Domaine St Maurice - UR 1052 - GAFL - CS 60094 84143 Montfavet, France ;

2 Syngenta Seeds 12 Chemin Hobit, 31790 Saint-Sauveur, France ;

3 Institut National de la Recherche Agronomique, UR0588, 45075 Orleans , France;

4 Syngenta Jealott Hill International Research Centre Bracknell, Berkshire RG42 6EY, UK;

5 Plant Research International, 6700 AA Wageningen, the Netherlands;

** present address: Boyce Thompson Institute, 533 Tower road, NY14850, Ithaca, USA;*

Modern breeding has impacted fruit quality and volatile content, modifying the organoleptic quality of tomato crop varieties. To decipher quality trait inheritance and improve it in modern tomato breeding programs, we characterized an association panel including 70 wild relative accessions (*S. pimpinellifolium*), 170 admixed accessions (*S.l. cerasiforme*) and 50 elite cultivars (*S. lycopersicum*). GWAS analysis was conducted using 12,000 SNP markers and a set of primary metabolism traits (sugars, organic and amino acids) as well as a broad range of volatile compounds. This study is the first in tomato reporting associations for a large set of volatiles at the genome scale. Metabolite content and genetic inheritance varied in a broad range over different genetic groups. Four genomic regions were detected highlighting clusters of associations for several metabolites. We found significant associations for 84 loci with a total of 17 traits including glucose, malate or phenylacetaldehyde. Some associations were consistent with previously published quantitative trait loci (sugars, guaiacol), while new loci were identified (phenylacetaldehyde). Local linkage disequilibrium analysis and allele mining allowed the identification of candidate genes to be functionally investigated. These results provide (1) a list of candidate loci and (2) a fast and efficient analytical approach for finding genetic variants that can be directly used for fruit quality improvement and deciphering the genetic architecture of complex traits.