



HAL
open science

Are cotyledonary somatic embryos of *Pinus pinaster* Ait. equivalent to the zygotic counterpart ?

Caroline Teyssier, Alexandre Morel, Jean-François Trontin, Françoise Corbineau, Anne-Marie Lomenech, Martine Beaufour, Isabelle Reymond, Claire Le Mette, Kévin Ader, Luc Harvengt, et al.

► To cite this version:

Caroline Teyssier, Alexandre Morel, Jean-François Trontin, Françoise Corbineau, Anne-Marie Lomenech, et al.. Are cotyledonary somatic embryos of *Pinus pinaster* Ait. equivalent to the zygotic counterpart ?. 3. International Conference of the IUFRO Working Party 2.09.02: Somatic Embryogenesis and Other Vegetative Propagation Technologies, Sep 2014, Vitoria-Gasteiz, Spain. 2014, 3rd International Conference of the IUFRO Working Party 2.09.02: Somatic Embryogenesis and Other Vegetative Propagation Technologies. Book of Abstracts. hal-01269493

HAL Id: hal-01269493

<https://hal.science/hal-01269493v1>

Submitted on 3 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



3rd International
Conference of the

IUFRO

Working Party
2.09.02

Somatic Embryogenesis and
Other Vegetative Propagation Technologies

BOOK OF ABSTRACTS

Woody Plant Production Integrating Genetic
and Vegetative Propagation Technologies

September 8 - 12, 2014. Vitoria-Gasteiz, Spain

EUSKO JAURLARITZA



GOBIERNO VASCO

EKONOMIAREN GARAPEN
ETA LEHIAKORTASUN SAILA

DEPARTAMENTO DE DESARROLLO
ECONÓMICO Y COMPETITIVIDAD

Eusko Jaurlaritzaren Argitalpen Zerbitzu Nagusia

Servicio Central de Publicaciones del Gobierno Vasco

Vitoria-Gasteiz, 2014

First published: July 2014

Edition: 140 copies

© Administration of the Autonomous Community of the Basque Country. Department of Economic Development and Competitiveness

Internet: www.euskadi.net

Publisher: Eusko Jaurlaritzaren Argitalpen Zerbitzua • Servicio Central de Publicaciones del Gobierno Vasco (Donostia-San Sebastián, 1 - 01010 Vitoria-Gasteiz)

Design and layout by: Alex Fernández Morán

Printed by: Gráficas Juaristi, S.L.

Legal Record: SS 1.000-2014

COMMITTEES

Organizing Committee

Yill-Sung Park, Natural Resources Canada-Canadian Forest Service, Fredericton, New Brunswick, Canada. yillsung.park@nrca-nrcan.gc.ca

Paloma Moncaleán, NEIKER-TECNALIA, Centro de Arkaute. 01080 Vitoria-Gasteiz. Spain. pmoncalean@neiker.net

Mariano Toribio, Instituto Madrileño de Investigación y Desarrollo Rural, Agrario y Alimentario (IMIDRA), Alcalá de Henares (Madrid), Spain. mariano.toribio@madrid.org

Jean-Francois Trontin, FCBA Technological Institute, Biotechnology and Advanced Forestry Department, Biotechnology Lab, Bordeaux, France. jean-francois.trontin@fcba.fr

Heung-Kyu Moon, Korea Forest Research Institute (KFRI), Department of Forest Genetic Resources, Suwon, Korea. mhkmoon@korea.kr

Jana Krajnakova, Mendel University in Brno, Faculty of Forestry and Wood Technology, Brno, Czech Republic. jana.krajnakova@uniud.it

Conference Executive Committee and Local Arrangements (NEIKER-Tecnalia)

Paloma Moncaleán

Itziar Montalbán

Iranzu Telletxea

Alex Fernández

Olatz Garcia-Mendiguren

Scientific Committee

Antonio Ballester (Spain)

Jorge Canhoto (Portugal)

Pramod Gupta (USA)

Cathy Hargreaves (New Zealand)

Marie-Anne Lelu-Walter (France)

Terezia Salaj (Slovak Republic)

Patrick von Aderkas (Canada)

Sara von Arnold (Sweden)

Tuija Aronen (Finland)

Carmen Diaz-Sala (Spain)

Yousry El-Kassaby (Canada)

Hely Hagman (Finland)

Krystyna Klimaszewska (Canada)

Scott Merkle (USA)

Heung-Kyu Moon (Korea)

Ana Vieitez (Spain)

Jan Bonga (Canada)



Are cotyledonary somatic embryos of *Pinus pinaster* Ait. equivalent to the zygotic counterpart?

Teyssier C.^{1*}, Morel A.¹, Trontin J-F.², Corbineau F.³, Lomenech A-M.⁴, Beaufour M.⁵, Reymond I.², Le Mett  C.¹, Ader K.¹, Harvengt L.², Cadene M.⁵, Label P.⁶, Lelu-Walter M-A.¹

¹INRA, UR 0588 UAGPF, 2163 Avenue de la Pomme de Pin, CS 4001, Ardon, 45075 Orl ans Cedex 2 (France).

²FCBA, P le BSA, 71 route d'Arcachon, 33610 Cestas (France).

³UPMC University Paris 06, UR5-EAC 7180 CNRS, Physiologie Cellulaire et Mol culaire des Plantes, Bo te Courrier 156, 4 place Jussieu, F-75005 Paris (France).

⁴Universit  de Bordeaux, Centre G nominique Fonctionnelle, Plateforme Prot ome, F-33000 Bordeaux, France.

⁵CNRS, UPR4301, Centre de Biophysique Mol culaire, Rue Charles Sadron, F-45100 Orl ans (France)

⁶INRA-Universit  Blaise Pascal, UMR_A 547 PIAF, 24 Avenue des Landais, 63171 Aubi re Cedex (France).

*Corresponding author: caroline.teyssier@orleans.inra.fr

In maritime pine, improved protocols are now available for the whole somatic embryogenesis process i.e. from initiation of embryogenic tissue to somatic plant regeneration. However, field trials established in France from somatic plant material have consistently revealed that somatic embryos (SEs) developed at a lower initial growth rate compared to control seedlings. A better understanding of SEs maturation is therefore required in order to produce high-quality, vigorous somatic plants. SEs are currently matured for 12 weeks to reach the cotyledonary stage before being germinated and converted to plantlets. Although regeneration success is highly dependent on SEs quality, the harvesting date is still determined from morphological features. This empirical method does not provide any accurate information about embryo quality with respect to storage compounds (proteins, carbohydrates). We first analyzed SEs matured for 10, 12 and 14 weeks by carrying out biological (dry weight, water content) and biochemical measurements (total protein and carbohydrate contents). No significant difference could be found between collection dates, suggesting that SE harvesting after 12 weeks maturation is appropriate. Cotyledonary SEs were then compared to various stages, from fresh to fully desiccated, in the development of cotyledonary zygotic embryos (ZEs). Using hierarchical ascendant cluster analysis, we demonstrated that cotyledonary SEs matured for 12 weeks are most similar to fresh cotyledonary ZEs sampled from late July to early August (as indicated by dry weight, water content, sucrose, RFOs content, RFO/Sucrose ratio) or at any time up to October (with respect to total protein). Both types of embryo exhibited similar carbohydrate and protein content and signatures. This high level of similarity (94.5%) was further supported by proteome profiling. Highly expressed proteins included storage, stress-related, late embryogenesis abundant (LEA) and energy metabolism proteins. By comparing overexpressed proteins in developing and cotyledonary SEs or ZEs, some (23 proteins) could be identified as candidate biomarkers for the late, cotyledonary stage. Of these, 18 belonged to five large families of proteins including five HSPs, four LEAs and two other stress-related proteins (aldose reductase, 6-phosphogluconate dehydrogenase), five storage proteins and two proteins involved in purine metabolism (adenosine kinase 2, SAM synthase).

This is the first report of useful generic protein markers for monitoring embryo development in maritime pine. Our results also suggest that improvements of SEs quality may be achieved if the current maturation conditions are refined.

Acknowledgements: This work was supported by grants from the French "Conseil R gional de la R gion Centre" (EMBRYOME project, contract 33639) and "Conseil R gional de la R gion Aquitaine" (EMBRYO2011 project, contract 09012579-045). Some equipment used at FCBA to get results presented in this work was funded by ANR-10-EQPX-16 XYLOFOREST. We gratefully acknowledge Gilles Bruneau, Francis Canlet, Sandrine Debille, Karine Durandeu, S verine Quoniou and Fran oise Wan-Fung (FCBA) for their valuable contribution to the collection of somatic and zygotic embryos and Pierre Alazard (FCBA) for cone sampling in the field. We acknowledge Marl ne Bailly for her technical assistance in electrophoresis.

Keywords: carbohydrates, embryo quality, maturation, proteomic, somatic embryogenesis, zygotic embryogenesis.

