



HAL
open science

An extension of the Tusnady inequality for general distributions

Ion Grama

► **To cite this version:**

Ion Grama. An extension of the Tusnady inequality for general distributions. The 14th Conference on Applied and Industrial Mathematics. (Satellite Conference of ICM 2006.) , The Romanian Society of Applied and Industrial Mathematics (ROMAI), Mathematical Society of the Republic of Moldova, Moldova State University, Tiraspol State University, Institute of Mathematics and Computer Science of the Academy of Sciences of Moldova and the Center for Education and Research in Mathematics and Computer Science at Moldova State University, Aug 2006, Chisinau, Moldova. pp.184-188. hal-01270357

HAL Id: hal-01270357

<https://hal.science/hal-01270357>

Submitted on 29 Jan 2017

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

An extension of the Tusnady inequality for general distributions

Ion Grama

Université de Bretagne Sud, France

`ion.grama@univ-ubs.fr`

Under mild assumptions an analog of Tusnady's inequality for conditional distributions of partial sums of independent nonidentically distributed r.v.'s is obtained. These type of inequalities provide very sharp bounds for couplings of random variables via quantile transformations, which are main tools for proving various strong invariance principles for sums of independent random variables. We refer to Komlós, Major and Tusnady (1975, 1976), Sakhanenko (1984), Zaitsev (1996), Grama, I. and Nussbaum, M. (2002), Carter and Pollard (2004) and to references therein for details. Recently quantile transformations have been applied for establishing a constructive asymptotic equivalence of statistical experiments, see Brown, Carter, Low and Zhang (2004).

Let (Ω, \mathcal{F}, P) be a probability space and λ be a real number satisfying $0 < \lambda < \infty$. Denote by $\mathfrak{D}(\lambda)$ the subset of all real valued random variables S on the probability space (Ω, \mathcal{F}, P) , which admits for some $n \geq 1$, a representation of the form $S = X_1 + \dots + X_n$, where X_1, \dots, X_n are independent r.v.'s, satisfying assumptions

A1 The r.v.'s X_1, \dots, X_n are of means zero and finite variances:

$$EX_i = 0, \quad 0 < EX_i^2 < \infty, \quad i = 1, \dots, n.$$

A2 The following Sakhanenko's condition holds true:

$$\lambda E |X_i|^3 \exp \{ \lambda |X_i| \} \leq EX_i^2, \quad i = 1, \dots, n.$$

Let $\mathbf{i} = \sqrt{-1}$ and ε, ν be real numbers, such that $0 < \varepsilon \leq 1/96$ and $0 < \nu < \infty$. Denote by $\mathfrak{D}_1(\lambda)$ the subset of all r.v.'s $S \in \mathfrak{D}(\lambda)$ which additionally satisfy the smoothness assumption

S1 The r.v. S is such that,

$$\sup_{|h| \leq \lambda} \int_{|t| > \varepsilon \lambda} \left| \frac{E \exp \{ (\mathbf{i}t + h)S \}}{E \exp \{ hS \}} \right| dt \leq \frac{\nu}{\lambda E S^2}.$$

Denote by $\mathfrak{D}_2(\lambda)$ the subset of all r.v.'s $S \in \mathfrak{D}(\lambda)$ which take discrete values $a + \Delta k$, $k \in \{0, \pm 1, \pm 2, \dots\}$ with maximal span Δ , where $0 \leq a < \Delta$ and $\lambda \Delta \leq \pi/\varepsilon$, and which additionally satisfy the assumption (S2)

S2 The r.v. S is such that,

$$\sup_{|h| \leq \lambda} \int_{\varepsilon \lambda < |t| \leq \pi/\Delta} \left| \frac{E \exp \{ (\mathbf{i}t + h)S \}}{E \exp \{ hS \}} \right| dt \leq \frac{\nu}{\lambda E S^2}.$$

Let us agree that \mathcal{X} is the range of a r.v. S in the class $\mathfrak{D}_1(\lambda)$ or $\mathfrak{D}_2(\lambda)$, i.e. $\mathcal{X} = R$, if $S \in \mathfrak{D}_1(\lambda)$ and $\mathcal{X} = \{a + \Delta k : k = 0, \pm 1, \pm 2, \dots\}$, if $S \in \mathfrak{D}_2(\lambda)$.

Assume that we are given three r.v.'s S_0, S_1 and S_2 with mean 0 and finite variance, so that S_1 and S_2 are independent and $S_0 = S_1 + S_2$, $S_i \in \mathfrak{D}(\lambda)$. Set $B_k^2 = \mathbb{E}S_k^2$, $k = 0, 1, 2$ and $B = B_1B_2/B_0$. Let $\Psi_k(h)$ be the cumulant generating function of the r.v. S_k , i.e.

$$\Psi_k(h) = \log \mathbb{E} \exp\{hS_k\}, \quad |h| \leq \lambda, \quad k = 0, 1, 2.$$

It is easy to see that $\Psi'_0(h)$ is strictly increasing, for $|h| \leq \lambda$. Let $H_0(x) = h$ be the solution of the equation $\Psi'_0(h) = x$, for x in the range of $\Psi'_0(h)$. Set $B = B_1B_2/B_0$,

$$\mu(y) = \Psi'_1(H_0(y)) = y - \Psi'_2(H_0(y)),$$

and

$$\sigma^2(y) = \Psi''_1(H_0(y))\Psi''_2(H_0(y))/\Psi''_0(H_0(y)).$$

Define the conditional distribution function

$$F(x|y) = P(S_1 \leq x | S_0 = y).$$

In the sequel c_i , $i = 0, 1, \dots$ denote positive absolute constants and θ_i , $i = 1, 2, \dots$ real numbers satisfying $|\theta_i| \leq 1$.

Theorem 1. *Assume that $S_0, S_1, S_2 \in \mathfrak{D}_1(\lambda)$ or $S_0, S_1, S_2 \in \mathfrak{D}_2(\lambda)$, for some $0 < \lambda < \infty$. Then, for any $x, y \in \mathcal{X}$ satisfying $|x| \leq c_0\lambda B^2$ and $|y| \leq c_0\lambda B^2$, we have*

$$F(x + \mu(y)|y) = \Phi\left(\frac{x}{\sigma(y)} + \theta_1 \frac{c_1}{\lambda\sigma(y)} \left(1 + \frac{x^2}{\sigma^2(y)}\right)\right),$$

provided $\lambda B \geq c_2$. Moreover, for $y \in \mathcal{X}$ satisfying $|y| \leq c_0\lambda B^2$, we have

$$\mu(y) = \beta_1 y + \theta_2 c_3 \frac{y^2}{\lambda B^2}, \quad \frac{1}{\sigma^2(y)} = \frac{1}{B^2} + \theta_3 c_4 \frac{|y|}{\lambda B^4},$$

where $\beta_1 = B_1^2/B_0^2$.

In the case of sums of i.i.d. r.v.'s we get the following result. Let ξ_1, \dots, ξ_n , $n \geq 1$ be a sequence of i.i.d. r.v.'s of means 0 and finite variances $\sigma^2 = \mathbb{E}\xi_i^2 < \infty$, obeying one of the two conditions (G1) or (G2) below:

G1 For some $0 < \lambda < \infty$ and $q \geq 1$,

$$\sup_{|h| \leq \lambda} \int_{-\infty}^{\infty} |\mathbb{E} \exp((it + h)\xi_i)|^q dt < \infty.$$

G2 The r.v. ξ_i takes values on the lattice $a + \Delta k$, $k \in \mathbb{Z}$ with maximal span Δ and for some $0 < \lambda < \infty$,

$$\mathbb{E} \exp(\lambda |\xi_i|) < \infty.$$

The common cumulant generating function of ξ_i -th is denoted by $\psi(t) = \log \mathbb{E} \exp(t\xi_i)$, for $|t| \leq \lambda$. Let $h(x)$, $|x| \leq \lambda/8$ be the inverse of the monotone

function $\psi'(t)$, $|t| \leq \lambda/4$. Assume that n_1, n_2 are positive integers such that $n_1 + n_2 = n$. Set $S_1 = \xi_1 + \dots + \xi_{n_1}$, $S_0 = \xi_1 + \dots + \xi_n$ and $F(x|y) = P(S_1 \leq x | S_0 = y)$.

Theorem 2. For any $|x| \leq c_0 \lambda n \sigma^2$ and $|y| \leq c_0 \lambda n \sigma^2$, we have

$$F(x + \mu(y)|y) = \Phi\left(\frac{x}{\sigma(y)} + \theta_1 \frac{c_1}{\lambda \sigma(y)} \left(1 + \frac{x^2}{\sigma^2(y)}\right)\right).$$

where

$$\mu(y) = \frac{n_1}{n}y, \quad \sigma^2(y) = \frac{n_1 n_2}{n} \psi''\left(h\left(\frac{y}{n}\right)\right).$$

Proofs of Theorems 1 and 2 rely on Lemmas 1 and 2 below, which are of independent interest.

Assume that we are given a real valued r.v. X , which has a density $p(x)$ w.r.t. Lebesgue measure on the real line. We shall impose conditions (C1-C2) below, where $\gamma_0, \dots, \gamma_4 > 0$ and $\sigma > 1$ denote real quantities not depending on x .

C1 For any $|x| \leq \gamma_0 \sigma$,

$$p(x) = \frac{1}{\sqrt{2\pi}} \exp\left(-\frac{x^2}{2} + \theta_1 \frac{\gamma_1}{2} \frac{x^3}{\sigma}\right) \left(1 + \theta_2 \gamma_2 \frac{1 + |x|}{\sigma}\right),$$

where $|\theta_i| \leq 1$, $i = 1, 2$.

C2 For any $|x| \leq \gamma_0 \sigma$ and any real z satisfying $zx \geq 0$,

$$p(x+z) \leq \gamma_3 \exp\left(-\frac{x^2}{2} + \frac{\gamma_1}{2} \frac{x^3}{\sigma} - \gamma_4 zx\right).$$

Lemma 1. Assume that the density $p(x)$ satisfies conditions (C1-C2). Then, for any x satisfying $|x| \leq c_0 \sigma$, we have

$$\Phi\left(x - c_1 \frac{1+x^2}{\sigma}\right) \leq P(X \leq x) \leq \Phi\left(x + c_1 \frac{1+x^2}{\sigma}\right),$$

provided $\sigma \geq c_2$, where c_0, c_1 and c_2 are constants depending only on $\gamma_0, \dots, \gamma_4$.

Let $\Delta_0 > 0$ and $\Delta \in (0, \Delta_0]$. Assume that we are given a r.v. X , which takes discrete values $a_k = a + k\Delta$, $k \in \{0, \pm 1, \pm 2, \dots\}$, with maximal span Δ and with $a \in [0, \Delta)$. In the sequel we shall impose conditions (D1-D2) below, where we assume that $\gamma_0, \dots, \gamma_4 > 0$ and $\sigma > 1$ denote real quantities not depending on values a_k , $k \in \{0, \pm 1, \pm 2, \dots\}$. For the sake of brevity we set $x_k = a_k/\sigma$.

D1 For any integer k satisfying $|x_k| \leq \gamma_0 \sigma$,

$$P(X = a_k) = \frac{\Delta}{\sqrt{2\pi}\sigma} \exp\left(-\frac{x_k^2}{2} + \theta_1 \frac{\gamma_1}{2} \frac{x_k^3}{\sigma}\right) \left(1 + \theta_2 \gamma_2 \frac{1 + |x_k|}{\sigma}\right).$$

D2 For any integers k, j satisfying $|x_k| \leq \gamma_0 \sigma$ and $(x_{k+j} - x_k)x_k \geq 0$,

$$P(X = a_{k+j}) \leq \gamma_3 \frac{\Delta}{\sigma} \exp\left(-\frac{x_k^2}{2} + \frac{\gamma_1}{2} \frac{x_k^3}{\sigma} - \gamma_4 (x_{k+j} - x_k)x_k\right).$$

Lemma 2. Assume that conditions (D1-D2) are satisfied. Then, for any x_k satisfying $|x_k| \leq c_0\sigma$,

$$\Phi\left(x_k - c_1 \frac{1 + x_k^2}{\sigma}\right) \leq P(X \leq a_k) \leq \Phi\left(x_k + c_1 \frac{1 + x_k^2}{\sigma}\right),$$

provided $\sigma \geq c_2$, where c_0, c_1 and c_2 are constants depending only on $\gamma_0, \dots, \gamma_4$ and Δ_0 .

Note that condition $\Delta \in (0, \Delta_0]$ is used to assure the uniformity of the assertion of the lemma in Δ .

1. Brown, L., Carter, A., Low, M. and Zhang, C.-H. *Equivalence theory for density estimation, Poisson processes and Gaussian white noise with drift.* Annals of statistics, **32** (2004) p. 2074-2097.

2. Carter, A. and Pollard, D. *Tusnady's inequality revisited.* Annals of statistics, **32** (2004) p. 2731-2741.

3. Grama I. and Nussbaum M. *A nonstandard Hungarian construction for partial sums.* Ann. I. H. Poincaré - PR, **38**, No. 6 (2002) p. 923-957.

4. Komlós, J., Major, P. and Tusnády, G. *An approximation of partial sums of independent rv's and the sample df. I.* Z. Wahrsch. verw. Gebiete, **32**, (1975). p. 111-131.

5. Komlós, J., Major, P. and Tusnády, G. *An approximation of partial sums of independent rv's and the sample df. II.* Z. Wahrsch. verw. Gebiete, **34** (1976). p. 33-58.

6. Sakhanenko, A. *The rate of convergence in the invariance principle for non-identically distributed variables with exponential moments.* Limit theorems for sums of random variables: Trudy Inst. Matem., Sibirsk. Otdel AN SSSR, Vol. 3 (1984). p. 3-49 (in Russian).

7. Zaitsev, A. *Bounds for quantiles of smooth conditional distributions and multidimensional invariance principle.* Sibirskii Matematicheskii Jurnal, **37** (1996). p. 807-831 (in Russian).