



HAL
open science

Empirical comparison of two methods for the Bayesian update of the parameters of probability distributions in a two-level hybrid probabilistic-possibilistic uncertainty framework for risk assessment

Nicola Pedroni, Enrico Zio, A Pasanisi, M Couplet

► To cite this version:

Nicola Pedroni, Enrico Zio, A Pasanisi, M Couplet. Empirical comparison of two methods for the Bayesian update of the parameters of probability distributions in a two-level hybrid probabilistic-possibilistic uncertainty framework for risk assessment. *ASCE-ASME Journal of Risk and Uncertainty in Engineering Systems, Part A: Civil Engineering*, 2015, pp.04015015. 10.1061/AJRUA6.0000848 . hal-01270598

HAL Id: hal-01270598

<https://hal.science/hal-01270598>

Submitted on 10 Feb 2016

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 Empirical comparison of two methods for the Bayesian update 2 of the parameters of probability distributions in a two-level 3 hybrid probabilistic-possibilistic uncertainty framework for 4 risk assessment

5 N. Pedroni¹, E. Zio², A. Pasanisi³, M. Couplet⁴

7 Abstract

8 *In this paper, we address the issue of updating in a Bayesian framework, the possibilistic*
9 *representation of the epistemically-uncertain parameters of (aleatory) probability distributions, as*
10 *new information (e.g., data) becomes available. Two approaches are considered: the first is based*
11 *on a purely possibilistic counterpart of the classical, well-grounded probabilistic Bayes' theorem;*
12 *the second relies on the hybrid combination of (i) Fuzzy Interval Analysis (FIA) to process the*
13 *uncertainty described by possibility distributions and (ii) repeated Bayesian updating of the*
14 *uncertainty represented by probability distributions.*

15 *The feasibility of the two methods is shown on a literature case study involving the risk-based*
16 *design of a flood protection dike.*

17 **Keywords:** hierarchical two-level uncertainty, Bayesian update, possibility distributions, fuzzy
18 interval analysis, possibilistic Bayes theorem, flood protection dike.
19
20

¹ Corresponding author. Assistant professor at the Chair of system science and the energetic challenge-Fondation EdF at CentraleSupélec, 3 Rue Joliot Curie – Plateau de Moulon, 91190, Gif-Sur-Yvette, France. E-mail: nicola.pedroni@supelec.fr.

² Full professor at the Chair of system science and the energetic challenge-Fondation EdF at CentraleSupélec, Grande Voie des Vignes, 92290, Chatenay-Malabry, France. E-mail: enrico.zio@ecp.fr. Also: Energy Department, Politecnico di Milano, Via Ponzio, 34/3 – 20133 Milano, Italy. E-mail: enrico.zio@polimi.it.

³ Project Manager at EDF R&D - EIFER, Emmy-Noether-Str. 11. 76131 Karlsruhe, Germany. E-mail address: alberto.pasanisi@edf.fr.

⁴ Research Engineer at Electricité de France, R&D, 6 Quai Watier, 78400, Chatou, France. E-mail address: mathieu.couplet@edf.fr.

21

22 **Introduction**

23 We consider a framework of uncertainty representation with two hierarchical levels [Limbourg and
24 de Rocquigny, 2010], in which risk analysis models of aleatory (i.e., random) events (e.g., failures)
25 contain parameters (e.g., probabilities, failure rates, ...) that are epistemically-uncertain, i.e., known
26 with poor precision due to lack of knowledge and information. Traditionally, both types of
27 uncertainty are represented by probability distributions [Apostolakis and Kaplan, 1981;
28 Apostolakis, 1990; NUREG-CR-6850, 2005; USNRC, 2002 and 2009; NASA, 2010] and Bayes' rule
29 is useful for updating the (probabilistic) epistemic uncertainty representation as new
30 information (e.g., data) becomes available [Bernardo and Smith, 1994; Siu and Kelly, 1998;
31 Lindley, 2000 and 2006; Bedford and Cooke, 2001; Atwood et al., 2003; Kelly and Smith, 2009 and
32 2011; Pasanisi et al., 2012].

33 However, in some situations, insufficient knowledge, information and data impair a probabilistic
34 representation of epistemic uncertainty. A number of alternative representation frameworks have
35 been proposed for such cases [Aven, 2010 and 2011; Aven and Steen, 2010; Aven and Zio, 2011;
36 Flage et al., 2009; Beer et al., 2013b and 2014b; Zhang et al., 2013], e.g., e.g., fuzzy set theory [Klir
37 and Yuan, 1995], fuzzy probabilities [Buckley, 2005; Beer, 2009b; Pannier et al., 2013], random set
38 theory [Molchanov, 2005], evidence theory [Ferson et al., 2003 and 2004; Helton et al., 2007 and
39 2008; Sentz and Ferson, 2002; Le Duy et al., 2013; Sallak et al., 2013], possibility theory (that can
40 be considered a special case of evidence theory) [Baudrit and Dubois, 2006; Baudrit et al., 2006 and
41 2008; Dubois, 2006; Dubois and Prade, 1988], probability bound analysis using probability boxes
42 (p-boxes) [Ferson and Ginzburg, 1996; Crespo et al., 2013; Mehl, 2013], interval analysis [Ferson
43 and Hajagos, 2004; Ferson and Tucker, 2006; Ferson et al., 2007 and 2010; Jalal-Kamali and
44 Kreinovich, 2013; Muscolino and Sofi, 2013; Zhang et al., 2013] and interval probabilities
45 [Weichselberger, 2000]; notice that most of these theories can be included within the general
46 common framework of *imprecise probabilities* [Kuznetsov, 1991; Walley, 1991; Kozine and

47 Filimonov, 2000; Kozine and Utkin, 2002; Coolen and Utkin, 2007; Beer and Ferson, 2013; Beer et
48 al., 2013a; Blockley, 2013; Reid, 2013; Sankararaman and Mahadevan, 2013].

49 In this paper, we adopt possibility distributions to describe epistemic uncertainty and address the
50 issue of updating, in a Bayesian framework, the possibilistic representation of the epistemically-
51 uncertain parameters of (aleatory) probability distributions. We take two approaches of literature.

52 The first is based on a purely possibilistic counterpart of the classical, well-grounded probabilistic
53 Bayes' theorem: it requires the construction of a possibilistic likelihood function which is used to
54 revise the prior possibility distributions of the uncertain parameters (determined on the basis of a
55 priori subjective knowledge and/or data) [Dubois and Prade, 1997; Lapointe and Bobee, 2000]. This
56 approach has been already applied by the authors for updating possibility distributions in [Pedroni
57 et al., 2014]. The second is a hybrid probabilistic-possibilistic method that relies on the use of
58 Fuzzy Probability Density Functions (FPDFs), i.e., PDFs with possibilistic (fuzzy) parameters. It is
59 based on the combination of (i) Fuzzy Interval Analysis (FIA) to process the uncertainty described
60 by possibility distributions and (ii) repeated Bayesian updating of the uncertainty represented by
61 probability distributions [Beer, 2009a; Stein and Beer, 2011; Stein et al., 2013; Beer et al., 2014a].

62 The objective (and the main contribution of the paper) is to compare the effectiveness of the two
63 methods. To the best of the authors' knowledge, this is the first time that the above mentioned
64 techniques are systematically compared with reference to risk assessment problems where hybrid
65 uncertainty is separated into two hierarchical levels. To keep the analysis simple and retain a clear
66 view of each step, the investigations are carried out with respect to a simple literature case study
67 involving the risk-based design of a flood protection dike [Pasanisi et al., 2009; Limbourg and de
68 Rocquigny, 2010]. In addition, different numerical indicators (e.g., cumulative distributions,
69 exceedance probabilities, percentiles, ...) are considered to perform a fair, quantitative comparison
70 between the methods and evaluate their rationale and appropriateness in relation to risk analysis.

71 Other methods have been proposed in the literature to revise, in a Bayesian framework, non-
72 probabilistic representations of epistemic uncertainty [Ferson, 2005]. In [Viertl, 1996, 1997, 1999,

73 2008a,b and 2011; Viertl and Hareter, 2004a,b; Viertl and Hule, 1991] a modification of the Bayes’
74 theorem is presented to account for the presence of fuzzy data and fuzzy prior PDFs: the approach
75 is similar to that employed by [Beer, 2009a; Stein and Beer, 2011; Stein et al., 2013] and considered
76 in this paper. In [Smets, 1993] a Generalized Bayes Theorem (GBT) is proposed within the
77 framework of evidence theory: this approach is applied by [Le-Duy et al., 2011] to update the
78 estimates of the failure rates of mechanical components in the context of nuclear Probabilistic Risk
79 Assessment (PRA). Finally, in [Walley, 1996; Bernard, 2005; Masegosa and Moral, 2014]
80 Imprecise Dirichlet Models (IDMs) are proposed for objective statistical inference from
81 multinomial data. In the IDM, prior or posterior uncertainty about a parameter is described by a set
82 of Dirichlet distributions, and inferences about events are summarized by lower and upper
83 probabilities. This model has been extended by [Quaeghebeur and de Cooman, 2005] to generalized
84 Bayesian inference from canonical exponential families and by [Walter and Augustin, 2009] with
85 the aim of handling prior-data conflicts.

86 The remainder of the paper is organized as follows. First, the representation of aleatory
87 (probabilistic) and epistemic (possibilistic) uncertainties in a “two-level” framework is provided;
88 then, the two methods employed in this paper for the Bayesian update of the possibilistic parameters
89 of aleatory probability distributions are described in details; after that, the case study concerning the
90 risk-based design of a flood protection dike is presented; in the following Section, the methods
91 described are applied to the case study: the results obtained are discussed and the two methods are
92 synthetically compared; finally, some conclusions are drawn in the last Section.

93 **Representation of aleatory and epistemic uncertainties in a two-level** 94 **framework: fuzzy random variables**

95 In all generality, we consider an uncertain variable Y , whose uncertainty is described by the
96 Probability Distribution Function (PDF) $p^Y(y|\theta)$, where $\theta = \{\theta_1, \theta_2, \dots, \theta_m, \dots, \theta_p\}$ is the vector of
97 the corresponding internal parameters. In a two-level framework, the parameters θ are themselves

98 affected by epistemic uncertainty [Limbourg and de Rocquigny, 2010]. In the present work, we
 99 describe these uncertainties by the (generally joint) possibility distribution $\pi^\theta(\theta)$ (it is
 100 straightforward to notice that in case the internal parameters $\{\theta_1, \theta_2, \dots, \theta_m, \dots, \theta_p\}$ are independent,
 101 then $\pi^\theta(\theta)$ is simply represented by a group of P separate, marginal and independent (i.e., non-
 102 interactive) possibility distributions, i.e., $\pi^\theta(\theta) = \{\pi^{\theta_1}(\theta_1), \pi^{\theta_2}(\theta_2), \dots, \pi^{\theta_m}(\theta_m), \dots, \pi^{\theta_p}(\theta_p)\}$). A
 103 random variable Y with possibilistic parameters θ is a particular case of a Fuzzy Random Variable
 104 (FRV), i.e., of a random variable whose values are not real, but rather fuzzy numbers [Féron, 1976;
 105 Kwakernaak, 1978; Puri and Ralescu, 1986; Baudrit et al., 2008; Couso and Sanchez, 2008]. The
 106 corresponding Fuzzy Probability Distribution Function (FPDF) is here indicated as $\tilde{p}^Y(y|\theta)$.
 107 For clarification by way of example, we may consider the generic uncertain variable Y described
 108 by a Gumbel PDF, i.e., $Y \sim p^Y(y|\theta) = Gum(\theta) = Gum(\theta_1, \theta_2) = Gum(\gamma, \delta) = p^Y(y|\gamma, \delta)$.
 109 Parameter $\delta = \theta_2$ (i.e., the scale parameter) is a fixed point-wise value ($\delta = \theta_2 = 100$), whereas
 110 parameter $\gamma = \theta_1$ (i.e., the location parameter) is epistemically-uncertain. By hypothesis, the only
 111 information available on $\gamma = \theta_1$ is that it is defined on interval $[a_\gamma, b_\gamma] = [900, 1300]$ and its most
 112 likely value is $c_\gamma = 1100$. Notice that such information is *not sufficient* for assigning a *single specific*
 113 probability distribution to describe the epistemic uncertainty in parameter $\gamma = \theta_1$. In facts, such
 114 scarce information is actually *compatible* with a *variety* of probability distributions (e.g., truncated
 115 normal, lognormal, triangular, ...). To address this issue, this limited state of knowledge about
 116 $\gamma = \theta_1$ is here described by a triangular possibility distribution $\pi^\gamma(\gamma)$ with core $c_\gamma = 1100$ and
 117 support $[a_\gamma, b_\gamma] = [900, 1300]$ (Fig. 1, top left column) [Baudrit and Dubois, 2006]. Indeed, this
 118 representation is *coherent* with the *information available*, as it can be demonstrated that such
 119 possibility distribution “encodes” the *family* of *all* the probability distributions with mode $c_\gamma = 1100$
 120 and support $[a_\gamma, b_\gamma] = [900, 1300]$ (obviously, this does *not* mean that the triangular possibility
 121 distribution is the *only one* with these characteristics, i.e., the only one able to encode such a

122 probability family). In other words, *one single* possibility distribution generates in practice a
 123 “*bundle*” of probability distributions with mode $c_\gamma = 1100$ and support $[a_\gamma, b_\gamma] = [900, 1300]$. The
 124 reader is referred to [Baudrit and Dubois, 2006; Couso et al., 2001; Dubois et al., 2004] for further
 125 technical details and a formal proof of these statements.

126 In order to provide an additional *practical* interpretation of the possibility distribution $\pi^\gamma(\gamma)$ of
 127 $\gamma = \theta_1$, we can define its α -cut sets $A_\alpha^\gamma = \{ \gamma : \pi^\gamma(\gamma) \geq \alpha \}$, with $0 \leq \alpha \leq 1$. For example, $A_{0.5}^\gamma =$
 128 $[1000, 1200]$ is the set of γ values for which the possibility function is greater than or equal to 0.5
 129 (dashed segment in Fig. 1, top left column). Notice that the α -cut set A_α^γ of parameter γ can be
 130 interpreted also as the $(1 - \alpha) \cdot 100\%$ Confidence Interval (CI) for γ , i.e., the interval such that
 131 $P[\gamma \in A_\alpha^\gamma] \geq 1 - \alpha$. For example, $A_0^\gamma = [900, 1300]$ is the $(1 - 0) \cdot 100\% = 100\%$ CI for γ , i.e., the
 132 interval that contains the “true” value of γ with certainty (solid segment in Fig. 1, top left column);
 133 $A_{0.5}^\gamma = [1000, 1200]$ ($\subset A_0^\gamma$) is the $(1 - 0.5) \cdot 100\% = 50\%$ CI (dashed segment in Fig. 1, top left
 134 column); $A_{0.8}^\gamma = [1050, 1150]$ ($\subset A_{0.5}^\gamma \subset A_0^\gamma$) is the $(1 - 0.8) \cdot 100\% = 20\%$ CI, and so on. In this
 135 view, the possibility distribution $\pi^\gamma(\gamma)$ can be interpreted as a set of *nested* CIs for parameter γ
 136 [Baudrit and Dubois, 2006; Couso et al., 2001; Dubois et al., 2004].

137 For each possibility (resp., confidence) level α (resp., $1 - \alpha$) in $[0, 1]$, a *family* of PDFs for Y ,
 138 namely $\{p^Y(y | \gamma, \delta)\}_\alpha$, can be generated by letting the epistemically-uncertain parameter γ range
 139 within the corresponding α -cut set A_α^γ , i.e., $\{p^Y(y | \gamma, \delta)\}_\alpha = \{p^Y(y | \gamma, \delta) : \gamma \in A_\alpha^\gamma, \delta = 100\}$. By way
 140 of example, Fig. 1, top right column, shows four PDFs belonging to the family $\{p^Y(y | \gamma, \delta)\}_{\alpha=0}$
 141 (solid lines) and four PDFs belonging to the family $\{p^Y(y | \gamma, \delta)\}_{\alpha=0.5}$ (dashed lines).

142 In the same way, a *bundle* of Cumulative Distribution Functions (CDFs) for Y , namely
 143 $\{F^Y(y | \gamma, \delta)\}_\alpha$, can be constructed by letting γ range within A_α^γ , i.e., $\{F^Y(y | \gamma, \delta)\}_\alpha =$
 144 $\{F^Y(y | \gamma, \delta) : \gamma \in A_\alpha^\gamma, \sigma = 100\}$. This *family* of CDFs (of level α) is bounded above and below by the

145 upper and lower CDFs, $\bar{F}_\alpha^Y(y)$ and $\underline{F}_\alpha^Y(y)$, defined as $\bar{F}_\alpha^Y(y) = \sup_{\gamma \in A_\alpha^Y} \{F^Y(y | \gamma, \delta = 100)\}$ and
 146 $\underline{F}_\alpha^Y(y) = \inf_{\gamma \in A_\alpha^Y} \{F^Y(y | \gamma, \delta = 100)\}$, respectively. Since $\pi^\gamma(\gamma)$ can be interpreted as a set of nested CIs
 147 for parameter γ (see above), it can be argued that the α -cuts of $\pi^\gamma(\gamma)$ induce also a *set of nested*
 148 *pairs* of CDFs $\{(F_\alpha^Y(y), \bar{F}_\alpha^Y(y)) : 0 \leq \alpha \leq 1\}$ which bound the “true” CDF $F^Y(y)$ of Y with confidence
 149 larger than or equal to $(1 - \alpha)$, i.e., $P[\underline{F}_\alpha^Y(y) \leq F^Y(y) \leq \bar{F}_\alpha^Y(y)] \geq 1 - \alpha$, with $0 \leq \alpha \leq 1$ [Baudrit et
 150 al., 2007 and 2008]. In passing, notice that the upper and lower CDFs (of level α), $\bar{F}_\alpha^Y(y)$ and
 151 $\underline{F}_\alpha^Y(y)$, can be referred to as the *plausibility* and *belief* functions (of level α) of the set $Z = (-\infty, y]$,
 152 i.e., $\bar{F}_\alpha^Y(y) = Pl_\alpha^Y(Z)$ and $\underline{F}_\alpha^Y(y) = Bel_\alpha^Y(Z)$, respectively. For illustration purposes, Fig. 1, bottom
 153 row, shows the bounding upper and lower CDFs of Y , $\bar{F}_\alpha^Y(y) = Pl_\alpha^Y(Z)$ and $\underline{F}_\alpha^Y(y) = Bel_\alpha^Y(Z)$, built
 154 in correspondence of the α -cuts of level $\alpha = 0$ (solid lines), 0.5 (dashed lines) and 1 (dot-dashed
 155 line) of the possibility distribution $\pi^\gamma(\gamma)$ of parameter γ (Fig. 1, top left column).
 156 Finally, the *set of nested pairs* of CDFs $\{(F_\alpha^Y(y), \bar{F}_\alpha^Y(y)) : 0 \leq \alpha \leq 1\} = \{(Bel_\alpha^Y(Z), Pl_\alpha^Y(Z)) : 0 \leq \alpha \leq 1\}$,
 157 $Z = (-\infty, y]$, can be synthesized into a *single pair* of plausibility and belief functions as
 158 $Pl^Y(Z) = \int_0^1 Pl_\alpha^Y(Z) d\alpha$ and $Bel^Y(Z) = \int_0^1 Bel_\alpha^Y(Z) d\alpha$, respectively (dotted lines in Fig. 1, bottom
 159 row): in other words, $Pl^Y(Z)$ and $Bel^Y(Z)$ are obtained by *averaging* the different nested
 160 plausibility and belief functions (i.e., $\{(Bel_\alpha^Y(Z), Pl_\alpha^Y(Z)) : 0 \leq \alpha \leq 1\}$) generated at different
 161 possibility levels $\alpha \in [0, 1]$ (i.e., by averaging the different contributions to the plausibility and
 162 belief functions produced by different α -cuts of the epistemic parameter γ). The plausibility and
 163 belief functions $Pl^Y(Z)$ and $Bel^Y(Z)$, $Z = (-\infty, y]$, are shown to represent the “*best bounds*” for the
 164 “true” CDF $F^Y(y)$ of the uncertain variable Y [Ralescu, 2002; Baudrit et al., 2007 and 2008; Couso
 165 et al., 2004; Couso and Dubois, 2009; Couso and Sanchez, 2011].

166 Further details about FRVs are not given here for the sake of brevity: the interested reader is
167 referred to the cited references.

168

169

Fig. 1

170 **Bayesian update of the possibilistic parameters of aleatory probability** 171 **distributions**

172 In this Section, we present the methods employed in this study for updating, in a Bayesian
173 framework, the possibilistic representation of the epistemically-uncertain parameters of (aleatory)
174 probability distributions, as *new* information/evidence (e.g., data) becomes available. In this view,
175 let $\pi^\theta(\boldsymbol{\theta})$ be the (joint) *prior* possibility distribution for the parameters $\boldsymbol{\theta} = [\theta_1, \theta_2, \dots, \theta_m, \dots, \theta_p]$ of
176 the PDF $p^Y(y|\boldsymbol{\theta})$ of variable Y (built on the basis of a priori subjective engineering knowledge
177 and/or data). For example, in the risk assessment context of this paper Y may represent the yearly
178 maximal water flow of a river described by a Gumbel distribution: thus, $Y \sim p^Y(y|\boldsymbol{\theta}) = Gum(\boldsymbol{\theta}) =$
179 $Gum(\theta_1, \theta_2) = Gum(\gamma, \delta) = p^Y(y|\gamma, \delta)$ and $\pi^\theta(\boldsymbol{\theta}) = \pi^{\gamma, \delta}(\gamma, \delta)$. Moreover, let
180 $\mathbf{y} = [y_1, y_2, \dots, y_k, \dots, y_D]$ be a vector of D *observed* pieces of data representing the *new*
181 information/evidence *available* for the analysis: referring to the example above, \mathbf{y} may represent a
182 vector of D values collected over a long period time (e.g., many years) of the yearly maximal water
183 flow of the river under analysis. The objective of the Bayesian analysis is to update the a priori
184 representation $\pi^\theta(\boldsymbol{\theta}) = \pi^{\gamma, \delta}(\gamma, \delta)$ of $\boldsymbol{\theta} = [\gamma, \delta]$ on the basis of the new evidence acquired, i.e., to
185 calculate the *posterior* possibility distribution $\pi^\theta(\boldsymbol{\theta}|\mathbf{y})$ (i.e., $\pi^{\gamma, \delta}(\gamma, \delta|\mathbf{y})$) of $\boldsymbol{\theta}$ after \mathbf{y} is
186 obtained.

187 In the present paper, two methods are considered to this aim: the purely possibilistic method and the
188 hybrid probabilistic and possibilistic approach.

189 ***Purely possibilistic approach***

190 The purely possibilistic method (hereafter also referred to as ‘Approach A’ for brevity) is based on
 191 a purely possibilistic counterpart of the classical, probabilistic Bayes’ theorem [Dubois and Prade,
 192 1997; Lapointe and Bobée, 2000]:

193
$$\pi^\theta(\boldsymbol{\theta} | \mathbf{y}) = \frac{\pi_L^\theta(\boldsymbol{\theta} | \mathbf{y}) \cdot \pi^\theta(\boldsymbol{\theta})}{\sup_{\boldsymbol{\theta}} \{\pi_L^\theta(\boldsymbol{\theta} | \mathbf{y}) \cdot \pi^\theta(\boldsymbol{\theta})\}}, \quad (1)$$

194 where $\pi_L^\theta(\boldsymbol{\theta} | \mathbf{y})$ is the possibilistic likelihood of the parameter vector $\boldsymbol{\theta}$ given the newly observed
 195 data \mathbf{y} , and quantities $\pi^\theta(\boldsymbol{\theta} | \mathbf{y})$ and $\pi^\theta(\boldsymbol{\theta})$ are defined above. Notice that $\sup_{\boldsymbol{\theta}} \{\pi_L^\theta(\boldsymbol{\theta} | \mathbf{y}) \cdot \pi^\theta(\boldsymbol{\theta})\}$ is a
 196 normalization factor such that $\sup_{\boldsymbol{\theta}} \{\pi^\theta(\boldsymbol{\theta} | \mathbf{y})\} = 1$, as required by possibility theory [Dubois, 2006].

197 It is worth mentioning that forms of the possibilistic Bayes’ theorem alternative to (1) can be
 198 constructed as a result of other definitions of the operation of ‘conditioning’ with possibility
 199 distributions: the reader is referred to [Dubois and Prade, 1997; Lapointe and Bobée, 2000] for
 200 technical details. In this paper, expression (1) has been chosen because “it satisfies desirable
 201 properties of the revision process and lead to continuous posterior distributions” [Lapointe and
 202 Bobée, 2000].

203 The possibilistic likelihood $\pi_L^\theta(\boldsymbol{\theta} | \mathbf{y})$ is here obtained by *transforming* the classical probabilistic

204 likelihood function $L^\theta(\boldsymbol{\theta} | \mathbf{y})$ through *normalization*, i.e., $\pi_L^\theta(\boldsymbol{\theta} | \mathbf{y}) = \frac{L^\theta(\boldsymbol{\theta} | \mathbf{y})}{\sup_{\boldsymbol{\theta}} \{L^\theta(\boldsymbol{\theta} | \mathbf{y})\}}$ [Anoop et al.,

205 2006] (obviously, $L^\theta(\boldsymbol{\theta} | \mathbf{y}) = \prod_{k=1}^D p^Y(y_k | \boldsymbol{\theta})$ in the case the observations $\{y_k: k = 1, 2, \dots, D\}$ are

206 independent and identically distributed). This choice has been made for the following main reasons:

- 207 i. the transformation is simple and can be straightforwardly applied to any distribution [Anoop
 208 et al., 2006];
- 209 ii. the resulting possibilistic likelihood is *very closely related* to the classical, purely
 210 probabilistic one (which is theoretically well-grounded) by means of the simple and direct

- 211 operation of normalization that preserves the “*original structure*” of the experimental
212 evidence;
- 213 iii. it can be easily verified that the resulting possibilistic likelihood keeps the *sequential* nature
214 of the updating procedure typical of the standard Bayes’ theorem;
- 215 iv. the operation of likelihood normalization finds also *theoretical* justifications in some recent
216 works of literature (see the brief discussion below) [Denoeux, 2014; Moral, 2014].

217 However, two considerations are in order with respect to this choice. First, it has to be admitted that
218 the resulting possibility distributions do not in general adhere to the probability-possibility
219 consistency principle [Dubois and Prade, 1980]. Second, it has to be remembered that the
220 probabilistic likelihood function $L^\theta(\boldsymbol{\theta} | \mathbf{y})$ is *not* a probability distribution: in this view, from a
221 rigorous mathematical viewpoint, speaking of probability/possibility transformation for it would be
222 wrong. On the other side, from the practical engineering viewpoint of interest to the present paper,
223 an operation of normalization can be performed (i.e., $L^\theta(\boldsymbol{\theta} | \mathbf{y}) / \int_{\theta} L^\theta(\boldsymbol{\theta} | \mathbf{y}) d\boldsymbol{\theta}$) in order to
224 “technically” provide it with the “properties” of a probability distribution function.

225 It is worth noting that other techniques of transformation of probability density functions into
226 possibility distributions exist, but the corresponding details are not given here for brevity sake: the
227 interested reader is referred to [Dubois et al., 1993, 2004 and 2008; Flage et al., 2010 and 2013] for
228 some proposed techniques, e.g., the principle of maximum specificity [Dubois et al., 1993] and the
229 principle of minimal commitment [Dubois et al., 2008]. Also, it has to be noticed that techniques
230 are also available to construct possibility distributions (and, thus, possibilistic likelihood functions)
231 *directly* from rough experimental *data* (i.e., without resorting to probability-possibility
232 transformations): see [Masson and Denoeux, 2006; Mauris, 2008; Hou and Yang, 2010; Serrurier
233 and Prade, 2011] for more details. Finally, for a thorough *theoretical* justification of a “possibilistic
234 vision” of the likelihood the reader is referred to: e.g., [Dubois et al., 1997], where possibility
235 measures are considered as the *supremum* of a *family* of likelihood functions; [Denoeux, 2014],

236 where the evidence about a parameter (after observing a piece of data) is represented by a
 237 *consonant* “likelihood based” belief function, whose contour function equals the *normalized*
 238 *likelihood function* (see above): in the paper, this is also rigorously derived from three basic
 239 principles, i.e., the likelihood principle [Edwards, 1992], compatibility with Bayes’ rule and the
 240 minimal commitment principle [Smets, 1993]; and finally [Moral, 2014], where the approach by
 241 [Denoeux, 2014] is discussed and the issue of representing likelihood information is taken from the
 242 point of view of imprecise probabilities.

243 ***Hybrid probabilistic and possibilistic approach***

244 The hybrid probabilistic and possibilistic method (hereafter also called ‘Approach B’ for brevity) is
 245 based on the construction of a Fuzzy Probability Distribution Function (FPDF) to be used as a
 246 ‘fictitious’ prior for the epistemically-uncertain parameters θ of the PDF $p^Y(y|\theta)$ of the uncertain
 247 variable Y : in other words, a fictitious (artificial) *probabilistic* function has to be ‘superimposed’
 248 onto the *purely possibilistic* prior $\pi^\theta(\theta)$ that has to be updated. In more detail, let $\tilde{p}^\theta(\theta|\varphi)$ be the
 249 fictitious (prior) FPDF of θ , constructed by the superimposition of an (arbitrarily selected)
 250 fictitious PDF $p^\theta(\theta|\varphi)$ and a vector of parameters φ described by a (properly selected) possibility
 251 distribution $\pi^\varphi(\varphi)$ (it is straightforward to notice that in the case parameters $\{\theta_m: m = 1, 2, \dots, P\}$

252 are independent, then $p^\theta(\theta|\varphi) = \prod_{m=1}^P p^{\theta_m}(\theta_m|\varphi_m)$, with $\varphi_m = [\phi_{m1}, \phi_{m2}, \dots, \phi_{mN_m}]$, $m = 1, 2, \dots,$

253 P ; also, $\pi^\varphi(\varphi)$ is expressed as $\pi^\varphi(\varphi) = \{\pi^{\varphi_1}(\varphi_1), \pi^{\varphi_2}(\varphi_2), \dots, \pi^{\varphi_m}(\varphi_m), \dots, \pi^{\varphi_P}(\varphi_P)\}$. In addition,

254 if also the possibilistic parameters $\phi_{m1}, \phi_{m2}, \dots, \phi_{mN_m}$ of the fictitious PDF $p^{\theta_m}(\theta_m|\varphi_m)$, $m = 1, 2,$

255 \dots, P , are independent, then $\pi^{\varphi_m}(\varphi_m) = \{\pi^{\varphi_{m1}}(\phi_{m1}), \pi^{\varphi_{m2}}(\phi_{m2}), \dots, \pi^{\varphi_{mN_m}}(\phi_{mN_m})\}$. As before, let $Y \sim$

256 $p^Y(y|\theta) = Gum(\gamma, \delta) = p^Y(y|\gamma, \delta)$, with $\theta = [\theta_1, \theta_2] = [\gamma, \delta]$ epistemically-uncertain. For

257 simplicity, we consider $\theta_1 = \gamma$ and $\theta_2 = \delta$ independent, such that $\pi^\theta(\theta) = \{\pi^\gamma(\gamma), \pi^\delta(\delta)\}$ and that

258 the (fictitious) PDF $p^\theta(\theta|\varphi)$ can be written as $\prod_{m=1}^P p^{\theta_m}(\theta_m|\varphi_m) = p^{\theta_1}(\theta_1|\varphi_1) \cdot p^{\theta_2}(\theta_2|\varphi_2) =$
259 $p^\gamma(\gamma|\varphi_\gamma) \cdot p^\delta(\delta|\varphi_\delta)$. By hypothesis, the analyst *arbitrarily* selects Normal distributions to
260 represent $p^\gamma(\gamma|\varphi_\gamma)$ and $p^\delta(\delta|\varphi_\delta)$, i.e., $p^\gamma(\gamma|\varphi_\gamma) = N(\varphi_\gamma) = N(\mu_\gamma, \sigma_\gamma)$ and $p^\delta(\delta|\varphi_\delta) =$
261 $N(\varphi_\delta) = N(\mu_\delta, \sigma_\delta)$, respectively (Normal and Uniform distributions are indicated as good choices
262 for $p^\theta(\theta|\varphi)$ by [Beer, 2009a; Stein and Beer, 2011; Stein et al., 2013]). Then, parameters
263 $\varphi_\gamma = [\mu_\gamma, \sigma_\gamma]$ and $\varphi_\delta = [\mu_\delta, \sigma_\delta]$ are represented by the possibility distributions
264 $\pi^{\varphi_\gamma} = \{\pi^{\mu_\gamma}(\mu_\gamma), \pi^{\sigma_\gamma}(\sigma_\gamma)\}$ and $\pi^{\varphi_\delta} = \{\pi^{\mu_\delta}(\mu_\delta), \pi^{\sigma_\delta}(\sigma_\delta)\}$ (by so doing, the fictitious prior FPDF
265 $\tilde{p}^\theta(\theta|\varphi) = \tilde{p}^{\theta_1}(\theta_1|\varphi_1) \cdot \tilde{p}^{\theta_2}(\theta_2|\varphi_2) = \tilde{p}^\gamma(\gamma|\varphi_\gamma) \cdot \tilde{p}^\delta(\delta|\varphi_\delta)$ is constructed). These possibility
266 distributions should be *properly* selected by the analyst so as to reflect as closely as possible the
267 structure of the ‘*real*’ prior possibility distribution $\pi^\theta(\theta) = \{\pi^\gamma(\gamma), \pi^\delta(\delta)\}$ that has to be updated:
268 for example, π^{φ_γ} and π^{φ_δ} could be identified by ‘imposing’ that the *expected value* of the FPDF
269 $\tilde{p}^\theta(\theta|\varphi)$ corresponds to the *real* prior possibility distribution $\pi^\theta(\theta)$, i.e., $E_\varphi[\tilde{p}^\theta(\theta|\varphi)] = \pi^\theta(\theta)$
270 or, in this case, $E_{\varphi_\gamma}[\tilde{p}^\gamma(\gamma|\varphi_\gamma)] = \pi^\gamma(\gamma)$ and $E_{\varphi_\delta}[\tilde{p}^\delta(\delta|\varphi_\delta)] = \pi^\delta(\delta)$.

271 In extreme synthesis, the method relies on the hybrid combination of (i) Fuzzy Interval Analysis
272 (FIA) to process the uncertainty described by possibility distributions and (ii) *repeated* Bayesian
273 updating of the uncertainty represented by probability distributions [Beer, 2009a; Stein and Beer,
274 2011; Stein et al., 2013]. In more details, the algorithm proceeds as follows:

- 275 1. set $\alpha = 0$;
- 276 2. select the α -cut A_α^φ of the possibility distribution $\pi^\varphi(\varphi)$ of vector φ of the parameters of the
277 (fictitious) prior PDF $p^\theta(\theta|\varphi)$;
- 278 3. letting the parameter vector φ range within the corresponding α -cut A_α^φ identified at step 2.
279 above, generate a family of (fictitious) prior PDFs $\{p^\theta(\theta|\varphi)\}_\alpha = \{p^\theta(\theta|\varphi): \varphi \in A_\alpha^\varphi\}$. This is

280 empirically done by (i) randomly or deterministically selecting a *finite* number T (e.g., $T =$
 281 100 in this paper) of parameter vectors $\boldsymbol{\varphi}_{l,\alpha}$, $l = 1, 2, \dots, T$, in A_α^θ and (ii) evaluating
 282 $p^\theta(\boldsymbol{\theta} | \boldsymbol{\varphi})$ in correspondence of these vectors, i.e., $\{p^\theta(\boldsymbol{\theta} | \boldsymbol{\varphi})\}_\alpha \approx \{p^\theta(\boldsymbol{\theta} | \boldsymbol{\varphi}_{l,\alpha}) : l = 1, 2, \dots, T\}$;

283 4. apply the classical probabilistic Bayes theorem to *each* (fictitious) prior PDF $p^\theta(\boldsymbol{\theta} | \boldsymbol{\varphi}_{l,\alpha})$
 284 generated at step 3. above to get the corresponding posterior PDF $p_{l,\alpha}^\theta(\boldsymbol{\theta} | \mathbf{y})$:

$$285 \quad p_{l,\alpha}^\theta(\boldsymbol{\theta} | \mathbf{y}) = \frac{L^\theta(\boldsymbol{\theta} | \mathbf{y}) \cdot p(\boldsymbol{\theta} | \boldsymbol{\varphi}_{l,\alpha})}{\int_{\boldsymbol{\theta}} L^\theta(\boldsymbol{\theta} | \mathbf{y}) \cdot p(\boldsymbol{\theta} | \boldsymbol{\varphi}_{l,\alpha}) d\boldsymbol{\theta}}, \quad l = 1, 2, \dots, T \quad (2)$$

286 This is equivalent to generating a family $\{p^\theta(\boldsymbol{\theta} | \mathbf{y})\}_\alpha$ of posterior PDFs for $\boldsymbol{\theta}$, i.e.,
 287 $\{p^\theta(\boldsymbol{\theta} | \mathbf{y})\}_\alpha \approx \{p_{l,\alpha}^\theta(\boldsymbol{\theta} | \mathbf{y}) : l = 1, 2, \dots, T\}$;

288 5. calculate the *expected value* of *each* posterior PDF $p_{l,\alpha}^\theta(\boldsymbol{\theta} | \mathbf{y})$ generated at step 4. above to
 289 obtain a *point estimate* $\boldsymbol{\theta}_{l,\alpha} | \mathbf{y}$ for the epistemically-uncertain parameter vector $\boldsymbol{\theta}$ [Stein et
 290 al., 2013]:

$$291 \quad \boldsymbol{\theta}_{l,\alpha} | \mathbf{y} = E_{\boldsymbol{\theta}}[p_{l,\alpha}^\theta(\boldsymbol{\theta} | \mathbf{y})], \quad l = 1, 2, \dots, T \quad (3)$$

292 6. take the hull enveloping the T point estimates $\boldsymbol{\theta}_{l,\alpha} | \mathbf{y}$, $l = 1, 2, \dots, T$, as the (P -dimensional)
 293 α -cut $A_\alpha^{\theta|\mathbf{y}}$ of the (joint P -dimensional) posterior possibility distribution $\pi^\theta(\boldsymbol{\theta} | \mathbf{y})$ of $\boldsymbol{\theta}$. By
 294 way of example and for illustration purposes, Fig. 2 shows the identification of a (two-
 295 dimensional) α -cut $A_\alpha^{\theta|\mathbf{y}} = A_\alpha^{\theta_1, \theta_2|\mathbf{y}}$ (solid line) as the contour enclosing $T = 20$ point estimates
 296 $\boldsymbol{\theta}_{l,\alpha} | \mathbf{y}$, $l = 1, 2, \dots, 20$ (dots), in the simplified case of $P = 2$ parameters θ_1 and θ_2 ;

297 7. if $\alpha < 1$, then set $\alpha = \alpha + \Delta\alpha$ (e.g., $\Delta\alpha = 0.05$ in this paper) and return to step 2. above;
 298 otherwise, stop the algorithm.

299 The (joint P -dimensional) posterior possibility distribution $\pi^\theta(\boldsymbol{\theta} | \mathbf{y})$ for the epistemically-uncertain
 300 parameter vector $\boldsymbol{\theta}$ is empirically constructed as the (discrete) collection of the α -cuts $A_\alpha^{\theta|y}$, $\alpha = 0,$
 301 $0.05, \dots, 0.95, 1$, found at step 6. above.

302
 303 It is worth noting that the application of both approaches A and B always produces a *joint P -*
 304 *dimensional* posterior possibility distribution $\pi^\theta(\boldsymbol{\theta} | \mathbf{y})$ (*whatever* the state of dependence between
 305 the priors), characterized by P -dimensional α -cuts $A_\alpha^{\theta|y}$, with $0 < \alpha < 1$: as a consequence, there is
 306 an *interactive dependence* between the values that parameters $\{\theta_m: m = 1, 2, \dots, P\}$ can take when
 307 ranging within a given α -cut $A_\alpha^{\theta|y}$: for example, in Fig. 2 it is impossible that parameter θ_1 takes on
 308 low values *and* parameter θ_2 takes on high values at the same time.

309 From $\pi^\theta(\boldsymbol{\theta} | \mathbf{y})$ it is straightforward to obtain the *marginal* posterior possibility distribution
 310 $\pi^{\theta_m}(\theta_m | \mathbf{y})$ for each parameter θ_m as $\pi^{\theta_m}(\theta_m | \mathbf{y}) = \max_{\theta_j \in \mathfrak{R}, j \neq m} \{\pi^\theta(\boldsymbol{\theta} | \mathbf{y})\}$, $\forall \theta_m \in \mathfrak{R}$, $m = 1, 2, \dots, P$

311 [Baudrit et al., 2006]: $\pi^{\theta_m}(\theta_m | \mathbf{y})$ is the projection of $\pi^\theta(\boldsymbol{\theta} | \mathbf{y})$ onto the m -th axis. The (one-
 312 dimensional) α -cut $A_\alpha^{\theta_m|y} = [\underline{\theta}_{m,\alpha} | y, \bar{\theta}_{m,\alpha} | y]$ of the marginal possibility distribution $\pi^{\theta_m}(\theta_m | \mathbf{y})$ is
 313 then related to the (P -dimensional) α -cut $A_\alpha^{\theta|y}$ of the joint possibility distribution $\pi^\theta(\boldsymbol{\theta} | \mathbf{y})$ by the
 314 following straightforward relation, i.e., $A_\alpha^{\theta_m|y} = [\underline{\theta}_{m,\alpha} | y, \bar{\theta}_{m,\alpha} | y] = [\min_{\theta \in A_\alpha^{\theta|y}} \{\theta_m\}, \max_{\theta \in A_\alpha^{\theta|y}} \{\theta_m\}]$. In this

315 view, notice that the use of the P -dimensional α -cut $A_{\alpha, Cart}^{\theta|y}$ constructed by the Cartesian product of
 316 the (one-dimensional) α -cuts $A_\alpha^{\theta_m|y}$ of the marginal distributions, $m = 1, 2, \dots, P$ (i.e., $A_{\alpha, Cart}^{\theta|y} = A_\alpha^{\theta_1|y}$
 317 $\times A_\alpha^{\theta_2|y} \times \dots \times A_\alpha^{\theta_m|y} \times \dots \times A_\alpha^{\theta_p|y}$) would (incorrectly) imply *independence* between the posterior
 318 estimates of the parameters $\{\theta_m: m = 1, 2, \dots, P\}$; however, since $A_{\alpha, Cart}^{\theta|y}$ completely *contains* $A_\alpha^{\theta|y}$
 319 (i.e., by definition $A_\alpha^{\theta|y} \subset A_{\alpha, Cart}^{\theta|y}$), then conservatism would be still guaranteed [Stein et al., 2013].

320 For illustration purposes and with reference to the example above, Fig. 2 shows also the two-

321 dimensional α -cut $A_{\alpha, Cart}^{\theta_1, \theta_2 | y}$ (dashed line) generated by the Cartesian product of the (one-dimensional)
 322 α -cuts $A_{\alpha}^{\theta_1 | y}$ and $A_{\alpha}^{\theta_2 | y}$, i.e., $A_{\alpha, Cart}^{\theta_1, \theta_2 | y} = A_{\alpha}^{\theta_1 | y} \times A_{\alpha}^{\theta_2 | y}$.

323

324

325

Fig. 2

326 The characteristics of the two approaches are summarized in Table 1. Notice that both methods
 327 relies on *arbitrary assumptions* about either the prior or the likelihood functions: in the purely
 328 possibilistic approach (A) the “original” possibilistic prior is employed, but a possibilistic
 329 likelihood function has to be constructed (e.g., by probability-possibility transformations, directly
 330 from rough experimental data and/or by resorting to the guidelines provided by [Dubois et al., 1997;
 331 Denoeux, 2014]); instead, in the hybrid method (B) the original probabilistic likelihood function is
 332 used, but a “fictitious” prior Fuzzy Probability Distribution Function needs to be identified by
 333 superimposing an *arbitrarily selected* probabilistic PDF onto the “original” possibilistic prior that
 334 has to be updated.

335

336

Table 1.

337 A final consideration is in order with respect to the two approaches here outlined. In the hybrid
 338 probabilistic-possibilistic framework of interest to the present paper, the knowledge a priori
 339 available on the parameters $\theta = [\theta_1, \theta_2, \dots, \theta_m, \dots, \theta_p]$ of a given (aleatory) probability model
 340 $p^Y(y|\theta)$ is described by the *prior* possibility distribution function $\pi^\theta(\theta)$. As detailed in the
 341 previous Section, the possibilistic approach is particularly suitable to address those situations where
 342 the information a priori available on θ is scarce and imprecise, i.e., *not sufficient* for assigning a
 343 *single specific* probability distribution to (describe the epistemic uncertainty in) θ . Actually, the
 344 *possibilistic* function $\pi^\theta(\theta)$ is in practice “equivalent” to the *family* of *all* those *probability*
 345 *distributions* (of possibly *different shapes*) that are *coherent* with the scarce information available
 346 on θ . On the other hand, it is worth mentioning that in a classical *purely probabilistic* framework,
 347 imprecision in prior information about θ can be also accounted for by means of the so-called

348 *Hierarchical Bayes* approach. Hierarchical Bayes is so-named because it utilizes *hierarchical* or
349 *multistage* prior distributions [Gelman, 2006; Gelman et al., 2008; Congdon, 2010; Kelly and
350 Smith, 2011; Chung et al., 2015; Shirley and Gelman, 2015]. To develop a hierarchical model for
351 θ , we need to specify a *first-stage prior* (say, $p^\theta(\theta|\varphi)$), which is often of a *particular functional*
352 *form*, often a conjugate prior. However, analysts find it difficult to express their uncertainty
353 numerically at all, much less as particular probability distributions. Thus, a higher-dimensional
354 model is defined to represent such (epistemic) uncertainty: in particular, we need to specify an
355 “additional” prior distribution (say, $p^\varphi(\varphi|\omega)$) on the first-stage parameters φ . Distribution
356 $p^\varphi(\varphi|\omega)$ is called the *second-stage prior*, or *hyper-prior*. This way of proceeding amounts to
357 generating a ‘parametric’ family $\{p^\theta(\theta|\varphi)\}$ of (first-stage prior) probability distributions for θ , all
358 obtained in correspondence of different possible values of the first-stage parameters φ (described by
359 hyper-prior $p^\varphi(\varphi|\omega)$). This method has been investigated in the field of social and behavioral
360 sciences with the *main* aim of treating hierarchical data with different levels of variables in the same
361 statistical model. For example, the hierarchical data for sociological survey analysis include
362 measurements from individuals with different historical, geographic, or economic variables. To this
363 end, the hierarchical modeling was proposed to account for the different grouping or times at which
364 data are measured [Gill, 2002]. In addition, a common application of hierarchical Bayes analysis in
365 the Probabilistic Risk Assessment (PRA) of nuclear power plants has been as a model of variability
366 among data sources, for example variability in Emergency Diesel Generator (EDG) performance
367 across different plants, or across time [Siu and Kelly, 1998; Atwood et al., 2003; Kelly and Smith,
368 2009]. Finally, similar analogy can be made for the collected measurements from a structure under
369 different ambient and environmental conditions. This framework has been recently implemented for
370 uncertainty quantification applications in structural dynamics [Behmanesh et al., 2015; Ballesteros
371 et al., 2014]. Although hierarchical Bayes can address the issue of imprecise prior information by
372 means of multi-level models, the following conceptual and practical considerations should be made

373 about its applicability: (i) in principle, one could define even higher dimensional models to
374 represent the uncertainty. Some analysts have attempted three- and even four-level models
375 [Jaworska and Aldenberg, 2000], but this hardly seems a workable solution when the complexity of
376 the analysis is the primary problem; (ii) in addition, “if concern about the uncertainty in parameters
377 can in principle drive the analysis to a higher level, one could fall into an insoluble infinite cascade”
378 [Ferson, 2005]; (iii) finally, both first- and second-stage priors, $p^\theta(\theta|\varphi)$ and $p^\varphi(\varphi|\omega)$,
379 respectively, are often of a *particular* functional form, which forces the analyst to make “overly-
380 optimistic” and excessively precise statements about parameter uncertainties, even when the
381 information and data available are scarce and/or vague, i.e., *not sufficient* for assigning a *single*
382 *specific* probability distribution. In such cases, non-probabilistic methods (e.g., intervals, possibility
383 distributions or Dempster-Shafer structures from evidence theory) have been shown to provide
384 reliable and robust results [Ferson et al., 2003; Beer et al., 2013a]. On the basis of considerations
385 (i)-(iii) above and given that the objective of the paper is the comparison of methods for the update
386 of *possibilistic* parameters (of aleatory probability distributions), Hierarchical Bayes approaches are
387 not considered in the present work.

388 **Case study: flood protection risk-based design**

389 The case study deals with the design of a protection dike in a residential area closely located to a
390 river with potential risk of floods. Two issues of concern are: (i) high construction and annual
391 maintenance costs of the dike; (ii) uncertainty in the natural phenomenon of flooding. Then, the
392 different design options must be evaluated within a flooding risk analysis framework accounting for
393 uncertainty.

394 ***The model***

395 The maximal water level of the river (i.e., the output variable of the model, Z_c) is given as a
396 function of several (and some uncertain) parameters (i.e., the inputs to the model) [Limbourg and de
397 Rocquigny, 2010]:

398 $Z_c = f(Q, Z_m, Z_v, K_s, B, L) = f(Y_1, Y_2, Y_3, Y_4, Y_5, Y_6)$ (4)

399 where: $Y_1 = Q$ is the yearly maximal water discharge [m^3/s]; $Y_2 = Z_m$ and $Y_3 = Z_v$ are the riverbed
 400 levels [m asl] at the upstream and downstream parts of the river under investigation, respectively; Y_4
 401 $= K_s$ is the Strickler friction coefficient; $Y_5 = B$ and $Y_6 = L$ are the width and length of the river
 402 part [m], respectively. Quantities $Y_5 = B$ ($= 300\text{m}$) and $Y_6 = L$ ($= 5000\text{m}$) are *constant* parameters,
 403 whereas quantities $Y_1 = Q$, $Y_2 = Z_m$, $Y_3 = Z_v$, $Y_4 = K_s$ are *uncertain* variables.

404 ***The input variables: physical description and representation of the associated***
 405 ***uncertainty***

406 The $n = 4$ input variables Y_i , $i = 1, 2, 3, 4$, are affected by aleatory and epistemic uncertainties. The
 407 aleatory part of the uncertainty is described by probability distributions of defined shape. The
 408 parameters of the probability distributions describing the aleatory uncertainty are themselves
 409 affected by epistemic uncertainty and represented in terms of possibility distributions.

410 **The yearly maximal water flow, $Y_1 = Q$**

411 The aleatory uncertainty in the yearly maximal water flow $Y_1 = Q$ is well described by a Gumbel
 412 probability distribution $p^q(q|\gamma, \delta) = Gum(\gamma, \delta) = \frac{1}{\delta} \exp\left[-\exp\left(\frac{\gamma-q}{\delta}\right)\right] \exp\left[\frac{\gamma-q}{\delta}\right]$ [Limbourg and
 413 de Rocquigny, 2010]. The extreme physical bounds on variable Q are [Limbourg and de
 414 Rocquigny, 2010]:

- 415 - $Q_{\min} = 10\text{m}^3/\text{s}$, which is a typical drought flow level (irrelevant within a flood study);
- 416 - $Q_{\max} = 10000\text{m}^3/\text{s}$, which is three times larger than the maximal flood ever occurred.

417 When Approach A is used, the prior possibility distributions $\pi^\gamma(\gamma)$ and $\pi^\delta(\delta)$ for the
 418 epistemically-uncertain parameters γ and δ are subjectively chosen as triangular functions $TR(a_\gamma,$
 419 $c_\gamma, b_\gamma)$ and $TR(a_\delta, c_\delta, b_\delta)$, respectively, with cores (i.e., preferred or most likely values) $c_\gamma = 955\text{m}^3/\text{s}$
 420 and $c_\delta = 600\text{m}^3/\text{s}$, and supports $[a_\gamma, b_\gamma] = [869, 1157]\text{m}^3/\text{s}$ and $[a_\delta, b_\delta] = [455, 660]\text{m}^3/\text{s}$,

421 respectively. When Approach B is employed, the fictitious prior FPDFs $\tilde{p}^\gamma(\gamma)$ and $\tilde{p}^\delta(\delta)$ are
422 subjectively chosen as normal probability distributions $p^\gamma(\gamma) = N(\mu_\gamma, \sigma_\gamma)$ and $p^\delta(\delta) =$
423 $N(\mu_\delta, \sigma_\delta)$, respectively, with possibilistic parameters $\mu_\gamma, \sigma_\gamma, \mu_\delta$ and σ_δ : the characteristics of the
424 corresponding triangular possibility distributions $\pi^{\mu_\gamma}(\mu_\gamma)$, $\pi^{\sigma_\gamma}(\sigma_\gamma)$, $\pi^{\mu_\delta}(\mu_\delta)$ and $\pi^{\sigma_\delta}(\sigma_\delta)$ are
425 summarized in Table 2 for brevity. The Bayesian update of these uncertainty representations (based
426 on prior subjective knowledge) is realized with the aid of a vector $\mathbf{y}_1 = [y_{11}, y_{12}, \dots, y_{1k}, \dots, y_{1D_1}]$ of
427 $D_1 = 149$ (independent and identically distributed – iid) values of the annual maximal flow of the
428 river, i.e., $\mathbf{y}_1 = \mathbf{q} = [q_1, q_2, \dots, q_k, \dots, q_{149}]$. The point estimates for γ and δ obtained by the
429 classical, purely probabilistic Maximum Likelihood Estimation (MLE) method are $\hat{\gamma}^{MLE} = 1013.21$
430 m^3/s and $\hat{\delta}^{MLE} = 558.21 \text{ m}^3/\text{s}$, respectively.

431 **The upstream riverbed level, $Y_2 = Z_m$**

432 The aleatory part of the uncertainty in the upstream riverbed level $Y_2 = Z_m$ is represented by a
433 normal distribution, i.e., $Z_m \sim p^{Z_m}(z_m | \mu_{Z_m}, \sigma_{Z_m}) = N(\mu_{Z_m}, \sigma_{Z_m})$ [Limbourg and de Rocquigny,
434 2010]. This distribution is truncated at the minimum and maximum physical bounds on Z_m , i.e.,
435 $Z_{m,\min} = 53.5 \text{ m}$ (given by plausible lower geomorphologic limits to erosion) and $Z_{m,\max} = 57 \text{ m}$
436 (given by plausible upper geomorphologic limits to sedimentation), respectively. In Approach A,
437 the prior possibility distributions $\pi^{\mu_{Z_m}}(\mu_{Z_m})$ and $\pi^{\sigma_{Z_m}}(\sigma_{Z_m})$ for μ_{Z_m} and σ_{Z_m} are chosen as
438 triangular functions (Table 2). In Approach B, the fictitious prior FPDFs $\tilde{p}^{\mu_{Z_m}}(\mu_{Z_m})$ and $\tilde{p}^{\sigma_{Z_m}}(\sigma_{Z_m})$
439 are chosen as normal PDFs $p^{\mu_{Z_m}}(\mu_{Z_m}) = N(\mu_{\mu_{Z_m}}, \sigma_{\mu_{Z_m}})$ and $p^{\sigma_{Z_m}}(\sigma_{Z_m}) = N(\mu_{\sigma_{Z_m}}, \sigma_{\sigma_{Z_m}})$ with
440 parameters described by the triangular possibilistic functions $\pi^{\mu_{\mu_{Z_m}}}(\mu_{\mu_{Z_m}})$, $\pi^{\sigma_{\mu_{Z_m}}}(\sigma_{\mu_{Z_m}})$,
441 $\pi^{\mu_{\sigma_{Z_m}}}(\mu_{\sigma_{Z_m}})$ and $\pi^{\sigma_{\sigma_{Z_m}}}(\sigma_{\sigma_{Z_m}})$ reported in Table 2. The Bayesian update of these uncertainty
442 representations is carried out using a vector $\mathbf{y}_2 = [y_{21}, y_{22}, \dots, y_{2k}, \dots, y_{2D_2}]$ of $D_2 = 29$ (iid) values

443 of the upstream riverbed level, i.e., $\mathbf{y}_2 = \mathbf{z}_m = [z_{m,1}, z_{m,2}, \dots, z_{m,k}, \dots, z_{m,29}]$. The point estimates $\hat{\mu}_{Z_m}^{MLE}$
 444 and $\hat{\sigma}_{Z_m}^{MLE}$ for μ_{Z_m} and σ_{Z_m} obtained by the MLE method are $\hat{\mu}_{Z_m}^{MLE} = 50.19$ m and $\hat{\sigma}_{Z_m}^{MLE} = 0.38$ m,
 445 respectively.

446 **The downstream riverbed level, $Y_3 = Z_v$**

447 As for $Y_2 = Z_m$, the aleatory part of the uncertainty in the downstream riverbed level $Y_3 = Z_v$ is
 448 represented by a normal distribution, i.e., $Z_v \sim p^{Z_v}(z_v | \mu_{Z_v}, \sigma_{Z_v}) = N(\mu_{Z_v}, \sigma_{Z_v})$, truncated at
 449 $Z_{v,\min} = 48$ m and $Z_{v,\max} = 51$ m. As before, the prior possibility distributions $\pi^{\mu_{Z_v}}(\mu_{Z_v})$ and
 450 $\pi^{\sigma_{Z_v}}(\sigma_{Z_v})$ used in Approach A are triangular functions (Table 2) and the FPDFs $\tilde{p}^{\mu_{Z_v}}(\mu_{Z_v})$ and
 451 $\tilde{p}^{\sigma_{Z_v}}(\sigma_{Z_v})$ employed in Approach B are normal PDFs $p^{\mu_{Z_v}}(\mu_{Z_v}) = N(\mu_{\mu_{Z_m}}, \sigma_{\mu_{Z_m}})$ and $p^{\sigma_{Z_v}}(\sigma_{Z_v}) =$
 452 $N(\mu_{\sigma_{Z_v}}, \sigma_{\sigma_{Z_v}})$ with parameters described by the triangular possibility distributions $\pi^{\mu_{Z_v}}(\mu_{Z_v})$,
 453 $\pi^{\sigma_{Z_v}}(\sigma_{Z_v})$, $\pi^{\mu_{\sigma_{Z_v}}}(\mu_{\sigma_{Z_v}})$ and $\pi^{\sigma_{\sigma_{Z_v}}}(\sigma_{\sigma_{Z_v}})$ of Table 2. These representations are updated by means of
 454 a vector $\mathbf{y}_3 = [y_{31}, y_{32}, \dots, y_{3k}, \dots, y_{3D_3}]$ of $D_3 = 29$ (iid) values of the downstream riverbed level,
 455 i.e., $\mathbf{y}_3 = \mathbf{z}_v = [z_{v,1}, z_{v,2}, \dots, z_{v,k}, \dots, z_{v,29}]$. The MLE estimates of the parameters are $\hat{\mu}_{Z_v}^{MLE} = 55.03$ m
 456 and $\hat{\sigma}_{Z_v}^{MLE} = 0.45$ m, respectively.

457 **The Strickler friction coefficient, $Y_4 = K_s$**

458 The Strickler friction coefficient $Y_4 = K_s$ is the most critical source of uncertainty because it is
 459 usually a simplification of a complex hydraulic model. The absolute physical limits of K_s are 5 and
 460 60, respectively [Limbourg and de Rocquigny, 2010]:

- 461 - $K_s < 5$ corresponds to an “extremely sinuous shape of the canal, with large dents and strong
 462 vegetation”;
- 463 - $K_s = 60$ corresponds to a “canal with smoothest earth surface, rectilinear, without any
 464 vegetation”.

465 The friction coefficient K_s is affected by random events modifying the river status (e.g., erosion,
466 sedimentation, ...): the corresponding variability is typically described by a normal distribution, i.e.,
467 $K_s \sim p^{K_s}(k_s | \mu_{K_s}, \sigma_{K_s}) = N(\mu_{K_s}, \sigma_{K_s})$ [Limbourg and de Rocquigny, 2010]. However, the
468 parameters of this normal distribution are difficult to estimate because data can only be obtained
469 through “indirect calibration characterized by significant uncertainty” [Limbourg and de
470 Rocquigny, 2010]: the uncertainty in these parameters is described by possibility distributions.
471 Details about the possibilistic functions $\pi^{\mu_{K_s}}(\mu_{K_s})$ and $\pi^{\sigma_{K_s}}(\sigma_{K_s})$ and the FPDFs $\tilde{p}^{\mu_{K_s}}(\mu_{K_s})$ and
472 $\tilde{p}^{\sigma_{K_s}}(\sigma_{K_s})$ used to represent the a priori knowledge on μ_{K_s} and σ_{K_s} in Approaches A and B,
473 respectively, are reported in Table 2. The Bayesian revision of these a priori representations is
474 performed by means of a vector $\mathbf{y}_4 = [y_{41}, y_{42}, \dots, y_{4k}, \dots, y_{4D_4}]$ of $D_4 = 5$ (iid) values of the
475 Strickler friction coefficient, i.e., $\mathbf{y}_4 = \mathbf{k}_s = [k_{s,1}, k_{s,2}, \dots, k_{s,k}, \dots, k_{s,5}]$. The MLE estimates of the
476 parameters are $\hat{\mu}_{K_s}^{MLE} = 27.8$ and $\hat{\sigma}_{K_s}^{MLE} = 5.26$, respectively.

477
478 **Table 2.**
479

480 Results

481 In order to simplify the notation, in what follows let θ be one of the uncertain parameters of the
482 PDFs of $Y_1 = Q$, $Y_2 = Z_m$, $Y_3 = Z_v$ and $Y_4 = K_s$, i.e., $\theta = \gamma, \delta, \mu_{Z_m}, \sigma_{Z_m}, \mu_{Z_v}, \sigma_{Z_v}, \mu_{K_s}$ or σ_{K_s} . Fig.
483 3 illustrates the possibility distributions of the epistemically-uncertain parameters of the aleatory
484 PDFs $p^Q(q | \gamma, \delta)$ (top row), $p^{Z_m}(z_m | \mu_{Z_m}, \sigma_{Z_m})$ (middle-top row), $p^{Z_v}(z_v | \mu_{Z_v}, \sigma_{Z_v})$ (middle-bottom
485 row) and $p^{K_s}(k_s | \mu_{K_s}, \sigma_{K_s})$ (bottom row) of the uncertain input variables $Y_1 = Q$, $Y_2 = Z_m$, $Y_3 = Z_v$
486 and $Y_4 = K_s$, respectively, of the model of the previous Section: in particular, the prior possibility
487 distributions $\pi^\theta(\theta)$ ($= \pi^\gamma(\gamma), \pi^\delta(\delta), \pi^{\mu_{Z_m}}(\mu_{Z_m}), \pi^{\sigma_{Z_m}}(\sigma_{Z_m}), \pi^{\mu_{Z_v}}(\mu_{Z_v}), \pi^{\sigma_{Z_v}}(\sigma_{Z_v}), \pi^{\mu_{K_s}}(\mu_{K_s})$
488 and $\pi^{\sigma_{K_s}}(\sigma_{K_s})$) are shown as solid lines, whereas the *marginal* posterior possibility distributions

489 $\pi^\theta(\theta | \mathbf{y})$ ($= \pi^\gamma(\gamma | \mathbf{q}), \pi^\delta(\delta | \mathbf{q}), \pi^{\mu_{Z_m}}(\mu_{Z_m} | \mathbf{z}_m), \pi^{\sigma_{Z_m}}(\sigma_{Z_m} | \mathbf{z}_m), \pi^{\mu_{Z_v}}(\mu_{Z_v} | \mathbf{z}_v), \pi^{\sigma_{Z_v}}(\sigma_{Z_v} | \mathbf{z}_v),$
490 $\pi^{\mu_{K_s}}(\mu_{K_s} | \mathbf{k}_s)$ and $\pi^{\sigma_{K_s}}(\sigma_{K_s} | \mathbf{k}_s)$) obtained by Approaches A and B using $D_1 = 149, D_2 = 29, D_3 =$
491 29 and $D_4 = 5$ pieces of data are shown in dashed and dot-dashed lines, respectively; the point
492 estimates $\hat{\theta}^{MLE}$ ($= \hat{\gamma}^{MLE}, \hat{\delta}^{MLE}, \hat{\mu}_{Z_m}^{MLE}, \hat{\sigma}_{Z_m}^{MLE}, \hat{\mu}_{Z_v}^{MLE}, \hat{\sigma}_{Z_v}^{MLE}, \hat{\mu}_{K_s}^{MLE}$ and $\hat{\sigma}_{K_s}^{MLE}$) produced by the
493 classical MLE method are also shown for comparison (dots).

494

495

Fig. 3.

496

497 From a mere visual and qualitative inspection of Fig. 3 it can be seen that both approaches are
498 suitable for revising the prior possibility distributions (based on a priori purely subjective
499 knowledge) by means of empirical data. In particular, it is evident that: (i) the most likely (i.e.,
500 preferred) values c_θ of the epistemically-uncertain parameters (i.e., those values in correspondence
501 of which the possibility function equals 1) are moved towards the MLE estimates $\hat{\theta}^{MLE}$ in all the
502 cases considered; (ii) the area S_θ underlying the corresponding possibility distributions is
503 significantly reduced: noting that this area is related to the imprecision in the knowledge of the
504 possibilistic parameter (i.e., the larger the area, the higher the imprecision), it can be concluded that
505 both approaches succeed in reducing the epistemic uncertainty. With respect to that, Table 3 reports
506 the most likely values c_θ and the areas S_θ underlying the (marginal) possibility distributions of the
507 uncertain parameter θ ($= \gamma, \delta, \mu_{Z_m}, \sigma_{Z_m}, \mu_{Z_v}, \sigma_{Z_v}, \mu_{K_s}$ and σ_{K_s}) before and after the Bayesian
508 update performed by Approaches A and B; the point estimates $\hat{\theta}^{MLE}$ obtained by the classical MLE
509 method are also reported for completeness. In addition, in order to perform a quantitative
510 comparison between the methods, two indicators are defined:

- 511 i. the relative absolute distance d_θ^{MLE} between the (posterior) most likely value c_θ of parameter
512 θ and the corresponding MLE estimate $\hat{\theta}^{MLE}$, i.e.:

513
$$d_{\theta}^{MLE} = \frac{|c_{\theta} - \hat{\theta}^{MLE}|}{\hat{\theta}^{MLE}}. \quad (5)$$

514 Obviously, the lower is d_{θ}^{MLE} the closer is the most likely value c_{θ} to the MLE estimate
 515 $\hat{\theta}^{MLE}$, i.e., the higher is the *strength* of the approach in updating the prior possibilistic
 516 distribution on the basis of newly available experimental evidence;

517 ii. the percentage relative difference R_{θ} between the areas underlying the possibility
 518 distribution of parameter θ before and after the Bayesian update, namely S_{θ}^{Prior} and $S_{\theta}^{Posterior}$,
 519 respectively:

520
$$R_{\theta} = \frac{S_{\theta}^{Prior} - S_{\theta}^{Posterior}}{S_{\theta}^{Prior}} \cdot 100. \quad (6)$$

521 In this case, the higher is R_{θ} , the higher is the reduction in the area (i.e., in the epistemic
 522 uncertainty) and, thus, the higher is the “updating strength” of the approach.

523
 524
 525

Table 3.

526 It is evident that the *strength* of Approach A in moving the most likely values c_{θ} towards the
 527 corresponding MLE estimates $\hat{\theta}^{MLE}$ is *always higher* than that of Approach B. Actually, the values
 528 of d_{θ}^{MLE} (5) produced by Approach A are 1.58-5.24 times *lower* than those generated by Approach
 529 B for all the parameters.

530 From the analysis of quantitative indicator R_{θ} (6) it can be seen that both methods succeed in
 531 reducing the area underlying the possibility distributions of the uncertain parameters: in particular,
 532 the percentage reduction R_{θ} ranges between 8.33% and 30.49% for approach A and between 3.89%
 533 and 33.01% for Approach B. In addition, as expected, the strength of *both* approaches in reducing
 534 epistemic uncertainty *decreases* with the *size* of the data set used to perform the Bayesian update.
 535 For example, the area S_{γ} underlying the possibility distribution of γ (Fig. 3, top left column) is
 536 reduced by 30.49–33.01% with the aid of a *large* data set of size $D_1 = 149$; on the contrary, the area

537 $S_{\sigma_{K_s}}$ underlying the possibility distribution of σ_{K_s} (Fig. 3, bottom right column) is reduced only by
538 3.89–8.76% by means of $D_4 = 5$ pieces of data. Moreover, it is interesting to note that the strength
539 of Approach B in reducing epistemic uncertainty is *slightly higher* than that of Approach A *only*
540 when the amount of available data is quite *large* (i.e., in the revision of the possibility distributions
541 of parameters γ and δ of the PDF of $Y_1 = Q$, by means of $D_1 = 149$ pieces of data): actually, the
542 values of R_γ and R_δ range within 25.56-30.49% and 28.74-33.01% for Approaches A and B,
543 respectively. In all the other cases, the power of Approach A in reducing epistemic uncertainty is
544 *higher* than that of Approach B and this difference becomes more and *more evident* as the size of
545 the data set *decreases*: in particular, for medium-sized data sets (i.e., in the revision of the
546 possibility distributions of the parameters μ_{Z_m} , σ_{Z_m} , μ_{Z_v} , σ_{Z_v} of the PDFs of $Y_2 = Z_m$ and $Y_3 = Z_v$
547 with $D_2 = D_3 = 29$ pieces of data) the values of R_θ (6) produced by Approaches A and B range
548 within 8.33-24.38% and 4.17-22.00%, respectively; instead, for small-sized data sets (i.e., in the
549 revision of the possibility distributions of the parameters μ_{K_s} and σ_{K_s} of the PDF of $Y_4 = K_s$ with
550 $D_4 = 5$ pieces of data) the values of R_θ produced by Approaches A and B range within 8.76-16.28%
551 and 3.89-10.08%, respectively. This is particularly evident in the estimation of the standard
552 deviation σ_{K_s} of K_s (Fig. 3, bottom right column): on one side, the posterior distribution produced by
553 the hybrid approach (B) seems not to be influenced by the revision process (actually, the most likely
554 value of the parameter, $c_{\sigma_{K_s}} = 6.72$, and the area underlying the corresponding posterior possibility
555 distribution, $S_{\sigma_{K_s}} = 3.95$, are quite close to those of the prior, i.e., 6.89 and 4.11, respectively); on
556 the other side, the posterior distribution generated by the purely possibilistic approach (A) is almost
557 centered on the point estimates obtained by the MLE method and the corresponding area is reduced
558 by about 9%.
559
560 Finally, it has to be remarked that for the sake of simplicity the quantitative analyses above have
561 been performed only on the *marginal* posterior possibility distributions of the uncertain parameters

562 $(\pi^\gamma(\gamma|\mathbf{q}), \pi^\delta(\delta|\mathbf{q}), \pi^{\mu_{Z_m}}(\mu_{Z_m}|\mathbf{z}_m), \pi^{\sigma_{Z_m}}(\sigma_{Z_m}|\mathbf{z}_m), \pi^{\mu_{Z_v}}(\mu_{Z_v}|\mathbf{z}_v), \pi^{\sigma_{Z_v}}(\sigma_{Z_v}|\mathbf{z}_v), \pi^{\mu_{K_s}}(\mu_{K_s}|\mathbf{k}_s)$
563 and $\pi^{\sigma_{K_s}}(\sigma_{K_s}|\mathbf{k}_s))$; in fact, as highlighted before, the posterior possibility distributions of the
564 uncertain parameters are *multi-dimensional* functions (and the posterior estimates of the parameters
565 are *dependent*). Only for illustration purposes, Fig. 4 shows the α -cuts of level $\alpha = 0.05, 0.50$ and
566 0.95 of the joint posterior possibility distributions $\pi^{\gamma,\delta}(\gamma,\delta|\mathbf{q}), \pi^{\mu_{Z_m},\sigma_{Z_m}}(\mu_{Z_m},\sigma_{Z_m}|\mathbf{z}_m),$
567 $\pi^{\mu_{Z_v},\sigma_{Z_v}}(\mu_{Z_v},\sigma_{Z_v}|\mathbf{z}_v)$ and $\pi^{\mu_{K_s},\sigma_{K_s}}(\mu_{K_s},\sigma_{K_s}|\mathbf{k}_s)$ of the parameters of the PDFs of $Y_1 = Q$ (top right
568 column), $Y_2 = Z_m$ (top left column), $Y_3 = Z_v$ (bottom left column) and $Y_4 = K_s$ (bottom right column),
569 respectively, produced by Approaches A (solid lines) and B (dashed lines).

570

571

Fig. 4.

572

573 The visual inspection of Fig. 4 confirms the results obtained by the quantitative analysis carried out
574 on the marginal distributions. For example, it can be seen that the areas of the α -cuts of level $\alpha =$
575 0.95 of $\pi^{\mu_{Z_v},\sigma_{Z_v}}(\mu_{Z_v},\sigma_{Z_v}|\mathbf{z}_v)$ and $\pi^{\mu_{K_s},\sigma_{K_s}}(\mu_{K_s},\sigma_{K_s}|\mathbf{k}_s)$ produced by Approach A are consistently
576 smaller than those generated by Approach B; the α -cuts of $\pi^{\mu_{Z_m},\sigma_{Z_m}}(\mu_{Z_m},\sigma_{Z_m}|\mathbf{z}_m)$ produced by
577 Approaches A and B are comparable in size, whereas the area of the α -cut of $\pi^{\gamma,\delta}(\gamma,\delta|\mathbf{q})$ obtained
578 by Approach B is slightly lower than the one produced by Approach A.

579

580 In order to show the effect that the reduction of the epistemic uncertainty in the distribution
581 parameters has on the uncertain input variables $Y_1 = Q, Y_2 = Z_m, Y_3 = Z_v$ and $Y_4 = K_s$, Fig. 5 reports
582 the upper and lower CDFs, $\bar{F}^{Y_j}(y_j)$ and $\underline{F}^{Y_j}(y_j)$ (i.e., the plausibility and belief functions,
583 $Pl^{Y_j}((-\infty, y_j])$ and $Bel^{Y_j}((-\infty, y_j])$, respectively), $j = 1, 2, 3, 4$, of $Y_1 = Q$ (top left column), $Y_2 = Z_m$
584 (top right column), $Y_3 = Z_v$ (bottom left column) and $Y_4 = K_s$ (bottom right column) before (solid
585 lines) and after the Bayesian update performed by Approaches A (dashed lines) and B (dot-dashed

586 lines). The calculation of CDFs is of great importance in risk assessment since they summarize the
587 uncertainty “contained” in the variables of interest.

588

589

Fig. 5

590

591 Obviously, the gap between the plausibility and belief functions $Pl^{Y_j}((-\infty, y_j]) = \bar{F}^{Y_j}(y_j)$ and
592 $Bel^{Y_j}((-\infty, y_j]) = \underline{F}^{Y_j}(y_j)$, $j = 1, 2, 3, 4$, is larger before the Bayesian update in all the cases
593 considered: in particular, the ‘prior’ CDFs (solid lines) completely envelop the ‘posterior’ ones
594 (dashed and dot-dashed lines). This larger gap is explained by the larger area contained under the
595 possibility distribution functions of the corresponding epistemically-uncertain parameters (actually,
596 as highlighted before, the larger the area, the higher the imprecision in the knowledge of the
597 possibilistic parameters).

598 Then, in order to provide a fair and quantitative comparison between the two approaches adopted,
599 the intervals $\left[\left[\bar{F}^{Y_j} \right]^{-1}(\beta), \left[\underline{F}^{Y_j} \right]^{-1}(\beta) \right]$ for the β -100-th percentiles Y_j^β , $\beta = 0.05, 0.50$ and 0.95 , of the
600 variables Y_j , $j = 1, 2, 3, 4$, are computed (Table 4). For example, analyzing Q^β , $\beta = 0.05, 0.50$ and
601 0.95 , it can be seen that the width of the intervals is reduced by 30.54–40.34% and 29.88–43.29%
602 by Approaches A and B, respectively: coherently with the results reported in Table 3, when a *large*
603 data set is available (i.e., $D_1 = 149$ in this case) the strength of Approach B in reducing the
604 epistemic uncertainty is *slightly* (i.e., 2-3%) *higher* than that of Approach A. On the contrary,
605 analyzing K_s^β , $\beta = 0.05, 0.50$ and 0.95 , it is evident that the width of the intervals is reduced of
606 25.30-30.80% and 5.74-19.45% by Approaches A and B, respectively: as highlighted before (Table
607 3), when a *small* data set is available (i.e., $D_4 = 5$ in this case) the power of Approach A in reducing
608 epistemic uncertainty is *consistently* (i.e., 15-20%) *higher* than that of Approach B. This is
609 confirmed *also* by the analysis of the quantiles of $Y_2 = Z_m$ and $Y_3 = Z_v$ (see Table 4).

610

611

Table 4.

612

613 Comparable conclusions can be drawn by the analysis of the upper and lower CDFs $\bar{F}^{Z_c}(z_c)$ and
614 $\underline{F}^{Z_c}(z_c)$ (i.e., the plausibility and belief functions, $Pl^{Z_c}((-\infty, z_c])$ and $Bel^{Z_c}((-\infty, z_c])$,
615 respectively), of the maximal water level of the river Z_c (i.e., the model output) obtained before
616 (Fig. 6, solid lines) and after the Bayesian update performed by Approaches A (Fig. 6, dashed lines)
617 and B (Fig. 6, dot-dashed lines). Notice that the distributions for Z_c have been obtained by
618 propagating the two-level mixed probabilistic and possibilistic uncertainty through the
619 mathematical model by means of a hybrid Monte Carlo (MC) and Fuzzy Interval Analysis (FIA)
620 approach. This method combines the MC technique [Kalos and Withlock, 1986] with the extension
621 principle of fuzzy set theory [Guyonnet et al., 2003; Kentel and Aral, 2004 and 2007] in two
622 hierarchical, repeated steps [Baudrit et al., 2008; Kentel and Aral, 2005; Moller and Beer, 2004 and
623 2008; Moller et al., 2003 and 2006; Pedroni and Zio, 2012; Pedroni et al., 2013]: the reader is
624 referred to the cited references for details. Again, in order to provide a fair comparison between the
625 two approaches employed, proper quantitative indicators are computed. The final goal of the case
626 study presented in the previous Section is to determine (i) the dike level necessary to guarantee a
627 given flood return period or (ii) the flood risk for a given dike level. With respect to issue (i) above,
628 the quantity of interest that is most relevant to the decision maker is the $\beta \cdot 100\%$ -th quantile of Z_c
629 (i.e., Z_c^β): this corresponds to the yearly maximal water level with a $\beta \cdot 100$ -year return period. With
630 respect to issue (ii) above, the quantity of interest that is most relevant to the decision maker is the
631 probability that the maximal water level of the river Z_c exceeds a given threshold z^* , i.e.,
632 $P[Z_c \geq z_c^*]$: in the present paper, $z_c^* = 55.5$ m (Table 5). Analyzing the intervals $[\bar{F}^{Z_c}]^{-1}(\beta)$,
633 $[\underline{F}^{Z_c}]^{-1}(\beta)$ for the percentiles Z_c^β , $\beta = 0.05, 0.50$ and 0.95 , and the intervals
634 $[1 - \bar{F}^{Z_c}(z_c^*), 1 - \underline{F}^{Z_c}(z_c^*)]$ for the exceedance probability $P[Z_c > z_c^*]$, it can be seen that their width is
635 reduced of 28.14–38.63% and 16.88–29.23% by Approaches A and B, respectively (i.e., the

636 strength of Approach A in reducing epistemic uncertainty is about 4-11% *higher* than that of
637 Approach B).

638

639

Fig. 6.

640

641

Table 5.

642

643 Several considerations are in order with respect to the results obtained. When the Bayesian update is
644 performed based on a data set of *large* size (e.g., > 100 in this case), the difference in the behavior
645 of the two approaches is quite *low*. This demonstrates that although the two methods are
646 conceptually and algorithmically quite different, in presence of a “*strong*” experimental evidence
647 they produce “*coherent*” results (i.e., posterior possibility distributions that bear the *same* overall
648 “uncertainty content”): this is a fair outcome since the results provided by the two methods are
649 expected to be more and more *similar* (i.e., more and more coherent with the experimental
650 evidence) as the size of the data set *increases* (experts and practitioners may find a similarity and
651 parallelism between these results and those obtained in purely probabilistic, graphical Bayesian
652 models [Gelman et al., 2004]; in particular, see references concerning approaches used to “borrow
653 strength” in (Hierarchical) Bayesian analyses [Atwood et al., 2003; Kelly and Smith, 2009 and
654 2011]). Instead, when the Bayesian update is performed on a data set of *small-medium* size (e.g., \approx
655 5-30 in this case), the *strength* of Approach A in reducing epistemic uncertainty is *significantly*
656 *higher* than that of Approach B. This can be explained as follows. In Approach A the *purely*
657 *possibilistic* likelihood (i.e., the function “containing” the experimental evidence available) has a
658 *direct* and strong influence on the *purely possibilistic* prior (actually, they are directly multiplied in
659 (1)); on the contrary, in Approach B the *purely probabilistic* likelihood has a direct influence *only*
660 on the (fictitious) *probabilistic* distributions that are superimposed onto the prior possibilistic
661 parameters: this “artificial” procedure may in practice “soften” (i.e., reduce) the effect of the newly
662 available information (i.e., of the data) in the revision of the possibilistic priors (obviously, this

663 effect is expected to be more evident if the amount of data, i.e., the *strength* of the experimental
664 evidence, is *small*). In such cases, embracing one method instead of the other may significantly
665 change the outcome of a decision making process in a risk assessment problem involving
666 uncertainties: this is of paramount importance in systems that are critical from the safety view point,
667 e.g., in the nuclear, aerospace, chemical and environmental fields.

668 Finally, it is absolutely important to acknowledge that even if the strength of one method (i.e., A in
669 this case) in reducing epistemic uncertainty is higher than that of the other one (i.e., B in this case),
670 this does *not necessarily* imply that one method is “*better*” or “*more effective*” than the other
671 *overall*. Actually, if on one side a consistent reduction in the epistemic uncertainty is in general
672 *desirable* in decision making processes related to risk assessment problems (since it significantly
673 *increases* the analyst’s *confidence* in the decisions), on the other side this reduction must be
674 *coherent* with the amount of *information available*. In this view, an objection may arise in the
675 present case: is the *remarkable* strength of Approach A in reducing epistemic uncertainty (with very
676 few pieces of data) *fully justified* by such a small amount of data? In other words, is this
677 considerable reduction of epistemic uncertainty coherent with the strength of the experimental
678 evidence or is it too optimistic? These issues will be thoroughly discussed in the following
679 dedicated Section.

680
681 Finally, in addition to the strength of the approaches in revising the (prior) possibilistic description
682 of the uncertain parameters of aleatory variables, also the *computational time* associated to the
683 methods has to be taken into account. Table 6 reports the computational time t_{comp} required by the
684 Bayesian update of all the parameters of the PDFs of $Y_1 = Q$, $Y_2 = Z_m$, $Y_3 = Z_v$ and $Y_4 = K_s$ performed
685 by Approach A and by Approach B (with $T = 100$ repetitions of the purely probabilistic Bayes’
686 theorem for each of the $N_\alpha = 21$ α -cuts analyzed). Obviously, the computational time required by
687 Approach B is approximately $T \cdot N_\alpha$ (i.e., $T \cdot N_\alpha = 100 \cdot 21 = 2100$ in this case) times larger than that of
688 Approach A. On the other hand, notice that since Approach B is based on several *repetitions* of the

689 purely probabilistic Bayes' theorem, if possible, *parallelization* could be in principle employed to
690 reduce the associated computational cost.

691
692
693

Table 6.

694 Discussion of the results and comparison of the approaches

695 The results of the comparisons performed in the previous Sections can be summarized as follows:

- 696 • both methods succeed in updating the possibilistic description of the epistemically-uncertain
697 parameters of (aleatory) probability distributions by means of data. This is highlighted by
698 the fact that in most cases the posterior possibility distributions produced by the two
699 approaches are significantly different from the corresponding priors. In particular:
 - 700 ▪ the most likely values of the parameters (i.e., those values in correspondence of
701 which the possibility function equals 1) are moved towards the point estimates of the
702 parameters obtained by the classical, purely probabilistic MLE method;
 - 703 ▪ the area underlying the posterior possibility distributions is consistently lower than
704 that of the priors: since this area is related to the imprecision in the knowledge of the
705 possibilistic parameter (i.e., the larger the area, the higher the imprecision), it can be
706 concluded that both approaches succeed in reducing the epistemic uncertainty in the
707 possibilistic parameters of the aleatory probability distributions. This is also
708 confirmed by the reduction of the gap between the upper and lower CDFs (i.e., the
709 plausibility and belief functions) of the corresponding aleatory variables;
- 710 • when the Bayesian update is performed using a data set of *large* size (e.g., > 100), the
711 *strength* of the two approaches in reducing the epistemic uncertainty is quite *similar*. By
712 way of example, in the case study considered Approaches A and B reduce the areas
713 underlying the possibility distributions of the uncertain parameters by 25.56-30.49% and
714 28.74-33.01%, respectively. This demonstrates that although the two methods are

715 conceptually and algorithmically quite different, in presence of a “strong” experimental
716 evidence they produce “coherent” results (i.e., possibility distributions that bear almost the
717 *same* overall “uncertainty content”): this is a fair outcome since the it is “desired” that both
718 methods provide results that become more and more similar (i.e., that become more and
719 more coherent with the experimental evidence) as the size of the data set increases;

- 720 • the strength of the purely possibilistic approach (A) in reducing epistemic uncertainty is
721 *consistently higher* than that of the hybrid one (B) in presence of *medium-* and *small-*sized
722 data sets (e.g., $\approx 5-30$) (which is often the case in the risk analysis of complex safety-critical
723 systems). For example, the width of the intervals for the quantiles of the variables of interest
724 is reduced by 25.30-30.80% and 5.74-19.45% by Approaches A and B, respectively. This
725 significantly different behavior is explained by the fact that in Approach A the (purely
726 possibilistic) likelihood has an immediate and strong influence on the (purely possibilistic)
727 prior (i.e., they are directly multiplied); on the contrary, in Approach B the (purely
728 probabilistic) likelihood has a direct influence *only* on the (fictitious) probabilistic function
729 that is superimposed onto the possibilistic parameter (subject to the Bayesian update): this
730 “artificial” procedure could in practice weaken the effect of the newly available information
731 (i.e., the data) in the revision of the possibilistic prior (this effect is expected to be more
732 evident if the amount of data, i.e., the *strength* of the experimental evidence, is small).

733 As highlighted above, the fact that the power of Approach A in reducing epistemic
734 uncertainty is higher than that of Approach B does *not necessarily* imply that one method is
735 “*better*” or “*more effective*” than the other *overall*. For example, by hypothesis the
736 remarkable strength of Approach A in reducing epistemic uncertainty by means of very few
737 pieces of data (e.g., 5-30 in this case) might *not* be coherent with the “real” strength of such
738 a scarce experimental evidence and could be accidentally due to some bias in the procedure.
739 With respect to that, it has to be admitted that the uncertainty reduction power of the purely
740 possibilistic approach (A) is strongly dependent on the shape of a *constructed* possibilistic

741 likelihood that could in principle *bias* the analysis. However: (i) in the present paper, this
742 possibilistic function is *very closely related* to the classical, purely probabilistic one (which
743 is theoretically well-grounded) by a simple and direct operation of *normalization* that
744 preserves the “original structure” of the experimental evidence; (ii) the operation of
745 normalization of the probabilistic likelihood finds also some *theoretical* justification in the
746 work by [Denoeux, 2014]; (iii) in general, a probability-to-possibility transformation
747 (properly performed according to the rules of possibility theory) *always* introduces
748 *additional artificial* epistemic uncertainty into the analysis, i.e., it does not artificially reduce
749 it (because it replaces a *single* probabilistic distribution by a *family* of distributions) [Dubois
750 et al., 1993, 2004 and 2008; Flage et al., 2010 and 2013]. On the basis of considerations (i)-
751 (iii) above, it seems unlikely that the purely possibilistic approach (A) may produce results
752 that are dangerously over-optimistic with respect to those of the hybrid one (B). On the other
753 hand, future research should be devoted to the study and development of rigorous,
754 generalized methods for Bayesian model comparison and validation in a *purely possibilistic*
755 framework, in order to complement and strengthen the conclusions drawn by means of the
756 metrics originally introduced in the present manuscript: in this light, techniques from the
757 classical, purely probabilistic field may serve as inspiring references [Gelman et al., 1996;
758 Bayarri and Berger, 1999 and 2000; Johnson Valen, 2004];

759 • the computational time required by the hybrid approach (B) is *consistently higher* than that
760 associated to the purely possibilistic one (A): this is explained by the necessity of repeatedly
761 applying many times the purely probabilistic Bayes’ theorem for *each* α -cut analyzed. More
762 precisely, the application of method A just requires *one single* evaluation of the purely
763 possibilistic Bayes’ formula; on the other hand, approach B entails repeating $T \cdot N_\alpha$ times the
764 classical probabilistic Bayes’ theorem. In this respect, notice that N_α (i.e., the number of α -
765 cuts processed) is typically of the order of 10 [Baudrit et al., 2006], whereas T (i.e., the
766 number of parameter values selected to explore *each* α -cut) cannot be in principle prescribed

767 a priori: however, it needs to be large enough to thoroughly explore *each dimension* of the
768 epistemic parameter space (*at least* 10 values should be selected for each parameter);

- 769 • both methods relies on *assumptions* about either the prior or the likelihood functions: in the
770 purely possibilistic approach (A) the “original” possibilistic prior is employed, but a
771 possibilistic likelihood function has to be constructed (e.g., by probability-possibility
772 transformations or directly from rough experimental data); instead, in the hybrid method (B)
773 the original probabilistic likelihood function is used, but a “fictitious” prior Fuzzy
774 Probability Distribution Function needs to be identified by superimposing an *arbitrarily*
775 *selected* probabilistic PDF onto the “original” possibilistic prior that has to be updated.

776
777 Based on these findings, the advantages and drawbacks of the two approaches are summarized in
778 the following Table 7.

779
780
781

Table 7.

782 **Conclusions**

783 In this paper, we have considered two methods for the Bayesian update of the possibilistic
784 parameters of aleatory probability distributions, with exemplification on a case study concerning the
785 risk-based design of a flood protection dike. The first method considered is based on a purely
786 possibilistic counterpart of the classical probabilistic Bayes’ theorem; the second is a hybrid
787 (probabilistic and possibilistic) method combining Fuzzy Interval Analysis and the classical
788 probabilistic Bayes’ theorem.

789 The findings of the work show that in general adopting different methods may generate different
790 results and possibly different decisions in risk problems involving uncertainties: this is of
791 paramount importance in systems that are critical from the safety viewpoint, e.g., in the nuclear,
792 aerospace, chemical and environmental fields.

793 In particular, on the basis of the results obtained, it seems advisable to suggest the use of the purely
794 possibilistic approach (instead of the hybrid one) for the following reasons: (i) its strength in
795 reducing epistemic uncertainty is significantly higher, in particular when the amount of available
796 data is small: this is important in decision making processes since reducing epistemic uncertainty
797 significantly increases the analyst confidence in the decisions; (ii) the computational time required
798 is consistently lower.

799 However, it has to be remarked that the construction of a possibilistic likelihood required by the
800 purely possibilistic method, although recently tackled in the literature, still represents an issue to be
801 *further* investigated from both the theoretical and practical viewpoint in order to avoid introducing
802 biases in the analysis and to suggest the application of the approach for real risk assessment
803 problems: with respect to that, future research will be devoted to the investigation of additional
804 methods developed to this aim. Also, future studies will be aimed at developing generalized
805 methods for Bayesian model comparison and validation in a *purely possibilistic* framework, in order
806 to complement and strengthen the conclusions drawn by means of the metrics originally introduced
807 in the present manuscript.

808

809

810

References

- 811 Anoop MB, Balaji Rao K, Gopalakrishnan S, 2006. Conversion of probabilistic information into
812 fuzzy sets for engineering decision analysis. *Computers and Structures*, 84, pp. 141-155.
- 813 Apostolakis, G.E., 1990. The concept of probability in safety assessment of technological systems.
814 *Science*, 250, 1359.
- 815 Apostolakis, G.E., S. Kaplan, 1981. Pitfalls in risk calculations. *Reliability Engineering*, Vol. 2,
816 Issue 2, pp. 135-145.
- 817 Atwood, C.L., JL. LaChance, H.F. Martz, D.L Anderson, M. Englehardt, D. Whitehead, T.
818 Wheeler, 2003. *Handbook of Parameter Estimation for Probabilistic Risk Assessment*.
819 NUREG/CR-6823, US Nuclear Regulatory Commission, Washington, DC; SAND2003-
820 3348P, Sandia National Laboratories, Albuquerque, New Mexico.
- 821 Aven, T., 2010. On the Need for Restricting the Probabilistic Analysis in Risk Assessments to
822 Variability. *Risk Analysis*, Vol. 30, Issue 3, pp. 354-360.
- 823 Aven, T., 2011. Interpretations of alternative uncertainty representations in a reliability and risk
824 analysis context. *Reliability Engineering & System Safety*, Volume 96, Issue 3, pp. 353-360.
- 825 Aven, E. Zio, 2011. Some considerations on the treatment of uncertainties in risk assessment for
826 practical decision making, *Reliability Engineering and System Safety*, Vol. 96, Issue 1, pp.
827 64-74.
- 828 Ballesteros, G., Angelikopoulos, P., Papadimitriou, C., Koumoutsakos, P., 2014. Bayesian
829 Hierarchical Models for Uncertainty Quantification in Structural Dynamics. *Vulnerability,*
830 *Uncertainty, and Risk: Quantification, Mitigation, and Management*, ASCE, pp. 1615-1624.
- 831 Baudrit, D. Dubois, 2006. Practical Representations of Incomplete Probabilistic Knowledge,
832 *Computational Statistics & Data Analysis*, Vol. 51, Issue 1, pp. 86-108.

833 Baudrit, I. Couso, D. Dubois, 2007. Joint propagation of probability and possibility in risk analysis:
834 toward a formal framework, *Internat. J. Approx. Reasoning*, Vol. 45, Issue 1, pp. 82–105.

835 Baudrit, D. Dubois, D. Guyonnet, 2006. Joint Propagation and Exploitation of Probabilistic and
836 Possibilistic Information in Risk Assessment. *IEEE Transactions on Fuzzy Systems*, Vol. 14,
837 Issue 5, pp. 593-608.

838 Baudrit, D. Dubois, N. Perrot, 2008. Representing parametric probabilistic models tainted with
839 imprecision, *Fuzzy Sets and System*, Vol. 159, Issue 15, pp.1913-1928.

840 Bayarri MJ, Berger, JO, 1999. Quantifying surprise in the data and model verification. In: Bernardo,
841 JM, et al., editors. *Bayesian statistics 6*. Oxford: Oxford University Press. p.53–82.

842 Bayarri, MJ, Berger, JO, 2000. P-values for composite null models. *J Am Stat Assoc*; 95: 1127–42.

843 Bedford, T., Cooke, R., 2001. *Probabilistic Risk Analysis. Foundations and Methods*. Cambridge:
844 Cambridge University Publishing Ltd.

845 Beer, M., 2009a. Engineering quantification of inconsistent information, *Int J Reliability and*
846 *Safety*, 3(1/2/3): 174–197.

847 Beer, M., 2009b. Fuzzy probability theory, in: *Encyclopedia of Complexity and Systems Science*,
848 vol.6, Springer, NewYork, pp.4047–4059.

849 Beer, M., and Ferson, S., 2013. Special issue of *Mechanical Systems and Signal Processing*
850 "Imprecise probabilities - What can they add to engineering analyses?". *Mechanical Systems*
851 *and Signal Processing*, 37(1-2), 1-3.

852 Beer, M., Ferson, S., and Kreinovich, V., 2013a. Imprecise probabilities in engineering analyses.
853 *Mechanical Systems and Signal Processing*, 37(1-2), 4-29.

854 Beer, M., Zhang, Y., Quek, S.T., Phoon, K.K., 2013b. Reliability analysis with scarce information:
855 Comparing alternative approaches in a geotechnical engineering context, *Structural Safety*,
856 Vol. 41, pp. 1–10.

857 Beer, M., DiazDelaO, F.A., Patelli, E., Au, S.K., 2014a. Conceptual comparison of Bayesian
858 approaches and imprecise probabilities. In: Topping, B.H.V.; Ivanyi, P. (eds.), Computational
859 Technology Reviews 9, 2014, 1-29, Saxe-Coburg Publications.

860 Beer, M., Kougioumtzoglou, I.A., Patelli, E., 2014b. Emerging Concepts and Approaches for
861 Efficient and Realistic Uncertainty Quantification. In: Frangopol, D.M.; Tsompanakis, Y.
862 (eds.), Maintenance and Safety of Aging Infrastructure, Book Series “Structures &
863 Infrastructures”, Vol 10, Chapter 5, 121–154, CRC Press, Taylor & Francis Group, Boca
864 Raton, London, New York, Leiden.

865 Behmanesh, I., Moaveni, B., Lombaert, G., Papadimitriou, C., 2015. Hierarchical Bayesian Model
866 Updating for Probabilistic Damage Identification. In: Atamturktur, HS, Moaveni, B.,
867 Papadimitriou, C., Schoenherr, T. Model validation and uncertainty quantification, Volume 3,
868 Conference Proceedings of the Society for Experimental Mechanics Series, pp. 55-66.

869 Bernard, J.M., 2005. An introduction to the imprecise Dirichlet model for multinomial data, *Int. J.*
870 *Approx. Reason.* 39, 123–150.

871 Bernardo, J M and Smith, A F M, 1994. *Bayesian Theory*. Chichester: Wiley.

872 Blockley, D., 2013. Analysing uncertainties: Towards comparing Bayesian and interval
873 probabilities. *Mechanical Systems and Signal Processing*, 37(1-2), 30-42.

874 Buckley, J.J., 2005. *Fuzzy probabilities — new approach and applications*, *Studies in Fuzziness and*
875 *Soft Computing*, vol.115, Springer, Berlin, Heidelberg.

876 Chung, Y., Gelman, A., Rabe-Hesketh, S., Liu, J., Dorie, V., 2015. Weakly Informative Prior for
877 Point Estimation of Covariance Matrices in Hierarchical Models. *Journal of Educational and*
878 *Behavioral Statistics*, Vol. 40, No. 2, pp. 136–157.

879 Congdon, P.D., 2010. *Applied Bayesian Hierarchical Methods*, CRC Press.

880 Coolen, F. P. A., and Utkin, L. V., 2007. Imprecise probability: a concise overview." *Risk,*
881 *reliability and societal safety: proceedings of the European safety and reliability conference*
882 *(ESREL)*, Stavanger, Norway, 1959-66.

883 Couso, I., Dubois, D., 2009. On the Variability of the Concept of Variance for Fuzzy Random
884 Variables. *Fuzzy Systems, IEEE Transactions on*, Volume: 17, Issue: 5, pp. 1070 – 1080.

885 Couso, I., Miranda, E., de Cooman, G., 2004. A Possibilistic Interpretation of the Expectation of a
886 Fuzzy Random Variable. In: *Soft Methodology and Random Information Systems, Series in*
887 *Advances in Soft Computing*, Volume 26, pp. 133-140, Springer Berlin Heidelberg.

888 Couso, I., S. Montes, P. Gil, 2001. The necessity of the strong alpha-cuts of a fuzzy set,
889 *International Journal of Uncertainty, Fuzziness and Knowledge-Based Systems*, 9, 249-262.

890 Couso, I., L. Sánchez, 2008. Higher order models for fuzzy random variables, *Fuzzy Sets and*
891 *Systems* 159, 237-258.

892 Couso, I., L. Sánchez, 2011. Upper and lower probabilities induced by a fuzzy random variable.
893 *Fuzzy Sets and Systems*, Volume 165, Issue 1, 16 February 2011, Pages 1–23.

894 Crespo, L. G., Kenny, S. P., and Giesy, D. P., 2013. Reliability analysis of polynomial systems
895 subject to p-box uncertainties. *Mechanical Systems and Signal Processing*, 37(1-2), 121-136.

896 Denoeux, T., 2014. Likelihood-based belief function: Justification and some extensions to low
897 quality data. *International Journal of Approximate Reasoning*, Volume 55, Issue 7, Pages 1535–
898 1547.

899 Dubois, D., 2006. Possibility Theory and Statistical Reasoning. *Computational Statistics and Data*
900 *Analysis*, Vol. 51, pp. 47-69.

901 Dubois, D., L. Foulloy, G. Mauris, H. Prade, 2004. Probability-Possibility Transformation,
902 Triangular Fuzzy Sets, and Probabilistic Inequalities. *Reliable Computing*, vol. 10, pp. 273-
903 297.

904 Dubois, D., H. Prade, 1980. *Fuzzy Sets and Systems: Theory and Applications*, Academic Press,
905 New York, NY.

906 Dubois, D., Prade, H., 1988. *Possibility Theory: An Approach to Computerized Processing of*
907 *Uncertainty*, New York, Plenum Press.

908 Dubois, D., Prade, H., 1997. Bayesian conditioning in possibility theory. *Fuzzy Sets and Systems*,
909 92, pp. 223-240.

910 Dubois, D, Prade, H & Sandri, S, 1993. On possibility/probability transformations. In: Lowen, R &
911 Roubens, M (eds). *Fuzzy Logic: State of the Art*. Dordrecht: Kluwer Academic Publishers.
912 pp. 103112.

913 Dubois D., Prade H., Smets P., 2008. A definition of subjective possibility. *International Journal of*
914 *Approximate Reasoning*, 2008, Vol. 48, pp. 352-364.

915 Dubois D., Moral, S., Prade H., 1997. A Semantics for Possibility Theory Based on Likelihoods.
916 *Journal of mathematical analysis and applications*, 205, pp. 359-380.

917 Edwards, A.W.F., 1992. *Likelihood*, expanded edition. The John Hopkins University Press,
918 Baltimore, USA.

919 Féron, R., 1976. Ensembles aléatoires flous, *CR. Acad. Sc. paris - Série A*, Vol 282, 903-906.

920 Ferson, S., 2005. Bayesian methods in risk assessment. Technical Report:
921 www.ramas.com/bayes.pdf.

922 Ferson, S. and Hajagos, J.G., 2004. Arithmetic with uncertain numbers: rigorous and (often) best
923 possible answers. *Reliability Engineering and System Safety*, 85, pp. 135-152.

924 Ferson, V. Kreinovich, L. Ginzburg, K. Sentz, D.S. Myers, 2003. Constructing probability boxes
925 and Dempster-Shafer structures, Sandia National Laboratories, Technical Report SAND2002-
926 4015, Albuquerque, New Mexico.

927 Ferson, V. Kreinovich, J. Hajagos, W. Oberkampf and L. Ginzburg, 2007. *Experimental*
928 *Uncertainty Estimation and Statistics for Data Having Interval Uncertainty*, Setauket, New
929 York 11733, Technical Report SAND2007-0939.

930 Ferson, R.B. Nelsen, J. Hajagos, D.J. Berleant, J. Zhang, W.T. Tucker, L.R. Ginzburg, W.L.
931 Oberkampf, 2004. Dependence in probabilistic modeling, Dempster-Shafer theory, and
932 probability bounds analysis, Technical Report SAND2004-3072, Albuquerque, New Mexico.

933 Ferson, W.T. Tucker, 2006. Sensitivity in risk analyses with uncertain numbers, Setauket, New
934 York 11733, Technical Report SAND2006-2801.

935 Ferson, P. Van den Brink, T.L. Estes, K. Gallagher, R. O'Connor, F. Verdonck, 2010. Bounding
936 uncertainty analyses. In: Application of uncertainty analysis to ecological risks of pesticides /
937 Warren-Hicks, W.J., A. Hart. Pensacola and Boca Raton, FL.: SETAC and CRC Press, 2010 -
938 ISBN 9781439807347.

939 Flage R., Aven T, Zio E., 2009. Alternative representations of uncertainty in reliability and risk
940 analysis – review and discussion. In: Martorell S, Guedes Soares C, Barnett J, editors.
941 Valencia, Spain, 22–25 September 2008 Safety, reliability and risk analysis. Theory, methods
942 and applications. Proceedings of the European safety and reliability conference 2009 (ESREL
943 2009). London: CRC Press, p. 2081–91.

944 Flage, P. Baraldi, E. Zio, T. Aven, 2010. Possibility-probability transformation in comparing
945 different approaches to the treatment of epistemic uncertainties in a fault tree analysis, in: B.
946 Ale, I.A. Papazoglu, E. Zio (Eds.), Reliability, Risk and Safety - Proceedings of the European
947 Safety and Reliability (ESREL) 2010 Conference, Rhodes, Greece, 5-9 September 2010, pp.
948 714-721, Taylor & Francis Group, London, United Kingdom, 2010, ISBN 978-0-415-60427-
949 7.

950 Flage, P. Baraldi, E. Zio, T. Aven, 2013. Probability and possibility-based representations of
951 uncertainty in fault tree analysis. Risk Analysis, 33(1), pp. 121-33.

952 Gelman A., et al., 2004. Bayesian data analysis. Second Ed. Chapman & Hall, CRC.

953 Gelman, A., 2006. Prior distributions for variance parameters in hierarchical models. Bayesian
954 Analysis, 1(3), pp. 515–533.

955 Gelman, A., Meng, XL, Stern, H., 1996. Posterior predictive assessment of model fitness via
956 realized discrepancies. Stat Sin: 733–806.

957 Gelman, A., van Dyk, D., Huang, Z., Boscardin, W.J., 2008. Using Redundant Parameterizations to
958 Fit Hierarchical Models. *Journal of Computational and Graphical Statistics*, Volume 17,
959 Number 1, Pages 95–122.

960 Gill, J., 2002. *Bayesian methods: a social and behavioral sciences approach*, Chapman & Hall/CRC,
961 Boca Raton, FL.

962 Guyonnet, D., Bourgine, B., Dubois, D., Fargier, H., Côme, B., Chilès, J.P., 2003. Hybrid approach
963 for addressing uncertainty in risk assessments. *Journal of the Environmental Engineering*
964 *Division*, ASCE 129, pp. 68–78.

965 Helton, J.D. Johnson, W.L. Oberkampf, C.J. Sallaberry, 2008. *Representation of Analysis Results*
966 *Involving Aleatory and Epistemic Uncertainty*, Sandia National Laboratories, Technical
967 Report SAND2008-4379, Albuquerque, New Mexico.

968 Helton, J.D. Johnson, W.L. Oberkampf, C.B. Storlie, 2007. A sampling-based computational
969 strategy for the representation of epistemic uncertainty in model predictions with evidence
970 theory, *Computer Methods in Applied Mechanics and Engineering*, 196: 3980–98.

971 Hou, Y. and Yang, B., 2010. Probability-Possibility Transformation for Small Sample Size Data.
972 2010 Seventh International Conference on Fuzzy Systems and Knowledge Discovery (FSKD
973 2010), pp. 1720-1724.

974 Jalal-Kamali, A., and Kreinovich, V., 2013. Estimating correlation under interval uncertainty.
975 *Mechanical Systems and Signal Processing*, 37(1-2), 43-53.

976 Jaworska, J.S., Aldenberg, T., 2000. Estimation of HC5 taking into account uncertainties of
977 individual dose response curves and species sensitivity distribution. Presentation at the 2000
978 Society for Risk Analysis annual meeting, Arlington, Virginia,
979 <http://www.riskworld.com/Abstract/2000/SRAam00/ab0ac164.htm>.

980 Johnson Valen, E., 2004. A Bayesian χ^2 test for goodness-of-fit. *Ann Stat*, 32: pp. 2361–84.

981 Kalos, M.H., P.A. Whitlock, 1986. *Monte Carlo methods*. Volume I: Basics, Wiley, New York,
982 NY.

983 Kelly, D.L., Smith, C.L, 2009. Bayesian inference in probabilistic risk assessment — The current
984 state of the art. *Reliability Engineering and System Safety*, 94, pp. 628–643.

985 Kelly, D.L., Smith, C.L, 2011. *Bayesian Inference for Probabilistic Risk Assessment: A
986 Practitioner’s Guidebook*. Springer-Verlag: London, UK.

987 Kentel, E., M.M. Aral, 2004. Probabilistic-fuzzy health risk modeling. *Stoch. Envir. Res. and Risk
988 Ass.*, Vol. 18: pp. 324–338.

989 Kentel, E., M.M. Aral, 2005. 2D Monte Carlo versus 2D Fuzzy Monte Carlo Health Risk
990 Assessment, *Internat. J. Stochastic Environ. Res. Risk Assess.*, 19, pp. 86–96.

991 Kentel, E., M.M. Aral, 2007. Risk tolerance measure for decision-making in fuzzy analysis: a health
992 risk assessment perspective. *Stoch. Environ Res. Ris. Assess.*, Vol. 21: pp. 405–417.

993 Klir, B. Yuan, 1995. *Fuzzy Sets and Fuzzy Logic: Theory and Applications*, Prentice-Hall, Upper
994 Saddle River, NJ.

995 Kozine, I., Filimonov, Y., 2000. Imprecise reliabilities: experiences and advances, *Reliab. Eng.
996 Syst. Saf.*, 67, 75–83.

997 Kozine, I. O., and Utkin, L. V., 2002. Processing unreliable judgements with an imprecise
998 hierarchical model. *Risk, Decision and Policy*, 7(03), 325-339.

999 Kuznetsov, V. P., 1991. Interval statistical models (in Russian), *Radio i Svyaz*, Moscow.

1000 Kwakernaak, H., 1978. Fuzzy Random Variables-I. Definitions and Theorems, *Information
1001 Sciences* 15, 1-29.

1002 Lapointe S. and Bobeè B., 2000. Revision of possibility distributions: A Bayesian inference pattern.
1003 *Fuzzy Sets and Systems*, 116, pp. 119-140.

1004 Le Duy, T.D., Vasseur, D., Couplet, M., Dieulle, L., Bérenguer, C., 2011. A study on updating
1005 belief functions for pa-rameter uncertainty representation in Nuclear Probabilistic Risk
1006 Assessment. In F. Coolin, G. De Cooman, T. Fetz, M. Oberguggenberger (Eds.), *Proceedings
1007 of the 7th Interna-tional Symposium on Imprecise Probability: Theories and Applications*, 25-
1008 28 July 2011, Innsbruck, Austria: 247-256. Innsbruck, Austria: SIPTA.

1009 Limbourg, P., E. de Rocquigny, 2010. Uncertainty analysis using evidence theory – confronting
1010 level-1 and level-2 approaches with data availability and computational constraints,
1011 Reliability Engineering and System Safety, Vol. 95, Issue 5, pp. 550-564.

1012 Lindley, D.V., 2000. The philosophy of statistics. The Statistician, 49(3): 293-337.

1013 Lindley, D. V., 2006. Understanding uncertainty, Wiley, Hoboken, NJ.

1014 Masegosa, A., S. Moral, 2014. Imprecise probability models for learning multinomial distributions
1015 from data. Applications to learning credal networks. International Journal of Approximate
1016 Reasoning, Volume 55, Issue 7, Pages 1548–1569.

1017 Masson, M.H. and Denoeux, T., 2006. Inferring a possibility distribution from empirical data.
1018 Fuzzy Sets and Systems, 157, pp. 319–340.

1019 Mauris, G., 2008. Inferring a Possibility Distribution from Very Few Measurements. In: D. Dubois
1020 et al. (Eds.), Soft Methods for Hand. Var. and Imprecision, ASC 48, pp. 92–99, Springer-
1021 Verlag, Berlin, Heidelberg.

1022 Mehl, C. H., 2013. P-boxes for cost uncertainty analysis. Mechanical Systems and Signal
1023 Processing, 37(1-2), 253-263.

1024 Molchanov, I., 2005. Theory of Random Sets, Springer, New York.

1025 Möller, B., M. Beer, 2004. Fuzzy Randomness: Uncertainty in Civil Engineering and
1026 Computational Mechanics, Springer, Berlin.

1027 Möller, B., M. Beer, 2008. Engineering computation under uncertainty – Capabilities of non-
1028 traditional models. Computers and Structures, Vol. 86, pp. 1024–1041.

1029 Moller, B., W. Graf, M. Beer, 2003. Safety assessment of structures in view of fuzzy randomness.
1030 Computers and Structures, Vol. 81, pp. 1567–1582.

1031 Moller, B., M. Beer, W. Graf, J.U. Sickert, 2006. Time-dependent reliability of textile-strengthened
1032 RC structures under consideration of fuzzy randomness. Computers and Structures, Vol. 84,
1033 pp. 585–603.

1034 Moral, S., 2014. Comments on “Likelihood-based belief function: Justification and some extensions
1035 to low-quality data” by Thierry Denoeux. *International Journal of Approximate Reasoning*,
1036 Volume 55, Issue 7, Pages 1591–1593.

1037 Muscolino, G., and Sofi, A., 2013. Bounds for the stationary stochastic response of truss structures
1038 with uncertain-but-bounded parameters. *Mechanical Systems and Signal Processing*, 37(1-2),
1039 163-181.

1040 NUREG-CR-6850, 2005. EPRI/NRC-RES Fire PRA methodology for nuclear power facilities,
1041 Volume 2: detailed methodology. US Nuclear Regulatory Commission.

1042 NASA, 2010. Risk-Informed Decision Making Handbook. NASA/SP-2010-576 – Version 1.0,
1043 April 2010.

1044 Pannier, S., Waurick, M., Graf, W., and Kaliske, M., 2013. Solutions to problems with imprecise
1045 data” - An engineering perspective to generalized uncertainty models. *Mechanical Systems
1046 and Signal Processing*, 37(1-2), 105-120.

1047 Pasanisi, A., E. de Rocquigny, N. Bousquet, E. Parent, 2009. Some useful features of the Bayesian
1048 setting while dealing with uncertainties in industrial practice, *Proceedings of the ESREL 2009
1049 Conference, Prague, Czech Republic, 7-10 September 2009*, pp. 1795-1802.

1050 Pasanisi, A., Keller, M., Parent, M., 2012. Estimation of a quantity of interest in uncertainty
1051 analysis: Some help from Bayesian decision theory. *Reliability Engineering and System
1052 Safety*, 100, pp. 93–101.

1053 Pedroni, N., Zio, E., 2012. Empirical comparison of methods for the hierarchical propagation of
1054 hybrid uncertainty in risk assessment, in presence of dependences. *International Journal of
1055 Uncertainty, Fuzziness and Knowledge-Based Systems*, Vol. 20, Issue 4, 2012, pp. 509-557.

1056 Pedroni, N., E. Zio, E. Ferrario, A. Pasanisi, M. Couplet, 2013. Hierarchical propagation of
1057 probabilistic and non-probabilistic uncertainty in the parameters of a risk model. *Computers
1058 and Structures*, Volume 126, pp. 199–213.

1059 Pedroni, N., E. Zio, A. Pasanisi, M. Couplet, 2014. Bayesian update of the parameters of probability
1060 distributions for risk assessment in a two-level hybrid probabilistic-possibilistic uncertainty
1061 framework. In: R.D.J.M. Steenbergen, P.H.A.J.M. van Gelder, S. Miraglia and A. C.W.M.
1062 Ton. Vrouwenvelder (Eds.), *Safety, Reliability and Risk Analysis, Beyond the Horizon,*
1063 *Proceedings of the European Safety and RELiability Conference (ESREL) 2013,* Amsterdam,
1064 The Netherlands, 29 September-2 October 2013, Taylor and Francis Group, London, UK, pp.
1065 3295–3302.

1066 Puri, M.L., D.A. Ralescu, 1986. Fuzzy random variables, *Journal of Mathematical Analysis and*
1067 *Applications* 114, 409-422.

1068 Quaeghebeur and de Cooman, 2005. Imprecise probability models for inference in exponential
1069 families. *Electronic Proceedings of the 4th International Symposium on Imprecise*
1070 *Probabilities and Their Applications,* Pittsburgh, Pennsylvania, 2005.

1071 Ralescu, D., 2002. Average level of a fuzzy set, in: *Statistical Modeling, Analysis and Management*
1072 *of Fuzzy Data,* eds. C. Bertoluzza, M.A. Gil, D.A. Ralescu (Springer, Heidelberg) pp. 119–
1073 126.

1074 Reid, S. G., 2013. Probabilistic confidence for decisions based on uncertain reliability estimates.
1075 *Mechanical Systems and Signal Processing,* 37(1-2), 229-239.

1076 Sankararaman, S., and Mahadevan, S., 2013. Distribution type uncertainty due to sparse and
1077 imprecise data. *Mechanical Systems and Signal Processing,* 37(1-2), 182-198.

1078 Sentz, S. Ferson, 2002. *Combination of Evidence in Dempster-Shafer Theory,* Sandia National
1079 Laboratories, Technical Report SAND 2002-0835, Albuquerque, New Mexico.

1080 Serrurier, M., and Prade, H., 2011. Maximum-Likelihood Principle For Possibility Distributions
1081 Viewed As Families Of Probabilities. 2011 IEEE International Conference on Fuzzy Systems
1082 June 27-30, 2011, Taipei, Taiwan, pp. 2987-2993.

1083 Shirley, K.E., Gelman, A., 2015. Hierarchical models for estimating state and demographic trends
1084 in US death penalty public opinion. *J. R. Statist. Soc. A,* 178(1), pp. 1-28.

- 1085 Siu, N., Kelly, D., 1998. Bayesian parameter estimation in probabilistic risk assessment. *Reliability*
1086 *Engineering and System Safety*, 62, pp. 89–116.
- 1087 Smets, P., 1993. Belief Functions: The Disjunctive Rule of Combination and the Generalized
1088 Bayesian Theorem. *International Journal of Approximate Reasoning*, 9, pp. 1-35.
- 1089 Stein M. & M. Beer, 2011. Bayesian quantification of inconsistent information. *Applications of*
1090 *Statistics and Probability in Civil Engineering – Faber, Köhler & Nishijima (eds)*, Taylor &
1091 Francis Group, London, ISBN 978-0-415-66986-3.
- 1092 Stein M., M. Beer, V. Kreinovich, 2013. Bayesian Approach for Inconsistent Information.
1093 *Information Sciences*: <http://dx.doi.org/10.1016/j.ins.2013.02.024>.
- 1094 USNRC, 2002. An approach for using probabilistic risk assessment in risk-informed decisions on
1095 plant-specific changes to the licensing basis. NUREG-1.174 – Revision 1, US Nuclear
1096 Regulatory Commission, Washington, DC.
- 1097 USNRC, 2009. Guidance on the Treatment of Uncertainties Associated with PRAs in Risk-
1098 Informed Decision Making. NUREG-1855, US Nuclear Regulatory Commission,
1099 Washington, DC.
- 1100 Viertl, R., 1996. *Statistical Methods for Non-Precise Data*. CRC Press, Boca Raton, New York,
1101 London, Tokyo.
- 1102 Viertl, R., 1997. On statistical inference for non precise data. *Environmetrics*, 8, pp. 541-568.
- 1103 Viertl, R., 1999. Statistics and integration of fuzzy functions. *Environmetrics*, 10, pp. 487-491.
- 1104 Viertl, R., 2008a. Foundations of Fuzzy Bayesian Inference. *Journal of Uncertain Systems*, 2(3), pp.
1105 187–191.
- 1106 Viertl, R., 2008b. Fuzzy Bayesian Inference. In: D. Dubois et al. (Eds.), *Soft Methods for Handling*
1107 *Variability and Imprecision*, ASC 48, pp. 10–15.
- 1108 Viertl, R., 2011. *Statistical Methods for Fuzzy Data*, Wiley, Chichester.
- 1109 Viertl, R. & Hareter, D., 2004a. Fuzzy information and imprecise probability. *ZAMM, Z. Angew.*
1110 *Math. Mech.*, 84 (10–11), pp. 731 – 739.

- 1111 Viertl, R. & Hareter, D., 2004b. Generalized Bayes-theorem for non-precise a-priori distribution,
1112 Metrika, 59(3), pp. 263–273.
- 1113 Viertl R. and H. Hule, 1991. On Bayes’ theorem for fuzzy data. Stat. Pap. 32.
- 1114 Walley, P., 1991. Statistical reasoning with imprecise probabilities, Chapman and Hall, New York.
- 1115 Walley, P., 1996. Inferences from multinomial data: learning about a bag of marbles (with
1116 discussion), J. R. Stat. Soc. B 58, 3–57.
- 1117 Walter, G., and Augustin, T., 2009. Imprecision and Prior-data Conflict in Generalized Bayesian
1118 Inference. Journal of Statistical Theory and Practice, Volume 3, No.1, pp. 255-271.
- 1119 Weichselberger, K., 2000. The theory of interval-probability as a unifying concept for uncertainty,
1120 Int. J. Approx. Reasoning, 24(2–3), 149–170.
- 1121 Zhang, H., Dai, H., Beer, M., and Wang, W., 2013. Structural reliability analysis on the basis of
1122 small samples: An interval quasi-Monte Carlo method. Mechanical Systems and Signal
1123 Processing, 37(1-2), 137-151.
- 1124
- 1125

1126

1127

Figure captions list

1128

1129 **Fig. 1. Top left:** triangular possibility distribution $\pi^\gamma(\gamma)$ of the epistemically-uncertain parameter

1130 γ of the Gumbel probability distribution of Y ; in evidence the α -cuts of level $\alpha = 0, 0.5$ and 1 .

1131 **Top right:** four PDFs belonging to the family $\{p^Y(y|\gamma, \delta)\}_{\alpha=0}$ and four PDFs belonging to the

1132 family $\{p^Y(y|\gamma, \delta)\}_{\alpha=0.5}$. **Bottom:** bounding upper and lower CDFs of Y , $\bar{F}_\alpha^Y(y) = Pl_\alpha^Y(Z)$ and

1133 $\underline{F}_\alpha^Y(y) = Bel_\alpha^Y(Z)$, $Z = (-\infty, y]$, built in correspondence of the α -cuts of level $\alpha = 0, 0.5$ and 1

1134 of $\pi^\gamma(\gamma)$; the plausibility and belief functions $Pl^Y(Z)$ and $Bel^Y(Z)$, $Z = (-\infty, y]$, are also

1135 shown

1136

1137 **Fig. 2.** Identification of the P -dimensional α -cut $A_\alpha^{\theta|y}$ of the posterior possibility distribution of θ as

1138 the hull enveloping $T = 20$ point estimates $\theta_{l,\alpha} | y$, $l = 1, 2, \dots, 20$ (dots), generated by the

1139 repeated application of the Bayes' theorem in the hybrid probabilistic-possibilistic approach

1140 (B), in the particular case of $P = 2$ parameters θ_1 and θ_2 . The corresponding α -cut $A_{\alpha, Cart}^{\theta_1, \theta_2 | y}$

1141 generated by the Cartesian product of the (one-dimensional) α -cuts $A_\alpha^{\theta_1 | y}$ and $A_\alpha^{\theta_2 | y}$ of the

1142 marginal possibility distributions is also shown

1143

1144 **Fig. 3.** Prior and posterior possibility distributions of the epistemically-uncertain parameters of the

1145 aleatory PDFs of $Y_1 = Q$ (top row, left and right column), $Y_2 = Z_m$ (middle-top row, left and

1146 right column), $Y_3 = Z_v$ (middle-bottom row, left and right column) and $Y_4 = K_s$ (bottom row,

1147 left and right column). The point estimates of the parameters obtained by the classical MLE

1148 method are also shown for comparison. Adapted and extended from [Pedroni et al., 2014]

1149

1150 **Fig. 4.** Exemplary α -cuts of level $\alpha = 0.05, 0.50$ and 0.95 of the joint posterior possibility
1151 distributions of the parameters of the PDFs of $Y_1 = Q$ (**top left**), $Y_2 = Z_m$ (**top right**), $Y_3 = Z_v$
1152 (**bottom left**) and $Y_4 = K_s$ (**bottom right**), produced by Approaches A (solid lines) and B
1153 (dashed lines)

1154

1155 **Fig. 5.** Upper and lower CDFs, $\bar{F}^{Y_j}(y_j)$ and $\underline{F}^{Y_j}(y_j)$, $j = 1, 2, 3, 4$, of the uncertain input variables
1156 $Y_1 = Q$ (**top left**), $Y_2 = Z_m$ (**top right**), $Y_3 = Z_v$ (**bottom left**) and $Y_4 = K_s$ (**bottom right**) before
1157 and after the Bayesian update performed by Approaches A and B

1158

1159 **Fig. 6.** Plausibility and belief functions, $Pl^{Z_c}((-\infty, z_c])$ and $Bel^{Z_c}((-\infty, z_c])$, of the maximal water
1160 level of the river Z_c before and after the Bayesian update performed by Approaches A and B.

1161 Adapted and extended from [Pedroni et al., 2014]


1162

1163

1164

1165

Figures and corresponding associated captions



1166

1167

1168 **Fig. 1. Top left:** triangular possibility distribution $\pi^\gamma(\gamma)$ of the epistemically-uncertain parameter

1169 γ of the Gumbel probability distribution of Y ; in evidence the α -cuts of level $\alpha = 0, 0.5$ and 1 .

1170 **Top right:** four PDFs belonging to the family $\{p^Y(y | \gamma, \delta)\}_{\alpha=0}$ and four PDFs belonging to the


1171 family $\{p^Y(y | \gamma, \delta)\}_{\alpha=0.5}$. **Bottom:** bounding upper and lower CDFs of Y , $\bar{F}_\alpha^Y(y) = Pl_\alpha^Y(Z)$ and

1172 $\underline{F}_\alpha^Y(y) = Bel_\alpha^Y(Z)$, $Z = (-\infty, y]$, built in correspondence of the α -cuts of level $\alpha = 0, 0.5$ and 1

1173 of $\pi^\gamma(\gamma)$; the plausibility and belief functions $Pl^Y(Z)$ and $Bel^Y(Z)$, $Z = (-\infty, y]$, are also

1174 shown

1175



1176

1177 **Fig. 2.** Identification of the P -dimensional α -cut $A_\alpha^{\theta|y}$ of the posterior possibility distribution of θ as

1178 the hull enveloping $T = 20$ point estimates $\theta_{l,\alpha} | y$, $l = 1, 2, \dots, 20$ (dots), generated by the

1179 repeated application of the Bayes' theorem in the hybrid probabilistic-possibilistic approach



1180 (B), in the particular case of $P = 2$ parameters θ_1 and θ_2 . The corresponding α -cut $A_{\alpha, Cart}^{\theta_1, \theta_2 | y}$

1181 generated by the Cartesian product of the (one-dimensional) α -cuts $A_\alpha^{\theta_1 | y}$ and $A_\alpha^{\theta_2 | y}$ of the



1182 marginal possibility distributions is also shown

1183



1184





1185



1186




1187





1188

1189 **Fig. 3.** *Prior and posterior possibility distributions of the epistemically-uncertain parameters of the*
1190 *aleatory PDFs of $Y_1 = Q$ (top row, left and right column), $Y_2 = Z_m$ (middle-top row, left and*
1191 *right column), $Y_3 = Z_v$ (middle-bottom row, left and right column) and $Y_4 = K_s$ (bottom row,*
1192 *left and right column). The point estimates of the parameters obtained by the classical MLE*
1193 *method are also shown for comparison. Adapted and extended from [Pedroni et al., 2014]*
1194


1195



1196





1197





1198 **Fig. 4.** Exemplary α -cuts of level $\alpha = 0.05, 0.50$ and 0.95 of the joint posterior possibility
 1199 distributions of the parameters of the PDFs of $Y_1 = Q$ (top left), $Y_2 = Z_m$ (top right), $Y_3 = Z_v$
 1200 (bottom left) and $Y_4 = K_s$ (bottom right), produced by Approaches A (solid lines) and B
 1201 (dashed lines)

1202

1203




1204



1205

1206 **Fig. 5.** Upper and lower CDFs, $\bar{F}^{Y_j}(y_j)$ and $\underline{F}^{Y_j}(y_j)$, $j = 1, 2, 3, 4$, of the uncertain input variables
 1207 $Y_1 = Q$ (top left), $Y_2 = Z_m$ (top right), $Y_3 = Z_v$ (bottom left) and $Y_4 = K_s$ (bottom right) before
 1208 and after the Bayesian update performed by Approaches A and B

1209



1210

1211 **Fig. 6.** Plausibility and belief functions, $Pl^{Z_c}((-\infty, z_c])$ and $Bel^{Z_c}((-\infty, z_c])$, of the maximal water

1212 level of the river Z_c before and after the Bayesian update performed by Approaches A and B.

1213 Adapted and extended from [Pedroni et al., 2014]

1214

1215

Tables

1216

	Purely possibilistic method (A)	Hybrid probabilistic-possibilistic method (B)
Prior	Possibility distribution	Fuzzy Probability Distribution Function
Likelihood	Possibility distribution: - Probability-possibility transformation in this paper (e.g., normalization: see also [Denoeux, 2014])	Probabilistic function
Posterior	Possibility distribution	Fuzzy Probability Distribution Function: - Expected value: possibility distribution

1217 *Table 1. Characteristics of the purely possibilistic (A) and the hybrid probabilistic and possibilistic*

1218

(B) approaches

1219

		Purely possibilistic method (Approach A)	
Aleatory uncertainty		Epistemic uncertainty (Purely possibilistic priors)	
Q	$Gum(\gamma, \delta)$	$\pi^\gamma(\gamma) = TR(a_\gamma, c_\gamma, b_\gamma) = TR(869, 955, 1157)$	
		$\pi^\delta(\delta) = TR(a_\delta, c_\delta, b_\delta) = TR(455, 600, 660)$	
Z_m	$N(\mu_{Z_m}, \sigma_{Z_m})$	$\pi^{\mu_{Z_m}}(\mu_{Z_m}) = TR(a_{\mu_{Z_m}}, c_{\mu_{Z_m}}, b_{\mu_{Z_m}}) = TR(54.78, 54.93, 55.28)$	
		$\pi^{\sigma_{Z_m}}(\sigma_{Z_m}) = TR(a_{\sigma_{Z_m}}, c_{\sigma_{Z_m}}, b_{\sigma_{Z_m}}) = TR(0.33, 0.51, 0.58)$	
Z_v	$N(\mu_{Z_v}, \sigma_{Z_v})$	$\pi^{\mu_{Z_v}}(\mu_{Z_v}) = TR(a_{\mu_{Z_v}}, c_{\mu_{Z_v}}, b_{\mu_{Z_v}}) = TR(49.98, 50.11, 50.40)$	
		$\pi^{\sigma_{Z_v}}(\sigma_{Z_v}) = TR(a_{\sigma_{Z_v}}, c_{\sigma_{Z_v}}, b_{\sigma_{Z_v}}) = TR(0.23, 0.45, 0.54)$	
K_s	$N(\mu_{K_s}, \sigma_{K_s})$	$\pi^{\mu_{K_s}}(\mu_{K_s}) = TR(a_{\mu_{K_s}}, c_{\mu_{K_s}}, b_{\mu_{K_s}}) = TR(21.37, 25.23, 34.23)$	
		$\pi^{\sigma_{K_s}}(\sigma_{K_s}) = TR(a_{\sigma_{K_s}}, c_{\sigma_{K_s}}, b_{\sigma_{K_s}}) = TR(1.16, 6.91, 9.37)$	
		Hybrid probabilistic-possibilistic method (Approach B)	
Aleatory uncertainty		Epistemic uncertainty (Fuzzy Probability Density Functions - FPDFs)	
		Probabilistic part	Possibilistic part
Q	$Gum(\gamma, \delta)$	$p^\gamma(\gamma) = N(\mu_\gamma, \sigma_\gamma)$	$\pi^{\mu_\gamma}(\mu_\gamma) = TR(a_{\mu_\gamma}, c_{\mu_\gamma}, b_{\mu_\gamma}) = TR(869, 955, 1157),$ $\pi^{\sigma_\gamma}(\sigma_\gamma) = TR(a_{\sigma_\gamma}, c_{\sigma_\gamma}, b_{\sigma_\gamma}) = TR(39, 52, 57)$
		$p^\delta(\delta) = N(\mu_\delta, \sigma_\delta)$	$\pi^{\mu_\delta}(\mu_\delta) = TR(a_{\mu_\delta}, c_{\mu_\delta}, b_{\mu_\delta}) = TR(455, 600, 660)$ $\pi^{\sigma_\delta}(\sigma_\delta) = TR(a_{\sigma_\delta}, c_{\sigma_\delta}, b_{\sigma_\delta}) = TR(25, 31, 43)$
Z_m	$N(\mu_{Z_m}, \sigma_{Z_m})$	$p^{\mu_{Z_m}}(\mu_{Z_m}) = N(\mu_{\mu_{Z_m}}, \sigma_{\mu_{Z_m}})$	$\pi^{\mu_{\mu_{Z_m}}}(\mu_{\mu_{Z_m}}) = TR(a_{\mu_{\mu_{Z_m}}}, c_{\mu_{\mu_{Z_m}}}, b_{\mu_{\mu_{Z_m}}}) = TR(54.78, 54.93, 55.28)$ $\pi^{\sigma_{\mu_{Z_m}}}(\sigma_{\mu_{Z_m}}) = TR(a_{\sigma_{\mu_{Z_m}}}, c_{\sigma_{\mu_{Z_m}}}, b_{\sigma_{\mu_{Z_m}}}) = TR(0.06, 0.09, 0.11)$
		$p^{\sigma_{Z_m}}(\sigma_{Z_m}) = N(\mu_{\sigma_{Z_m}}, \sigma_{\sigma_{Z_m}})$	$\pi^{\mu_{\sigma_{Z_m}}}(\mu_{\sigma_{Z_m}}) = TR(a_{\mu_{\sigma_{Z_m}}}, c_{\mu_{\sigma_{Z_m}}}, b_{\mu_{\sigma_{Z_m}}}) = TR(0.33, 0.51, 0.58)$ $\pi^{\sigma_{\sigma_{Z_m}}}(\sigma_{\sigma_{Z_m}}) = TR(a_{\sigma_{\sigma_{Z_m}}}, c_{\sigma_{\sigma_{Z_m}}}, b_{\sigma_{\sigma_{Z_m}}}) = TR(0.03, 0.04, 0.06).$
Z_v	$N(\mu_{Z_v}, \sigma_{Z_v})$	$p^{\mu_{Z_v}}(\mu_{Z_v}) = N(\mu_{\mu_{Z_v}}, \sigma_{\mu_{Z_v}})$	$\pi^{\mu_{\mu_{Z_v}}}(\mu_{\mu_{Z_v}}) = TR(a_{\mu_{\mu_{Z_v}}}, c_{\mu_{\mu_{Z_v}}}, b_{\mu_{\mu_{Z_v}}}) = TR(49.98, 50.11, 50.40)$ $\pi^{\sigma_{\mu_{Z_v}}}(\sigma_{\mu_{Z_v}}) = TR(a_{\sigma_{\mu_{Z_v}}}, c_{\sigma_{\mu_{Z_v}}}, b_{\sigma_{\mu_{Z_v}}}) = TR(0.04, 0.08, 0.10),$
		$p^{\sigma_{Z_v}}(\sigma_{Z_v}) = N(\mu_{\sigma_{Z_v}}, \sigma_{\sigma_{Z_v}})$	$\pi^{\mu_{\sigma_{Z_v}}}(\mu_{\sigma_{Z_v}}) = TR(a_{\mu_{\sigma_{Z_v}}}, c_{\mu_{\sigma_{Z_v}}}, b_{\mu_{\sigma_{Z_v}}}) = TR(0.23, 0.45, 0.54)$ $\pi^{\sigma_{\sigma_{Z_v}}}(\sigma_{\sigma_{Z_v}}) = TR(a_{\sigma_{\sigma_{Z_v}}}, c_{\sigma_{\sigma_{Z_v}}}, b_{\sigma_{\sigma_{Z_v}}}) = TR(0.02, 0.04, 0.08)$
K_s	$N(\mu_{K_s}, \sigma_{K_s})$	$p^{\mu_{K_s}}(\mu_{K_s}) = N(\mu_{\mu_{K_s}}, \sigma_{\mu_{K_s}})$	$\pi^{\mu_{\mu_{K_s}}}(\mu_{\mu_{K_s}}) = TR(a_{\mu_{\mu_{K_s}}}, c_{\mu_{\mu_{K_s}}}, b_{\mu_{\mu_{K_s}}}) = TR(21.37, 25.23, 34.23)$ $\pi^{\sigma_{\mu_{K_s}}}(\sigma_{\mu_{K_s}}) = TR(a_{\sigma_{\mu_{K_s}}}, c_{\sigma_{\mu_{K_s}}}, b_{\sigma_{\mu_{K_s}}}) = TR(0.50, 2.80, 3.79)$
		$p^{\sigma_{K_s}}(\sigma_{K_s}) = N(\mu_{\sigma_{K_s}}, \sigma_{\sigma_{K_s}})$	$\pi^{\mu_{\sigma_{K_s}}}(\mu_{\sigma_{K_s}}) = TR(a_{\mu_{\sigma_{K_s}}}, c_{\mu_{\sigma_{K_s}}}, b_{\mu_{\sigma_{K_s}}}) = TR(1.16, 6.91, 9.37)$ $\pi^{\sigma_{\sigma_{K_s}}}(\sigma_{\sigma_{K_s}}) = TR(a_{\sigma_{\sigma_{K_s}}}, c_{\sigma_{\sigma_{K_s}}}, b_{\sigma_{\sigma_{K_s}}}) = TR(0.10, 0.86, 2.64)$

1221 **Table 2.** Characteristics and parameters of the prior distributions of the uncertain variables $Y_1 =$
 1222 Q , $Y_2 = Z_m$, $Y_3 = Z_v$, and $Y_4 = K_s$ used in Approaches A and B, respectively

1223

1224

Marginal possibility distributions update								
Variable	θ	MLE $\hat{\theta}^{MLE}$	Most likely value, $c_\theta (d_\theta^{MLE})$			Area, $S_\theta (R_\theta)$		
			Prior	Posterior A	Posterior B	Prior	Posterior A	Posterior B
Q [m^3/s]	γ	1013.21	955.55	1002.70 (0.010)	990.42 (0.023)	144.65	100.55 (30.49)	96.90 (33.01)
	δ	558.48	599.15	566.35 (0.014)	581.40 (0.041)	103.35	76.94 (25.56)	73.65 (28.74)
Z_m [m]	μ_{Z_m}	55.03	54.93	55.00 (5.45e-4)	54.98 (9.09e-4)	0.25	0.190 (24.00)	0.195 (22.00)
	σ_{Z_m}	0.45	0.51	0.47 (0.044)	0.50 (0.111)	0.12	0.110 (8.33)	0.115 (4.17)
Z_v [m]	μ_{Z_v}	50.19	50.11	50.17 (3.98e-4)	50.15 (7.97e-4)	0.21	0.165 (21.43)	0.170 (19.05)
	σ_{Z_v}	0.38	0.45	0.39 (0.026)	0.43 (0.132)	0.16	0.121 (24.38)	0.125 (21.88)
K_s [l]	μ_{K_s}	27.80	25.24	26.95 (0.031)	26.43 (0.049)	6.45	5.40 (16.28)	5.80 (10.08)
	σ_{K_s}	5.26	6.89	5.54 (0.053)	6.72 (0.278)	4.11	3.75 (8.76)	3.95 (3.89)

1226 **Table 3.** Most likely values c_θ of the parameters $\theta = \gamma, \delta, \mu_{Z_m}, \sigma_{Z_m}, \mu_{Z_v}, \sigma_{Z_v}, \mu_{K_s}$ and σ_{K_s} of
1227 the aleatory PDFs of the input variables $Y_1 = Q, Y_2 = Z_m, Y_3 = Z_v$ and $Y_4 = K_s$ and areas S_θ
1228 underlying the corresponding (marginal) possibility distributions before and after the
1229 Bayesian update performed by Approaches A and B. The point estimates $\hat{\theta}^{MLE}$ obtained by
1230 the classical MLE method are also shown for comparison together with the values of the
1231 quantitative indicators d_θ^{MLE} (5) and R_θ (6) (in parentheses). Adapted and extended from
1232 [Pedroni et al., 2014]
1233

INPUT VARIABLES				
Variable	Indicator	Prior	Purely possibilistic approach (A) (% width reduction)	Hybrid approach (B) (% width reduction)
$Y_1 = Q$	$Q^{0.05}$	[284.8, 480.9]	[329.3, 446.3] (0.4033)	[322.9, 434.1] (0.4329)
	$Q^{0.50}$	[1110.5, 1295.5]	[1143.6, 1269.1] (0.3216)	[1145.5, 1263.7] (0.3610)
	$Q^{0.95}$	[2478.2, 2917.3]	[2553.4, 2857.9] (0.3065)	[2571.1, 2879.0] (0.2987)
$Y_2 = Z_m$	$Z_m^{0.05}$	[53.98, 54.36]	[54.09, 54.37] (0.2631)	[54.06, 54.36] (0.2105)
	$Z_m^{0.50}$	[54.85, 55.12]	[54.92, 55.09] (0.4074)	[54.93, 55.12] (0.2592)
	$Z_m^{0.95}$	[55.56, 56.02]	[55.66, 55.96] (0.3478)	[55.67, 55.99] (0.3043)
$Y_3 = Z_v$	$Z_v^{0.05}$	[49.22, 49.64]	[49.35, 49.64] (0.3095)	[49.29, 49.62] (0.2142)
	$Z_v^{0.50}$	[50.04, 50.29]	[50.10, 50.25] (0.4000)	[50.10, 50.27] (0.3199)
	$Z_v^{0.95}$	[50.63, 51.09]	[50.72, 51.00] (0.3913)	[50.72, 51.06] (0.2608)
$Y_4 = K_s$	$K_s^{0.05}$	[10.95, 22.33]	[12.75, 21.25] (0.2530)	[11.15, 20.38] (0.1889)
	$K_s^{0.50}$	[23.49, 31.15]	[24.43, 29.73] (0.3080)	[24.05, 30.22] (0.1945)
	$K_s^{0.95}$	[32.66, 44.14]	[33.01, 41.40] (0.2691)	[33.32, 44.14] (0.0574)

1235 **Table 4.** Intervals $\left[\left[\bar{F}^{Y_j} \right]^{-1}(\beta), \left[\underline{F}^{Y_j} \right]^{-1}(\beta) \right]$ for the β -100-th percentiles Y_j^β , $\beta = 0.05, 0.50$ and 0.95 ,
 1236 of the input variables Y_j , $j = 1, 2, 3, 4$, before and after the Bayesian update performed by
 1237 Approaches A and B. The percentage reduction in the width of the intervals produced by
 1238 Approaches A and B is also shown in parentheses for comparison purposes

1239
 1240

1241

OUTPUT VARIABLE Z_c				
Variable	Indicator	Prior	Purely possibilistic approach (A) (% width reduction)	Hybrid approach (B) (% width reduction)
Z_c	$Z_c^{0.05}$	[50.70, 51.67]	[50.90, 51.56] (0.3196)	[50.84, 51.53] (0.2887)
	$Z_c^{0.50}$	[52.16, 53.46]	[52.38, 53.23] (0.3462)	[52.36, 53.28] (0.2923)
	$Z_c^{0.95}$	[54.13, 56.44]	[54.21, 55.87] (0.2814)	[54.28, 56.20] (0.1688)
	$P[Z_c > z_c^*]$	[0.8908, 0.9946]	[0.9284, 0.9921] (0.3863)	[0.9103, 0.9946] (0.1879)

1242 **Table 5.** Intervals $\left[[Pl^{Z_c}]^{-1}(\beta), [Bel^{Z_c}]^{-1}(\beta) \right]$ for the β -100-th percentiles Z_c^β , $\beta = 0.05, 0.50$ and
 1243 0.95 , of the maximal water level of the river Z_c (i.e., the model output) and intervals
 1244 $[1 - \bar{F}^{Z_c}(z_c^*), 1 - \underline{F}^{Z_c}(z_c^*)]$ for the exceedance probability $P[Z_c > z_c^* = 55.5m]$, before and
 1245 after the Bayesian update performed by Approaches A and B. The percentage reduction in the
 1246 width of the intervals produced by Approaches A and B is also shown in parentheses for
 1247 comparison purposes. Adapted and extended from [Pedroni et al., 2014]

1248

1249

	Approach A	Approach B ($T = 100, N_\alpha = 21$)
Computational time t_{comp} [s] on a Intel® Core™2 Duo CPU E7600 @ 3.06 and 3.07 GHz	0.73	1536.65

1250

Table 6. Computational time t_{comp} required by the Bayesian update of all the parameters of the

1251

PDFs of $Y_1 = Q$, $Y_2 = Z_m$, $Y_3 = Z_v$ and $Y_4 = K_s$ performed by Approach A and by Approach B

1252

with $T = 100$ repetitions of the purely probabilistic Bayes' theorem for each of the $N_\alpha = 21$ α -

1253

cuts analyzed

1254

	Purely possibilistic approach (A)	Hybrid probabilistic-possibilistic approach (B)
Advantages	<ul style="list-style-type: none"> - High strength in updating the possibilistic parameters independently on the size of the data set available - Computationally cheap 	<ul style="list-style-type: none"> - Well-established and rigorous theoretical framework (i.e., probabilistic Bayes' theorem)
Drawbacks	<ul style="list-style-type: none"> - Necessity to "build" a possibilistic likelihood (e.g., by probability-possibility transformations, normalization or directly from rough experimental data) 	<ul style="list-style-type: none"> - Necessity to "build" a (fictitious) Fuzzy Probability Distribution Functions as a prior - High strength in updating the possibilistic parameters only when the size of the data set available is large (e.g., > 100) - Computationally burdensome

1256 *Table 7. Advantages and drawbacks of the purely possibilistic (A) and the hybrid probabilistic and*
1257 *possibilistic (B) approaches*