



**HAL**  
open science

# A REMARK ON VANISHING CYCLES WITH TWO STRATA

Dung Trang Lê, David Massey

► **To cite this version:**

Dung Trang Lê, David Massey. A REMARK ON VANISHING CYCLES WITH TWO STRATA. Comptes Rendus. Mathématique, 2012, 350, pp.217 - 220. hal-01271388

**HAL Id: hal-01271388**

**<https://hal.science/hal-01271388>**

Submitted on 11 Feb 2016

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# A REMARK ON VANISHING CYCLES WITH TWO STRATA

LÊ DŨNG TRÁNG AND DAVID B. MASSEY

ABSTRACT. Suppose that the critical locus  $\Sigma$  of a complex analytic function  $f$  on affine space is, itself, a space with an isolated singular point at the origin  $\mathbf{0}$ , and that the Milnor number of  $f$  restricted to normal slices of  $\Sigma - \{\mathbf{0}\}$  is constant. Then, the general theory of perverse sheaves puts severe restrictions on the cohomology of the Milnor fiber of  $f$  at  $\mathbf{0}$ , and even more surprising restrictions on the cohomology of the Milnor fiber of generic hyperplane slices.

## 1. SETTINGS

Let  $\mathcal{U}$  be an open neighborhood of the origin in  $\mathbb{C}^{n+1}$ , and  $f : (\mathcal{U}, \mathbf{0}) \rightarrow (\mathbb{C}, 0)$  be a complex analytic function. Let  $(X, \mathbf{0})$  denote the germ of the complex analytic hypersurface defined by this function.

The Milnor fiber,  $F_{\mathbf{0}}$ , of  $f$  at the origin has been a fundamental object in the study of the local, ambient topology of  $(X, \mathbf{0})$  since the appearance of the foundational work by Milnor in [11]. In [11], Milnor proves, among other things, that, if  $f$  has an isolated critical point at  $\mathbf{0}$ , then the homotopy-type of  $F_{\mathbf{0}}$  is that of a finite one-point union, a *bouquet*, of  $n$ -spheres, where the number of spheres is given by the *Milnor number*,  $\mu_{\mathbf{0}}(f)$ .

It is natural to consider the question of what can be said about the homotopy-type, or even cohomology, of  $F_{\mathbf{0}}$  in the case where the dimension of the critical locus (at the origin),  $s := \dim_{\mathbf{0}} \Sigma f$ , is greater than 0.

One of the first general results along these lines was due to M. Kato and Y. Matsumoto in [4] who proved that, in the case the critical locus of the function  $f$  at the origin is  $s$ , the Milnor fiber of  $f$  at the origin is  $(n - s - 1)$ -connected.

Another general, more computational, result was obtained by the first author, in [5], where it is shown that, up to homotopy, the Milnor fiber of  $f$  is obtained from the Milnor fiber of a generic hyperplane restriction  $f|_H$  by attaching  $(\Gamma_{f,H} \cdot X)_{\mathbf{0}}$   $n$ -cells, where  $(\Gamma_{f,H} \cdot X)_{\mathbf{0}}$  is the intersection number of the relative polar curve  $\Gamma_{f,H}$  with the hypersurface  $X$ . In fact, the result of [4] can be obtained directly from [5] (see [2]).

A particular case of the main result of [5] is when the polar curve is empty (or, zero, as a cycle), so that the intersection number above is zero, and the Milnor fiber of  $f$  and of  $f|_H$  have the same homotopy-type: that of a bouquet of  $(n - 1)$ -spheres.

If  $\Sigma f$  is smooth and 1-dimensional, it is trivial to show that  $\Gamma_{f,H}$  being empty is equivalent to the Milnor number of the isolated critical point of generic transverse hyperplane sections being constant along  $\Sigma f$ . In fact, if  $\Sigma f$  is 1-dimensional, one can show, using [6], that  $\Gamma_{f,H}$  being empty is equivalent to  $\Sigma f$  is smooth and the Milnor number of the isolated critical point of generic transverse hyperplane sections being constant along  $\Sigma f$ . Thus, constant transverse Milnor number implies the constancy of the cohomology of the Milnor fiber  $F_{\mathbf{p}}$  of  $f$  at points  $\mathbf{p}$  along  $\Sigma f$ .

If  $\Sigma f$  is smooth, of arbitrary dimension  $s$ , then, proceeding inductively from the 1-dimensional case, one obtains that, if the generic  $s$ -codimensional transverse slices of  $f$  have constant Milnor

---

These notes were written with the help of the Research fund of Northeastern University.

number along  $\Sigma f$ , then the reduced cohomology of the Milnor fiber  $F_{\mathbf{p}}$ , of  $f$  at  $\mathbf{p}$ , is constant along  $\Sigma f$ , and is concentrated in the single degree  $n - s$ .

What if  $\Sigma f$  is smooth, of dimension  $s$ , and the generic  $s$ -codimensional transverse slices of  $f$  have constant Milnor number on  $\Sigma f - \{\mathbf{0}\}$ , but, perhaps, the transverse slice at  $\mathbf{0}$  has a different (necessarily higher) Milnor number? If  $s \geq 2$ , then, it follows from Proposition 1.31 of [9] that, in fact, the Milnor number of the  $s$ -codimensional transverse slices of  $f$  have constant Milnor number on all of  $\Sigma f$ , i.e., there can be no jump in the transverse Milnor numbers at isolated points on a smooth critical locus of dimension at least 2. The remaining case where  $s = 1$  was addressed by the authors in [7].

In this brief paper, we address the case where:

- (1)  $\Sigma f - \{\mathbf{0}\}$  is smooth near  $\mathbf{0}$ ;
- (2)  $s \geq 3$ ;
- (3) the Milnor number of a transverse slice of codimension  $s$  of the hypersurface  $f^{-1}(0)$  is constant along  $\Sigma f - \{\mathbf{0}\}$  near  $\mathbf{0}$ ; and
- (4) the intersection of  $\Sigma$  with a sufficiently small sphere  $S_\varepsilon$  centered at  $\mathbf{0}$  is  $(s-2)$ -connected.

Under these hypotheses, we have:

**Theorem 1.1.** *The Milnor fiber  $F_{\mathbf{0}}$  of  $f$  at  $\mathbf{0}$  can have non-zero cohomology only in degrees  $0$ ,  $n - s$ ,  $n - 1$  and  $n$ .*

**Corollary 1.2.** Suppose that  $s \geq 4$  and, for a generic hyperplane  $H$ , the real link  $S_\varepsilon \cap \Sigma \cap H$  of  $\Sigma \cap H$  at  $\mathbf{0}$  is  $(s-3)$ -connected. Then, the Milnor fiber  $F_H$  of  $f|_H$  at  $\mathbf{0}$  can have non-zero cohomology only in degrees  $0$ ,  $n - s$  and  $n - 1$ .

## 2. AN EXACT SEQUENCE

Let  $\mathbb{Z}_{\mathcal{U}}^\bullet$  be the constant sheaf on  $\mathcal{U}$  with stalks isomorphic to the ring of integers  $\mathbb{Z}$ . If  $\phi_f$  is the functor of vanishing cycles of  $f$ , we know (see, e.g., [3], Theorem 5.2.21) that the complex  $\phi_f[-1]\mathbb{Z}_{\mathcal{U}}^\bullet[n+1]$  is a perverse sheaf (see, e.g., [1] p. 9) on  $f^{-1}(0)$ . Let  $\mathbf{P}^\bullet$  denote the restriction of this sheaf to its support  $\Sigma$ , which is the set of critical points of  $f$  inside  $f^{-1}(0)$ .

We know that, for all  $x \in \Sigma$ , we have

$$\mathbb{H}^{-k}(\mathbb{B}(x) \cap \Sigma; \mathbf{P}^\bullet) \cong H^{-k}(\mathbf{P}^\bullet)_x \cong \tilde{H}^{n-k}(F_x; \mathbb{Z}),$$

where  $F_x$  is the Milnor fiber of  $f$  at  $x$  and  $\mathbb{B}(x)$  is a sufficiently small ball (open or closed, with non-zero radius) of  $\mathbb{C}^{n+1}$  centered at  $x$ . Let  $\mathbb{B}^*(x) = \mathbb{B}(x) - \{x\}$ .

Then, we have the exact sequence in hypercohomology:

$$\begin{aligned} \rightarrow \mathbb{H}^{-k}(\mathbb{B}(x) \cap \Sigma, \mathbb{B}^*(x) \cap \Sigma; \mathbf{P}^\bullet) &\rightarrow \mathbb{H}^{-k}(\mathbb{B}(x) \cap \Sigma; \mathbf{P}^\bullet) \\ &\rightarrow \mathbb{H}^{-k}(\mathbb{B}^*(x) \cap \Sigma; \mathbf{P}^\bullet) \rightarrow \mathbb{H}^{-k+1}(\mathbb{B}(x) \cap \Sigma, \mathbb{B}^*(x) \cap \Sigma; \mathbf{P}^\bullet) \rightarrow \end{aligned}$$

Since  $\mathbf{P}^\bullet$  is perverse, using the cosupport condition (see e.g. [1] p. 9):

$$\mathbb{H}^{-k+1}(\mathbb{B}(x) \cap \Sigma, \mathbb{B}^*(x) \cap \Sigma; \mathbf{P}^\bullet) = 0$$

for  $-k+1 < 0$ . The support condition (see loc. cit.) leads to:

$$H^k(\mathbb{B}(x) \cap \Sigma, \mathbf{P}^\bullet) \cong \tilde{H}^{n+k}(F_x; \mathbb{Z}) = 0$$

for  $k > 0$ . Therefore,

$$\tilde{H}^{n-k}(F_x; \mathbb{Z}) \cong \mathbb{H}^{-k}(\mathbb{B}(x) \cap \Sigma; \mathbf{P}^\bullet) \cong \mathbb{H}^{-k}(\mathbb{B}^*(x) \cap \Sigma; \mathbf{P}^\bullet)$$

for  $-k+1 < 0$  and:

$$\tilde{H}^k(F_x; \mathbb{Z}) = 0$$

for  $k > n$ .

### 3. TOPOLOGICAL HYPOTHESES

Throughout the remainder of this paper, we assume, as in the introduction, that:

- (1)  $s \geq 3$  (and  $\Sigma f$  might be singular at  $\mathbf{0}$ ).
- (2) There is an open neighborhood  $\mathcal{U}$  of the origin  $\mathbf{0}$ , such that the Milnor number of a transverse slice of codimension  $s$  of the hypersurface  $f^{-1}(0)$  is constant along the singular set  $\Sigma \cap \mathcal{U} (= \Sigma f \cap \mathcal{U})$  of  $X \cap \mathcal{U}$  outside of  $\mathbf{0}$ , and equal to  $\mu$ .
- (3) The intersection of  $\Sigma$  with a sufficiently small sphere  $S_\varepsilon$  centered at  $\mathbf{0}$  is  $(s-2)$ -connected.

Note that (1) and (3) imply, in particular, that  $S_\varepsilon \cap \Sigma$  is simply-connected. Also (2) implies that  $(\Sigma - \{0\}) \cap \mathcal{U} = (\Sigma f - \{0\}) \cap \mathcal{U}$  is smooth.

As we discussed in the introduction, without the language of sheaves, the assumption on the constancy of the Milnor number of  $f$ , restricted to a normal slice to  $\Sigma$ , is equivalent to saying that our shifted, restricted vanishing cycle complex  $\mathbf{P}^\bullet|_{\Sigma - \{0\}}$  is locally constant, with stalk cohomology  $\mathbb{Z}^\mu$  concentrated in degree  $-s$ . (The technical details of the sheaf result are non-trivial; see Theorem 6.9 of [9] and Corollary 3.14 of [10].) As  $\mathbb{B}^*(\mathbf{0}) \cap \Sigma$  is homotopy-equivalent to  $S_\varepsilon \cap \Sigma$ , which is simply-connected, it follows that  $\mathbf{P}^\bullet|_{\mathbb{B}^*(\mathbf{0}) \cap \Sigma}$  is isomorphic to the shifted constant sheaf  $(\mathbb{Z}^\mu)^\bullet_{\mathbb{B}^*(\mathbf{0}) \cap \Sigma}[s]$ .

This implies that

$$\mathbb{H}^{-k}(\mathbb{B}^*(\mathbf{0}) \cap \Sigma; \mathbf{P}^\bullet) \cong H^{-k+s}(\mathbb{B}^*(0) \cap \Sigma; \mathbb{Z}^\mu) \cong H^{-k+s}(S_\varepsilon \cap \Sigma; \mathbb{Z}^\mu).$$

Thus, as  $S_\varepsilon \cap \Sigma$  is  $(s-2)$ -connected, we have:

$$\mathbb{H}^{-s}(\mathbb{B}^*(\mathbf{0}) \cap \Sigma; \mathbf{P}^\bullet) \cong H^0(S_\varepsilon \cap \Sigma; \mathbb{Z}^\mu) \cong \mathbb{Z}^\mu,$$

and, if  $2 \leq k \leq s-1$ :

$$\mathbb{H}^{-k}(\mathbb{B}^*(\mathbf{0}) \cap \Sigma; \mathbf{P}^\bullet) \cong H^{s-k}(S_\varepsilon \cap \Sigma; \mathbb{Z}^\mu) = 0.$$

### 4. PROOFS

Combining the results from the previous two sections, we find that, if the real link of the critical locus  $\Sigma$  at  $\mathbf{0}$  is  $(s-2)$ -connected and  $s \geq 3$ , then we have for the Milnor fiber  $F$  of  $f$  at  $\mathbf{0}$ :

$$\tilde{H}^{n-s}(F; \mathbb{Z}) \cong H^0(S_\varepsilon \cap \Sigma; \mathbb{Z}^\mu) \cong \mathbb{Z}^\mu;$$

$$\tilde{H}^{n-k}(F; \mathbb{Z}) = 0, \text{ if } 2 \leq k \leq s-1.$$

$$\tilde{H}^k(F; \mathbb{Z}) = 0, \text{ for } k \leq n-s-1, \text{ because of the result of [4]}$$

$$\tilde{H}^k(F; \mathbb{Z}) = 0, \text{ for } k > n, \text{ because of the support condition.}$$

This proves the theorem.

Suppose now that, in addition to our other hypotheses,  $s \geq 4$  and, for generic hyperplanes  $H$ ,  $S_\varepsilon \cap \Sigma \cap H$  is  $(s-3)$ -connected. Then,  $f|_H$  satisfies the hypotheses of the theorem, except that  $n$  is replaced by  $n-1$  and  $s$  is replaced by  $s-1$ . Thus, for the Milnor fiber  $F_H$ :

$$\tilde{H}^{n-s}(F_H; \mathbb{Z}) \cong \mathbb{Z}^\mu;$$

$$\tilde{H}^k(F_H; \mathbb{Z}) = 0, \text{ if } k \neq n-2, n-1.$$

However, by the main result of [5], the Milnor fiber  $F$  is obtained from the Milnor fiber  $F_H$  by attaching cells in dimension  $n$ . Hence,  $\tilde{H}^{n-2}(F_H; \mathbb{Z}) \cong \tilde{H}^{n-2}(F; \mathbb{Z})$ , which we know is 0. This proves the corollary.

## 5. WHEN THE CRITICAL LOCUS IS AN ICIS

Assume that the critical locus  $\Sigma$  of  $f$  is an isolated complete intersection singularity (ICIS) of dimension  $s \geq 4$ .

For an ICIS, the real link  $S_\varepsilon \cap \Sigma$  is  $(s - 2)$ -connected (see [8]). In addition, for a generic hyperplane  $H$ , the critical locus of  $f|_H$ , which equals  $\Sigma \cap H$ , will also be an ICIS, but now of dimension  $s - 1$ . Thus,  $S_\varepsilon \cap \Sigma \cap H$  is  $((s - 1) - 2)$ -connected. Therefore, we are in the situation that we have considered above.

In his preprint [12] M. Shubladze asserts that if the singular locus  $\Sigma$  of  $f$  is a complete intersection with isolated singularity at  $\mathbf{0}$  of dimension  $\geq 3$  and the Milnor number for transverse sections is 1 along  $\Sigma \setminus \{\mathbf{0}\}$ , the Milnor number of  $f$  at 0 has cohomology possibly  $\neq 0$  only in dimensions 0,  $n - s$  and  $n$ .

The results above show that, under the hypothesis of M. Shubladze, one obtains in a general way that the cohomology of the Milnor fiber of  $f$  at  $\mathbf{0}$  is possibly  $\neq 0$  in dimension 0,  $n - s$ ,  $n - 1$  and  $n$ , and a similar result as the one of M. Shubladze in dimension 0,  $n - s$ ,  $n - 1$  for the cohomology of the Milnor fiber of  $f$  restricted to a general hyperplane section if  $\dim \Sigma \geq 4$ .

Shubladze's result would follow immediately from our corollary, if it were true that every function such as that studied by Shubladze can be obtained as a generic hyperplane restriction of a function satisfying the same hypotheses. We cannot easily prove or disprove this result.

6. WHAT IF  $S_\varepsilon \cap \Sigma$  IS A HOMOLOGY SPHERE?

One might also wonder what happens if the real link of  $\Sigma$  is  $(s - 1)$ -connected. This would, in fact, imply that  $S_\varepsilon \cap \Sigma$  is a homology sphere. In this case, our earlier exact sequence immediately yields that  $\tilde{H}^{n-1}(F; \mathbb{Z}) = 0$ .

A special case of  $S_\varepsilon \cap \Sigma$  being a homology sphere would occur if  $\Sigma$  were smooth. However, in this case, when  $s \geq 2$ , Proposition 1.31 of [9] implies that the Milnor number cannot change at  $\mathbf{0}$ , i.e., we have a smooth  $\mu$ -constant family, and so the non-zero cohomology of  $F$  occurs only in degrees 0 and  $n - s$ .

## REFERENCES

- [1] A. A. Beilinson, J. Bernstein, P. Deligne, Faisceaux pervers, *Astrisque* **100** (1982), Société Mathématique de France, Paris.
- [2] D. Cheniot, Lê Dũng Tráng, Remarque sur les cycles évanouissants d'une hypersurface analytique à singularité non isolée, *C. R. Acad. Sci. Paris Sr. A - B* **277** (1973), A599 - A600.
- [3] A. Dimca, *Sheaves in Topology*, Springer-Verlag, 2004.
- [4] Mitsuyoshi Kato, Yukio Matsumoto, On the connectivity of the Milnor fiber of a holomorphic function at a critical point. *Manifolds Tokyo 1973 (Proc. Internat. Conf., Tokyo, 1973)*, pp. 131-136. Univ. Tokyo Press, Tokyo, 1975.
- [5] Lê D. T., Calcul du nombre de cycles évanouissants d'une hypersurface complexe, *Ann. Inst. Fourier (Grenoble)* **23**, 261-270, 1973.
- [6] Lê D. T. Une application d'un théorème d'A'Campo à l'équisingularité, *Nederl. Akad. Wetensch. Proc. Ser. A* **76**, *Indag. Math.* **35** (1973), 403 - 409.
- [7] Lê D. T. and D. Massey, Hypersurface Singularities and Milnor Equisingularity, *Pure and Appl. Math. Quart.*, special issue in honor of Robert MacPherson's 60th birthday, vol. **2**, no. **3**, 893-914, 2006.
- [8] E. Looijenga, *Isolated Singular Points on Complete Intersection*, Cambridge Univ. Press, 1984.
- [9] D. Massey, *Lê Cycles and Hypersurface Singularities*, *Lecture Notes in Mathematics* **1615**, Springer-Verlag, 1995.
- [10] D. Massey, Vanishing Cycles and Thom's  $a_f$  Condition, *Bull. London Math. Soc.*, **39** (4), 591-602, 2007.
- [11] J. Milnor, *Singular Points of Complex Hypersurfaces*, *Annals of Math. Studies* **77**, Princeton Univ. Press, 1968.
- [12] M. Shubladze, Singularities with critical locus an complete intersection and transversal type  $A_1$ , Preprint 2010-16, Max Planck Institute for Mathematics preprints, <http://www.mpim-bonn.mpg.de/preblob/4141>

CMI, UNIVERSITÉ DE PROVENCE, F-13453 MARSEILLE CEDEX 13, FRANCE  
*E-mail address:* `ledt@ictp.it`

DEPARTMENT OF MATHEMATICS, NORTHEASTERN UNIVERSITY, BOSTON, MA 02115, USA  
*E-mail address:* `d.massey@neu.edu`