



**HAL**  
open science

# ON A LINEAR RUNS AND TUMBLES EQUATION

Stéphane Mischler, Q Weng

► **To cite this version:**

Stéphane Mischler, Q Weng. ON A LINEAR RUNS AND TUMBLES EQUATION. 2016. hal-01272429v1

**HAL Id: hal-01272429**

**<https://hal.science/hal-01272429v1>**

Preprint submitted on 10 Feb 2016 (v1), last revised 9 Sep 2016 (v3)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# ON A LINEAR RUNS AND TUMBLES EQUATION

S. MISCHLER, Q. WENG

ABSTRACT. We consider a linear runs and tumbles equation in dimension  $d \geq 1$  for which we establish the existence of a unique positive and normalized steady state as well as its asymptotic stability, improving similar results obtained by Calvez et al. [5] in dimension  $d = 1$ . Our analysis is based on the Krein-Rutman theory revisited in [18] together with some new moment estimates for proving confinement mechanism as well as dispersion, multiplier and averaging lemma arguments for proving some regularity property of suitable iterated averaging quantities.

Version of February 10, 2016

## CONTENTS

1. Introduction and main result	1
2. Well-posedness and exponential weighted $L^1$ estimate	4
3. The stationary state problem	7
4. Asymptotic stability of the stationary state	12
References	21

## 1. INTRODUCTION AND MAIN RESULT

**1.1. The “runs and tumbles” equation in chemotaxis.** In the present paper we are interested in a kinetic evolution PDE coming from the modeling of chemotactic movement of cells in the presence of self-produced chemical substance. The so-called *runs and tumbles* model of Othmer-Dunbar-Alt [1, 19, 20, 8]

$$(1.1) \quad \partial_t f = \mathcal{L}f = -v \cdot \nabla_x f + \int_{\mathcal{V}} \{K' f' - K f\} dv'$$

describes the evolution of the distribution function of a microorganisms density  $f = f(t, x, v) \geq 0$  which at time  $t \geq 0$  and at position  $x \in \mathbb{R}^d$  move with the velocity  $v \in \mathcal{V}$ . At a microscopic description level, microorganisms move in straight line with their own velocity  $v$  which changes accordingly to a jump process of parameter  $K = K(x, v, v') \geq 0$ . Here and below, we used the shorthands  $f' = f(t, x, v')$  and  $K' = K(x, v', v)$ . For the sake of simplicity, we assume that  $\mathcal{V} \subset \mathbb{R}^d$  is the centered ball with unit volume ( $\mathcal{V} := B(0, V_0)$ ) with  $V_0$  chosen such that  $|\mathcal{V}| = 1$ ). We complement the evolution PDE (1.1) with an initial condition

$$(1.2) \quad f(0, \cdot) = f_0 \quad \text{on} \quad \mathbb{R}^d \times \mathcal{V}.$$

---

2010 *Mathematics Subject Classification.* 35B40, 35Q92, 47D06, 92C17.

*Key words and phrases.* Kinetic equations, velocity-jump processes, chemotaxis, stationary state, asymptotic stability, hypocoercivity.

At least formally, for any multiplier  $\varphi = \varphi(x, v)$ , we have

$$\frac{d}{dt} \int f \varphi = \int f v \cdot \nabla_x \varphi + \int f K \int_{\mathcal{V}} \{\varphi' - \varphi\}.$$

In particular, choosing  $\varphi \equiv 1$  in the above identity, we see that the total mass is conserved and we may assume that it is normalized to the unit, namely

$$(1.3) \quad \langle\langle f(t, \cdot) \rangle\rangle = \langle\langle f_0 \rangle\rangle = 1, \quad t \geq 0,$$

where for functions  $g = g(x, v)$  and  $h = h(v)$ , we define

$$\langle h \rangle = \int_{\mathcal{V}} h \, dv, \quad \langle\langle g \rangle\rangle = \int_{\mathbb{R}^d} \langle g(x, \cdot) \rangle \, dx.$$

The precise form of the turning kernel  $K$  depends upon the space concentration  $S$  of a chemical agent: microorganisms have the tendency to move to where the chemical concentration is higher. More specifically, we assume that the turning kernel is given by

$$(1.4) \quad K = K[S](v) := 1 - \chi \Phi(v \cdot \nabla_x S(x)), \quad \chi \in (0, 1), \quad \Phi(y) = \text{sign}(y),$$

where the sign function is defined by  $\Phi(y) = -1$  if  $y < 0$  and  $\Phi(y) = 1$  if  $y > 0$ . In other words, the turning kernel  $K$  takes the two values  $1 \pm \chi$  depending on the velocity direction of the microorganism with respect to the gradient of the chemical concentration.

When the chemical agent is produced by the microorganisms themselves, it is usually assumed to be given as the solution to the damped Poisson equation

$$(1.5) \quad -\Delta S + S = \varrho := \int_{\mathcal{V}} f \, dv,$$

so that the evolution of the microorganisms density  $f$  is given by the coupled system of equations (1.1)-(1.4)-(1.5). We refer the reader interested by the well-posedness issue for related models to the review paper [4] and the references quoted therein. We also refer to [13] for related modelling considerations. Concerning the qualitative behaviour of the solutions it seems that the unique available information is the mass conservation (1.3). One of the main difficulty comes from the fact that both equations (1.1) and (1.5) are invariant by translations so that the expected confinement mechanism seems to be hard to prove.

On the other way round, one can see that for a given solution  $(f, S)$  of (1.1)-(1.4)-(1.5) and for any rotation  $R \in SO(d)$  the couple  $(f_R, S_R)$  is also a solution of the same equations, where we have set  $f_R(x, v) := f(Rx, Rv)$  and  $S_R(x) := S(Rx)$ . In particular, an invariant by rotations initial datum  $f_0$  gives rise to an invariant by rotations solution  $(f, S)$ .

More specifically, we may observe that in the case when  $S$  is radially symmetric and strictly decreasing in the position variable (which is the case if the density  $\varrho$  satisfies the same properties thanks to the maximum principle), we have

$$-\Phi(v \cdot \nabla_x S) = -\Phi(-v \cdot x) = \text{sign}(x \cdot v),$$

and thus the associated turning kernel writes

$$(1.6) \quad K = K(x, v) := 1 + \chi \zeta, \quad \chi \in (0, 1), \quad \zeta = \zeta(x, v) = \text{sign}(x \cdot v).$$

Such a kernel has been introduced in [5] and the associated (now linear!) evolution equation (1.1)-(1.6) has then been analyzed in dimension  $d = 1$ : the existence of a unique (positive and normalized) steady state has been established and its asymptotic exponential stability has been proved.

**1.2. The linear “runs and tumbles” equation.** The main purpose of the present work is to provide an alternative approach to study the linear “runs and tumbles” (linear RT) equation (1.1)-(1.6) which makes possible to generalize the analysis of [5] to any dimension  $d \geq 1$ . In order to state our main result, we introduce some notations and the functional framework we will work with.

First, we denote by  $m$  some weight function which is either a polynomial or an exponential

$$(1.7) \quad m(x) = \langle x \rangle^k, \quad k > 0, \quad \text{or} \quad m(x) = \exp(\gamma \langle x \rangle), \quad \gamma \in (0, \gamma^*),$$

for some positive constant  $\gamma^*$  which will be defined latter, and where  $\langle x \rangle^2 = 1 + |x|^2$ . To a given weight  $m$  we define the associated rate function  $\Theta_m$  and weight function  $\omega$  by

$$\begin{aligned} \Theta_m(t) &:= \langle t \rangle^{-\ell}, \quad \ell \in (0, k), & \omega &= 1 & \text{if} & \quad m = \langle x \rangle^k; \\ \Theta_m(t) &= e^{at}, \quad a < 0, & \omega &= m & \text{if} & \quad m = e^{\gamma \langle x \rangle}. \end{aligned}$$

Finally for given weight function  $m = m(x) : \mathbb{R}^d \rightarrow \mathbb{R}_+$  and exponent  $1 \leq p \leq \infty$ , we define the associated weighted Lebesgue space  $L^p(m)$  and weighted Sobolev space  $W^{1,p}(m)$ , through their norms

$$(1.8) \quad \|f\|_{L^p(m)} := \|mf\|_{L^p}, \quad \|f\|_{W^{1,p}(m)} := \|mf\|_{W^{1,p}}.$$

We use the shorthands  $L_k^p = L^p(m)$ , when  $m = \langle x \rangle^k$ , and  $H^1(m) = W^{1,2}(m)$ . We write  $a \lesssim b$  if there exists a positive constant  $C$  such that  $a \leq Cb$ .

**Theorem 1.1.** *There exists  $\gamma^* > 0$  and there exists a unique positive, invariant by rotations and normalized stationary state*

$$\begin{aligned} 0 < G &\in L^1(m_0) \cap L^\infty, \quad \langle\langle G \rangle\rangle = 1, \\ -v \cdot \nabla_x G &+ \int_{\mathcal{V}} \{K'G' - KG\} dv' = 0, \end{aligned}$$

where  $m_0$  stands for the exponential weight function  $m_0(x) := \exp(\gamma_* \langle x \rangle)$ . Moreover, for any weight function  $m$  satisfying (1.7) and for any  $0 \leq f_0 \in L^1(m)$ , there exists a unique solution  $f \in C([0, \infty); L^1(m))$  to the equation (1.1)-(1.6) associated to the initial datum  $f_0$  and

$$(1.9) \quad \|f(t) - \langle\langle f_0 \rangle\rangle G\|_{L^1(\omega)} \leq \Theta_m(t) \|f_0\|_{L^1(m)}, \quad \forall t \geq 0,$$

where  $\omega$  and  $\Theta_m$  are defined just above.

The present result generalizes to any dimension  $d \geq 1$  similar results ([5, Theorem 2.1] and [5, Proposition 1]) obtained by Calvez et al. in dimension  $d = 1$ . As pointed out in [5], the main novelty and mathematical interest of the model lie in the fact that the confinement is achieved by a biased velocity jump process, where the bias replaces the confining acceleration field which is the classical confinement mechanism for Boltzmann and Fokker-Planck models, see for instance [23, 16, 7] and the references quoted therein.

Our strategy is different from the one used in [5] and it is adapted from the Krein-Rutman theory developed in [18] for positive semigroups which do not fulfill the classical compactness assumption on the associated resolvent but have a nice splitting structure. The main difficulty is then to get suitable estimates on some related operators and semigroups.

The first step consists in proving a weighted  $L^1$  bound which brings out the confinement mechanism. That is the main new bound which is in the spirit of weighted  $L^p$  estimates

obtained for performing similar spectral analysis in [12, 18, 16] for Boltzmann, growth-fragmentation and kinetic Fokker-Planck models.

Next, in order to apply the Krein-Rutman theory of [18], we adapt to our context a series of regularity properties known for solutions to the free transport equation. We successively use a dispersion argument as introduced by Bardos and Degond [2] for providing better integrability in the position variable (transfer of integrability from the velocity variable to the position variable), next a multiplier method in the spirit of Lions-Perthame multiplier (see [21, 14, 22]) for improving again the integrability estimate in the position variable near the origin and finally a space variable averaging lemma in the spirit of the variant [6, 3] of the classical time and space averaging lemma of Golse et al [11, 10]. It is worth emphasizing that the needed regularity estimate is not obtained using an abstract hypo-coercivity operator as in [7, 5] nor using an iterated averaging lemma as in [12] (which allows to transfer regularity from the velocity variable to the position variable thanks to a suitable commutator and the associated “gliding norms”) but using the more classical (and more robust) averaging lemma.

Finally, it is worth pointing out that it is not clear how to use the above analysis in order to make any progress in the understanding of the nonlinear equation (1.1)-(1.4)-(1.5). In particular, we have not been able to prove that the chemical agent density  $S$  is decaying with respect to the radial variable  $|x|$  and thus that  $G$  is also a stationary state of the nonlinear equation (1.1)-(1.4)-(1.5), as one can expect by making an analogy with the case when the velocity set  $\mathcal{V}$  is replaced by  $\bar{\mathcal{V}} := S^{d-1}$ . Indeed, in that case, we may observe that for  $\bar{G}(x, v) = g(|x|) = C e^{-\chi|x|}$ , we have

$$v \cdot \nabla_x \bar{G} = v \cdot \frac{x}{|x|} g'(|x|) = -\chi \operatorname{sign}(x \cdot v) g(|x|) = \int_{\bar{\mathcal{V}}} \{K' \bar{G} - K \bar{G}\} dv',$$

so that we have exhibited an explicit (unique, positive and unit mass) stationary state  $\bar{G}$ . The associated macroscopic density  $\bar{\varrho}$  is then decaying and thus also the associated chemical agent density  $\bar{S}$  (thanks to the maximum principle applied to the elliptic equation (1.5)). It turns out then that  $G$  is also a stationary state of the nonlinear equation (1.1)-(1.4)-(1.5).

Let us end the introduction by describing the plan of the paper. In Section 2, we mainly present the weighted  $L^1$  estimate which highlights the confinement mechanism of the model. In Section 3, we introduce a first splitting of the generator in order to prove the existence (and next the uniqueness) of a positive stationary state. Finally, in Section 4, we introduce a second splitting of the generator which enjoys better smoothness properties and for which we can use the Krein-Rutman theory revisited in [18] and conclude to the asymptotic stability of the stationary state.

**Acknowledgments.** We thank V. Clavez for fruitful discussions which have been a source of the present work. The research leading to this paper was (partially) funded by the French “ANR blanche” project Kibord: ANR-13-BS01-0004.

## 2. WELL-POSEDNESS AND EXPONENTIAL WEIGHTED $L^1$ ESTIMATE

We first state a well-posedness result concerning the linear RT equation (1.1)-(1.6), whose very classical proof is skipped, and we recall some notations and definition.

**Lemma 2.1.** *For any  $f_0 \in L^p(m)$ ,  $1 \leq p \leq \infty$ , there exists a unique weak (distributional) solution  $f \in C([0, T]; L^1_{loc}) \cap L^\infty(0, T; L^p(m))$ ,  $\forall T \in (0, \infty)$ , to the linear RT equation (1.1)-(1.6) which furthermore satisfies:*

- (1)  $f(t, \cdot) \geq 0$  for any  $t \geq 0$  if  $f_0 \geq 0$  (preservation of positivity);  
(2)  $\langle\langle f(t, \cdot) \rangle\rangle = \langle\langle f_0 \rangle\rangle$  for any  $t \geq 0$  if  $L^p(m) \subset L^1$  (mass conservation);  
(3)  $f \in C(\mathbb{R}_+; L^p(m))$  and we may associate to  $\mathcal{L}$  a continuous semigroup  $S_{\mathcal{L}}$  in  $L^p(m)$  by setting  $S_{\mathcal{L}}(t)f_0 := f(t, \cdot)$  for any  $t \geq 0$  and  $f_0 \in L^p(m)$ . Here the continuity has to be understood in the strong (norm) topology sense when  $p \in [1, \infty)$  and in the weak  $\sigma(L^\infty, L^1)^*$  topology sense when  $p = \infty$ .

In the sequel, we will thus associate to the generator  $\mathcal{L}$  (and next to the related generators  $\mathcal{B}_0, \mathcal{B}_1, \mathcal{B}, \dots$ ) a semigroup  $S_{\mathcal{L}}$  and to any initial datum  $f_0 \in L^p(m)$  we will denote by  $f(t) = f_{\mathcal{L}}(t) = S_{\mathcal{L}}(t)f_0$  the solution associated to the related abstract evolution equation, which is then also a distributional, weak and renormalized solution to the corresponding PDE equation. More precisely, the above function  $f$  satisfies the linear RT equation (1.1) in the following renormalized sense

$$\frac{d}{dt} \int_{\mathbb{R}^d \times \mathcal{V}} \beta(f) \varphi = \int_{\mathbb{R}^d \times \mathcal{V}} \beta(f) v \cdot \nabla_x \varphi + \int_{\mathbb{R}^d \times \mathcal{V}} f K \int_{\mathcal{V}} [\varphi' \beta(f') - \varphi \beta'(f)]$$

for any (renormalizing) Lipschitz function  $\beta : \mathbb{R} \rightarrow \mathbb{R}$  and for any (test) function  $\varphi \in C_c^1(\mathbb{R}^d \times \mathcal{V})$ . It is worth mentioning that we deduce the preservation of positivity property by just choosing  $\beta(s) = s_+$ ,  $\varphi = 1$ , in the above identity and next using the Gronwall lemma. When  $L^p(m) \subset L^1$ , the uniqueness result follows from choosing  $\beta(s) = |s|$  and  $\varphi = 1$  in the above identity. The existence part can be achieved by combining the characteristics method for the free transport equation and a perturbation by bounded operators argument. In the sequel, we will denote  $\mathcal{B}(X, Y)$  the space of bounded linear operators from a Banach space  $X$  into a Banach space  $Y$ , and we write  $\mathcal{B}(X) = \mathcal{B}(X, X)$ . Another way to prove existence and uniqueness consists in using the Hille-Yosida theorem for maximal dissipative unbounded operators. We recall for futur references that we say that an unbounded operator  $\mathcal{L}$  with dense domain  $D(\mathcal{L}) \subset X$  is  $a$ -dissipative,  $a \in \mathbb{R}$ , if

$$\forall f \in D(\mathcal{L}), \exists f^* \in J_f \quad \langle f^*, \mathcal{L}f \rangle_{X', X} \leq a \|f\|_X^2,$$

where  $J_f$  denotes the (nonempty) dual set

$$J_f := \{g \in X'; \langle g, f \rangle_{X', X} = \|g\|_{X'}^2 = \|f\|_X^2\}.$$

We now establish a uniform in time exponential weighted  $L^1$  estimate, which is one of the cornerstone arguments of the proof of our main theorem.

**Lemma 2.2.** *There exists a constant  $\gamma^* > 0$  and for any  $\gamma \in (0, \gamma^*)$  there exist a weight function  $\tilde{m}$  such that  $\tilde{m}(x) \sim e^{\gamma(x)}$  as  $x \rightarrow \infty$  and a constant  $C \in (0, \infty)$  such that the solution  $f$  to the linear RT equation with initial datum  $f_0 \in L^1(m)$  satisfies*

$$(2.1) \quad \int |f(t)| \tilde{m} \leq \max\left(C \int |f_0|, \int |f_0| \tilde{m}\right), \quad \forall t \geq 0.$$

*In particular, the semigroup  $S_{\mathcal{L}}$  is bounded in  $L^1(m)$ .*

*Proof of Lemma 2.2.* We define the dual operator  $\mathcal{L}^*$  by

$$\int (\mathcal{L}^* \varphi) f = \int (\mathcal{L} f) \varphi, \quad \forall \varphi \in W^{1, \infty}(\mathbb{R}^d \times \mathcal{V}), f \in C_c(\mathbb{R}^d \times \mathcal{V}),$$

so that

$$(\mathcal{L}^* \varphi)(x, v) := v \cdot \nabla_x \varphi + K \int_{\mathcal{V}} \{\varphi' - \varphi\} dv'.$$

For a given  $\gamma > 0$ , we compute

$$\mathcal{L}^* e^{\gamma\langle x \rangle} = \gamma(v \cdot \frac{x}{\langle x \rangle}) e^{\gamma\langle x \rangle},$$

and next

$$\begin{aligned} \mathcal{L}^* \left[ (v \cdot \frac{x}{\langle x \rangle}) e^{\gamma\langle x \rangle} \right] &= v \cdot \nabla_x [(v \cdot \frac{x}{\langle x \rangle}) e^{\gamma\langle x \rangle}] - K (v \cdot \frac{x}{\langle x \rangle}) e^{\gamma\langle x \rangle} \\ &= \left( \frac{|v|^2}{\langle x \rangle} - \frac{(v \cdot x)^2}{\langle x \rangle^3} + \gamma \frac{(v \cdot x)^2}{\langle x \rangle^2} - (v \cdot \frac{x}{\langle x \rangle}) - \chi \frac{|v \cdot x|}{\langle x \rangle} \right) e^{\gamma\langle x \rangle}. \end{aligned}$$

Defining

$$(2.2) \quad V_1 := \int_{\mathcal{V}} |v'_1| dv',$$

we finally have

$$\begin{aligned} \mathcal{L}^* \left[ \frac{|v \cdot x|}{\langle x \rangle} e^{\gamma\langle x \rangle} \right] &= v \cdot \nabla_x \left[ \frac{|v \cdot x|}{\langle x \rangle} e^{\gamma\langle x \rangle} \right] + (1 + \chi\zeta) \left( V_1 \frac{|x|}{\langle x \rangle} - \frac{|v \cdot x|}{\langle x \rangle} \right) e^{\gamma\langle x \rangle} \\ &= \zeta \left( \frac{|v|^2}{\langle x \rangle} - \frac{|v \cdot x|^2}{\langle x \rangle^3} + \gamma \frac{|v \cdot x|^2}{\langle x \rangle^2} \right) e^{\gamma\langle x \rangle} \\ &\quad + \left( (1 + \chi\zeta) V_1 \frac{|x|}{\langle x \rangle} - (1 + \chi\zeta) \frac{|v \cdot x|}{\langle x \rangle} \right) e^{\gamma\langle x \rangle}. \end{aligned}$$

Consider  $\beta, \gamma \in (0, 1)$  small enough such that the weight function

$$(2.3) \quad \tilde{m} := \left( 1 + \gamma(v \cdot \frac{x}{\langle x \rangle}) - \beta \frac{|v \cdot x|}{\langle x \rangle} \right) e^{\gamma\langle x \rangle}$$

satisfies

$$(2.4) \quad e^{\gamma\langle x \rangle} (1 - \delta) \leq \tilde{m} \leq (1 + \delta) e^{\gamma\langle x \rangle},$$

for some  $\delta \in (0, 1)$ , which is possible because  $\mathcal{V}$  is a bounded set. Gathering the previous identities, we find

$$\begin{aligned} (\mathcal{L}^* \tilde{m}) e^{-\gamma\langle x \rangle} &= \gamma \left( \frac{|v|^2}{\langle x \rangle} - \frac{(v \cdot x)^2}{\langle x \rangle^3} + \gamma \frac{(v \cdot x)^2}{\langle x \rangle^2} - \chi \frac{|v \cdot x|}{\langle x \rangle} \right) \\ &\quad - \beta \zeta \left( \frac{|v|^2}{\langle x \rangle} - \frac{|v \cdot x|^2}{\langle x \rangle^3} + \gamma \frac{|v \cdot x|^2}{\langle x \rangle^2} \right) - \beta \left( (1 + \chi\zeta) V_1 \frac{|x|}{\langle x \rangle} - (1 + \chi\zeta) \frac{|v \cdot x|}{\langle x \rangle} \right) \\ &\leq \gamma V_0^2 \left( \frac{1}{\langle x \rangle} + \gamma \right) + \beta V_0^2 \left( \frac{2}{\langle x \rangle} + \gamma \right) + \beta(1 + \chi) \frac{|v \cdot x|}{\langle x \rangle} \\ &\quad - \gamma\chi \frac{|v \cdot x|}{\langle x \rangle} - \beta(1 - \chi) V_1 \frac{|x|}{\langle x \rangle} \\ &\leq [V_0^2(\gamma + 2\beta) + \beta(1 - \chi)V_1] \frac{1}{\langle x \rangle} + [\gamma^2 V_0^2 + \beta\gamma V_0^2 - \beta(1 - \chi)V_1], \end{aligned}$$

and thus

$$(2.5) \quad (\mathcal{L}^* \tilde{m}) e^{-\gamma\langle x \rangle} \leq \frac{C}{\langle x \rangle} - 2\alpha,$$

by choosing  $\beta(1 + \chi) = \gamma\chi$  and  $\gamma > 0$  small enough so that  $2\alpha := \beta(1 - \chi)V_1 - \gamma^2 V_0^2 - \beta\gamma V_0^2 > 0$ . Observing that

$$\left( \frac{C}{\langle x \rangle} - 2\alpha \right) e^{\gamma\langle x \rangle} \leq C e^{\gamma\langle R \rangle} \mathbf{1}_{B(0,R)} - \alpha(1 + \delta) e^{\gamma\langle x \rangle} \leq A - \alpha\tilde{m},$$

for some constant  $A > 0$ , we have proved

$$\mathcal{L}^* \tilde{m} \leq A - \alpha \tilde{m}, \quad \alpha < 0.$$

We consider now  $f$  the solution to the linear RT equation (1.1) associated to  $f_0 \in L^1(m)$ . Denoting  $g := |f|$ , we deduce from the above inequality that

$$\frac{d}{dt} \int g \tilde{m} = \int (\mathcal{L}f) (\text{sign}f) \tilde{m} \leq \int (\mathcal{L}g) \tilde{m} \leq A \int g - \alpha \int g \tilde{m}.$$

As a consequence, (2.1) holds with  $C := A/\alpha$ .  $\square$

### 3. THE STATIONARY STATE PROBLEM

We introduce two generators  $\mathcal{B}_0$  and  $\mathcal{B}_1$  in the next paragraphs and we study the regularity properties of the associated semigroups. We then use these estimates in order to prove the existence of a stationary state.

**3.1. The operator  $\mathcal{B}_0$  and the associated semigroup  $S_{\mathcal{B}_0}$ .** We define  $\mathcal{B}_0$  by

$$\mathcal{B}_0 f := -v \cdot \nabla_x f - K f.$$

**Lemma 3.1.** *There exist  $\gamma^* > 0$  and  $a^* < 0$ , such that for any  $1 \leq p \leq \infty$ ,  $m = e^{\gamma \langle x \rangle}$ ,  $\gamma \in [0, \gamma^*)$ , there holds*

$$\|S_{\mathcal{B}_0}(t)\|_{L^p(m) \rightarrow L^p(m)} \lesssim e^{at}, \quad \forall a > a^*.$$

*Proof of Lemma 3.1.* We consider a solution  $f = S_{\mathcal{B}_0}(t)f_0$  to the evolution equation associated to  $\mathcal{B}_0$  and we compute

$$\begin{aligned} \frac{d}{dt} \int |f|^p m^p &= \int |f|^p p [v \cdot \nabla_x m - K m] m^{p-1} \\ &\leq p [V_0 \gamma + \chi - 1] \int |f|^p m^p, \end{aligned}$$

from which we easily conclude thanks to the Gronwall lemma.  $\square$

For any  $\varphi \in L_{xv}^\infty$ , we define  $A = A_\varphi : L_x^q L_v^1 \rightarrow L_x^q$  by

$$A f = A_\varphi f(x) := \int_{\mathcal{V}} \varphi(x, v') f(x, v') dv'.$$

Given some Banach spaces  $X_i$  and two functions  $u \in L^1(\mathbb{R}_+; \mathcal{B}(X_2, X_3))$ ,  $v \in L^1(\mathbb{R}_+; \mathcal{B}(X_1, X_2))$ , we define the convolution function  $u * v \in L^1(\mathbb{R}_+; \mathcal{B}(X_1, X_3))$  by

$$(u * v)(t) := \int_0^t u(t-s) v(s) ds.$$

We also define  $u^{*n}$  by  $u^{*1} = u$  and  $u^{*n} = u^{*(n-1)} * u$  for  $n \geq 2$ .

**Lemma 3.2.** *There exists  $a^* < 0$  such that for any  $\varphi \in L_{xv}^\infty$  and  $m = e^{\gamma \langle x \rangle}$ ,  $\gamma \in [0, \gamma^*)$ , there holds*

$$(3.1) \quad \|A_\varphi S_{\mathcal{B}_0}(t)\|_{L_x^1 L_v^\infty(m) \rightarrow L_x^\infty L_v^\infty} \lesssim t^{-d} e^{at}, \quad \forall t > 0, \forall a > a^*.$$

*As a consequence, there exists  $n \in \mathbb{N}^*$  ( $n = d + 2$  is suitable) such that*

$$(3.2) \quad \|(A_\varphi S_{\mathcal{B}_0})^{(*n)}(t)\|_{L_{xv}^1(m) \rightarrow L_{xv}^\infty} \lesssim e^{at}, \quad \forall t \geq 0, \forall a > a^*.$$



*Proof of Lemma 3.2.* We split the proof into two steps.

*Step 1.* We adapt the classical dispersion result of Bardos and Degond [2]. We denote  $f(t) := S_{\mathcal{B}_0}(t)f_0$  the solution to the damped transport equation

$$\partial_t f + v \cdot \nabla_x f + Kf = 0, \quad f(0) = f_0.$$

The characteristics method gives the representation formula

$$(S_{\mathcal{B}_0}(t)f_0)(x, v) = f_0(x - vt, v) e^{-\int_0^t K(x-vs, v) ds}.$$

We then have

$$\rho(t, x) := (AS_{\mathcal{B}_0}(t)f_0)(x) = \int_{\mathcal{V}} \varphi(x, v_*) f_0(x - v_*t, v_*) e^{-\int_0^t K(x-v_*s, v_*) ds} dv_*.$$

Using that  $K(x, v) \geq (1 - \chi)$ , we deduce

$$\begin{aligned} |\rho(t, x)| &\leq e^{-(1-\chi)t} \int_{\mathcal{V}} |\varphi(x, v_*)| \left[ \sup_{w \in \mathcal{V}} |f_0|(x - v_*t, w) \right] dv_* \\ &\leq e^{-(1-\chi)t} \|\varphi(x, \cdot)\|_{L_v^\infty} \int_{\mathcal{V}} \left[ \sup_{w \in \mathcal{V}} |f_0|(y, w) \right] \frac{dy}{t^d}, \end{aligned}$$

and we conclude with

$$\|AS_{\mathcal{B}_0}(t)f_0\|_{L_{xv}^\infty} \lesssim \frac{e^{-(1-\chi)t}}{t^d} \|f_0\|_{L_x^1 L_v^\infty},$$

which in particular implies (3.1).

*Step 2.* From Lemma 3.1, for  $r = 1$  and  $r = \infty$ , we clearly have

$$\|AS_{\mathcal{B}_0}(t)\|_{L_x^r L_v^\infty(m^{1/r}) \rightarrow L_x^r L_v^\infty(m^{1/r})} \lesssim e^{at}.$$

Gathering that estimate with (3.1), we may repeat the proof of [17, Proposition 2.2] (see also [16, Lemma 2.4] or [15]) and we get

$$(3.3) \quad \|(AS_{\mathcal{B}_0})^{*(d+1)}(t)\|_{L_x^1 L_v^\infty(m) \rightarrow L_x^\infty L_v^\infty} \lesssim e^{at}.$$

Observing that

$$(3.4) \quad \|(AS_{\mathcal{B}_0})(t)\|_{L_x^1 L_v^1(m) \rightarrow L_x^1 L_v^\infty(m)} \lesssim e^{at},$$

thanks to Lemma 3.1 and  $A : L_{xv}^1 \rightarrow L_x^1 L_v^\infty$ , we conclude to (3.2) by taking  $n = d + 2$ .  $\square$

**3.2. The operator  $\mathcal{B}_1$  and the associated semigroup  $S_{\mathcal{B}_1}$ .** We define  $\mathcal{B}_1$  by

$$(3.5) \quad \mathcal{B}_1 f := -v \cdot \nabla_x f - Kf + (1 - \phi_R) \int_{\mathcal{V}} K' f' dv',$$

where we have defined the truncation functions  $\phi_R(x) := \phi(x/R)$  for a given  $\phi \in \mathcal{D}(\mathbb{R}^d)$  which is radially symmetric and satisfies  $\mathbf{1}_{B(0,1)} \leq \phi \leq \mathbf{1}_{B(0,2)}$ .

**Lemma 3.3.** *There exist  $\gamma^* > 0$ ,  $a^* < 0$  and  $C \geq 1$  such that for  $R$  large enough and  $m = e^{\gamma \langle x \rangle}$ ,  $\gamma \in (0, \gamma^*)$ , the semigroup  $S_{\mathcal{B}_1}$  satisfies the following growth estimate*

$$\|S_{\mathcal{B}_1}(t)\|_{L^1(m) \rightarrow L^1(m)} \leq C e^{a^* t}, \quad \forall t \geq 0.$$

*Proof of Lemma 3.3.* We observe that the dual operator  $\mathcal{B}_1^*$  writes

$$\mathcal{B}_1^* \varphi = \mathcal{L}^* \varphi - \phi_R K \varphi.$$

Defining the modified weight function  $\tilde{m}$  as in (2.3) and using the inequalities (2.4) and (2.5), we get

$$\begin{aligned} \mathcal{B}_1^* \tilde{m} &\leq \left( \frac{C}{\langle x \rangle} - 2\alpha \right) e^{-\gamma \langle x \rangle} - \phi_R K \tilde{m} \\ &\leq \left( \frac{C}{\langle x \rangle} - 2\alpha \right) e^{-\gamma \langle x \rangle} - \mathbf{1}_{B_R(0)} (1 - \chi) (1 - \delta) e^{-\gamma \langle x \rangle} \\ &\leq -\alpha \tilde{m}, \end{aligned}$$

for  $\gamma > 0$  small enough and  $R > 1$  large enough, because  $C = \mathcal{O}(\gamma)$  and  $\delta = \mathcal{O}(\gamma)$ . We then have proved that  $\mathcal{B}_1$  is dissipative in  $L^1(\tilde{m})$  and we immediately conclude.  $\square$

**Lemma 3.4.** *For the same constants  $\gamma^* > 0$  and  $a^* < 0$  as defined in Lemma 3.3, for any  $1 \leq p \leq \infty$  and  $m = e^{\gamma \langle x \rangle}$ ,  $\gamma \in (0, \gamma^*)$ , the semigroup  $S_{\mathcal{B}_1}$  satisfies the growth estimate*

$$(3.6) \quad \|S_{\mathcal{B}_1}(t)\|_{L^1(m) \cap L^p(m^{1/p}) \rightarrow L^1(m) \cap L^p(m^{1/p})} \leq C e^{at}, \quad \forall t \geq 0, \quad \forall a > a^*.$$

*Proof of Lemma 3.4.* We set  $X := L^1(m) \cap L^\infty$  and we write

$$\mathcal{B}_1 = \mathcal{B}_0 + \mathcal{A}_0^c,$$

with  $\mathcal{A}_0^c = A_\psi$ ,  $\psi := (1 - \phi_R)K(x, v)$ , and then the iterated Duhamel formula

$$S_{\mathcal{B}_1} = \{S_{\mathcal{B}_0} + \dots + S_{\mathcal{B}_0} * (\mathcal{A}_0^c S_{\mathcal{B}_0})^{(*n)}\} + S_{\mathcal{B}_0} * (\mathcal{A}_0^c S_{\mathcal{B}_0})^{(*n)} * \mathcal{A}_0^c S_{\mathcal{B}_1} =: U_1 + U_2,$$

with  $n = d + 2$ . From Lemma 3.1, (3.2) in Lemma 3.2 and Lemma 3.3, we easily deduce that

$$\|U_2\|_{L^1(m) \rightarrow L^\infty} \leq \|S_{\mathcal{B}_0}\|_{L^\infty \rightarrow L^\infty} * \|(\mathcal{A}_0^c S_{\mathcal{B}_0})^{(*n)}\|_{L^1 \rightarrow L^\infty} * \|\mathcal{A}_0^c S_{\mathcal{B}_1}\|_{L^1 \rightarrow L^1} \lesssim e^{at},$$

for any  $a > a^*$ . We have similarly the same decay estimate on the remainder term  $U_1$  in  $\mathcal{B}(X)$  by just using Lemma 3.1. We deduce that (3.6) holds for  $p = \infty$ . We conclude that (3.6) holds for any  $1 \leq p \leq \infty$  by interpolating that first estimate in  $L^\infty$  together with the estimate established in Lemma 3.3.  $\square$

**Lemma 3.5.** *For the same constants  $\gamma^* > 0$  and  $a^* < 0$  as defined in Lemma 3.3, for any  $\varphi \in L_{xv}^\infty$  and  $m = e^{\gamma \langle x \rangle}$ ,  $\gamma \in [0, \gamma^*)$ , there holds*

$$(3.7) \quad \|A_\varphi S_{\mathcal{B}_1}(t)\|_{L_x^1 L_v^\infty(m) \rightarrow L_{xv}^\infty} \lesssim t^{-d} e^{at}, \quad \forall t \geq 0, \quad \forall a > a^*.$$

*As a consequence, for  $n \in \mathbb{N}^*$  large enough ( $n = d + 2$  is suitable), there holds*

$$(3.8) \quad \|(A_\varphi S_{\mathcal{B}_1})^{(*n)}(t)\|_{L^1(m) \rightarrow L^\infty} \lesssim e^{at}, \quad \forall t \geq 0, \quad \forall a > a^*.$$

*Proof of Lemma 3.5.* With the notation of the proof of Lemma 3.4, we write

$$A_\varphi S_{\mathcal{B}_1} = A_\varphi S_{\mathcal{B}_0} + A_\varphi S_{\mathcal{B}_0} * \mathcal{A}_0^c S_{\mathcal{B}_1},$$

and we immediately conclude that (3.7) holds putting together (3.1) and the fact that  $\mathcal{A}_0^c S_{\mathcal{B}_1}$  has the appropriate decay rate in  $\mathcal{B}(L^1(m); L_x^1 L_v^\infty(m))$  thanks to Lemma 3.3. Introducing the notations  $X_1 := L_x^1 L_v^\infty(m)$  and  $X_\infty := L_x^1 L_v^\infty(m) \cap L_{xv}^\infty$ , we then easily see from (3.6) and (3.7) that

$$\|A_\varphi S_{\mathcal{B}_1}\|_{X_p \rightarrow X_q} \lesssim \Theta_{p,q}(t) e^{at}, \quad \forall a > a^*,$$

for  $(p, q) = (1, 1), (1, \infty), (\infty, \infty)$ , with  $\Theta_{1,1} = \Theta_{\infty, \infty} = 1$  and  $\Theta_{1, \infty} = t^{-d}$ . Repeating again the proof of [17, Proposition 2.2], we deduce

$$\|(A_\varphi S_{\mathcal{B}_1})^{(*n-1)}(t)\|_{X_1 \rightarrow X_\infty} \lesssim e^{at},$$

and we then conclude by using that  $A_\varphi S_{\mathcal{B}_1}$  has the appropriate decay rate in  $\mathcal{B}(L^1(m); L_x^1 L_v^\infty(m))$  thanks to Lemma 3.3.  $\square$

**3.3. Existence of a steady state.** We establish the existence of a steady state thanks to a fixed point argument. We fix an exponential weight  $m := e^{\gamma(x)}$ ,  $\gamma \in (0, \gamma^*)$ . We define the Banach space  $X := L^1(m) \cap L^\infty$  as well as

$$\forall f \in X, \quad \|f\| := \sup_{t \geq 0} \|S_{\mathcal{L}}(t)f\|_X.$$

**Lemma 3.6.** *The quantity  $\| \cdot \|$  defines a norm on  $X$  which is equivalent to its usual norm  $\| \cdot \|_X$ . For  $C \geq 1$  large enough, the set*

$$\mathcal{C} := \left\{ f \in L^1(m) \cap L^\infty; f \geq 0, \langle f \rangle = 1, f_R = f, \forall R \in SO(d), \|f\| \leq C \right\}$$

*is a weakly  $*$  compact and convex set of  $X$  which is furthermore left invariant by the action of the flow associated to the linear RT equation (1.1)-(1.6). As a consequence, there exists at least one invariant element  $G \in \mathcal{C}$ , which is nothing but a stationary state for the linear RT equation which fulfills all the properties listed in the statement of Theorem 1.1.*

*Proof of Lemma 3.6.* We split the proof into two steps.

*Step 1.* We split the operator  $\mathcal{L}$  as

$$\mathcal{L} = \mathcal{A}_1 + \mathcal{B}_1, \quad \mathcal{A}_1 := A_\psi, \quad \psi := \phi_R K(x, v),$$

and with the same integer  $n$  as in Lemma 3.5 we write the iterated Duhamel formula

$$S_{\mathcal{L}} = \{S_{\mathcal{B}_1} + \dots + S_{\mathcal{B}_1} * (\mathcal{A}_1 S_{\mathcal{B}_1})^{(n*)}\} + S_{\mathcal{B}_1} * (\mathcal{A}_1 S_{\mathcal{B}_1})^{(n*)} * \mathcal{A}_1 S_{\mathcal{L}} =: V_1 + V_2.$$

For the first term and thanks to Lemma 3.4, for some constant  $K_1 \geq 1$ , we have

$$\|V_1(t)f_0\|_X \leq \sum_{\ell=0}^n \|S_{\mathcal{B}_1} * (\mathcal{A}_1 S_{\mathcal{B}_1})^{(\ell*)} f_0\|_X \leq K_1 \|f_0\|_X.$$

For the second term, for some constant  $K_2 \geq 1$ , we have

$$\|V_2(t)\|_{L^1(m) \rightarrow X} \leq \|S_{\mathcal{B}_1}\|_{X \rightarrow X} * \|(\mathcal{A}_1 S_{\mathcal{B}_1})^{(n*)}\|_{L^1(m) \rightarrow X} * \|\mathcal{A}_1 S_{\mathcal{L}}\|_{L^1(m) \rightarrow L^1(m)} \leq K_2,$$

using both (3.6) and (3.8). All together, we have proved

$$\|S_{\mathcal{L}}(t)f_0\|_X \lesssim \|f_0\|_X, \quad \forall t \geq 0.$$

As a consequence, the norm  $\| \cdot \|$  is equivalent to the standard one on  $X$ .

*Step 2.* We now define  $C := \|g_0\|$  for a given  $g_0 \in X$  which is also a probability measure. With this choice, we see that the set  $\mathcal{C}$  is not empty. Moreover, thanks to Lemma 2.1, the flow is continuous for the  $L^1$  norm and preserves positivity and total mass. By construction, we see that for any  $f_0 \in \mathcal{C}$  and  $t \geq 0$ , we have

$$\|S_{\mathcal{L}}(t)f_0\| = \sup_{s \geq t} \|S_{\mathcal{L}}(s)f_0\|_{L^p(m)} \leq \sup_{s \geq 0} \|S_{\mathcal{L}}(s)f_0\|_{L^p(m)} = \|f_0\| \leq C.$$

All together, the set  $\mathcal{C}$  is clearly invariant by the flow  $S_{\mathcal{L}}$ .

Because  $\mathcal{C}$  is also compact for the weak topology of  $L^1 \cap L^\infty$ , thanks to a standard variant of the Brouwer-Schauder-Tychonoff fixed point theorem (see for instance [9, Theorem 1.2]), we obtain the existence of an invariant element  $G$  for the linear RT flow which furthermore belongs to  $\mathcal{C}$ , in other words

$$\exists G \in \mathcal{C} \text{ such that } S_{\mathcal{L}}(t)G = G, \quad \forall t \geq 0.$$

As a consequence, we have  $G \in D(\mathcal{L}) \setminus \{0\}$  and  $\mathcal{L}G = 0$ , so that  $G$  is a stationary state for the linear RT equation.  $\square$

**3.4. Uniqueness of the stationary state.** In this section we prove a strong maximal principle on the operator  $-\mathcal{L}$ . The uniqueness of the normalized and positive steady state then follows using classical arguments. We skip the proof and refer for instance to [18, Step 4, proof of Theorem 5.3] for details.

**Lemma 3.7.** *The operator  $-\mathcal{L}$  satisfies the following version of the strong maximum principle: for any given  $0 \leq g \in L^2(m)$  and  $\mu \in \mathbb{R}$ , there holds*

$$g \in D(\mathcal{L}) \setminus \{0\} \text{ and } (-\mathcal{L} + \mu)g \geq 0 \text{ imply } g > 0$$

*Proof of Lemma 3.7.* We consider  $g$  as in the above statement and we prove that it is a positive function in several steps.

*Step 1.* Defining  $M := 1 + \chi + \mu \in \mathbb{R}$  and  $m := 1 - \chi > 0$ , we see that

$$\begin{aligned} v \cdot \nabla_x g + M g &\geq v \cdot \nabla_x g + K g \\ &\geq \int_{\mathcal{V}} K' g' dv' \geq m \varrho, \quad \varrho := \int_{\mathcal{V}} g' dv'. \end{aligned}$$

Integrating the above inequality along the free transport characteristics, we get

$$(3.9) \quad g(x, v) \geq m \int_0^t \varrho(x - v s) e^{-M s} ds + g(x - v t, v) e^{-M t}, \quad \forall t \geq 0.$$

*Step 2.* In particular, because the second term at the RHS is nonnegative, we may keep only the contribution of the first term and we get

$$\begin{aligned} \varrho(x) &\geq m \int_{1/2}^1 \int_{\mathcal{V}} \varrho(x - v s) e^{-M s} ds dv \\ &\geq \kappa \int_{\mathcal{V}/2} \varrho(x + w) dw, \quad \kappa > 0. \end{aligned}$$

Since  $g \geq 0$  and  $g \not\equiv 0$ , we also have  $\varrho \geq 0$  and  $\varrho \not\equiv 0$ , and there exists  $x_0 \in \mathbb{R}^d$  and  $r > 0$ , small enough, such that  $\langle \varrho \mathbf{1}_{B(x_0, r)} \rangle = \alpha > 0$  and  $B(0, 2r) \subset \mathcal{V}$ , or in other words  $B(x_0, r) \subset x + \mathcal{V}/2$  for any  $x \in B(x_0, r)$ . As a consequence, we have

$$(3.10) \quad \varrho \geq \alpha_0 \mathbf{1}_{B(x_0, r)}, \quad \alpha_0 := \kappa \alpha.$$

*Step 3.* Observing that for any  $x \in \mathbb{R}^d$ , there exists a small ball  $B \subset \mathcal{V}$  and some times  $\tau_1 > \tau_0 > 0$  such that  $x - sv \in B(x_0, r/2)$  for any  $v \in B$  and  $s \in (\tau_0, \tau_1)$ , we may argue as above and we get

$$\begin{aligned} \varrho(x) &\geq m \int_{\tau_0}^{\tau_1} \int_B \varrho(x - v s) e^{-M s} ds dv \\ &\geq \alpha_x := m |B| (\tau_1 - \tau_0) e^{-M \tau_1} \alpha_0 > 0. \end{aligned}$$

Finally, using (3.9) again, we deduce

$$g(x, v) \geq m \int_0^1 \varrho(x - v s) e^{-M s} ds > 0,$$

which concludes the proof.  $\square$

## 4. ASYMPTOTIC STABILITY OF THE STATIONARY STATE

**4.1. A new splitting.** In all this section, excepted in paragraph 4.5, we fix an exponential weight function  $m = e^{\gamma(x)}$ , with  $\gamma \in (0, \gamma^*)$  and  $\gamma^* > 0$  defined in Lemma 3.3, and we define the Banach space  $X := L^1(m) \cap L^2(m^{1/2})$ . We also introduce the splitting

$$(4.1) \quad \mathcal{L} = \mathcal{A} + \mathcal{B}, \quad \mathcal{A}f := \int_{\mathcal{V}} K'_{R, \delta_i} f' dv',$$

where

$$K_{R, \delta_i} = \phi_{\delta_2, R}(x) \psi_{\delta_1}(v) K_{\delta_3}(x, v), \quad K_{\delta_3}(x, v) = 1 + \chi \zeta_{\delta_3}(x \cdot v),$$

for some real numbers  $R > 1$ ,  $\delta_1, \delta_2, \delta_3 \in (0, 1)$  to be fixed, and where we have defined the truncation functions  $\phi_\lambda(z) := \phi(z/\lambda)$ ,  $\phi \in \mathcal{D}(\mathbb{R}^d)$  radially symmetric,  $\mathbf{1}_{B(0,1)} \leq \phi \leq \mathbf{1}_{B(0,2)}$ , and then  $\phi_{\delta, R}(x) := \phi_R(x) - \phi_\delta(x)$ ,  $\psi_\delta(v) := 1 - \phi_\delta(V_0 - |v|) - \phi_\delta(v)$ , as well as a regularized sign function  $\zeta_\delta \in C^\infty(\mathbb{R})$  which is odd, increasing and satisfies  $\zeta_\delta(s) = 1$  for any  $s \geq \delta$ .

We establish that  $S_{\mathcal{B}}$  and  $\mathcal{A}S_{\mathcal{B}}$  enjoy suitable decay estimate (Section 4.2) and regularity estimate (Section 4.3) from which we deduce the asymptotic stability in  $X$  thanks to a semigroup version of the Krein-Rutman theorem (Section 4.4) and next the asymptotic stability in any exponential and polynomial weighted  $L^1$  space by using an extension argument (Section 4.5).

**4.2. Decay estimates for the semigroup  $S_{\mathcal{B}}$ .**

**Proposition 4.1.** *For the same constant  $a^* < 0$  as defined in Lemma 3.3, there holds*

$$(4.2) \quad \|S_{\mathcal{B}}(t)\|_{X \rightarrow X} \lesssim e^{at}, \quad \forall t \geq 0, \quad \forall a > a^*.$$

*Proof of Proposition 4.1.* We split the proof into five steps.

*Step 1. Norms and splitting.* Inspired by [12, Proposition 5.15] and the moment trick introduced in [14], we define the three norms  $\|\cdot\|_X$ ,  $\|\|\cdot\|\|$  and  $N(\cdot)$  in the following way

$$\begin{aligned} \|f\|_X^2 &:= \|f\|_{L^1(m)}^2 + \|f\|_{L^2(m^{1/2})}^2, \\ \|\|f\|\|^2 &:= \eta_2 \|f\|_X^2 + \int_0^\infty \|S_{\mathcal{B}_1}(\tau)f\|_X^2 d\tau, \\ N(f)^2 &:= \eta_1 \|f\|_{L^2(\tilde{m}_0^{1/2})}^2 + \|\|f\|\|^2, \end{aligned}$$

for some constants  $\eta_1, \eta_2 \in (0, 1)$  to be fixed and where  $\tilde{m}_0$  is the weight function

$$\tilde{m}_0 := \left(1 - \frac{x}{|x|^{1/2}} \cdot \frac{v}{|v|}\right) \phi_{1/2}(x),$$

so that  $0 \leq \tilde{m}_0 \leq 2\phi_1$ . Thanks to the decay estimate of Lemma 3.4, one easily sees that these three norms are equivalent.

We fix  $f_0 \in X$  and we define  $f(t) = f_{\mathcal{B}}(t) = S_{\mathcal{B}}(t)f_0$  the associated trajectory along the action of the semigroup  $S_{\mathcal{B}}$ . In order to prove (4.2), we compute

$$(4.3) \quad T := \frac{1}{2} \frac{d}{dt} N(f_{\mathcal{B}}(t))^2 = \eta_1 T_1 + \eta_2 T_2 + T_3,$$

where  $T_i$  are the contributions of the terms involved in the definition of the norm  $N$  that we compute separately. For latter references, we introduce the splitting of  $\mathcal{B}$  as

$$\mathcal{B} := \mathcal{B}_1 + \mathcal{A}_1^c + \mathcal{A}_2^c + \mathcal{A}_3^c,$$

where  $\mathcal{B}_1$  is defined in section 3.2 with  $R \geq 1$  large enough so that Lemma 3.3 and Lemma 3.4 hold true, and we have set

$$\begin{aligned}\mathcal{A}_1^c f &= \phi_R(x) \int_{\mathcal{V}} K' f' \psi_{\delta_1}^c(v') dv' \\ \mathcal{A}_2^c f &= \phi_{\delta_2}(x) \int_{\mathcal{V}} K' f' \psi_{\delta_1}(v') dv' \\ \mathcal{A}_3^c f &= \phi_{\delta_2, R}(x) \int_{\mathcal{V}} K_{\delta_3}^c(x \cdot v') f' \psi_{\delta_1}(v') dv',\end{aligned}$$

with  $\psi_{\delta_1}^c := 1 - \psi_{\delta_1}$ ,  $\phi_R^c := 1 - \phi_R$ ,  $K_{\delta}^c = \chi \zeta_{\delta}^c$ ,  $\zeta_{\delta}^c = \zeta - \zeta_{\delta}$ . We shall also denote

$$\mathcal{A}_{123}^c := \mathcal{A}_1^c + \mathcal{A}_1^c + \mathcal{A}_2^c \quad \text{and} \quad \mathcal{A}_{0123}^c := \mathcal{A}_0^c + \mathcal{A}_1^c + \mathcal{A}_1^c + \mathcal{A}_2^c,$$

where we recall that  $\mathcal{A}_0^c$  has been defined during the proof of Lemma 3.4.

*Step 2. Contribution of the term  $T_1$ .* We prove that

$$(4.4) \quad T_1 := \frac{1}{2} \frac{d}{dt} \|f_{\mathcal{B}}(t)\|_{L^2(\tilde{m}_0^{1/2})}^2 \leq -\frac{1}{4} \|f_{\mathcal{B}}(t)\|_{L^2(\tilde{m}_1^{1/2})}^2 + C_1 \|f_{\mathcal{B}}(t)\|_X^2,$$

for some positive constant  $C_1$  and where  $\tilde{m}_1$  is the weight function defined by

$$\tilde{m}_1(x, v) := \frac{|v|}{|x|^\alpha} \phi_{1/2}(x).$$

In order to prove (4.4), we first observe that

$$T_1 = (\mathcal{B} f_{\mathcal{B}}(t), f_{\mathcal{B}}(t))_{L^2(\tilde{m}_0^{1/2})},$$

and next we compute the RHS by splitting it in several pieces. For any  $f \in X$ , we have

$$\begin{aligned}(\mathcal{B}f, f)_{L^2(\tilde{m}_0^{1/2})} &= (-v \cdot \nabla_x f, f)_{L^2(\tilde{m}_0^{1/2})} + (\mathcal{A}_{0123}^c f - Kf, f)_{L^2(\tilde{m}_0^{1/2})} \\ &=: T_{1,1} + T_{1,2}.\end{aligned}$$

We compute the first key term. Performing one integration by parts, we find

$$\begin{aligned}T_{1,1} &= \int (-v \cdot \nabla_x f) f \left(1 - \frac{x}{|x|^{1/2}} \cdot \frac{v}{|v|}\right) \phi_{1/2}(x) dx dv \\ &= \frac{1}{2} \int f^2 \left\{ \left[ \frac{1}{2} \left(\frac{x}{|x|} \cdot \frac{v}{|v|}\right)^2 - 1 \right] \frac{|v|}{|x|^{1/2}} \phi_{1/2} + \left(1 - \frac{x}{|x|^{1/2}} \cdot \frac{v}{|v|}\right) (v \cdot \nabla_x \phi_{1/2}) \right\} dx dv \\ &= -\frac{1}{4} \int f^2 \frac{|v|}{|x|^{1/2}} \phi_{1/2} dx dv + \tilde{T}_{1,1}\end{aligned}$$

with  $\tilde{T}_{1,1} \lesssim \|f\|_{L^2}^2$ . For the remainder term, we also easily have  $|T_{1,2}| \lesssim \|f\|_{L^2}^2$  from what (4.4) immediately follows.

*Step 3. Contribution of the term  $T_2$ .* We prove that

$$(4.5) \quad T_2 := \frac{1}{2} \frac{d}{dt} \|f_{\mathcal{B}}\|_X^2 \leq C_2 \|f_{\mathcal{B}}\|_X^2,$$

for some positive constant  $C_2$ . We have

$$\begin{aligned}T_2 &= \frac{1}{2} \frac{d}{dt} \|f_{\mathcal{B}}\|_{L^2(m)}^2 + \frac{1}{2} \frac{d}{dt} \|f_{\mathcal{B}}\|_{L^1(m)}^2 \\ &= (\mathcal{B} f_{\mathcal{B}}, f_{\mathcal{B}})_{L^2(m)} + \langle \mathcal{B} f_{\mathcal{B}}, \text{sign} f_{\mathcal{B}} \rangle_{L^1(m), L^\infty} \|f_{\mathcal{B}}\|_{L^1(m)} \\ &=: T_{2,1} + T_{2,2},\end{aligned}$$

with

$$T_{2,1} := (\mathcal{B}_0 f_{\mathcal{B}}, f_{\mathcal{B}})_{L^2(m)} + \langle \mathcal{B}_0 f_{\mathcal{B}}, \text{sign} f_{\mathcal{B}} \rangle_{L^1(m), L^\infty} \|f_{\mathcal{B}}\|_{L^1(m)} \leq 0,$$

because  $\mathcal{B}_0$  is dissipative in both  $L^1(m)$  and  $L^2(m^{1/2})$  from Lemma 3.1, and with

$$\begin{aligned} T_{2,2} &:= (\mathcal{A}_{0123}^c f_{\mathcal{B}}, f_{\mathcal{B}})_{L^2(m)} + \langle \mathcal{A}_{123}^c f_{\mathcal{B}}, \text{sign} f_{\mathcal{B}} \rangle_{L^1(m), L^\infty} \|f_{\mathcal{B}}\|_{L^1(m)}^2 \\ &\leq \|\mathcal{A}_{0123}^c f_{\mathcal{B}}\|_{L^2(m)} \|f_{\mathcal{B}}\|_{L^2(m)} + \|\mathcal{A}_{123}^c f_{\mathcal{B}}\|_{L^1(m)} \|f_{\mathcal{B}}\|_{L^1(m)}^2 \\ &\lesssim \|f_{\mathcal{B}}\|_X^2, \end{aligned}$$

because  $|\mathcal{A}_i f| \leq \mathcal{A}|f|$  for any  $i \in \{1, 2, 3\}$  and  $\mathcal{A} \in \mathcal{B}(L^p(m), L^p(m))$  for any  $p \in \{1, 2\}$ .

*Step 4. Contribution of the term  $T_3$ .* We prove that for any  $\eta_1 \in (0, 1)$ , we can find  $\delta_1, \delta_2, \delta_3 \in (0, 1)$  and  $R \geq 1$  such that the associated operator  $\mathcal{B}$  satisfies

$$(4.6) \quad T_3 := \frac{1}{2} \frac{d}{dt} \int_0^\infty \|S_{\mathcal{B}_1}(\tau) f_{\mathcal{B}}(t)\|_X^2 d\tau \leq -\frac{3}{8} \|f_{\mathcal{B}}(t)\|_X^2 + \frac{\eta_1}{4} \|f_{\mathcal{B}}(t)\|_{L^2(\tilde{m}_1^{1/2})}^2.$$

We split the term  $T_3$  as

$$\begin{aligned} T_3 &= \frac{1}{2} \frac{d}{dt} \int_0^\infty \|S_{\mathcal{B}_1}(\tau) f_{\mathcal{B}}(t)\|_{L^1(m)}^2 d\tau + \frac{1}{2} \frac{d}{dt} \int_0^\infty \|S_{\mathcal{B}_1}(\tau) f_{\mathcal{B}}(t)\|_{L^2(m^{1/2})}^2 d\tau \\ &=: T_{3,1} + T_{3,2}. \end{aligned}$$

For the first term, we compute

$$\begin{aligned} T_{3,1} &= \int_0^\infty \frac{1}{2} \frac{d}{dt} \|S_{\mathcal{B}_1}(\tau) f_{\mathcal{B}}(t)\|_{L^1(m)}^2 d\tau \\ &= \int_0^\infty \langle S_{\mathcal{B}_1}(\tau) \mathcal{B} f_{\mathcal{B}}(t), \text{sign}(S_{\mathcal{B}_1}(\tau) f_{\mathcal{B}}(t)) \rangle_{L^1(m), L^\infty} \|S_{\mathcal{B}_1}(\tau) f_{\mathcal{B}}(t)\|_{L^1(m)} d\tau \\ &= \int_0^\infty \langle \mathcal{B}_1 S_{\mathcal{B}_1}(\tau) f_{\mathcal{B}}(t), \text{sign}(S_{\mathcal{B}_1}(\tau) f_{\mathcal{B}}(t)) \rangle_{L^1(m), L^\infty} \|S_{\mathcal{B}_1}(\tau) f_{\mathcal{B}}(t)\|_{L^1(m)} d\tau \\ &\quad + \int_0^\infty \langle S_{\mathcal{B}_1}(\tau) \mathcal{A}_{123}^c f_{\mathcal{B}}(t), \text{sign}(S_{\mathcal{B}_1}(\tau) f_{\mathcal{B}}(t)) \rangle_{L^1(m), L^\infty} \|S_{\mathcal{B}_1}(\tau) f_{\mathcal{B}}(t)\|_{L^1(m)} d\tau \\ &=: T_{3,1,1} + T_{3,1,2}. \end{aligned}$$

On the one hand, we observe that

$$T_{3,1,1} = \int_0^\infty \frac{1}{2} \frac{d}{d\tau} \|S_{\mathcal{B}_1}(\tau) f_{\mathcal{B}}(t)\|_{L^1(m)}^2 d\tau = -\frac{1}{2} \|f_{\mathcal{B}}(t)\|_{L^1(m)}^2,$$

where in the last line we have used that  $S_{\mathcal{B}_1} f_0$  has the nice decay estimate (3.6) in the space  $L^1(m)$  because  $f_0 \in X$ . On the other hand, using again the decay estimate (3.6), for any  $\varepsilon > 0$ , we have

$$\begin{aligned} T_{3,1,2} &= \int_0^\infty \|S_{\mathcal{B}_1}(\tau) \mathcal{A}_{123}^c f_{\mathcal{B}}(t)\|_{L^1(m)} \|S_{\mathcal{B}_1}(\tau) f_{\mathcal{B}}(t)\|_{L^1(m)} d\tau \\ &\leq \frac{1}{2\varepsilon} \int_0^\infty \|S_{\mathcal{B}_1}(\tau) \mathcal{A}_{123}^c f_{\mathcal{B}}(t)\|_{L^1(m)}^2 d\tau + \frac{\varepsilon}{2} \int_0^\infty \|S_{\mathcal{B}_1}(\tau) f_{\mathcal{B}}(t)\|_{L^1(m)}^2 d\tau \\ &\lesssim \frac{1}{\varepsilon} \|\mathcal{A}_{123}^c f_{\mathcal{B}}(t)\|_X^2 + \varepsilon \|f_{\mathcal{B}}(t)\|_X^2. \end{aligned}$$

We may treat the second term in a similar way, using in particular the fact that  $S_{\mathcal{B}_1} f_0$  has the nice decay estimate (3.6) in the space  $L^2(m^{1/2})$  because  $f_0 \in X$ . Gathering the two

resulting estimates and taking  $\varepsilon > 0$  small enough, we get

$$(4.7) \quad T_3 \leq -\frac{7}{16} \|f_{\mathcal{B}}(t)\|_X^2 + \widetilde{T}_3, \quad \widetilde{T}_3 \lesssim \|\mathcal{A}_{123}^c f_{\mathcal{B}}(t)\|_X^2.$$

In order to conclude, we compute the contributions  $\|\mathcal{A}_i^c f_{\mathcal{B}}(t)\|_X^2$  for any  $i \in \{1, 2, 3\}$ .

On the one hand, using the Cauchy-Schwarz inequality, for any  $p \in \{1, 2\}$ ,  $R > 2$  and  $\delta_1 \in (0, 1)$ , we have

$$\begin{aligned} \|\mathcal{A}_1^c f\|_{L^p}^p(m^{1/p}) &= \int \left| \phi_R(x) \int_{\mathcal{V}} K' f' \psi_{\delta_1}^c(v') dv' \right|^p m dx dv \\ &\leq 2^p m(2R) \int \left[ \int_{\mathcal{V}} |f'|^2 dv' \right]^{p/2} \left[ \int_{\mathcal{V}} (\mathbf{1}_{|v'| \leq 2\delta_1} + \mathbf{1}_{V_0 - |v'| \leq 2\delta_1}) dv' \right]^{p/2} dx dv \\ &\lesssim m(2R) \delta_1^{p/2} \|f\|_{L^2}^{p/2}. \end{aligned}$$

Similarly, when furthermore  $\delta_2 \in (0, 1/4)$ , we have

$$\begin{aligned} \|\mathcal{A}_2^c f\|_{L^p}^p(m^{1/p}) &= \int \left| \phi_{\delta_2}(x) \int_{\mathcal{V}} K' f' \psi_{\delta_1}(v') dv' \right|^p m dx dv \\ &\leq 2^p m(2) \left[ \int_{B(0,1) \times \mathcal{V}} \mathbf{1}_{|x| \leq 2\delta_2} |f'|^2 \mathbf{1}_{|v'| \geq \delta_1} dv' dx \right]^{p/2} \\ &\lesssim \left[ \frac{\delta_2^{1/2}}{\delta_1} \right]^{p/2} \|f\|_{L^2(m_1^{1/2})}^p. \end{aligned}$$

Finally and similarly again, when furthermore  $\delta_3 \in (0, 1/2)$ , observing that

$$0 \leq K_{\delta_3}^c(x, v') = \chi (\zeta - \zeta_{\delta_3})(x \cdot v') \leq \chi \mathbf{1}_{|x \cdot v'| \leq \delta_3},$$

we have

$$\begin{aligned} \|\mathcal{A}_3^c f\|_{L^p}^p(m^{1/p}) &= \int \left| \phi_{\delta_2, R}(x) \int K_{\delta_3}^c(x \cdot v') f' \psi_{\delta_1}(v') dv' \right|^p m dx dv \\ &\leq m(2R) \chi^p \int \left[ \int_{\mathcal{V}} |f'|^2 dv' \right]^{p/2} \left[ \int_{\mathcal{V}} \mathbf{1}_{|x \cdot v'| \leq 2\delta_3} dv' \right]^{p/2} \mathbf{1}_{|x| \geq \delta_2} dx dv \\ &\lesssim m(2R) \left[ \int f^2 dv dx \right]^{p/2} \left[ \text{meas}\{v \in \mathcal{V}; |v_1| \leq \delta_3/\delta_2\} \right]^{p/2} \\ &\lesssim m(2R) \frac{\delta_3^{p/2}}{\delta_2^{p/2}} \|f\|_{L^2}^p. \end{aligned}$$

All these estimates together, we get

$$(4.8) \quad \|\mathcal{A}_{123}^c f_{\mathcal{B}}(t)\|_X^2 \lesssim m(2R) \delta_1 \|f\|_X^2 + \frac{\delta_2^{1/2}}{\delta_1} \|f\|_{L^2(m_1^{1/2})}^2 + m(2R) \frac{\delta_3}{\delta_2} \|f\|_X^2.$$

We thus obtain (4.6) by just gathering (4.7) and (4.8) and by choosing  $\delta_1, \delta_2, \delta_3 > 0$  adequately.

*Step 5. Conclusion.* From estimates (4.4), (4.5) and (4.6), we have

$$\begin{aligned} T &\leq \eta_1 C_1 \|f_{\mathcal{B}}\|_X^2 + \eta_2 C_2 \|f_{\mathcal{B}}\|_X^2 - \frac{3}{8} \|f_{\mathcal{B}}\|_X^2 \\ &\leq -\frac{1}{4} \|f_{\mathcal{B}}\|_X^2 \lesssim -N(f_{\mathcal{B}})^2, \end{aligned}$$

by choosing  $\eta_1, \eta_2 > 0$  small enough. We have proved that  $\mathcal{B}$  is  $-1/4$ -dissipative for the norm  $N$  and thus (4.2) follows.  $\square$



**4.3. Some regularity associated to  $\mathcal{AS}_{\mathcal{B}}$ .** In this section we show that the family of operators  $\mathcal{AS}_{\mathcal{B}}$  satisfies a regularity and growth estimate that we express in terms of the abstract Sobolev space  $X_{\mathcal{B}}^{1/2}$  defined as the usual 1/2 interpolated space between  $X$  and the domain

$$X_{\mathcal{B}}^1 = D(\mathcal{B}) := \{f \in X; \mathcal{B}f \in X\}$$

endowed with the graph norm.

**Proposition 4.2.** *For the same constant  $a^* < 0$  as defined in Lemma 3.3, the family of operators  $\mathcal{AS}_{\mathcal{B}}$  satisfies*

$$(4.9) \quad \int_0^\infty \|\mathcal{AS}_{\mathcal{B}}(t)\|_{X \rightarrow Y}^2 e^{-2a^*t} dt < \infty,$$

with

$$Y := \{f \in L^2(\mathbb{R}^d \times \mathcal{V}); \text{supp } f \subset B(0, R) \times \mathcal{V}, f \in H^{1/2}\}.$$

The proof is mainly a consequence of Bouchut-Desvillettes' version [3, Theorem 2.1] (see also [6] for a related discrete version) of the classical averaging Lemma initiated in the famous articles of Golse et al. [11, 10]. We give in step 1 below a simpler, more accurate and more adapted version of [3, Theorem 2.1] for which we sketch the proof for the sake of completeness. During the proof, we will use the following classical trace result.

**Lemma 4.3.** *There exists a constant  $C_d \in (0, \infty)$  such that for any  $\phi \in H^{d/2}(\mathbb{R}^d)$  and any  $u \in \mathbb{R}^d$ ,  $|u| = 1$ , the function  $\phi_u(s) := \phi(su)$  satisfies*

$$\|\phi_u\|_{L^2(\mathbb{R})} \leq C_d \|\phi\|_{H^{d/2}(\mathbb{R}^d)} = C_d \left( \int_{\mathbb{R}^d} |\check{F}\phi|^2(w) \langle w \rangle^d dw \right)^{1/2},$$

where  $\check{F}$  stands for the (inverse) Fourier transform operator.

*Proof of Proposition 4.2.* We split the proof into two steps.

*Step 1.* We consider the damped free transport equation

$$(4.10) \quad \partial_t f = \mathcal{T}f := -v \cdot \nabla_x f - f, \quad f|_{t=0} = f_0,$$

and we denote by  $S_{\mathcal{T}}(t)$  the associated semigroup defined through the characteristics formula

$$(4.11) \quad [S_{\mathcal{T}}(t)f_0](x, v) := f(t, x, v) = f_0(x - vt, v) e^{-t}.$$

We claim that for any  $\varphi \in L^2(\mathcal{V})$ , there holds

$$(4.12) \quad \int_0^\infty \|A_\varphi S_{\mathcal{T}}(t)\|_{L_{x,v}^2 \rightarrow H_x^{1/2}}^2 e^{2t} dt \lesssim \|\varphi\|_{L^2(\mathcal{V})}.$$

For a given function  $h$  which depends on the  $x$  variable or on the  $(x, v)$  variable, we denote by  $\hat{h}$  its Fourier transform on the  $x$  variable and by  $\mathcal{F}h$  its Fourier transform on both variables  $x$  and  $v$ . We fix  $f_0 \in L^2(\mathbb{R}^d \times \mathcal{V})$  and  $\varphi \in L^\infty(\mathbb{R}^d)$ , we denote by  $f$  the solution to the free transport equation (4.10) and by  $\rho$  the average function

$$\rho(t, x) := \int_{\mathbb{R}^d} f(t, x, v) \varphi(v) dv = [A_\varphi S_{\mathcal{T}}(t)f_0](x).$$

In Fourier variables, the free transport equation (4.10) writes

$$\partial_t \hat{f} + iv \cdot \xi \hat{f} - \hat{f} = 0, \quad \hat{f}|_{t=0} = \hat{f}_0,$$

so that

$$\hat{f}(t, \xi, v) = e^{iv \cdot \xi t - t} \hat{f}_0(\xi, v)$$

and

$$\hat{\rho}(t, \xi) = \int_{\mathbb{R}^d} e^{iv \cdot \xi t - t} \hat{f}_0(\xi, v) \varphi(v) dv = \mathcal{F}(f_0 \varphi)(\xi, t\xi) e^{-t}.$$

We deduce

$$\int_0^\infty |\hat{\rho}(t, \xi)|^2 e^{2t} dt \leq \int_{\mathbb{R}} |\mathcal{F}(f_0 \varphi)(\xi, t\xi)|^2 dt.$$

Performing one change of variable, introducing the notation  $\sigma_\xi = \xi/|\xi|$  and using Lemma 4.3, we deduce

$$\begin{aligned} \int_{\mathbb{R}} |\mathcal{F}(f_0 \varphi)(\xi, t\xi)|^2 dt &= \frac{1}{|\xi|} \int_{\mathbb{R}} |\mathcal{F}(f_0 \varphi)(\xi, s \sigma_\xi)|^2 ds \\ &\lesssim \frac{1}{|\xi|} \int_{\mathbb{R}^d} |(\hat{f}_0 \varphi)(\xi, w)|^2 \langle w \rangle^d dw. \end{aligned}$$

Thanks to Plancherel identity, we then obtain

$$\int_0^\infty \int_{\mathbb{R}^d} |\xi| |\hat{\rho}(t, \xi)|^2 d\xi e^{2t} dt \lesssim \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} |(f_0 \varphi)(x, w)|^2 \langle w \rangle^d dw dx = \|\varphi\|_{L^2_{d/2}}^2 \|f_0\|_{L^2_{xv}}^2,$$

which ends the proof (4.12).

*Step 2.* We show a similar estimate on  $\mathcal{AS}_T(t)$ . Using that  $K_{R, \delta_i} \in C_c^\infty(\mathbb{R}^d \times \mathbb{R}^d)$ ,  $\text{supp } K_{R, \delta_i} \subset B(0, 2R) \cap B(0, V'_0)$ ,  $V'_0 \in (0, V_0)$ , we may expand it as a Fourier series

$$K_{R, \delta_i}(x, v) = \sum_{k, \ell \in \mathbb{Z}^d} a_{k, \ell} e^{ix \cdot k} e^{iv \cdot \ell} \vartheta(v), \quad \forall (x, v) \in \mathcal{Q},$$

$\mathcal{Q} := \{x \in \mathbb{R}^d, v \in \mathbb{R}^d; |x_i| \leq 2R, |v_i| \leq V_0, \forall i = 1, \dots, d\}$ , for a truncation function  $\vartheta \in C^\infty(\mathbb{R}^d)$ ,  $\text{supp } \vartheta \subset B(0, V_0)$ ,  $\vartheta \equiv 1$  on  $B(0, V'_0)$  and with fast decaying Fourier coefficients

$$|a_{k, \ell}| \lesssim \langle k \rangle^{-2d-4} \langle \ell \rangle^{-2d-2}.$$

From the definition of  $\mathcal{A}$  and denoting  $f(t) = S_T(t) f_0$  for some  $f_0 \in L^2(\mathbb{R}^d \times \mathcal{V})$ , we may then write

$$(\mathcal{AS}_T(t) f_0)(x) = \sum_{k, \ell \in \mathbb{Z}^d} a_{k, \ell} e^{ix \cdot k} \rho_\ell(t, x), \quad \rho_\ell(t, x) := \int_{\mathcal{V}} f(t, x, v) e^{iv \cdot \ell} \vartheta(v) dv.$$

On the one hand, from Step 1, we have

$$(4.13) \quad \sup_{\ell \in \mathbb{Z}^d} \int_0^\infty \|\rho_\ell(t, \cdot)\|_{H^{1/2}}^2 e^{2t} dt \lesssim \|f_0\|_{L^2}^2.$$

On the other hand, we denote  $e_k(x) := e^{ix \cdot k}$  and we define the mapping

$$U(\rho_\ell) := \sum_{k, \ell \in \mathbb{Z}^d} a_{k, \ell} e_k \rho_\ell.$$

From Cauchy-Schwarz inequality and Fubini Theorem, we have

$$\begin{aligned}
& \int_0^\infty \|U(\rho_\ell)(t, \cdot)\|_{L^2(B_{2R})}^2 e^{2t} dt \leq \\
& \leq \int_0^\infty \int_{B_{2R}} \left( \sum_{k,\ell} |a_{k,\ell}|^2 \langle k \rangle^{d+1} \langle \ell \rangle^{d+1} \right) \left( \sum_{k,\ell} |\rho_\ell|^2 \langle k \rangle^{-d-1} \langle \ell \rangle^{-d-1} \right) e^{2t} dx dt \\
& \lesssim \sum_{k,\ell} \langle k \rangle^{-d-1} \langle \ell \rangle^{-d-1} \int_0^\infty \int_{B_{2R}} |\rho_\ell|^2 e^{2t} dt dx \\
& \lesssim \sup_{\ell \in \mathbb{Z}^d} \int_0^\infty \|\rho_\ell(t, \cdot)\|_{L^2(B_R)}^2 e^{2t} dt.
\end{aligned}$$

Using furthermore that

$$\nabla_x U(\rho_\ell) = \sum_{k,\ell \in \mathbb{Z}^d} a_{k,\ell} (ik) e_k \rho_\ell + \sum_{k,\ell \in \mathbb{Z}^d} a_{k,\ell} e_k \nabla_x \rho_\ell,$$

we find similarly

$$\int_0^\infty \|\nabla_x U(\rho_\ell)(t, \cdot)\|_{L^2(B_{2R})}^2 e^{2t} dt \lesssim \sup_{\ell \in \mathbb{Z}^d} \int_0^\infty \|\rho_\ell(t, \cdot)\|_{H^1(B_R)}^2 e^{2t} dt.$$

Observing that

$$\{g \in L^2(\mathbb{R}^d \times \mathcal{V}); \text{supp } g \subset B(0, 2R) \times \mathcal{V}, \nabla_x g \in L^2\} \subset X_{\mathcal{B}}^1,$$

both estimates together and an interpolation argument yield

$$(4.14) \quad \int_0^\infty \|U(\rho_\ell)(t, \cdot)\|_{X_{\mathcal{B}}^{1/2}}^2 e^{2t} dt \lesssim \sup_{\ell \in \mathbb{Z}^d} \int_0^\infty \|\rho_\ell(t, \cdot)\|_{H^{1/2}(B_R)}^2 e^{2t} dt.$$

Gathering estimates (4.13) and (4.14), we have established

$$(4.15) \quad \int_0^\infty \|\mathcal{A}S_{\mathcal{T}}(t)f_0\|_{X_{\mathcal{B}}^{1/2}}^2 e^{2t} dt \lesssim \|f_0\|_{L^2}^2.$$

*Step 3. Conclusion.* We split  $\mathcal{B}$  as  $\mathcal{B} = \mathcal{T} + \mathcal{C}$ . The Duhamel formula writes

$$S_{\mathcal{B}} = S_{\mathcal{T}} + S_{\mathcal{T}} * \mathcal{C}S_{\mathcal{B}},$$

from which we deduce

$$\mathcal{A}S_{\mathcal{B}} = \mathcal{A}S_{\mathcal{T}} + \mathcal{A}S_{\mathcal{T}} * \mathcal{C}S_{\mathcal{B}}.$$

Using that identity and estimates (4.2) and (4.15), we obtain (4.9).  $\square$

**4.4. A first asymptotic stability estimate in  $X$ .** In order to apply the semigroup version [18, Theorem 5.3] of the Krein-Rutman theorem, we list below some properties satisfied by the operators  $\mathcal{L}$ ,  $\mathcal{A}$  and  $\mathcal{B}$ .

*Fact 1.* There exists  $a^* < 0$  such that for any  $a > a^*$  and  $\ell \in \mathbb{N}$ , the following growth estimate holds

$$t \mapsto \|S_{\mathcal{B}} * (\mathcal{A}S_{\mathcal{B}})^{(*\ell)}(t)\|_{\mathcal{B}(X)} e^{-at} \in L^\infty(\mathbb{R}_+).$$

That is an immediate consequence of Proposition 4.1 and  $\mathcal{A} \in \mathcal{B}(X)$ .

*Fact 2.* For the same value  $a^* < 0$ , there exists  $Y \subset X_{\mathcal{L}}^s$ ,  $s \in (0, 1/2)$ , with compact embedding such that for any  $a > a^*$  the following growth estimate holds

$$t \mapsto \|\mathcal{A}S_{\mathcal{B}}(t)\|_{\mathcal{B}(X,Y)} e^{-at} \in L^1(\mathbb{R}_+).$$

That is an immediate consequence of Proposition 4.2 together with the fact that

$$\{f \in L^2(\mathbb{R}^d \times \mathcal{V}); \text{supp } f \subset B(0, R) \times \mathcal{V}, f \in H^1\} \subset X_{\mathcal{L}}^1$$

and an interpolation argument.

*Fact 3.* The semigroup  $S_{\mathcal{L}}$  is positive, the operator  $-\mathcal{L}$  satisfies the strong maximum principle as stated in Lemma 3.7 and  $\mathcal{L}$  satisfies the following Kato's inequality

$$(4.16) \quad (\text{sign } f)\mathcal{L}f := \frac{1}{2|f|}(f\mathcal{L}\bar{f} + \bar{f}\mathcal{L}f) \leq \mathcal{L}|f|,$$

for any complex valued function  $f \in X + iX$ . For proving the last inequality, we just compute

$$\begin{aligned} \frac{1}{2|f|}(f\mathcal{L}\bar{f} + \bar{f}\mathcal{L}f) &= -v \cdot \nabla|f| - K|f| + \frac{1}{2|f|}\left(f \int_{\mathcal{V}} K' \bar{f}' + \bar{f} \int_{\mathcal{V}} K' f'\right) \\ &\leq -v \cdot \nabla|f| - K|f| + \int_{\mathcal{V}} K'|f'| = \mathcal{L}|f|. \end{aligned}$$

*Fact 4.* The mass conservation property writes  $\mathcal{L}^*1 = 0$ , so that  $0 > a^*$  and 0 is an eigenvalue for the dual problem associated to a positive dual eigenfunction.

Gathering these above facts, we may then apply [18, Theorem 5.3], or more exactly we may repeat the proof of [18, Theorem 5.3] with minor and straightforward adaptations (we refer to [15] where these slight modifications are performed), in order to obtain that 0 is a (algebraically) simple eigenvalue, that there exists a spectral gap between this largest eigenvalue 0 and the remainder part of the spectrum and that a quantitative (partial but principal) spectral mapping theorem holds true. More precisely, we have the following asymptotic estimate: there exists  $\alpha \in (a^*, 0)$  such that

$$(4.17) \quad \|S_{\mathcal{L}}(t)\Pi^{\perp}f_0\|_X \lesssim e^{at} \|f_0\|_X, \quad \forall f_0 \in X, \forall a > \alpha, \forall t \geq 0,$$

where we have set  $\Pi^{\perp} := I - \Pi$  and  $\Pi f_0 := \langle\langle f_0 \rangle\rangle G$ .

**4.5. Asymptotic stability estimate in weighted  $L^1$  spaces.** We first consider the exponential weight  $m(x) := \exp(\gamma\langle x \rangle)$  with  $\gamma \in (0, \gamma^*)$  and  $\gamma^* > 0$  identified in Lemma 2.2. Iterating the Duhamel formula, we may write

$$S_{\mathcal{L}}\Pi^{\perp} = \Pi^{\perp}\{\mathcal{S}_{\mathcal{B}_1} + \dots + \mathcal{S}_{\mathcal{B}_1} * (\mathcal{A}_1\mathcal{S}_{\mathcal{B}_1})^{N-1}\} + (S_{\mathcal{L}}\Pi^{\perp}) * (\mathcal{A}_1\mathcal{S}_{\mathcal{B}_1})^N,$$

with  $N = d + 2$ . From Lemma 3.3 and (3.8) we have  $\mathcal{S}_{\mathcal{B}_1} * (\mathcal{A}_1\mathcal{S}_{\mathcal{B}_1})^{\ell} : L^1(m) \rightarrow L^1(m)$  with rate  $e^{at}$  for any  $\ell \in \{0, \dots, N-1\}$  and  $(\mathcal{A}_1\mathcal{S}_{\mathcal{B}_1})^N : L^1(m) \rightarrow X$  with rate  $e^{at}$ . Using that  $S_{\mathcal{L}}\Pi^{\perp} : X \rightarrow X \subset L^1(m)$  with rate  $e^{at}$  from (4.17) and gathering all the preceding decay estimates, we conclude that (1.9) holds in  $L^1(m)$ .

We next consider the polynomial weight  $m(x) := \langle x \rangle^k$  with  $k \in (0, \infty)$ . We begin with a decay estimate on the semigroup  $S_{\mathcal{B}_1}$ .

**Lemma 4.4.** *For any  $k > \ell > 0$ , the semigroup  $S_{\mathcal{B}_1}$  satisfies the following growth estimate*

$$\|S_{\mathcal{B}_1}\|_{L_k^1 \rightarrow L_{\ell}^1} \lesssim \langle t \rangle^{-(k-\ell)}, \quad \forall t \geq 0.$$

*Proof of Lemma 4.4.* Recalling that the dual operator  $\mathcal{L}^*$  has been defined in the proof of Lemma 2.2, for any  $q > 0$ , we compute

$$\mathcal{L}^*\langle \gamma x \rangle^q = q\gamma(v \cdot x)\langle \gamma x \rangle^{q-2},$$

$$\begin{aligned}\mathcal{L}^*(v \cdot x)\langle \gamma x \rangle^{q-2} &= v \cdot \nabla_x [(v \cdot x)\langle \gamma x \rangle^{q-2}] - q(v \cdot x)\langle \gamma x \rangle^{q-2} \\ &= \left( |v|^2 - (v \cdot x) - \chi |v \cdot x| \right) \langle \gamma x \rangle^{q-2} + (q-2)\gamma(v \cdot x)^2 \langle \gamma x \rangle^{q-4}.\end{aligned}$$

We then compute

$$\begin{aligned}\mathcal{L}^*|v \cdot x|\langle \gamma x \rangle^{q-2} &= v \cdot \nabla_x [|v \cdot x|\langle \gamma x \rangle^{q-2}] + (1 + \chi\zeta) \left( V_1 |x|\langle \gamma x \rangle^{q-2} - |v \cdot x|\langle \gamma x \rangle^{q-2} \right) \\ &= \left( |v|^2 \frac{v \cdot x}{|v \cdot x|} + (1 + \chi\zeta)V_1|x| - (1 + \chi\zeta)|v \cdot x| \right) \langle \gamma x \rangle^{q-2} \\ &\quad + (q-2)\gamma|v \cdot x|(v \cdot x)\langle \gamma x \rangle^{q-4},\end{aligned}$$

where we recall that  $V_1$  has been defined in (2.2). We consider  $\beta, \gamma \in (0, 1)$  to be fixed later such that the weight function

$$\tilde{m}_q := \langle \gamma x \rangle^q + q\gamma(v \cdot x)\langle \gamma x \rangle^{q-2} - q\beta|v \cdot x|\langle \gamma x \rangle^{q-2}$$

satisfies

$$(1 - \delta)\langle \gamma x \rangle^q \leq \tilde{m}_q \leq (1 + \delta)\langle \gamma x \rangle^q,$$

for some  $\delta \in (0, 1)$ . Gathering the previous estimates, there holds

$$\begin{aligned}\mathcal{B}_1^* \tilde{m}_q &= \mathcal{L}^* \tilde{m}_q - \mathcal{A}_1^* \tilde{m}_q = \mathcal{L}^* \tilde{m}_q - (1 + \chi\zeta)\phi_R \int_{\mathcal{V}} \tilde{m}_q dv \\ &= q\gamma|v|^2 \langle \gamma x \rangle^{q-2} + q(q-2)\gamma(v \cdot x)^2 \langle \gamma x \rangle^{q-4} - q\gamma\chi|v \cdot x|\langle \gamma x \rangle^{q-2} \\ &\quad - q\beta|v|^2 \frac{v \cdot x}{|v \cdot x|} \langle \gamma x \rangle^{q-2} - q(q-2)\beta\gamma|v \cdot x|(v \cdot x)\langle \gamma x \rangle^{q-4} \\ &\quad - q\beta(1 + \chi\zeta)V_1|x|\langle \gamma x \rangle^{q-2}\phi_R^c + q\beta(1 + \chi\zeta)|v \cdot x|\langle \gamma x \rangle^{q-2} \\ &\quad - (1 + \chi\zeta)\langle \gamma x \rangle^q \phi_R,\end{aligned}$$

and then

$$\begin{aligned}\mathcal{B}_1^* \tilde{m}_q &\leq q\gamma V_0^2 \langle \gamma x \rangle^{q-2} + q|q-2|\gamma V_0^2 |x|^2 \langle \gamma x \rangle^{q-4} - q\gamma\chi|v \cdot x|\langle \gamma x \rangle^{q-2} \\ &\quad + q\beta V_0^2 \langle \gamma x \rangle^{q-2} + q|q-2|\beta\gamma V_0^2 |x|^2 \langle \gamma x \rangle^{q-2} \\ &\quad - q\beta(1 - \chi)V_1|x|\langle \gamma x \rangle^{q-2} + q\beta(1 + \chi)|v \cdot x|\langle \gamma x \rangle^{q-2} \\ &\quad - (1 - \chi)\langle \gamma x \rangle^{q-1}\phi_R + q\beta(1 - \chi)V_1\langle \gamma x \rangle^{q-1}\phi_R \\ &\leq \left( qV_0^2(\gamma + \beta) + q\beta(1 - \chi)V_1 \right) \langle \gamma x \rangle^{q-2} + q|q-2|V_0^2\gamma(1 + \beta)|x|^2 \langle \gamma x \rangle^{q-4} \\ &\quad - (1 - \chi)(1 - q\beta V_1)\langle \gamma x \rangle^q \phi_R - q\beta(1 - \chi)V_1\langle \gamma x \rangle^{q-1} \\ &\leq \left( \frac{C_1}{\langle \gamma x \rangle} - C_2\phi_R - q\beta(1 - \chi)V_1 \right) \langle \gamma x \rangle^{q-1}.\end{aligned}$$

Choosing  $\beta(1 + \chi) = \gamma\chi$  with  $\gamma > 0$  small enough and  $R \geq 1$  large enough, and observing that  $C_1 = O(\gamma)$ ,  $C_2 \geq (1 - \chi)/2$  as  $\gamma \rightarrow 0$ , we deduce

$$\mathcal{B}_1^* \tilde{m}_q \leq -\frac{q\beta(1 - \chi)V_1}{2} \langle \gamma x \rangle^{q-1} \lesssim -\langle x \rangle^{q-1}.$$

We denote  $f_{\mathcal{B}_1}(t) := S_{\mathcal{B}_1}(t)f_0$  for some  $0 \leq f_0 \in L_k^1$  and then  $\widetilde{M}_q = \langle\langle f_{\mathcal{B}_1} \tilde{m}_q \rangle\rangle$ ,  $M_q = \langle\langle f_{\mathcal{B}_1} \langle x \rangle^q \rangle\rangle$ , so that

$$(4.18) \quad \widetilde{M}_q \lesssim M_q \lesssim \widetilde{M}_q.$$

From the above inequality, we get

$$(4.19) \quad \frac{d}{dt} \widetilde{M}_q = \int f(\mathcal{B}_1^* \tilde{m}_q) \lesssim -M_{q-1},$$

and in particular

$$(4.20) \quad \widetilde{M}_k(t) \leq \widetilde{M}_k(0), \quad \forall t \geq 0.$$

A classical interpolation inequality together with (4.18) and (4.20) give

$$M_\ell(t) \leq M_{\ell-1}(t)^\theta M_k(t)^{1-\theta} \lesssim M_{\ell-1}(t)^\theta M_k(0)^{1-\theta},$$

with  $\theta \in (0, 1)$  such that  $\ell = \theta(\ell - 1) + (1 - \theta)k$ . Coming back to (4.19), we get

$$\frac{d}{dt} \widetilde{M}_\ell \lesssim -M_k(0)^{-1/\alpha} \widetilde{M}_\ell^{1+1/\alpha}$$

where

$$\alpha := \frac{1}{(1/\theta) - 1} = k - \ell.$$

Integrating the above differential inequality, we obtain

$$M_\ell(t) \lesssim \frac{M_k(0)}{t^\alpha}, \quad \forall t > 0,$$

and we conclude gathering that last inequality with (4.20).  $\square$

In order to establish the asymptotic stability in  $L^1(m)$  for a polynomial weight  $m$ , we write

$$\mathcal{S}_{\mathcal{L}} \Pi^\perp = \Pi^\perp \mathcal{S}_{\mathcal{B}_1} + (\mathcal{S}_{\mathcal{L}} \Pi^\perp) * (\mathcal{A}_1 \mathcal{S}_{\mathcal{B}_1}).$$

Introducing the exponential weight  $m_0 := e^{\langle x \rangle}$ , we observe that  $\Pi^\perp \mathcal{S}_{\mathcal{B}_1} : L^1(m) \rightarrow L^1$  and  $\mathcal{A}_1 \mathcal{S}_{\mathcal{B}_1} L^1(m) \rightarrow L^1(m_0)$ , with rate  $\langle t \rangle^{-\ell}$  for any  $\ell \in (0, k)$  from Lemma 4.4. Because we have already established that  $\mathcal{S}_{\mathcal{L}} \Pi^\perp : L^1(m_0) \rightarrow L^1$  with rate  $e^{at}$  for any  $a \in (a^*, 0)$ , we immediately conclude that (1.9) holds in  $L^1(m)$ .

## REFERENCES

- [1] ALT, W. Biased random walk models for chemotaxis and related diffusion approximations. *J. Math. Biol.* 9, 2 (1980), 147–177.
- [2] BARDOS, C., AND DEGOND, P. Global existence for the Vlasov-Poisson equation in 3 space variables with small initial data. *Ann. Inst. H. Poincaré Anal. Non Linéaire* 2, 2 (1985), 101–118.
- [3] BOUCHUT, F., AND DESVILLETES, L. Averaging lemmas without time Fourier transform and application to discretized kinetic equations. *Proc. Roy. Soc. Edinburgh Sect. A* 129, 1 (1999), 19–36.
- [4] BOURNAVEAS, N., AND CALVEZ, V. A review of recent existence and blow-up results for kinetic models of chemotaxis. *Can. Appl. Math. Q.* 18, 3 (2010), 253–265.
- [5] CALVEZ, V., RAOUL, G., AND SCHMEISER, C. Confinement by biased velocity jumps: aggregation of *escherichia coli*. *Kinet. Relat. Models* 8, 4 (2015), 651–666.
- [6] DESVILLETES, L., AND MISCHLER, S. About the splitting algorithm for Boltzmann and B.G.K. equations. *Math. Models Methods Appl. Sci.* 6, 8 (1996), 1079–1101.
- [7] DOLBEAULT, J., MOUHOT, C., AND SCHMEISER, C. Hypocoercivity for linear kinetic equations conserving mass. *Trans. Amer. Math. Soc.* 367, 6 (2015), 3807–3828.
- [8] ERBAN, R., AND OTHMER, H. G. From individual to collective behavior in bacterial chemotaxis. *SIAM J. Appl. Math.* 65, 2 (2004/05), 361–391 (electronic).
- [9] ESCOBEDO, M., MISCHLER, S., AND RODRIGUEZ RICARD, M. On self-similarity and stationary problem for fragmentation and coagulation models. *Ann. Inst. H. Poincaré Anal. Non Linéaire* 22, 1 (2005), 99–125.
- [10] GOLSE, F., LIONS, P.-L., PERTHAME, B., AND SENTIS, R. Regularity of the moments of the solution of a transport equation. *J. Funct. Anal.* 76, 1 (1988), 110–125.
- [11] GOLSE, F., PERTHAME, B., AND SENTIS, R. Un résultat de compacité pour les équations de transport et application au calcul de la limite de la valeur propre principale d’un opérateur de transport. *C. R. Acad. Sci. Paris Sér. I Math.* 301, 7 (1985), 341–344.
- [12] GUALDANI, M., MISCHLER, S., AND MOUHOT, C. Factorization for non-symmetric operators and exponential H-Theorem. arxiv:1006.5523.

- [13] JAMES, F., AND VAUCHELET, N. Chemotaxis: from kinetic equations to aggregate dynamics. *NoDEA Nonlinear Differential Equations Appl.* 20, 1 (2013), 101–127.
- [14] LIONS, P.-L., AND PERTHAME, B. Lemmes de moments, de moyenne et de dispersion. *C. R. Acad. Sci. Paris Sér. I Math.* 314, 11 (1992), 801–806.
- [15] MISCHLER, S. Semigroups in Banach spaces - factorization approach for spectral analysis and asymptotic estimates. *In preparation.*
- [16] MISCHLER, S., AND MOUHOT, C. Exponential stability of slowly decaying solutions to the kinetic-Fokker-Planck equation. Preprint arxiv:1412.7487, 2015.
- [17] MISCHLER, S., QUIÑINAO, C., AND TOUBOUL, J. On a kinetic fitzhugh-nagumo model of neuronal network. hal-01108872.
- [18] MISCHLER, S., AND SCHER, J. Spectral analysis of semigroups and growth-fragmentation equations. Preprint arXiv:1310.7773, 2013.
- [19] OTHMER, H. G., DUNBAR, S. R., AND ALT, W. Models of dispersal in biological systems. *J. Math. Biol.* 26, 3 (1988), 263–298.
- [20] OTHMER, H. G., AND STEVENS, A. Aggregation, blowup, and collapse: the ABCs of taxis in reinforced random walks. *SIAM J. Appl. Math.* 57, 4 (1997), 1044–1081.
- [21] PERTHAME, B. Global existence to the BGK model of Boltzmann equation. *J. Differential Equations* 82, 1 (1989), 191–205.
- [22] PERTHAME, B. Time decay, propagation of low moments and dispersive effects for kinetic equations. *Comm. Partial Differential Equations* 21, 3-4 (1996), 659–686.
- [23] VILLANI, C. Hypocoercivity. *Mem. Amer. Math. Soc.* 202, 950 (2009), iv+141.

STÉPHANE MISCHLER

UNIVERSITÉ PARIS-DAUPHINE  
CEREMADE, UMR CNRS 7534  
PLACE DU MARÉCHAL DE LATTRE DE TASSIGNY 75775 PARIS CEDEX 16  
FRANCE

E-MAIL: [mischler@ceremade.dauphine.fr](mailto:mischler@ceremade.dauphine.fr)

QILONG WENG

UNIVERSITÉ PARIS-DAUPHINE  
CEREMADE, UMR CNRS 7534  
PLACE DU MARÉCHAL DE LATTRE DE TASSIGNY 75775 PARIS CEDEX 16  
FRANCE

E-MAIL: [weng@ceremade.dauphine.fr](mailto:weng@ceremade.dauphine.fr)