

# Generalized guidance equation for peaked quantum solitons and effective gravity

Thomas Durt

# ► To cite this version:

Thomas Durt. Generalized guidance equation for peaked quantum solitons and effective gravity. EPL - Europhysics Letters, 2016, 114, pp.10004. 10.1209/0295-5075/114/10004 . hal-01272636v2

# HAL Id: hal-01272636 https://hal.science/hal-01272636v2

Submitted on 11 Apr 2016  $\,$ 

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Generalized guidance equation for peaked quantum solitons and effective gravity.

## Thomas Durt<sup>1</sup>

### Abstract

Bouncing oil droplets have been shown to follow de Broglie-Bohm like trajectories and at the same time they exhibit attractive and repulsive pseudo-gravitation. We propose a model aimed at rendering account of these phenomenological observations. It inspires, in a more speculative approach, a toy model for quantum gravity. In studies concerning quantum wave mechanics, de Broglie-Bohm (dB-B) trajectories [10, 3, 17] remained during many years a rather confidential and academic topic, but they regained interest since they were realized in the lab. with artificial macroscopic systems, the bouncing oil droplets or walkers [4, 9], which were shown experimentally to follow dB-B like quantum trajectories. For instance, when the walker passes through one slit of a two-slit device, it undergoes the influence of its "pilot-wave" passing through the other slit, in such a way that, after averaging over many dB-B like trajectories, the interference pattern typical of a double-slit experiment is restored, despite of the fact that each walker passes through only one slit. Pseudogravitational interaction has also been reported between two droplets. In [9] for instance we can read: ... We find that, depending on the value of d, (d represents here the impact parameter of the collision) the interaction is either repulsive or attractive. When repulsive, the drops follow two approximately hyperbolic trajectories. When attractive, there is usually a mutual capture of the two walkers into an orbital motion similar to that of twin stars ...

Our study was motivated by the double solution program [2, 10, 11] of Louis de Broglie, according to which the quantum wave function, solution of the linear Schrödinger equation does not contain all elements of reality concerning the quantum system. In de Broglie's view, particles would consist of a peaked concentration of energy that would coexist with the linear wave function, and the latter would guide the former according to the de Broglie-Bohm guidance equation. Our work consists of an attempt to realize de Broglie's program by adding another ingredient to it, which is that we suppose the existence of a non-linear self focusing potential (2) of gravitational nature which prohibits the spreading of the peaked soliton (solitary wave) representing the particle. Therefore we assume that the system evolves in time according to the non-linear equation (1).

$$i\hbar\frac{\partial\Psi(t,\mathbf{x})}{\partial t} = -\hbar^2\frac{\Delta\Psi(t,\mathbf{x})}{2m} + V^L(t,\mathbf{x})\Psi(t,\mathbf{x}) + V^{NL}(\Psi)\Psi(t,\mathbf{x}), \quad (1)$$

 $<sup>^1 \</sup>rm Aix$  Marseille Université, CNRS, Centrale Marseille, Institut Fresnel (UMR 7249),13013 Marseille, France.email: thomas.durt@centrale-marseille.fr

where  $V^{L}(t, \mathbf{x})$  represents the usual external potentials, for instance it could represent the coupling to external electro-magnetic fields. It is thus represented by a self-adjoint operator linearily acting on the Hilbert space.  $V_{NL}$  represents a self focusing non-linearity. Contrary to  $V^{L}$  it depends on  $\Psi$ . In order to distinguish short range and long range we shall consider in the rest of the paper the Yukawa like screened S-N potential:

$$V^{NL}(\Psi) \equiv -K \int d^3 x' \left(\frac{|\Psi(t, \mathbf{x}')|^2}{|\mathbf{x} - \mathbf{x}'|}\right) e^{-\frac{|\mathbf{x} - \mathbf{x}'|}{\lambda}},\tag{2}$$

where K is a positive coupling constant and  $\lambda$  is a characteristic length to be defined soon. The Schrödinger-Newton (S-N) potential is in turn defined as follows:  $V_{SN}^{NL}(\Psi)((t, \mathbf{x}) \equiv -Gm^2 \int d^3x' (\frac{|\Psi(t, \mathbf{x}')|^2}{|\mathbf{x} - \mathbf{x}'|})$ ; (1) has been abundantly studied in the literature in presence of the S-N potential and in absence of external potential ( $V^L = 0$ ). In particular, when  $V^L = 0$ , (1) possesses a "ground-state" solution of the form  $\psi(\mathbf{x}, t) = e^{-\frac{iE_g t}{\hbar}} \phi_{NLg}(\mathbf{x})$ ;  $\phi_{NLg}$  is thus a solution of

$$\frac{-\hbar^2}{2m} \Delta \phi_{NLg}(\mathbf{x}) + Gm^2 \int d^3 y \, \frac{\left|\phi_{NLg}(\mathbf{y})\right|^2}{\left|\mathbf{x} - \mathbf{y}\right|} \, \phi_{NLg}(\mathbf{x}) \\ = E_g \phi_{NLg}(\mathbf{x}) \,, \tag{3}$$

which was studied in astrophysics and is known under the name of the *Choquard* equation [18, 5]. In [18], Lieb showed that the energy functional

$$E(\phi) = \frac{\hbar^2}{2m} \int d^3x \left| \nabla \phi(\mathbf{x}) \right|^2$$

$$- \frac{Gm^2}{2} \int d^3x \int d^3y \frac{\left| \phi(\mathbf{y}) \right|^2}{|\mathbf{x} - \mathbf{y}|} \left| \phi(\mathbf{x}) \right|^2,$$
(4)

is minimized by a unique solution  $\phi_{NLg}(\mathbf{x})$  of the Choquard equation (3) for a given  $L_2$  norm  $N(\phi)$ .

As is well-known, the Choquard equation possesses well-defined scaling properties [5, 25]; in particular, we may scale its solution in such a way that in the limit where the  $L_2$  norm goes to plus infinity, the extent goes to zero, and the energy to minus infinity. Numerical studies also established that  $\phi_{NLg}$  has a "hump", quasi-gaussian shape [5, 25]. In the case of quantum particles the non-linear potential  $V^{NL}$  aims represents in our eyes a short-range self-focusing interaction of gravitational nature which localizes the wave function  $\Psi$ , analogous to the Poincaré pressure [20] aimed at ensuring the cohesion of electrons in presence of Coulombian self-repulsion. In the rest of the paper we shall identify the quantum particles with these self-collapsed wave packets (solitons). Our aim is to show that when several particles are present there appears a long range effective gravitational interaction between them. We explicitly established a distinction between gravitation at short and long spatial scale in our model, with the aim of gaining information about the short scale behaviour by fitting the long range effective gravitational potential with Newton gravitation. In the case of bouncing droplets, the non-linearity mimicks the complicate mechanism which ensures their stability. That we consider droplets or quantum systems,  $\lambda$  is intentionally chosen to be larger than the size of the object under interest (particle or droplet), in such a way that the solitonic ground state of (1) when  $V^L = 0$  is essentially the solution  $\phi_{NLg}(\mathbf{x})$  of (3). If we consider quantum particles like the electron or the proton, we choose K to be  $Gm^2$ ; in the case of droplets K ought to be calibrated in function of the experimental set-up.

The first real novelty of our paper is that we assume that the solution of (1) obeys the factorisation ansatz:

$$\Psi = \left(\frac{1}{A_L(t, \mathbf{x_0}(\mathbf{t}))}\right) \Psi_L(t, \mathbf{x}) \cdot \phi_{NLg}(\mathbf{x} - \mathbf{x_0}(t)) e^{-iE_g t/\hbar},\tag{5}$$

where we introduce the pilot wave  $\Psi_L$ , of amplitude  $A_L$  and phase  $\varphi_L$ :  $\Psi_L(t, \mathbf{x}) \equiv A_L(t, \mathbf{x}) e^{i\varphi(t, \mathbf{x})}$  and  $\mathbf{x}_0$  the position of the barycentre of the soliton  $\phi_{NLg}$ . We also suppose to begin with that  $\mathbf{x}_0$  follows a de Broglie-Bohm trajectory:

$$\mathbf{x}_{0}(t) = \mathbf{x}_{0}(t=0) + \int_{0}^{t} dt \mathbf{v}_{dB-B},$$
(6)

where  $\mathbf{v}_{dB-B}$  represents the dB-B velocity derived from the pilot wave  $\Psi_L$  evaluated at the position of the barycentre of  $\Psi$ :

$$\mathbf{v}_{dB-B} = \frac{\hbar}{m} \nabla \varphi_L(\mathbf{x}_0(t), t). \tag{7}$$

Our goal is to study the feedback of the soliton on the pilot wave. We shall thus study in the rest of the paper solutions of (1,2), under the constraints (5,6,7).

The second novelty of our paper is that contrary to the majority of the papers that can be found in the literature concerning the S-N equation (1), we assume, for reasons that we shall clarify in the core of the paper, that whenever we consider elementary particles the size of the solution of (1,5) is very small, actually of the order of  $Gm/c^2$ . In the case of an electron for instance this size is of the order of  $10^{-55}$  meter. This means [5] that the energy (4) of the self-collapsed ground state is of the order of  $-mc^2$ . In our approach, the external potentials are thus treated as a perturbation, contrary to the standard approach, where the non-linearity is treated as a weak perturbation [22, 6]. The non-linearity is so strong in our case that it forces the state of the system to behave as a solitary wave, a soliton of extremely small size, solution of the so-called Choquard equation (3) for which the spread gets compensated by the non-linearity. The wave function of the system is thus supposedly self-collapsed to begin with, since arbitrary long times. This marks a difference with previous studies which rather focused on the collapse process itself [12, 19, 25, 5, 6]. We consider situations where  $A_L$  and  $\varphi_L$  smoothly vary in space and time compared to  $\phi_{NLg}$  so that in good approximation (5) reads

$$\Psi = e^{i\varphi_L} \cdot \phi_{NLg}(\mathbf{x} - \mathbf{x_0}(t))e^{-iE_g t/\hbar},\tag{8}$$

which explicitly shows that  $\Psi$  is a constant  $L_2$  norm function, in accordance with the norm-preserving character of the evolution equation (1). In this paper, we shall apply these ideas to a system consisting of two particles in order to derive an effective long range gravitational attraction between the particles. We also compare our predictions to droplets phenomenology and make in this case new predictions that are likely to be tested in the lab.

#### 1 Effective gravitation

To simplify the treatment we shall first consider distinguishable particles  ${\cal A}$  and  ${\cal B}$  and suppose that the wave function of the full system reads

A and *D* and suppose that the wave function of the full system reads  $\Psi(\mathbf{x},t) = \Psi_L(\mathbf{x},t) \cdot \left(\frac{\phi_{NLg}^{A}(\mathbf{x},t)}{A_L(\mathbf{x_0}^{A},t)} + \frac{\phi_{NLg}^{B}(\mathbf{x},t)}{A_L(\mathbf{x_0}^{B},t)}\right) \text{ where we introduced the short notation}$  $<math display="block">\phi_{NLg}^{A(B)}(\mathbf{x},t) \equiv \phi_{NLg}(\mathbf{x} - \mathbf{x_0}^{A(B)}(t))e^{-iE_gt/\hbar}, \text{ with } \mathbf{x_0}^{A} \text{ and } \mathbf{x_0}^{B} \text{ the positions of the barycentres of the solitons assigned to the two particles.}$  $In virtue of (8), at the locations <math>\mathbf{x_0}^{A(B)}, V^{NL}(\Psi) = V^{NL}(\phi_{NLg}), \text{ so that}$  $\phi_{NLq}^{A(B)}(\mathbf{x},t)$  satisfies the constraint

$$i\hbar \cdot \frac{\partial \phi_{NLg}^{A(B)}}{\partial t} = -\frac{\hbar^2}{2m} \cdot \Delta \phi_{NLg}^{A(B)} + V^{NL}(\Psi) \phi_{NLg}^{A(B)}$$
$$-\frac{\hbar^2}{m} \cdot i \nabla \varphi_L(t, \mathbf{x_0^{A(B)}}) \cdot \nabla \phi_{NLg}^{A(B)}$$
(9)

Multiplying (9) by  $\Psi_L(\mathbf{x}, t)$ , and substracting the result from (1) we find, in the low-dBB-velocity regime where the temporal derivative of  $A_L(\mathbf{x_0^{A(B)}}, t)$  is small compared to other contributions

$$\left(\frac{\phi_{NLg}^{A}(\mathbf{x},t)}{A_{L}(\mathbf{x_{0}^{A}},t)} + \frac{\phi_{NLg}^{B}(\mathbf{x},t)}{A_{L}(\mathbf{x_{0}^{B}},t)}\right)\left(i\hbar \cdot \frac{\partial\Psi_{L}(t,\mathbf{x})}{\partial t} + \frac{\hbar^{2}}{2m}\Delta\Psi_{L}(t,\mathbf{x}) - V^{L}(t,\mathbf{x})\Psi_{L}(t,\mathbf{x})\right) = -\frac{\hbar^{2}}{m}\left(\nabla\Psi_{L}(t,\mathbf{x}) + \Psi_{L}(t,\mathbf{x}) \cdot i\nabla\varphi_{L}(t,\mathbf{x_{0}})\right) \cdot \nabla\left(\frac{\phi_{NLg}^{A}(\mathbf{x},t)}{A_{L}(\mathbf{x_{0}^{A}},t)} + \frac{\phi_{NLg}^{B}(\mathbf{x},t)}{A_{L}(\mathbf{x_{0}^{B}},t)}\right).$$
(10)

We shall only consider situations where the positions  $\mathbf{x}_0^{\mathbf{A}}$  and  $\mathbf{x}_0^{\mathbf{B}}$  are not too close to each other e.g. larger than  $\lambda$ ; then we may neglect the overlap between the two solitons, so that, dividing (10) by  $\left(\frac{\phi_{NLg}^{A}(\mathbf{x},t)}{A_{L}(\mathbf{x_{0}^{A}},t)} + \frac{\phi_{NLg}^{A}(\mathbf{x_{0}^{A}},t)}{\Phi_{L}(\mathbf{x_{0}^{A}},t)}\right)$  $\frac{\phi^B_{NLg}(\mathbf{x},t)}{A_L(\mathbf{x_0^B},t)}\big)$  we find

$$i\hbar \cdot \frac{\partial \Psi_L(t, \mathbf{x})}{\partial t} + \frac{\hbar^2}{2m} \Delta \Psi_L(t, \mathbf{x}) - V^L(t, \mathbf{x}) \Psi_L(t, \mathbf{x}) = (-\frac{\hbar^2}{m} (\nabla \Psi_L(t, \mathbf{x}) + \Psi_L(t, \mathbf{x}) \cdot i \nabla \varphi_L(t, \mathbf{x_0})) \cdot (\frac{\nabla \phi^A_{NLg}(\mathbf{x}, t)}{\phi^A_{NLg}(\mathbf{x}, t)} + \frac{\nabla \phi^B_{NLg}(\mathbf{x}, t)}{\phi^B_{NLg}(\mathbf{x}, t)}),$$
(11)

which looks like the linear Schrödinger equation with source terms. At some distance from the soliton  $\phi_{NLg}$  decreases exponentially, because  $\phi_{NLg}$  is a negative energy state with radial symmetry. Moreover  $\frac{\nabla_{x,y,z}\phi_{NLg}(\mathbf{x},t)}{\phi_{NLg}(\mathbf{x},t)}$  is odd in x, y, z around x around x to be the last of the state o is odd in x, y, z around  $x_0, y_0, z_0$ . It behaves thus along the x, y, z axis passing through  $(x_0, y_0, z_0)$  as a kink with a s-like shape --/--. In the plane orthogonal to this axis its tail decreases like  $(x-x_0)/\sqrt{(x-x_0)^2 + (y-y_0)^2 + (z-z_0)^2}$ , but we should not worry about the tails because when several particles are isotropically distributed, the tails cancel each other and the shape of the kink is rather --/-. The integration over space of the source term proportional to  $\Psi_L$  is equal to zero. Integrating the source term

proportional to  $\nabla \Psi_L$  by part we find that everything happens as if the source terms in equation (10) were proportional to Dirac 3-deltas, and we may write (10) in the form

$$i\hbar \cdot \frac{\partial \Psi_L(t, \mathbf{x})}{\partial t} + \frac{\hbar^2}{2m} \Delta \Psi_L(t, \mathbf{x}) - V^L(t, \mathbf{x}) \Psi_L(t, \mathbf{x})$$
$$= -4\pi \frac{\hbar^2}{2m} \Psi_L(t, \mathbf{x}) \cdot (L_A \delta^{Dirac}(\mathbf{x} - \mathbf{x_0^A}) + L_B \delta^{Dirac}(\mathbf{x} - \mathbf{x_0^B})),$$
(12)

where L is a length of the order of the size (extent) of the particle. We incorporated in  $L_{A(B)}$  factors of the order of unity, in order to simplify the computations.

In a perturbative or bootstrap approach, we shall now treat (12) as a generalised Poisson equation with a source term proportional to a solution of the homogeneous (linear Schrödinger) equation without sources; moreover we impose a factorisable solution in the form

 $\Psi_L^{inhom}(t, \mathbf{x}) = \Psi_L^{hom}(t, \mathbf{x})(-\phi^G(t, \mathbf{x})/c^2)$ , then (12) reads

$$(-\phi^{G}(t,\mathbf{x})/c^{2})(i\hbar \cdot \frac{\partial \Psi_{L}^{hom}(t,\mathbf{x})}{\partial t} + \frac{\hbar^{2}}{2m}\Delta\Psi_{L}^{hom}(t,\mathbf{x})$$
$$-V^{L}(t,\mathbf{x})) + \frac{\hbar^{2}}{m}\nabla\Psi_{L}^{hom}(t,\mathbf{x})\cdot\nabla(-\phi^{G}(t,\mathbf{x})/c^{2})$$
$$+(\Psi_{L}^{hom}(t,\mathbf{x}))(i\hbar \cdot \frac{\partial(-\phi^{G}(t,\mathbf{x})/c^{2})}{\partial t} + \frac{\hbar^{2}}{2m}\Delta(-\phi^{G}(t,\mathbf{x})/c^{2})) = -4\pi\frac{\hbar^{2}}{2m}\cdot\Psi_{L}^{hom}(t,\mathbf{x})\cdot$$
$$(L_{A}\delta^{Dirac}(\mathbf{x}-\mathbf{x_{0}^{A}}) + L_{B}\delta^{Dirac}(\mathbf{x}-\mathbf{x_{0}^{B}})).$$
(13)

The three first terms of (13) are equal to zero, because  $\Psi_L^{hom}$  is solution of the linear Schrödinger equation without sources. It is consistent, in the low-dBB-velocity regime, to neglect

 $\nabla \Psi_L^{hom}(t, \mathbf{x}) \cdot \nabla (-\phi^G(t, \mathbf{x})/c^2)$  as well as  $\frac{\partial (-\phi^G(t, \mathbf{x})/c^2)}{\partial t}$  as we shall explain soon. Then  $\phi^G$  obeys the Poisson equation

$$\Delta(\phi^G(t, \mathbf{x})/c^2) = 4\pi \cdot (L_A \delta^{Dirac}(\mathbf{x} - \mathbf{x_0^A}) + L_B \delta^{Dirac}(\mathbf{x} - \mathbf{x_0^B}))$$
(14)

Making use of the well-known properties of the Green functions associated to the Laplace equation, it is easy to check that the solution of (14) is

$$\phi^{G}(t,\mathbf{x})/c^{2} = -\left(\frac{L_{A}}{|\mathbf{x} - \mathbf{x_{0}^{A}}|} + \frac{L_{B}}{|\mathbf{x} - \mathbf{x_{0}^{B}}|}\right).$$
(15)

 $\phi^G(t, \mathbf{x})/c^2$  looks thus like an effective (here repulsive) gravitational potential<sup>2</sup>. We check a posteriori that we were in right to neglect certain corrections proportional to the dB-B velocities.

In order to fit  $\phi^G$  with a Newtonian potential in the long range domain, let us consider  $H_L$  the linear Hamiltonian of the system:

<sup>&</sup>lt;sup>2</sup>Note that the singularities of this expression in  $\mathbf{x_0^A}$  and  $\mathbf{x_0^B}$  are artificial, they result from our discretisation procedure.

 $H_L = H_L^A + H_L^B + m_A c^2 + m_B c^2 + V_L^{AB}, \text{ including a possible coupling}$ between the A and B systems. In the non-relativistic regime,  $H_L^{A(B)} << m_{A(B)}c^2$ ,  $H_L^{AB} << m_{A(B)}c^2$ ,  $|i\hbar \cdot \frac{\partial \Psi}{\partial t}| << m_{A(B)}c^2|\Psi_L|$ . Note that  $\Psi_L(t, \mathbf{x})$ , the solution of (12) obeys  $\Psi_L = \Psi_L^{hom} + \Psi^{inhom} =$ 

 $(1 - \phi^G(t, \mathbf{x})/c^2)\Psi_L^{hom}$ . Let us from now on neglect self-interactions which were already taken account in  $V^{NL}$ . Requiring that  $\Psi_L^{hom} = \Psi_L/(1 - \psi_L)$  $\phi^G/c^2$ ) satisfies the homogeneous equation  $H_L \Psi_L^{hom} = i\hbar \cdot \frac{\partial \Psi_L^{hom}}{\partial t}$  results,  $\phi^{G'/c^*}$ ) satisfies the homogeneous equation  $H_L \Psi_L^{Gam} = i\hbar \cdot \frac{1}{\partial t}$  results, provided we only consider the dominating contribution, in the replacement in  $H^{A(B)}$  of  $m_{A(B)}c^2$  by  $m_{A(B)}c^2(m_{A(B)}c^2)\phi^G_{A(B)}(t, \mathbf{x})/c^2$ . At this level, an effective gravitational interaction between A and B emerges which satisfies  $\phi^G_{A(B)}(t, \mathbf{x})/c^2 = -(\frac{L_{B(A)}}{|\mathbf{x_0^{B(A)}}-\mathbf{x_0^{A(B)}}|})$ . Requiring the equivalence of this effective potential with the Newto-nian gravitational potential  $\phi^{Newton} = -\frac{Gm_Am_B}{|\mathbf{x_0^{B(A)}}-\mathbf{x_0^{A(B)}}|}$  imposes the con-ctroint:

straint:

$$L_{A(B)} = \frac{G \cdot m_{A(B)}}{2c^2},$$
(16)

as announced in the introduction.

Disregarding the self-interactions, the netto contribution of the perturbation is thus, when (16) is satisfied, to add  $-Gm_Am_B(\frac{1}{|\mathbf{x}_A^A-\mathbf{x}_B^B|})$  to the total energy.

If A and B represent identical particles, we impose that the fermionic or bosonic character of the full wave  $\Psi$  is expressed at the level of the linear component only. The non-linear components behave thus like bosons and we get

$$\Psi(1,2) = \Psi_L(t,\mathbf{x_1},\mathbf{x_2}) \cdot (\frac{1}{A_L}) \cdot S \cdot \phi^A_{NLg}(t,\mathbf{x_1})\phi^B_{NLg}(t,\mathbf{x_2}))$$
$$\approx e^{i\varphi_L(t,\mathbf{x_1},\mathbf{x_2})} \cdot S \cdot \phi^A_{NLg}(t,\mathbf{x_1})\phi^B_{NLg}(t,\mathbf{x_2})),$$

where  $\phi_{NLg}^{A(B)}(t, \mathbf{x_i}) = \phi_{NLg}(\mathbf{x_i} - \mathbf{x_0^{A(B)}}(t))e^{-iE_gt/\hbar}$  while S is a bosonic symmetrization operator:  $2S \cdot \phi_{NLg}^A(t, \mathbf{x_1})\phi_{NLg}^B(t, \mathbf{x_2})) = \phi_{NLg}^A(t, \mathbf{x_1})\phi_{NLg}^B(t, \mathbf{x_2}) + \phi_{NLg}^A(t, \mathbf{x_2})\phi_{NLg}^B(t, \mathbf{x_1}).$ The barycentres  $\mathbf{x_0^{A(B)}}$  of the particles move according to the two

particles dB-B guidance equation

$$\mathbf{v}_{\mathbf{i}drift} = \frac{\hbar}{m} \nabla_{\mathbf{i}} \varphi_L(\mathbf{x_0^{A(B)}}(t), \mathbf{x_0^{B(A)}}(t), t), (i = 1, 2)$$
(17)

In the present case, (1) and (17) are dynamical equations in the  $3N{=}6$ dimensional configuration space (here there are N = 2 particles). Therefore  $\Psi_L(1,2) = A_L e^{i\varphi_L}$ , where  $A_L$  and  $\varphi_L$  depend on  $\mathbf{x}^1$  and  $\mathbf{x}^2$  instantaneously. The system is therefore likely to exhibit a non-local [16] behaviour in presence of entanglement [3].

We shall only consider situations where  $\mathbf{x}_0^{\mathbf{A}}$  and  $\mathbf{x}_0^{\mathbf{B}}$  do not overlap. By computations very similar to those already performed in the case of indistingusihable particles, we find that an effective gravitational interaction appears, symmetrized in  $\mathbf{x}_1$  and  $\mathbf{x}_2$ .

$$\phi^{G}(t, \mathbf{x})/c^{2} = -L\Sigma_{i=1,2}\left(\frac{1}{|\mathbf{x}_{i} - \mathbf{x}_{0}^{\mathbf{A}}|} + \frac{1}{|\mathbf{x}_{i} - \mathbf{x}_{0}^{\mathbf{B}}|}\right)$$
(18)

Fitting this expression with the Newton interaction provides as before the constraint  $L = \frac{G \cdot m}{2c^2}$  that we may also rewrite in the form  $mc^2 = \frac{G \cdot m^2}{2L}$  in accordance with the results announced in the introduction and in particular with the value of the energy functional (4) which is of the order of  $-mc^2$  in this case. Relying on previous stability analyses of the S-N equation [1, 25, 5], it is worth noting that to destabilize the soliton would require a positive energy of the order of  $mc^2$  which confirms our interpretation in which the soliton and the particle are the same object.

# 2 Droplets, dB-B trajectories, and gravitation

dB-B trajectories [10, 3, 17] have been studied in relation with the measurement problem, but they are often considered as conceptual tools rather than empirical realities, among other reasons because the dB-B dynamics is often considered to be an ad hoc reinterpretation of the standard quantum theory. In the last decade, however, they were realized in the lab. with artificial macroscopic systems, the bouncing oil droplets or walkers [4, 9], which were shown to follow dB-B like quantum trajectories.

Our model seemingly catches some fundamental properties of droplets in the sense that it explains the appearance of an effective gravitational interaction in presence of dB-B trajectories, as they were observed directly in droplets phenomenology (and only there).

Of course, droplets are complicated, hydrodynamical, macroscopic systems, but in order to simplify the treatment we shall from now on identify the droplets with the self focused solitons of our model  $\phi_{NL}$  and the liquid on which they are floating with the pilot wave  $\Psi_L$ , which implies that the state of the system is represented by  $\Psi(\mathbf{x},t) = \Psi_L(\mathbf{x},t) \cdot \left(\frac{\phi_{NL}^{0(A)}(\mathbf{x},t)}{A_L(\mathbf{x_0^0},t)} + \frac{\phi_{NL}^{0(B)}(\mathbf{x},t)}{A_L(\mathbf{x_0^0},t)}\right)$ , where A and B refer to the presence of two droplets. It is then easy to repeat the same treatment as before, provided the two solitons stay away from each other. As has been reported in [8], stationary waves are present, resulting from the forcing frequency imposed to the container at frequency  $f_0$ . The stationary waves are characterized by Faraday wave lengths  $\lambda_F$  in good agreement with the values computed from the dispersion relation, at the frequency  $f_F$ , where the Faraday frequency  $f_F$  is equal to half the forcing frequency  $f_0$ .

In order to simulate the propagation of vibrations in the container, let us modelize, in a first step, the "Schrödinger" linear equation describing wave propagation in the container by the d' Alembert equation

$$\left(\frac{1}{v^2}\frac{\partial^2}{\partial t^2} - \Delta\right)\Psi_L = 0,\tag{19}$$

with  $v = \lambda_F \cdot f_F$ . Of course d'Alembert equation is not dispersive, whereas the liquid of the container is a dispersive medium, but we shall assume in first approximation that the velocity of propagation smoothly depends on the frequency, in the spectral domain under interest. This remark matters, because, as noted in [8] two frequencies play a role here,  $f_0$  and  $f_F = f_0/2$ .

When two droplets are present we must solve an inhomogeneous d' Alembert equation with a source term  $4\pi(L_A \cdot \delta(\mathbf{x} - \mathbf{x_0^A}) + L_B \cdot \delta(\mathbf{x} - \mathbf{x_0^B}))$ . Repeating the same process as for the inhomogeneous Schrödinger equation, we predict the appearance of a pseudo-gravitational interaction  $\phi^{PG}$  which obeys

$$\left(\frac{1}{v^2}\frac{\partial^2}{\partial t^2} - \Delta\right)\phi^{PG} = -4\pi(L_A \cdot \delta(\mathbf{x} - \mathbf{x_0^A}) + L_B \cdot \delta(\mathbf{x} - \mathbf{x_0^B})).$$

Due to the forcing, we shall replace d' Alembert equation by Helmholtz equation

$$\left(\frac{(2\pi)^2 f_0^2}{v^2} + \Delta\right) \phi^{PG} = (k_0^2 + \Delta) \phi^{PG} =$$

$$4\pi (L_A \cdot \delta(\mathbf{x} - \mathbf{x_0^A}) + L_B \cdot \delta(\mathbf{x} - \mathbf{x_0^B})),$$
(20)

where  $k_0 \equiv 2\pi/\lambda_0$ .

(20) has been studied by us in another context [14], and, in the case of a punctual source, its Green function is not 1/r but  $\cos(kr)/r$  which leads as shown in [14] to the appearance of attractive and repulsive gravitational zones periodically distributed among the punctual sources.

Henceforth we predict the appearance of a pseudo-gravitational field  $\phi^{PG} = -v^2 (L_A(\cos(k_0 |\mathbf{x} - \mathbf{x_0^A}|) / |\mathbf{x} - \mathbf{x_0^A}|) + L_B(\cos(k_0 |\mathbf{x} - \mathbf{x_0^B}|) / |\mathbf{x} - \mathbf{x_0^B}|)).$ 

Disregarding self-interactions, the sum of the potential energy undergone by A (due to B) with the potential energy undergone by B (due to A) is thus equal to

 $-v^2(\cos(k_0|\mathbf{x}^{\mathbf{B}} - \mathbf{x}_0^{\mathbf{A}}|)/|\mathbf{x}^{\mathbf{B}} - \mathbf{x}_0^{\mathbf{A}}|)(M_BL_A + M_AL_B)$ . Our model explains in this way the appearance of repulsive and attractive gravitation in droplets phenomenology, in connection with the presence of dB-B trajectories.

In the same vein, we explain the appearance of a pseudo-quantisation rule, self-adapting to the forcing frequency, and similar to the one observed in [7] according to which orbital radii obey  $d_n^{orb} = (n/2 - \epsilon)\lambda_F$ , with n a positive integer, where we made use of the fact that the Faraday frequency is one halve of the forcing frequency  $f_0$ . In our eyes this effect is not specifically quantum, it is rather related to the very unfamiliar topology of the attractive and repulsive (!) gravitational basins (in case of circular orbits we conjecture that the aforementioned quantisation rule imposes that after one closed circular orbit each soliton keeps the Faraday phase that it possessed before the revolution). Let us now propose a toy model for quantum gravity, inspired by droplet phenomenology.

## 3 Minimal coupling to gravity

Coming back to the single particle case, replacing  $\Psi_L^{hom}$  by  $\Psi_L^{hom}/(1-\phi^G/c^2)$  in the Schrödinger equation delivers the modified Schrödinger equation

$$i\hbar \cdot \frac{\partial}{\partial t} \left( \frac{\Psi_L(t, \mathbf{x})}{(1 - \phi^G/c^2)} \right) = -\frac{\hbar^2}{2m} \Delta \left( \frac{\Psi_L(t, \mathbf{x})}{(1 - \phi^G/c^2)} \right)$$
(21)  
+ $V^L(t, \mathbf{x}) \left( \frac{\Psi_L(t, \mathbf{x})}{(1 - \phi^G/c^2)} \right) + mc^2 \left( \frac{\Psi_L(t, \mathbf{x})}{(1 - \phi^G/c^2)} \right),$ 

to be interpreted according to us in a perturbative approach, in terms of the small parameter  $\phi^G/c^2$ , replacing  $1/(1 - \phi^G/c^2)$  by  $1 + \phi^G/c^2 + \phi^G/c^2$ 

 $(\phi^G/c^2)^2 + \dots$  There appears thus post-Newtonian corrections proportional to  $\nabla \Psi_L(t, \mathbf{x}) \cdot \nabla (-\phi^G(t, \mathbf{x})/c^2)$  as well as  $\frac{\partial (-\phi^G(t, \mathbf{x})/c^2)}{\partial t}$  and  $\phi^G/c^2$ . It is not in the scope of our paper to study these corrections<sup>3</sup>.

We shall emit the hypothesis that the minimal coupling condition that consists of replacing  $\Psi_L$  by  $\Psi_L/(1 - \phi^G/c^2)$  in the linear Schrödinger equation is general, and that  $\phi^G$  is a field in "real", 3-dimensional space, equal to the product of the gravitational fields generated by all species of indistinguishable elementary particles. In a regime where all these contributions  $\phi^k_G$  are small compared to  $c^2$ , we find that  $\pi_k(1 - \phi^k_G/c^2) \approx$  $(1 - \sum_k \phi^k_G/c^2)$ , which expresses the universal character of gravitation.

If we had chosen to begin with to describe the system with the relativistic Klein-Gordon equation (that we write in the free case here for simplicity but the result is still valid in the presence of external potentials e.g. electro-magnetic potentials)

$$\left(c^{2}\left(\frac{1}{c^{2}}\frac{\partial^{2}}{\partial t^{2}}-\Delta\right)-\frac{m^{2}c^{2}}{\hbar^{2}}\right)\Psi_{L}=0$$
(22)

then, by a treatment similar to the one performed in the previous sections we would have found finally that the gravitational potential obeys the inhomogeneous d' Alembert equation:

$$-c^{2}\left(\frac{1}{c^{2}}\frac{\partial^{2}}{\partial t^{2}}-\Delta\right)\phi_{G} = 4\pi G\rho.$$
(23)

We find so that all gravitational waves move at the (same) speed of light in vacuum.

Beginning with the Dirac equation

$$\beta i\hbar \partial_t \Psi(t, \mathbf{x}) - \beta \alpha c \frac{\hbar}{i} \nabla \Psi(t, \mathbf{x}) = mc^2 \Psi(t, \mathbf{x}) + V_{NL} \Psi(t, \mathbf{x}),$$

where  $\alpha$  and  $\beta$  represent Dirac matrices and imposing the ansatz  $\Psi = \Psi_{\mathbf{L}}(t, \mathbf{x}) \cdot \phi_{NL}(t, \mathbf{x})$  where  $\Psi_{\mathbf{L}}(t, \mathbf{x})$  a Dirac 4-spinor and  $\phi_{NL}(t, \mathbf{x})$  is a Lorentz scalar moving at the velocity  $\mathbf{v}_{DiracdB-B} = \Psi_{\mathbf{L}}(t, \mathbf{x})^{\dagger} \alpha \Psi_{\mathbf{L}}(t, \mathbf{x}) / \Psi_{\mathbf{L}}(t, \mathbf{x})^{\dagger} \Psi_{\mathbf{L}}(t, \mathbf{x})$ , in accordance with the dB-B guidance equation [23, 17], we find again by similar methods as those of previous sections the inhomogeneous Dirac equation

$$i\hbar\partial_t \Psi_{\mathbf{L}}(t,\mathbf{x}) - \alpha c \frac{\hbar}{i} \nabla \Psi_{\mathbf{L}}(t,\mathbf{x}) - mc^2 \beta \Psi_{\mathbf{L}}(t,\mathbf{x}) = -\frac{\hbar}{i} \alpha c (\Psi_{\mathbf{L}}(t,\mathbf{x}) - \Psi_{\mathbf{L}}(t,\mathbf{x_0})) \frac{\nabla \phi_{NLg}(t,\mathbf{x})}{\phi_{NLg}(t,\mathbf{x})}$$
(24)

Multiplying (24) by  $(i\hbar\partial_t + \alpha c\frac{\hbar}{i}\nabla + mc^2\beta)$  we get  $-\hbar^2 c^2 (\frac{1}{c^2}(\frac{\partial^2}{\partial t^2} - \Delta) + \frac{m^2 c^2}{\hbar^2})\Psi_{\mathbf{L}}(t,\mathbf{x}) = (i\hbar\partial_t + \alpha c\frac{\hbar}{i}\nabla + m\beta)\frac{(-)\hbar}{i}\alpha c(\Psi_{\mathbf{L}}(t,\mathbf{x}) - \Psi_{\mathbf{L}}(t,\mathbf{x_0}))\frac{\nabla\phi_{NLg}(t,\mathbf{x})}{\phi_{NLg}(t,\mathbf{x})}$ 

Applying the same reasonings as in the previous section, making use of (16), and  $\alpha_k^2 = 1$ , still in the limit of slowly moving bodies, we find (23). Together with

 $i\hbar\partial_t \frac{\Psi_{\mathbf{L}}(t,\mathbf{x})}{(1-\phi_G/c^2)} - \alpha c \frac{\hbar}{i} \bigtriangledown \frac{\Psi_{\mathbf{L}}(t,\mathbf{x})}{(1-\phi_G/c^2)} - mc^2 \beta \frac{\Psi_{\mathbf{L}}(t,\mathbf{x})}{(1-\phi_G/c^2)} = 0,$ 

and the Dirac-dB-B guidance relation  $\mathbf{v}_{DiracdB-B} = \Psi_{\mathbf{L}}(t, \mathbf{x})^{\dagger} \alpha \Psi_{\mathbf{L}}(t, \mathbf{x}) / \Psi_{\mathbf{L}}(t, \mathbf{x})^{\dagger} \Psi_{\mathbf{L}}(t, \mathbf{x})$  our model is self-consistent.

<sup>&</sup>lt;sup>3</sup>These two corrections are reminiscent of the standard coupling in electro-magnetism in which we replace  $i\hbar \frac{\partial}{\partial \mu}$  by  $i\hbar \frac{\partial}{\partial \mu} - eA_{\mu}$ . Here we replace  $i\hbar \frac{\partial}{\partial \mu}$  by  $i\hbar \frac{\partial}{\partial \mu} + i\frac{\hbar}{c^2} \frac{\partial \phi_G}{\partial \mu}$ . Formally, the gravitational potential looks like a purely imaginary gauge field. This is not so amazing having in mind the role played by rescalings in our approach.

## 4 Conclusions

Treating particles/droplets as incompressible solitons which obey dB-B dynamics<sup>4</sup>, the spread of which is compensated by a self focusing nonlinear interaction, we predicted the appearance of an effective gravitation. In the case of elementary particles, our model makes it possible, fitting the effective gravitation with Newtonian gravitation to estimate the extent of the particle (16). In the case of droplets, it delivers precise predictions about the pseudo-gravitational interaction. Our model is quite unorthodox. We found our main inspiration in old fashioned concepts such as the Poincaré pressure [20] or de Broglie's double solution program [10], and we did not make any kind of reference to curved space-time. This is maybe an open door for quantum gravity, maybe not. At this level, we simply do not know. In order to falsify our model it suffices in principle to test its post-Newtonian predictions, which is beyond the scope of our paper. According to us, alternative approaches to gravity could maybe help to understand apparent modifications of Newton's equation in the solar system [21] and in galactic rotation curves as well [14] but this is another story. In the meanwhile, our study suggests that dB-B trajectories and effective gravitation could be observed with collective self-collapsed assemblies of cold atoms [24], which has never been done so far. In any case, a virtue of our model is that it has some retrodictive power in the sense that it explains the appearance, in presence of dB-B trajectories, of a pseudo-gravitational interaction in droplets phenomenology, attractive and repulsive as well. Moreover we predict that the interaction scales like  $v^2(L_A M_B + L_B M_A) cos(k_0 r_{AB})/r_{AB}$ , which can possibly be tested in the lab.. There are two unspecified parameters in our model, at this level, which are the exact ratios between  $L_{A(B)}$  and the sizes of the droplets. We expect those to be of the order of unity however, and they can easily be evaluated experimentally.

## Acknowledgments

This work benefitted from the support of grants 21326 and 60230 from the John Templeton Foundation and the FQXI project FQXi-RFP-1506. Thanks to C. Champenois and R. Willox for kind comments.

## References

- E. Ruiz Arriola and J. Soler. A Variational Approach to the Schrödinger-Poisson System: Asymptotic Behaviour, Breathers, and Stability. J. Stat. Phys., 103(5-6):1069–1105, 2001.
- [2] G. Bacciagaluppi and A. Valentini. Quantum Theory at the Crossroads: Reconsidering the 1927 Solvay Conference. Cambridge University Press (2010), and quant-ph/0609184.

<sup>&</sup>lt;sup>4</sup>In [15] we explain how we were brought to impose (5,6,7) that in the present paper we treat as a postulate. Actually in [15] we studied the feedback of the pilot wave  $\Psi_L$  on the soliton or particle  $\phi_{NL}$ , postulating to begin with that (1,2) is satisfied and also that  $\Psi = \Psi_L \cdot \phi_{NL}$ , while  $\Psi_L$  is a solution of the homogeneous Schrödinger equation; we argued that this feedback results in conditions (5,6,7) in the appropriate regime. [15] is complementary to the present paper in the sense that here we study the feedback of the solitons on the pilot wave.

- [3] D. Bohm. A Suggested Interpretation of the Quantum Theory in Terms of "Hidden" Variables. I and II. Phys. Rev., 85(2):166–179, 180–193 (1952).
- [4] J.W.M. Busch. Pilot-Wave Hydrodynamics. Ann. Rev. Fluid Mech., 49:269-292 (2015).
- [5] S. Colin, T. Durt, and R. Willox. Can quantum systems succumb to their own (gravitational) attraction? *Class. Quantum Grav.*, 31:245003 (2014).
- [6] S. Colin, T. Durt, and R. Willox. Crucial tests of macrorealist and semi-classical gravity models with freely falling mesoscopic nanospheres. preprint (2014) arXiv:1402.5653v3.
- [7] Y. Couder, S. Protière, E. Fort and A. Boudaoud. Dynamical phenomena: walking and orbiting droplets. *Nature* 8; 437 (7056), 208 (2005).
- [8] Y. Couder and E. Fort. Single-Particle Diffraction and Interference at a Macroscopic Scale. *Phys. Rev. Lett.*, 97:15410 (2006).
- [9] Y. Couder, A. Boudaoud, S. Protière and E. Fort. Walking droplets, a form of wave-particle duality at macroscopic scale? *Europhysics News* Vol. 41, No. 1, 14-18 (2010).
- [10] L. de Broglie. Une tentative d'interprétation causale et non linéaire de la mécanique ondulatoire: la théorie de la double solution. Paris: Gauthier- Villars, 1956. English translation: Nonlinear wave mechanics: A causal interpretation. Elsevier, Amsterdam 1960.
- [11] L. de Broglie. Interpretation of quantum mechanics by the double solution theory Annales de la Fondation Louis de Broglie, 12, 4, 1987, english translation from a paper originally published in the book Foundations of Quantum Mechanics- Rendiconti della Scuola Internazionale di Fisica Enrico Fermi, IL Corso, B. d' Espagnat ed. Academic Press N.Y. 1972.
- [12] L. Diósi. Gravitation and quantum-mechanical localization of macroobjects. *Phys. Lett. A*, 105:199–202 (1984).
- [13] T. Durt. Characterisation of an entanglement-free evolution. Zeit. fur Naturf., 59 A:425, 2004. (2001).
- [14] T. Durt. Estimate of the weight of empty space based on astronomical observations quant/ph arXiv:1302.0835 (2013).
- [15] T. Durt. Generalized guidance equation for peaked quantum solitons: the single particle case. quant/ph arxiv 1602.03133 (2016).
- [16] A. Einstein, B. Podolsky, and N. Rosen: Can Quantum-Mechanical Description of Physical Reality be Considered Complete? *Phys. Rev.*, 47, 777 (1935).
- [17] P.R.. Holland. The Quantum Theory of Motion (Cambridge University Press, 1993)
- [18] E.H. Lieb. Existence and Uniqueness of the Minimizing Solution of Choquard's Nonlinear Equation. *Studies in Applied Mathematics*, 57:93–105 (1977).
- [19] R. Penrose. On Gravity's Role in Quantum State Reduction. General Relativity and Gravitation, 28(5):581–600, 1996.
- [20] H. Poincaré Sur la dynamique de l'électron Rendiconti del Circolo matematico di Palermo, 21,129-176 (1906).

- [21] S. Reynaud and M-T Jaekel. Gravity tests in the solar system and the pioneer anomaly. Mod. Phys. Letters A, 20, 14 (2005).
- [22] L. Smolin. Quantum fluctuations and inertia. Physics Letters A, 113:408, (1986).
- [23] T. Takabayasi Relativistic hydrodynamics of the Dirac matter. Suppl. Prog. Theor. Phys. 4, 1-80 (1957).
- [24] K.E. Strecker, G.B. Partridge, and R.G. Hulet. Formation and propagation of matter-wave soliton trains. *Nature*. 417 150 (2002).
- [25] J. R. van Meter. Schrödinger-Newton "collapse" of the wave-function. Class. Quant. Grav., 28:215013, 2011.