



Stationary measures and phase transition for a class of probabilistic cellular automata

Paolo Dai Pra, Pierre-Yves Louis, Sylvie Roelly

► To cite this version:

Paolo Dai Pra, Pierre-Yves Louis, Sylvie Roelly. Stationary measures and phase transition for a class of probabilistic cellular automata. ESAIM: Probability and Statistics, 2002, 6, pp.89-104. <10.1051/ps:2002004>. <hal-01277058v2>

HAL Id: hal-01277058

<https://hal.science/hal-01277058v2>

Submitted on 25 Apr 2016

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons CC BY-NC-ND 4.0 - Attribution - Non-commercial use - No Derivative Works - International License

STATIONARY MEASURES AND PHASE TRANSITION FOR A CLASS OF PROBABILISTIC CELLULAR AUTOMATA

PAOLO DAI PRA¹, PIERRE-YVES LOUIS² AND SYLVIE ROELLY^{3, 4}

Abstract. We discuss various properties of Probabilistic Cellular Automata, such as the structure of the set of stationary measures and multiplicity of stationary measures (or phase transition) for reversible models.

1991 Mathematics Subject Classification. 60G60 ; 60J10 ; 60K35 ; 82C20 ; 82B26.

Received December 4, 2001. Revised March 11, 2002.

1. INTRODUCTION

Probabilistic Cellular Automata (PCA) are discrete-time Markov chains on a product space S^Λ (*configuration space*) whose transition probability is a product measure. In this paper, S is assumed to be a finite set (*spin space*), and Λ (set of *sites*) a subset, finite or infinite, of \mathbb{Z}^d . The fact that the transition probability $P(d\sigma|\sigma')$, $\sigma, \sigma' \in S^\Lambda$, is a product measure means that all spins $\{\sigma_i : i \in \Lambda\}$ are simultaneously and independently updated (*parallel updating*). This transition mechanism differs from the one in the most common Gibbs samplers (e.g. [3, 8]), where only one site is updated at each time step (*sequential updating*).

Several properties of PCA's, mainly of general and qualitative nature, have been investigated ([5, 7, 14, 18, 23]). As far as we know, however, sharper properties like e.g. rate of convergence to equilibrium or use of parallel dynamics in perfect sampling, have not yet been investigated. PCA's are hard to analyze mainly for the following reason. Suppose Λ is a finite subset of \mathbb{Z}^d , and let μ be a given probability on S^Λ . To fix ideas, we may think of μ as a finite volume Gibbs measure for a given interaction and assigned boundary conditions. It is simple to construct Markov chains on S^Λ with sequential updating which have μ as reversible measure. Transition probabilities are given in simple form in terms of μ , and reversibility immediately implies that μ is an stationary measure for the dynamics. Quite differently, for a given μ , there is no general recipe to construct a PCA for which μ is stationary. In particular, there exists Gibbs measures on $S^{\mathbb{Z}^2}$ such that no PCA admits them as stationary measures (Theorem 4.2 in [5]).

Despite of this discouraging starting point, other aspects of PCA's make them interesting stochastic models, and motivate further investigation.

Keywords and phrases: Probabilistic Cellular Automata, stationary measure, Gibbs measure

¹ Dipartimento di Matematica Pura e Applicata, Università di Padova, Via Belzoni 7, 35131 Padova, Italy e-mail: daipra@math.unipd.it

² Laboratoire de Statistique et Probabilités, F.R.E. C.N.R.S. 2222, U.F.R. de Mathématiques, Université Lille 1, 59655 Villeneuve d'Ascq Cedex, France, e-mail: louis@lps.univ-lille1.fr

³ Weierstraß-Institut für Angewandte Analysis und Stochastik, Mohrenstraße 39, 10117 Berlin, Germany, e-mail: roelly@wias-berlin.de

⁴ Permanent address: CMAP, UMR CNRS 7641 École Polytechnique, 91128 Palaiseau Cedex France.

- (1) For simulation and sampling, PCA's are natural stochastic algorithms for parallel computing. At least in some simple models (see Section 3) it is interesting to evaluate their performance versus algorithms with sequential updating. This will be the subject of a forthcoming paper.
- (2) In opposition to dynamics with sequential updating, it is simple to define PCA's in infinite volume without passing to continuous time. One may try to study, for instance, convergence to equilibrium in infinite volume, or in finite volume uniformly in the volume size. Although some perturbative methods are available (see [20] Chapter 7, [17, 18]), a theory corresponding the one in ([21]) in continuous time, is yet to be developed.
- (3) PCA's that are reversible with respect to a Gibbs measure μ have been completely characterized in [11]. In particular it has been shown that only a small class of Gibbs measures may be reversible for a PCA. For such PCA's one can investigate metastable behavior. A first step in this direction is done in [2].

The present paper is a small step toward a better understanding of PCA's. Our objective is first to present some links between the sets of reversible, resp. stationary, resp. Gibbs measures for general PCA's. We then illustrate these results on a particular class of reversible PCA's already introduced in [2].

More precisely it was proved in [11] that for PCA's possessing a reversible Gibbs measure w.r.t. a potential Φ , all reversible measures are gibbsian w.r.t the same potential. We prove a similar statement on the set of stationary measures : For a general PCA, if one shift invariant stationary measure is Gibbsian for a potential Φ , then all shift invariant stationary measures are Gibbsian w.r.t. the same potential Φ (see Proposition 2.2). This induce that for a class of local, shift invariant, non-degenerated, reversible PCA the reversible measures coincide with the Gibbsian stationary ones (Remark 3.1).

Applying this general statements to the class of PCA's considered in [2], one can do explicit a stationary measure which is in fact Gibbsian w.r.t. a certain potential Φ we write down (cf Proposition 3.2); we show that, for sufficiently small values of the temperature parameter, *phase transition* occurs, that is there are several Gibbs measures w.r.t. Φ . At least in certain cases, existence of phase transition would follow from general expansion arguments, like Pirogov-Sinai theory. We have preferred here, however, to use "softer" contour arguments. The understanding of the right notion of contour for a specific model is in any case useful in many respects (percolation, block dynamics, ...).

However, unlike what happens with sequential updating, not all these Gibbs measures need to be stationary for the infinite volume PCA, the non-stationary ones being periodic with period two. To conclude, we exhibit a Gibbs measure which is not stationary for the associated PCA.

2. SHIFT INVARIANT PROBABILISTIC CELLULAR AUTOMATA

Let S be a finite set. For $\sigma \in S^{\mathbb{Z}^d}$, $\sigma = (\sigma_i)_{i \in \mathbb{Z}^d}$, and $\Lambda \subset \mathbb{Z}^d$, we let $\sigma_\Lambda \in S^\Lambda$ its restriction to Λ . Sometimes, when no confusion arises, we omit the index Λ in σ_Λ .

A time-homogeneous Markov chain on S^Λ is determined, in law, by its transition probabilities $P_\Lambda(d\sigma|\eta)$. If $P_\Lambda(d\sigma|\eta)$ is a product measure, as a probability measure on S^Λ , then we say that the Markov chain is a Probabilistic Cellular Automaton. More explicitly

$$P_\Lambda(d\sigma|\eta) = \otimes_{i \in \Lambda} P_i(d\sigma_i|\eta),$$

and

$$P_i(\sigma_i = s|\eta) \equiv p_i(s|\eta), \quad s \in S. \tag{1}$$

In the case $\Lambda = \mathbb{Z}^d$, we omit the index Λ in $P_\Lambda(d\sigma|\eta)$. In this case, we say that a PCA is *shift invariant* if, for every $i \in \mathbb{Z}^d$, $s \in S$, $\eta \in S^{\mathbb{Z}^d}$, we have

$$p_i(s|\eta) = p_0(s|\theta_i\eta),$$

where θ_i is the shift in \mathbb{Z}^d : $(\theta_i\eta)_j = \eta_{i+j}$ for every $j \in \mathbb{Z}^d$. A shift invariant PCA is said to be *local* if, for each $s \in S$, the map $\eta \rightarrow p_0(s|\eta)$ is local, i.e. it depends on a finite number of components of η .

From now on, all PCA's we consider in this paper satisfy the non degeneration condition :

$$p_0(s|\eta) > 0, \quad \forall s \in S, \eta \in S^{\mathbb{Z}^d}.$$

This means that we are dealing with dynamics which can not contain a deterministic component.

In this paper we are mostly interested in stationary measures for PCA's. For this purpose we recall the notion of Gibbs measure on $S^{\mathbb{Z}^d}$. A shift invariant potential Φ is a family $\{\Phi_\Lambda : \Lambda \subset \mathbb{Z}^d, |\Lambda| < +\infty\}$ of maps $\Phi_\Lambda : S^\Lambda \rightarrow \mathbb{R}$ with the properties

i. For all $i \in \mathbb{Z}^d$, $\Lambda \subset \mathbb{Z}^d$ finite:

$$\Phi_{\Lambda+i} = \Phi_\Lambda \circ \theta_i.$$

ii.

$$\sum_{\Lambda \ni 0} \|\Phi_\Lambda\|_\infty < +\infty.$$

Here and later $|\Lambda|$ denotes the cardinality of Λ . Letting $H_\Lambda(\sigma) = \sum_{A \cap \Lambda \neq \emptyset} \Phi_A(\sigma)$ and choosing $\tau \in S^{\mathbb{Z}^d}$, also write $H_\Lambda^\tau(\sigma_\Lambda) = H_\Lambda(\sigma_\Lambda \tau_{\Lambda^c})$, where $\sigma_\Lambda \tau_{\Lambda^c}$ is the element of $S^{\mathbb{Z}^d}$ which coincides with σ on Λ and with τ on Λ^c . The finite volume Gibbs measure on S^Λ with boundary condition τ is given by

$$\mu_\Lambda^\tau(\sigma_\Lambda) = \frac{\exp[-H_\Lambda^\tau(\sigma_\Lambda)]}{Z_\Lambda^\tau},$$

where Z_Λ^τ is the normalization factor. A probability measure μ on $S^{\mathbb{Z}^d}$ is said to be Gibbsian for the potential Φ , and we write $\mu \in \mathcal{G}(\Phi)$ if for every $\Lambda \subset \mathbb{Z}^d$ finite and $\sigma \in S^{\mathbb{Z}^d}$

$$\mu(\{\eta : \eta_\Lambda = \sigma_\Lambda\} | \eta_{\Lambda^c} = \tau_{\Lambda^c}) = \mu_\Lambda^\tau(\sigma_\Lambda)$$

for μ -a.e. τ . If μ is *shift-invariant*, i.e. $\mu \circ \theta_i = \mu$ for all $i \in \mathbb{Z}^d$, then we write $\mu \in \mathcal{G}_s(\Phi)$. More generally, we let \mathcal{P} (resp. \mathcal{P}_s) be the set of probability measures (resp. shift-invariant probability measures) on $S^{\mathbb{Z}^d}$.

Given $\Lambda \subset \mathbb{Z}^d$, we denote by \mathcal{F}_Λ the σ -field on $S^{\mathbb{Z}^d}$ generated by the projection $\sigma \rightarrow \sigma_\Lambda$. For $\nu \in \mathcal{P}$, $\pi_\Lambda \nu$ is the restriction of ν to \mathcal{F}_Λ . We will use, for $\nu, \mu \in \mathcal{P}$, the notion of local relative entropy:

$$h_\Lambda(\nu|\mu) = \sum_{\sigma_\Lambda} \pi_\Lambda \nu(\sigma_\Lambda) \log \frac{\pi_\Lambda \nu(\sigma_\Lambda)}{\pi_\Lambda \mu(\sigma_\Lambda)} \quad (2)$$

with $\Lambda \subset \mathbb{Z}^d$ finite, and of specific relative entropy

$$h(\nu|\mu) = \limsup_{\Lambda \uparrow \mathbb{Z}^d} \frac{1}{|\Lambda|} h_\Lambda(\nu|\mu) \quad (3)$$

where in the limit above Λ varies over hypercubes centered in the origin. It is easily seen that $0 \leq h(\nu|\mu) \leq +\infty$. In the case of $\mu \in \mathcal{G}_s(\Phi)$ for a potential Φ , in (2) $\pi_\Lambda \mu(\sigma_\Lambda)$ can be replaced by $\mu_\Lambda^\tau(\sigma_\Lambda)$, for an arbitrary τ , without changing the limit in (3). Moreover, for $\mu \in \mathcal{G}_s(\Phi)$ and $\nu \in \mathcal{P}_s$, the limsup in (3) is actually a limit. In this case the Gibbs variational principle states that, for $\nu \in \mathcal{P}_s$, $h(\nu|\mu) = 0$ if and only if $\nu \in \mathcal{G}_s(\Phi)$; so $h(\nu|\mu)$ represents a notion of (pseudo-) distance of ν from $\mathcal{G}_s(\Phi)$.

We now define a corresponding notion of specific relative entropy for transition probabilities, that will be used to measure distance between two dynamics. Let $P(d\sigma|\eta)$ and $Q(d\sigma|\eta)$ two transition probabilities on $S^{\mathbb{Z}^d}$, and $\nu \in \mathcal{P}$. We define

$$\mathcal{H}_\nu(P|Q) = \limsup_{\Lambda \uparrow \mathbb{Z}^d} \frac{1}{|\Lambda|} \int h_\Lambda(P(\cdot|\eta)|Q(\cdot|\eta)) \nu(d\eta).$$

Clearly $\mathcal{H}_\nu(P|Q) \geq 0$. By conditioning to σ the joint law $Q_\nu(d\sigma, d\eta) \equiv P(d\sigma|\eta)\nu(d\eta)$ we obtain the backward transition probability, that we denote by $\hat{P}_\nu(d\eta|\sigma)$. We also let $P\nu(d\sigma)$ be given by

$$P\nu(A) = \int P(A|\eta)\nu(d\eta)$$

for $A \subset S^{\mathbb{Z}^d}$ measurable. If $P\nu = \nu$ we say that ν is stationary for $P(d\sigma|\eta)$.

Our first result concerns the entropy production for a PCA (cf. [4]). The corresponding result in continuous time has appeared in [9].

Proposition 2.1. *Suppose μ is a stationary measure for a shift invariant, local PCA with transition probability $P(d\sigma|\eta)$. If μ is also a shift invariant Gibbs measure w.r.t. a certain potential Φ ($\mu \in \mathcal{G}_s(\Phi)$), then, for any shift invariant measure ν ,*

$$h(\nu|\mu) - h(P\nu|\mu) = \mathcal{H}_\nu(\hat{P}_\nu|\hat{P}_\mu).$$

In particular, if $\nu \in \mathcal{G}_s(\Phi)$, then $P\nu \in \mathcal{G}_s(\Phi)$, that is the set of shift-invariant Gibbs measures w.r.t. the potential Φ is stable under the action of this PCA dynamics.

Proof. Let Λ be a finite subset of \mathbb{Z}^d , and consider

$$P_\Lambda(\sigma|\eta) = \prod_{i \in \Lambda} p_i(\sigma_i|\eta).$$

This expression depends on the restriction of η to a neighborhood of Λ , that we denote by $\bar{\Lambda}$.

Consider now the measure $Q_\nu(d\sigma, d\eta)$ defined above. For $A, B \subset \mathbb{Z}^d$ with A finite, we denote by $Q_\nu(\sigma_A|\eta_B)$ the restriction to the σ -field generated by the projection $(\sigma, \eta) \rightarrow \sigma_A$ of the measure Q conditioned to the σ -field generated by the projection $(\sigma, \eta) \rightarrow \eta_B$. So, e.g., $P_\Lambda(\sigma|\eta) = Q_\nu(\sigma_\Lambda|\eta_{\mathbb{Z}^d}) \equiv Q_\nu(\sigma_\Lambda|\eta)$, independently of ν . Similarly, $\hat{Q}_\nu(\eta_A|\sigma_B)$ denotes the time-reversed conditioning, so that

$$\pi_\Lambda \hat{P}_\nu(\eta_\Lambda|\sigma) = \hat{Q}_\nu(\eta_\Lambda|\sigma). \quad (4)$$

For $C \subset \mathbb{Z}^d$ we will also use conditionings of the form

$$\hat{Q}(\eta_A|\sigma_B, \eta_C),$$

with the obvious meaning.

A simple computation, using the fact that $P\mu = \mu$, yields

$$\begin{aligned} h_{\bar{\Lambda}}(\nu|\mu) - h_\Lambda(P\nu|\mu) &= \\ &= \sum_{\sigma_\Lambda} \pi_\Lambda(P\nu)(\sigma_\Lambda) \sum_{\eta_{\bar{\Lambda}}} \hat{Q}_\nu(\eta_{\bar{\Lambda}}|\sigma_\Lambda) \log \frac{\hat{Q}_\nu(\eta_{\bar{\Lambda}}|\sigma_\Lambda)}{\hat{Q}_\mu(\eta_{\bar{\Lambda}}|\sigma_\Lambda)} \\ &= E^Q \left[\log \frac{\hat{Q}_\nu(\eta_{\bar{\Lambda}}|\sigma_\Lambda)}{\hat{Q}_\mu(\eta_{\bar{\Lambda}}|\sigma_\Lambda)} \right]. \end{aligned}$$

Since

$$h(\nu|\mu) - h(P\nu|\mu) = \lim_{\Lambda \uparrow \mathbb{Z}^d} \frac{1}{|\Lambda|} [h_{\bar{\Lambda}}(\nu|\mu) - h_\Lambda(P\nu|\mu)],$$

then the conclusion follows provided we show (see (4))

$$\lim_{\Lambda \uparrow \mathbb{Z}^d} \frac{1}{|\Lambda|} E^Q \left[\log \frac{\hat{Q}_\nu(\eta_{\bar{\Lambda}}|\sigma_\Lambda)}{\hat{Q}_\nu(\eta_{\bar{\Lambda}}|\sigma)} \right] = 0 \quad (5)$$

and

$$\lim_{\Lambda \uparrow \mathbb{Z}^d} \frac{1}{|\Lambda|} E^Q \left[\log \frac{\hat{Q}_\mu(\eta_\Lambda | \sigma_\Lambda)}{\hat{Q}_\mu(\eta_\Lambda | \sigma)} \right] = 0. \quad (6)$$

Note that (6) is a special case of (5).

Let now λ^\otimes be the probability measure on $S^{\mathbb{Z}^d}$ obtained by taking the infinite product of the uniform measure λ in S . We denote by $\lambda^\otimes(\sigma_\Lambda)$ the projection of λ^\otimes on \mathcal{F}_Λ . Let also $\{i_1, \dots, i_{|\bar{\Lambda}|}\}$ be the lexicographic ordering of the elements of $\bar{\Lambda}$; define $\bar{\Lambda}_k = \{i_1, \dots, i_k\}$ for $1 \leq k \leq |\bar{\Lambda}|$, and $\bar{\Lambda}_0 = \emptyset$. By the chain rule for conditional measures

$$\log \frac{\hat{Q}(\eta_{\bar{\Lambda}} | \sigma_\Lambda)}{\lambda^\otimes(\eta_{\bar{\Lambda}})} = \sum_{k=1}^{|\bar{\Lambda}|} \log \frac{\hat{Q}(\eta_{i_k} | \sigma_\Lambda, \eta_{\bar{\Lambda}_{k-1}})}{\lambda(\eta_{i_k})}. \quad (7)$$

Moreover, by shift invariance of Q

$$E^Q \left[\log \frac{\hat{Q}(\eta_{i_k} | \sigma_\Lambda, \eta_{\bar{\Lambda}_{k-1}})}{\lambda(\eta_{i_k})} \right] = E^Q \left[\log \frac{\hat{Q}(\eta_0 | \sigma_{\theta_{-i_k}\Lambda}, \eta_{\theta_{-i_k}\bar{\Lambda}_{k-1}})}{\lambda(\eta_0)} \right]. \quad (8)$$

Let $\mathbb{Z}_-^d = \{i \in \mathbb{Z}^d : i \prec 0\}$, where “ \prec ” is the lexicographic order. By the Shannon-Breiman-McMillan Theorem ([1]), for every $\epsilon > 0$ there are $A \subset \mathbb{Z}^d$, $B \subset \mathbb{Z}_-^d$ finite such that if $A \subset V$ and $B \subset W \subset \mathbb{Z}_-^d$ then

$$\left| E^Q \left[\log \frac{\hat{Q}(\eta_0 | \sigma_V, \eta_W)}{\lambda(\eta_0)} \right] - E^Q \left[\log \frac{\hat{Q}(\eta_0 | \sigma_A, \eta_B)}{\lambda(\eta_0)} \right] \right| < \epsilon. \quad (9)$$

Note that, if we take Λ large enough and $i_k \in \Lambda$ is far enough from the boundary of Λ , then $A \subset \theta_{-i_k}\Lambda$, and $B \subset \theta_{-i_k}\bar{\Lambda}_{k-1}$. For the other values of $i_k \in \bar{\Lambda}$,

$$E^Q \left[\log \frac{\hat{Q}(\eta_0 | \sigma_{\theta_{-i_k}\Lambda}, \eta_{\theta_{-i_k}\bar{\Lambda}_{k-1}})}{\lambda(\eta_0)} \right] \leq \log |S|,$$

which is the upper bound for the entropy of any probability measure is S with respect to λ . Summing all up

$$\lim_{\Lambda \uparrow \mathbb{Z}^d} \frac{1}{|\Lambda|} E^Q \left[\log \frac{\hat{Q}(\eta_{\bar{\Lambda}} | \sigma_\Lambda)}{\lambda^\otimes(\eta_{\bar{\Lambda}})} \right] = E^Q \left[\log \frac{\hat{Q}(\eta_0 | \sigma_{\mathbb{Z}^d}, \eta_{\mathbb{Z}_-^d})}{\lambda(\eta_0)} \right]. \quad (10)$$

Exactly in the same way one shows that

$$\lim_{\Lambda \uparrow \mathbb{Z}^d} \frac{1}{|\Lambda|} E^Q \left[\log \frac{\hat{Q}(\eta_{\bar{\Lambda}} | \sigma)}{\lambda^\otimes(\eta_{\bar{\Lambda}})} \right] = E^Q \left[\log \frac{\hat{Q}(\eta_0 | \sigma_{\mathbb{Z}^d}, \eta_{\mathbb{Z}_-^d})}{\lambda(\eta_0)} \right]. \quad (11)$$

Thus (10) and (11) establish (5).

Next result shows that the measures in \mathcal{P}_s for which the entropy production is zero are exactly those in $\mathcal{G}_s(\Phi)$. This result goes back to [10], where it has been proved for reversible systems in continuous time. The assumption of reversibility has been dropped in [12]. In discrete-time, the proof for a special class of reversible PCA is given in [11], Proposition 1. In the generality given here, the first proof was contained (but unpublished) in one of the authors' PhD Thesis ([4]). Later, a proof using general entropy arguments was given in [19]. In this paper we have preferred to emphasize the fact that the following result comes from the precise entropy production formula presented in Proposition 2.1.

Proposition 2.2. *Under the same assumptions of Proposition 2.1, suppose $\nu \in \mathcal{P}_s$ is such that*

$$h(\nu|\mu) = h(P\nu|\mu) \quad (12)$$

(in particular, this happens when ν is stationary). Then $\nu \in \mathcal{G}_s(\Phi)$.

Proof. By what seen in Proposition 2.1, (12) amounts to

$$\mathcal{H}_\nu(\hat{P}_\nu|\hat{P}_\mu) = 0. \quad (13)$$

We now adapt a classical argument for Gibbs measures (see e.g. [22], Th. 7.4). Let V be a fixed hypercube and, for $k > 0$,

$$\partial_k V = \{i \in V^c : \text{dist}(i, V) \leq k\},$$

where $\text{dist}(\cdot)$ is the Euclidean distance. Take, now, a hypercube $\Lambda_{m,k}$ that is obtained as disjoint union of m^d translates of $V \cup \partial_k V$, say

$$\Lambda_{m,k} = \cup_{i=1}^{m^d} W_{i,k},$$

where $W_{i,k} = T_i(V \cup \partial_k V)$, and T_i is a suitable translation. We also write $V_i = T_i V$. Defining, for $i \in \{1, \dots, m^d\}$

$$B_{i,k} = W_{i,k} \setminus V_i$$

we have (we use the notations introduced in the proof of Proposition 2.1)

$$\log \frac{\hat{Q}_\nu(\eta_{\Lambda_{m,k}}|\sigma)}{\hat{Q}_\mu(\eta_{\Lambda_{m,k}}|\sigma)} = \sum_{i=1}^{m^d} \log \frac{\hat{Q}_\nu(\eta_{V_i}|\eta_{B_{i,k}}, \sigma)}{\hat{Q}_\mu(\eta_{V_i}|\eta_{B_{i,k}}, \sigma)} + \log \frac{\hat{Q}_\nu(\eta_{B_{1,k}}|\sigma)}{\hat{Q}_\mu(\eta_{B_{1,k}}|\sigma)}.$$

By positivity of relative entropy:

$$E^Q \left[\log \frac{\hat{Q}_\nu(\eta_{B_{1,k}}|\sigma)}{\hat{Q}_\mu(\eta_{B_{1,k}}|\sigma)} \right] \geq 0$$

so that

$$E^Q \left[\log \frac{\hat{Q}_\nu(\eta_{\Lambda_{m,k}}|\sigma)}{\hat{Q}_\mu(\eta_{\Lambda_{m,k}}|\sigma)} \right] \geq \sum_{i=1}^{m^d} E^Q \left[\log \frac{\hat{Q}_\nu(\eta_{V_i}|\eta_{B_{i,k}}, \sigma)}{\hat{Q}_\mu(\eta_{V_i}|\eta_{B_{i,k}}, \sigma)} \right]. \quad (14)$$

By translation invariance of Q :

$$E^Q \left[\log \frac{\hat{Q}_\nu(\eta_{V_i}|\eta_{B_{i,k}}, \sigma)}{\hat{Q}_\mu(\eta_{V_i}|\eta_{B_{i,k}}, \sigma)} \right] = E^Q \left[\log \frac{\hat{Q}_\nu(\eta_V|\eta_{T_i^{-1}B_{i,k}}, \sigma)}{\hat{Q}_\mu(\eta_V|\eta_{T_i^{-1}B_{i,k}}, \sigma)} \right]. \quad (15)$$

Moreover, since $B_{i,k} \uparrow V_i^c$ as $k \uparrow +\infty$, using again the Shannon-Breiman-McMillan Theorem, for each $\epsilon > 0$ we can choose k large enough so that

$$\left| E^Q \left[\log \frac{\hat{Q}_\nu(\eta_V|\eta_{T_i^{-1}B_{i,k}}, \sigma)}{\hat{Q}_\mu(\eta_V|\eta_{T_i^{-1}B_{i,k}}, \sigma)} \right] - E^Q \left[\log \frac{\hat{Q}_\nu(\eta_V|\eta_{V^c}, \sigma)}{\hat{Q}_\mu(\eta_V|\eta_{V^c}, \sigma)} \right] \right| \leq \epsilon. \quad (16)$$

Summing up (14), (15) and (16), we get

$$\frac{1}{m^d} E^Q \left[\log \frac{\hat{Q}_\nu(\eta_{\Lambda_{m,k}}|\sigma)}{\hat{Q}_\mu(\eta_{\Lambda_{m,k}}|\sigma)} \right] \geq E^Q \left[\log \frac{\hat{Q}_\nu(\eta_V|\eta_{V^c}, \sigma)}{\hat{Q}_\mu(\eta_V|\eta_{V^c}, \sigma)} \right] - \epsilon. \quad (17)$$

But m^d is proportional to $|\Lambda_{m,k}|$, so, by (13)

$$\lim_{m \rightarrow +\infty} \frac{1}{m^d} E^Q \left[\log \frac{\hat{Q}_\nu(\eta_{\Lambda_{m,k}}|\sigma)}{\hat{Q}_\mu(\eta_{\Lambda_{m,k}}|\sigma)} \right] = 0.$$

Thus, since ϵ is arbitrary, (17) yields

$$E^Q \left[\log \frac{\hat{Q}_\nu(\eta_V|\eta_{V^c}, \sigma)}{\hat{Q}_\mu(\eta_V|\eta_{V^c}, \sigma)} \right] = 0$$

that, by elementary properties of relative entropy, implies

$$\hat{Q}_\nu(\eta_V|\eta_{V^c}, \sigma) = \hat{Q}_\mu(\eta_V|\eta_{V^c}, \sigma) \quad Q - \text{a.s.} \quad (18)$$

At this point we use Proposition 3.2 in [13], which implies that if (18) holds for a $\mu \in \mathcal{G}_s(\Phi)$, then $\nu(\eta_V|\eta_{V^c}) = \mu(\eta_V|\eta_{V^c})$ a.s. and then $\nu \in \mathcal{G}_s(\Phi)$ too. This completes the proof.

3. A CLASS OF REVERSIBLE DYNAMICS

In this section we introduce a class of reversible PCA's we will be dealing with in the rest of the paper, and give some general results on their stationary measures, resp. reversible measures. Let us remember that a PCA P is called reversible if there exists at least one probability measure μ such that the Markov process with initial law μ and dynamics P is reversible.

We choose $S = \{-1, 1\}$ as spin space. Consider a function $k : \mathbb{Z}^d \rightarrow \mathbb{R}$ that is of finite range, i.e. there exists $R > 0$ such that $k(i) = 0$ for $|i| > R$, and symmetric, i.e. $k(i) = k(-i)$ for every $i \in \mathbb{Z}^d$ (this last assumption being necessary to assure the reversibility of the PCA, cf [11]). Moreover, let $\tau \in \{-1, 1\}^{\mathbb{Z}^d}$ be a fixed configuration, that will play the role of boundary condition. For $\Lambda \subset \mathbb{Z}^d$, we define the transition probability $P_\Lambda^\tau(d\sigma|\eta) = \otimes_{i \in \Lambda} P_i^\tau(d\sigma_i|\eta)$ by

$$P_i^\tau(\sigma_i = s|\eta) = p_i(s|\tilde{\eta}) = \frac{1}{2} \left[1 + s \tanh(\beta \sum_{j \in \mathbb{Z}^d} k(i-j)\tilde{\eta}_j + \beta h), \right] \quad (19)$$

where $\tilde{\eta} = \eta_\Lambda \tau_{\Lambda^c}$; $h \in \mathbb{R}$, $\beta > 0$ are given parameters. According to [11], this particular form of p_i is indeed the most general one for a shift invariant non degenerate local PCA on $\{-1, 1\}^{\mathbb{Z}^d}$.

In the case Λ is a hypercube, we can also consider periodic boundary conditions. The associated transition probability is denoted by P_Λ^{per} . In general, when Λ is finite, we write $P_\Lambda^\tau(\sigma|\eta)$ in place of $P_\Lambda^\tau(\{\sigma\}|\eta)$. In the case $\Lambda = \mathbb{Z}^d$, the boundary condition τ plays no role, and will be omitted.

In the rest of this section we establish some simple facts about stationary measures for these PCA's.

Proposition 3.1. *Let $\Lambda \subset \mathbb{Z}^d$ finite, and $\tau \in \{-1, 1\}^{\mathbb{Z}^d}$. Then the finite volume PCA with transition probability $P_\Lambda^\tau(\sigma|\eta)$ has a unique stationary measure ν_Λ^τ given by*

$$\nu_\Lambda^\tau(\sigma) = \frac{1}{W_\Lambda^\tau} \prod_{i \in \Lambda} e^{\beta h \sigma_i} \cosh \left[\beta \sum_{j \in \mathbb{Z}^d} k(i-j)\tilde{\sigma}_j + \beta h \right] e^{\beta \sigma_i \sum_{j \in \Lambda^c} k(i-j)\tau_j},$$

where, as before, $\tilde{\sigma} = \sigma_\Lambda \tau_{\Lambda^c}$, and W_Λ^τ is the normalization. Moreover, ν_Λ^τ is reversible for P_Λ^τ .

Proof. It is clear that $P_\Lambda^\tau(\sigma|\eta) > 0 \forall \sigma, \eta$, so that the Markov chain with transition probability P_Λ^τ has a unique stationary measure. Thus, we only have to show that ν_Λ^τ is reversible, i.e.

$$P_\Lambda^\tau(\sigma|\eta)\nu_\Lambda^\tau(\eta) \equiv P_\Lambda^\tau(\eta|\sigma)\nu_\Lambda^\tau(\sigma). \quad (20)$$

Observe that, since $\sigma_i \in \{-1, 1\}$, P_Λ^τ may be written in the form

$$P_\Lambda^\tau(\sigma|\eta) = \prod_{i \in \Lambda} \frac{e^{\beta \sigma_i (\sum_j k(i-j) \tilde{\eta}_j + h)}}{2 \cosh \left(\beta \sum_j k(i-j) \tilde{\eta}_j + \beta h \right)}.$$

Thus (20) amounts to

$$\sum_{i \in \Lambda} \sum_{j \in \mathbb{Z}^d} \sigma_i \tilde{\eta}_j k(i-j) + \sum_{i \in \Lambda} \sum_{j \notin \Lambda} \eta_i \tau_j k(i-j) = \sum_{i \in \Lambda} \sum_{j \in \mathbb{Z}^d} \eta_i \tilde{\sigma}_j k(i-j) + \sum_{i \in \Lambda} \sum_{j \notin \Lambda} \sigma_i \tau_j k(i-j)$$

which is easily checked.

The above result on stationary measures for PCA's in finite volume, has an immediate consequence in infinite volume.

Proposition 3.2. *Let τ be any fixed boundary condition, and μ be any limit point of ν_Λ^τ as $\Lambda \uparrow \mathbb{Z}^d$. Then μ is reversible for the infinite volume PCA defined in (19), and μ is Gibbsian for the shift-invariant potential Φ given by*

$$\begin{aligned} \Phi_{\{i\}}(\sigma_i) &= -\beta h \sigma_i \\ \Phi_{U_i}(\sigma_{U_i}) &= -\log \cosh \left[\beta \sum_j k(i-j) \sigma_j + \beta h \right] \\ \Phi_\Lambda(\sigma_\Lambda) &= 0 \text{ otherwise,} \end{aligned} \quad (21)$$

where $U_i = \{j : k(i-j) \neq 0\}$, that is finite by assumption.

Proof. Note that the finite volume Gibbs measure for Φ is

$$\mu_\Lambda^\tau(\sigma) = \frac{1}{Z_\Lambda^\tau} \prod_{i: \text{dist}(i, \Lambda) \leq R} \cosh \left[\beta \sum_j k(i-j) \tilde{\sigma}_j + \beta h \right] e^{\beta h \sigma_i},$$

that differs from ν_Λ^τ only for boundary terms (and for the renormalization constant). The fact that the limit of ν_Λ^τ is Gibbsian for Φ follows therefore from general facts on Gibbs measures ([6]). The reversibility of μ for the infinite volume PCA is obtained as follows. Let $f : \{-1, 1\}^{\mathbb{Z}^d} \times \{-1, 1\}^{\mathbb{Z}^d} \rightarrow \mathbb{R}$ be a function which is local in both variables. For Λ large enough, reversibility of ν_Λ^τ yields

$$\sum_{\sigma, \tau} P_\Lambda^\tau(\sigma|\eta) \nu_\Lambda^\tau(\eta) f(\sigma, \eta) = \sum_{\sigma, \tau} P_\Lambda^\tau(\eta|\sigma) \nu_\Lambda^\tau(\sigma) f(\sigma, \eta). \quad (22)$$

Note that, for Λ large enough, the boundary condition τ in P_Λ^τ does not play any role in (22). Thus, letting $\Lambda \uparrow \mathbb{Z}^d$ in (22) obtaining

$$\int P(d\sigma|\eta) \mu(d\eta) f(\sigma, \eta) = \int P(d\eta|\sigma) \mu(d\sigma) f(\sigma, \eta), \quad (23)$$

that establishes reversibility of μ .

Instead of fixed boundary conditions, one can choose periodic boundary conditions. In this case, the finite volume measure defined by

$$\nu_{\Lambda}^{per}(\sigma) = \frac{1}{W_{\Lambda}^{per}} \prod_{i \in \Lambda} \cosh \left[\beta \sum_{j \in \mathbb{Z}^d} k(i-j) \tilde{\sigma}_j + \beta h \right] e^{\beta h \sigma_i}$$

where $\tilde{\sigma}$ is the periodic continuation of σ , is the unique stationary reversible measure for P_{Λ}^{per} . Remark that, in opposition to fixed boundary conditions, we now have that $\nu_{\Lambda}^{per} = \mu_{\Lambda}^{per}$, which means that the finite volume stationary measure for the finite volume PCA is equal to the local specification of the associated Gibbs measure.

Moreover, the following result gives a complete description of the links between the set of reversible measures for the PCA P (which will be denoted by \mathcal{R}), the set of stationary ones denoted by \mathcal{S} , the set $\mathcal{G}(\Phi)$ of Gibbs measures with respect to the potential Φ defined by (21), and their respective intersections with the set of shift-invariant measures : $\mathcal{R}_s, \mathcal{S}_s, \mathcal{G}_s(\Phi)$.

Proposition 3.3. *The reversible measures for the PCA P defined in (19) are exactly those Gibbs measures w.r.t. Φ given in (21) which are also stationary :*

$$\mathcal{R} = \mathcal{S} \cap \mathcal{G}(\Phi). \quad (24)$$

Moreover, the subset of shift invariant reversible measures is equal to the set of shift invariant stationary measures :

$$\mathcal{R}_s = \mathcal{S}_s. \quad (25)$$

Proof. The proof of the first assertion is based on the following proposition proved in [11] :

Let P be a non degenerate local reversible PCA. Each reversible measure μ for P is Gibbs w.r.t. a certain potential Φ_P . Reciprocally, any Gibbs measure w.r.t. Φ_P is either a reversible measure for P or periodic of period two.

Since obviously $\mathcal{R} \subset \mathcal{S}$, the abovementioned proposition implies $\mathcal{R} \subset \mathcal{S} \cap \mathcal{G}(\Phi)$. For the reciprocal inclusion, since stationary measures can not be 2-periodic, a stationary Gibbsian measure is necessarily a reversible one.

To prove the second assertion, note that by Proposition 3.2 and Remark 3, $\mathcal{S}_s \cap \mathcal{G}_s(\Phi) \ni \mu^{per}$. Thus Proposition 2.2 applies, that is : $\mathcal{S}_s \subset \mathcal{G}_s(\Phi)$. On the other hand, from the first assertion: $\mathcal{R}_s = \mathcal{S}_s \cap \mathcal{G}_s(\Phi)$. Then $\mathcal{R}_s = \mathcal{S}_s$.

Remark 3.1. The proof of Proposition 3.3 doesn't use the specific form of the PCA P . So equalities (24) and (25) hold as soon as Proposition 2.2 and the abovementioned result of [11] apply, that is for the general class of local, shift invariant, non degenerate reversible PCA dynamics on $S^{\mathbb{Z}^d}$ for any S finite.

4. PHASE TRANSITION

In this section we show that for some reversible PCA it is indeed the case that not all Gibbs measures for the potential in (21) are stationary. We treat those PCA defined in (19) for which $k(i) = 0$ for $|i| > 1$ (id est $R=1$), $h = 0$ and $d = 2$. Besides β , there are three parameters in the game: $k(0), k(e_1)$ and $k(e_2)$, where e_1, e_2 are the basis vectors in \mathbb{R}^2 . The first result concerns the existence of phase transition for the potential Φ .

Proposition 4.1. *Assume $k(e_1) \neq 0, k(e_2) \neq 0$. Then there exists $\beta_c \in (0, +\infty)$ such that for $\beta > \beta_c$ $|\mathcal{G}(\Phi)| > 1$.*

Proof. We divide the proof into different cases, depending on the signes of $k(0), k(e_1), k(e_2)$. Note that the transformation $k(\cdot) \rightarrow -k(\cdot)$ leaves invariant the potential Φ .

Case 1: $k(0) \geq 0, k(e_1) > 0, k(e_2) > 0$.

For a given square $\Lambda \subset \mathbb{Z}^2$, let $Cl_m(\Lambda) = \{i \in \mathbb{Z}^d : \text{dist}(i, \Lambda) \leq m\}$. Consider a fixed configuration $\sigma \in \{-1, +1\}^{\mathbb{Z}^2}$ such that $\sigma_i \equiv +1$ for $i \notin \Lambda$ ($\sigma_{\Lambda^c} \equiv +1$). Moreover let $\mathbb{Z}_*^2 = \mathbb{Z}^2 + (1/2, 1/2)$. We recall

the classical notion of *Peierls contour* associated to σ . We say that the segment joining two nearest neighbors $a, b \in \mathbb{Z}_*^2$ is marked if this segment separates two nearest neighbors $i, j \in \mathbb{Z}^2$ for which $\sigma_i \sigma_j = -1$. Marked segments form a finite family of closed, non self-intersecting, piecewise linear curves, that we call Peierls contours. Each segment of a contour γ separates two nearest neighbors whose spins have different signes (they necessarily belong to $Cl_1(\Lambda)$). If i, j are nearest neighbors separated by γ and $\sigma_i = -1$ we write $i \in \partial^- \gamma$ and $j \in \partial^+ \gamma$. We call the union of the sets of sites $\partial^- \gamma$ and $\partial^+ \gamma$ the *boundary of the contour* γ . For each $i \in \Lambda$ for which $\sigma_i = -1$, there is a *minimal Peierls contour* γ around i , i.e. such that i is in the interior of the closed curve γ .

This notion of minimal contour is the one used for the Ising model. Here we have to modify it as follows. Two Peierls contours γ, γ' are called *adjacent* if their boundaries have a common point. We say that two Peierls contours γ, γ' communicates if they belong to a sequence of Peierls contours $\gamma_1, \dots, \gamma_n$ such that for all k , γ_k and γ_{k+1} are adjacent. The relation of communicating is an equivalence relation. We call simply *contour* the union of the Peierls contours in an equivalence class. The minimal contour around i with $\sigma_i = -1$ is the one formed by the equivalence class which contains the minimal Peierls contour around i . The boundary (∂^+ or ∂^-) of a contour is simply the union of the boundaries of the Peierls contours that form it (see Fig. 4).

Let now μ_Λ^+ be the finite volume Gibbs measure with $+$ boundary condition, that we write as follows:

$$\mu_\Lambda^+(\sigma) = \frac{1}{\mathbb{Z}_\Lambda^+} \prod_{i \in Cl_1(\Lambda)} \frac{\cosh(\beta \sum_j k(i-j)\sigma_j^+)}{\cosh(\beta \sum_j k(i-j))} \quad \text{with } \sigma^+ = \sigma_\Lambda(+1)_{\Lambda^c}.$$

We have modified the normalization for later convenience. A given $\sigma^+ \in \{-1, +1\}^{\mathbb{Z}^2}$ corresponds, as described above, to a collection of contours $\Gamma = \{c_1, \dots, c_m\}$. Each contour c_i is a union of Peierls contours. Peierls contours belonging to different c_i 's do not communicate. We can write:

$$\mu_\Lambda^+(\sigma) = \frac{1}{\mathbb{Z}_\Lambda^+} \prod_{k=1}^m F(c_k),$$

where

$$F(c_k) = \prod_{i \in \partial c_k} \frac{\cosh(\beta \sum_j k(i-j)\sigma_j)}{\cosh(\beta \sum_j k(i-j))}$$

and $\partial c_k = \partial^+ c_k \cup \partial^- c_k$. Observing that if $\sigma_0 = -1$ then there is a contour around 0, we have:

$$\mu_\Lambda^+(\sigma_0 = -1) = \frac{1}{\mathbb{Z}_\Lambda^+} \sum_{c_1 \text{ around } 0} F(c_1) \sum_{\Gamma \ni c_1} F(\Gamma \setminus c_1),$$

where, for $\Gamma = c_1 \cup c_2 \cup \dots \cup c_m$, we let $F(\Gamma \setminus c_1) = \prod_{k=2}^m F(c_k)$. Note that, if Γ is a contour, $\Gamma \setminus c_1$ is also a contour, that corresponds to the configuration obtained by flipping all the spins -1 inside c_1 in the configuration associated to Γ . It follows that

$$\sum_{\Gamma \ni c_1} F(\Gamma \setminus c_1) \leq Z_\Lambda^+ \equiv \sum_{\Gamma} F(\Gamma),$$

and therefore

$$\mu_\Lambda^+(\sigma_0 = -1) \leq \sum_{c_1 \text{ around } 0} F(c_1). \quad (26)$$

Now note that if c_1 is a contour and $i \in \partial c_1$, then the spins $\sigma_i, \sigma_{i \pm e_1}, \sigma_{i \pm e_2}$ do not have the same sign, so that

$$\frac{\cosh(\beta \sum_j k(i-j)\sigma_j)}{\cosh(\beta \sum_j k(i-j))} \leq \frac{\cosh(\beta A)}{\cosh(\beta B)},$$

where $B = \sum_j k(j)$, A is the maximum value of $|\sum_j k(i-j)\sigma_j|$ for σ such that $\sigma_0, \sigma_{\pm e_1}, \sigma_{\pm e_2}$ do not have the same sign, and therefore $A < B$. Thus, we have to compare for a contour c_1 , the cardinal of its boundary $|\partial c_1|$




FIGURE 1. Example of a configuration σ on $Cl_2(\Lambda)$ such that $\sigma_0 = -1$ and $\sigma_{\Lambda^c} \equiv +1$. Drawing of its corresponding contours : γ_1 is the *minimal Peierls contour* around the origin ; $(\gamma_1 \cup \gamma_2 \cup \gamma_3 \cup \gamma_4)$ is the *minimal contour* around the origin (*i.e.* the equivalence class of γ_1) ; $\{\gamma_5, \gamma_6\}, \{\gamma_7\}, \{\gamma_8, \gamma_9, \gamma_{10}\}, \{\gamma_{11}\}, \{\gamma_{12}\}$ are the other equivalence classes.

with its length denoted by $l(c_1)$. But remark that to any point of ∂c_1 correspond at most 4 marked segments on c_1 . So, $l(c_1) \leq 4|\partial c_1|$, and we have

$$F(c_1) \leq \left[\frac{\cosh(\beta A)}{\cosh(\beta B)} \right]^{|\partial c_1|} \leq \left[\frac{\cosh(\beta A)}{\cosh(\beta B)} \right]^{l(c_1)/4}.$$

On the other hand, for a given length l , it is easily checked that the number of contours around 0 of length l is bounded by $l^3 3^{l-1}$. Thus, by (26),

$$\mu_{\Lambda}^+(\sigma_0 = -1) \leq \sum_{l \geq 0} l^3 3^{l-1} \left[\frac{\cosh(\beta A)}{\cosh(\beta B)} \right]^{l/4}$$

that goes to zero as $\beta \uparrow +\infty$. Thus, taking β large enough and letting $L \uparrow \mathbb{Z}^d$ in μ_{Λ}^+ , we construct a Gibbs measure μ for which $\mu^+(\sigma_0 = -1) < 1/2$. Symmetrically, taking minus boundary conditions, we obtain a Gibbs measure μ^- for which $\mu^-(\sigma_0 = -1) > 1/2$, and this proves phase transition.

Case 2: $k(0) < 0$, $k(e_1) > 0$, $k(e_2) > 0$.

Define

$$k^*(i) = \begin{cases} k(i) & \text{for } i \neq 0 \\ -k(0) & \text{for } i = 0, \end{cases}$$

and let Φ^* be the associated potential. Consider also the map $T : \{-1, 1\}^{\mathbb{Z}^2} \rightarrow \{-1, 1\}^{\mathbb{Z}^2}$ given by

$$(T\sigma)_i = \begin{cases} \sigma_i & \text{for } i \in \mathbb{Z}_e^2 \\ -\sigma_i & \text{for } i \in \mathbb{Z}_o^2. \end{cases}$$

To stress dependence on the potential Φ we write $\mu_{\Lambda, \Phi}^{\tau}$ for μ_{Λ}^{τ} . It is easily seen that

$$\mu_{\Lambda, \Phi}^{\tau}(\sigma) = \mu_{\Lambda, \Phi^*}^{T\tau}(T\sigma),$$

so that the map $\mu \rightarrow \mu \circ T$ is a bijection between $\mathcal{G}(\Phi)$ and $\mathcal{G}(\Phi^*)$. The conclusion follows from the fact that $|\mathcal{G}(\Phi^*)| > 1$, as seen in case 1.

Case 3: $k(0) \geq 0$, $k(e_1) > 0$, $k(e_2) < 0$.

This case is treated as case 2, with the following choices:

$$k^*(i) = \begin{cases} k(i) & \text{for } i \neq e_2 \\ -k(e_2) & \text{for } i = e_2, \end{cases}$$

and

$$(T\sigma)_i = \begin{cases} \sigma_i & \text{for } i = (x, y) \text{ with } y \text{ even} \\ -\sigma_i & \text{otherwise.} \end{cases}$$

the proof is now completed.

Remark 4.1. The special case $k(0) = 0$ was already treated in [11] example 2 (for $k(e_1) = k(e_2) = 1$), where a remarkable relation with Ising model was pointed out. We recall here in some more generality the principal steps of the argumentation :

let $\mathbb{Z}_o^2 = \{(x, y) \in \mathbb{Z}^2 : x + y \text{ is odd}\}$, $\mathbb{Z}_e^2 = \mathbb{Z}^2 \setminus \mathbb{Z}_o^2$ and, similarly, $\Lambda_o = \Lambda \cap \mathbb{Z}_o^2$, $\Lambda_e = \Lambda \cap \mathbb{Z}_e^2$. Note that since $k(0) = 0$, σ_{Λ_o} and σ_{Λ_e} are independent under μ_{Λ}^{τ} , i.e. $\mu_{\Lambda}^{\tau} = \mu_{\Lambda_e}^{\tau} \otimes \mu_{\Lambda_o}^{\tau}$. Consider the following anisotropic Ising model on $\{-1, 1\}^{\Lambda}$:

$$\rho_{\Lambda}^{\tau}(\sigma) = \frac{1}{N_{\Lambda}^{\tau}} \exp \left[\beta \sum_{i \in \Lambda} (k(e_1) \sigma_i \tilde{\sigma}_{i+e_1} + k(e_2) \sigma_i \tilde{\sigma}_{i+e_2}) \right],$$

where N_Λ^τ is the normalization and $\tilde{\sigma} = \sigma_\Lambda \tau_{\Lambda^c}$. Restricting this measure to the sites in Λ_e we obtain

$$\begin{aligned} \pi_{\Lambda_e} \rho_\Lambda^\tau(\sigma_{\Lambda_e}) &= \sum_{\sigma_{\Lambda_o}} \rho_\Lambda^\tau(\sigma) \\ &= \frac{2}{N_\Lambda^\tau} \prod_{i \in \Lambda_o} \cosh \left[\beta \sum_j k(i-j) \tilde{\sigma}_j \right] \\ &= \mu_{\Lambda_o}^\tau(\sigma_{\Lambda_e}). \end{aligned}$$

Therefore, phase transition for $\mathcal{G}(\Phi)$ follows from phase transition for the Ising model: since $\rho^-(\sigma_0 = +1) < \frac{1}{2} < \rho^+(\sigma_0 = +1)$, the restrictions $\pi_{\Lambda_e} \rho^-$ and $\pi_{\Lambda_e} \rho^+$ are different, and then

$$(\mu_{\mathbb{Z}_o^2}^+ = \pi_{\Lambda_e} \rho^+) \neq (\pi_{\Lambda_e} \rho^- = \mu_{\mathbb{Z}_o^2}^-).$$

We now show that, in certain cases, there are elements in $\mathcal{G}(\Phi)$ that are not stationary.

Proposition 4.2. *Suppose $k(0) \leq 0$, $k(e_1) < 0$, $k(e_2) < 0$, and let μ^+ be the Gibbs measure corresponding to plus boundary conditions. Suppose β is large enough so that $\mu^+ \neq \mu^-$. Then μ^+ is not stationary.*

Proof. We first observe that the transformation $k(\cdot) \rightarrow -k(\cdot)$ do not change the elements of $\mathcal{G}(\Phi)$, but it does change the dynamics. We recall few basic notions on stochastic ordering. Given $\sigma, \eta \in \{-1, 1\}^{\mathbb{Z}^2}$, we say that $\sigma \leq \eta$ if $\sigma_i \leq \eta_i$ for every $i \in \mathbb{Z}^2$. Monotonicity of functions $\{-1, 1\}^{\mathbb{Z}^2} \rightarrow \mathbb{R}$ is defined with respect to this partial order. Finally, for ν, μ probabilities on $\{-1, 1\}^{\mathbb{Z}^2}$, we say that $\nu \leq \mu$ if $\int f d\nu \leq \int f d\mu$ for every increasing f .

The key observation consists in the fact that, under our assumptions on $k(\cdot)$, the transition probability $P(d\sigma|\eta)$ is decreasing, i.e.

$$\nu \leq \mu \text{ implies } P\nu \geq P\mu.$$

This follows from the facts that $p_0(1|\eta)$ is decreasing in η , while $p_0(-1|\eta)$ is increasing in η (see [14] or [16] for details). Let now μ^0 be a limit point of the sequence ν_Λ^{per} defined in Remark 3. By using the criterion in [15], Th. II 2.9, it is easy to check that $\nu_\Lambda^{per} \leq \nu_\Lambda^+$ for every Λ , and so $\mu^0 \leq \mu^+$. Moreover, $0 = \mu^0(\sigma_0 = -1) < \frac{1}{2} < \mu^+(\sigma_0 = -1)$. So $\mu^0 \neq \mu^+$. On the other hand, by Proposition 3.2, μ^0 is stationary. Therefore $P\mu^+ \leq P\mu^0 = \mu^0 < \mu^+$, which completes the proof.

: P.-Y. Louis thanks, for their kind hospitality, the Mathematics' Departement of Padova University and the Interacting Random Systems group of Weierstrass Institute for Applied Analysis and Stochastics in Berlin, where part of this work was done.

REFERENCES

- [1] Barron A.R., *The strong ergodic theorem for densities: generalized Shannon-McMillan-Breiman theorem*, Ann. Prob. 13, no. 4 (1985), 1292-1303.
- [2] Bigelis S., Cirillo E.N.M., Lebowitz J.L., Speer E.R., *Critical droplets in metastable states of probabilistic cellular automata.*, Phys. Rev. E (3) 59 (1999), no. 4, 3935-3941.
- [3] Brémaud P., *Markov chains. Gibbs fields, Monte Carlo simulation, and queues*, Texts in Applied Mathematics, 31. Springer-Verlag, New York, 1999.
- [4] Dai Pra P., Ph.D. Thesis, Rutgers University, 1992.
- [5] Dawson, D. A., *Synchronous and asynchronous reversible Markov systems*, Canad. Math. Bull. 17 (1974/75), no. 5, 633-649.
- [6] Georgii H-O., *Gibbs measures and phase transitions*, de Gruyter Studies in Mathematics, 9. Walter de Gruyter & Co., Berlin, 1988.
- [7] Goldstein S., Kuik R., Lebowitz J.L., Maes C., *From PCAs to equilibrium systems and back.*, Comm. Math. Phys. 125 (1989), no. 1, 71-79.
- [8] Guyon X., *Champs aléatoires sur un réseau. Modélisations, statistique et applications.*, Techniques stochastiques, Masson, Paris, 1992.
- [9] Handa K., *Entropy production per site in (nonreversible) spin-flip processes*, J. Statist. Phys. 83 (1996), no. 3-4, 555-571.
- [10] Holley R., *Free energy in a Markovian model of a lattice spin system*, Comm. Math. Phys. 23 (1971), 87-99.
- [11] Kozlov, O., Vasilyev, N., *Reversible Markov chains with local interaction*, Multicomponent random systems, pp. 451-469, Adv. Probab. Related Topics, 6, Dekker, New York, 1980.
- [12] Künsch, H., *Nonreversible stationary measures for infinite interacting particle systems*, Z. Wahrsch. Verw. Gebiete 66 (1984), no. 3, 407-424.
- [13] Künsch H., *Time reversal and stationary Gibbs measures*, Stochastic Processes and their Applications 17 (1984), 159-166.
- [14] Lebowitz J.L., Maes C., Speer E.R., *Statistical mechanics of probabilistic cellular automata.*, J. Statist. Phys. 59 (1990), no. 1-2, 117-170.
- [15] Liggett T.M., *Interacting particle systems*. 276. Springer-Verlag, New York-Berlin, 1985.
- [16] Lopez F.J., Sanz G., *Stochastic comparisons for general probabilistic cellular automata*, Stat. Probab. Letters 46 (2000), 401-410.
- [17] Maes C., Shlosman S.B., *Ergodicity of probabilistic cellular automata: a constructive criterion*. Comm. Math. Phys. 135 (1991), no. 2, 233-251.
- [18] Maes C., Shlosman S.B., *When is an interacting particle system ergodic?*, Comm. Math. Phys. 151 (1993), no. 3, 447-466.
- [19] Maes C., Vande Velde K., *The (non-) Gibbsian nature of states invariant under stochastic transformations*, Physica A 206, (1994), 587-603.
- [20] Malyshev V.A., Minlos R.A., *Gibbs random fields, Cluster expansions*, Mathematics and its Applications 44, Kluwer Academic Publishers, Dordrecht, 1991.
- [21] Martinelli F., *Lectures on Glauber dynamics for discrete spin models*, Lectures on probability theory and statistics (Saint-Flour, 1997), 93-191, Lecture Notes in Math. 1717, Springer, Berlin, 1999.
- [22] Preston C., *Random fields*, Lecture Notes in Mathematics, Vol. 534. Springer-Verlag, Berlin-New York, 1976.
- [23] Toom A.L., Vasilyev N.B., Stavskaya O.N., Mityushin L.G., Kurdyumov G.L., Pirogov S.A., *Discrete local Markov systems*, in Stochastic Cellular Systems: ergodicity, memory, morphogenesis, edited by R.L. Dobrushin, V.I. Kryukov, A.L. Toom, Manchester University Press, Manchester, 1990, 1-182.
- [24] Vasilyev N.B., *Bernoulli and Markov stationary measures in discrete local interactions*, Locally interacting systems and their applications in biology (Pushchino, 1976), R.L. Dobrushin, V.I. Kryukov and A.L. Toom eds., Lecture Notes in Mathematics 653, Springer, Berlin 1978.