



HAL
open science

On the monodromy conjecture for non-degenerate hypersurfaces *

Alexander Esterov, Ann Lemahieu, Kiyoshi Takeuchi

► **To cite this version:**

Alexander Esterov, Ann Lemahieu, Kiyoshi Takeuchi. On the monodromy conjecture for non-degenerate hypersurfaces *. 2016. hal-01279910v1

HAL Id: hal-01279910

<https://hal.science/hal-01279910v1>

Preprint submitted on 27 Feb 2016 (v1), last revised 13 Mar 2022 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

On the monodromy conjecture for non-degenerate hypersurfaces ^{*}

Alexander ESTEROV [†], Ann LEMAHIEU [‡] and Kiyoshi TAKEUCHI [§]

December 22, 2014

Abstract

Recently the second author and Van Proeyen proved the monodromy conjecture on topological zeta functions for all non-degenerate surface singularities. In this paper, we obtain some higher-dimensional analogues of their results. First we study configurations of B_1 -pyramid facets which produce fake poles. Secondly, we introduce fully supermodular functions which are useful to find eigenvalues of monodromies. Finally, we obtain a result which would be useful to treat the monodromy conjecture for non-degenerate hypersurfaces whose Newton polyhedron is not convenient. In particular, we prove the conjecture for non-degenerate hypersurface singularities in \mathbb{C}^4 under some additional assumptions.

1 Introduction

Over the fields \mathbb{R} and \mathbb{C} it is well-known that the poles of the local zeta function associated to a polynomial f are contained in the set of roots of the Bernstein-Sato polynomial and integer shifts of them. By a celebrated theorem of Kashiwara and Malgrange, this implies that for any such pole $s_0 \in \mathbb{Q}$ the complex number $\exp(2\pi i s_0) \in \mathbb{C}$ is an eigenvalue of the monodromies of the complex hypersurface defined by f . Igusa predicted a similar beautiful relationship between the poles of p -adic integrals and the complex monodromies. This is now called the monodromy conjecture (see Denef [4] and Nicaise [25] for excellent reviews on this subject). Later in [6] Denef and Loeser introduced the local topological zeta function $Z_{\text{top},f}(s)$ associated to f and proposed a weaker version of the monodromy

***2010 Mathematics Subject Classification:** 14M25, 32S40, 32S60, 35A27

[†]National Research University Higher School of Economics

Faculty of Mathematics NRU HSE, 7 Vavilova 117312 Moscow, Russia, Email: aesterov@hse.ru

This study (research grant No 14-01-0152) is supported by The National Research University–Higher School of Economics Academic Fund Program in 2014/2015. Partially supported by RFBR grant 13-01-00755 and the Dynasty Foundation fellowship.

[‡]Laboratoire Paul Painlevé, Université de Lille 1, Cité Scientifique, 59655, Villeneuve d’Ascq Cedex, France, Email: ann.lemahieu@math.univ-lille1.fr

This research is partially supported by MCI-Spain grant MTM2010-21740-C02 and the ANR ‘SUSI’ project (ANR-12-JS01-0002-01).

[§]Institute of Mathematics, University of Tsukuba, 1-1-1, Tennodai, Tsukuba, Ibaraki, 305-8571, Japan. E-mail: takemicro@nifty.com

conjecture. For important contributions to this Denef-Loeser conjecture, see Loeser [19], [20], Artal Bartolo-Cassou-Noguès-Luengo-Melle Hernández [2], Veys [30] etc. Recently the second author and Van Proeyen [17] proved it for all non-degenerate surface singularities. The aim of this paper is to exploit to what extent the results of [17] hold true in higher dimensions. In Section 3, as a straightforward generalization of the notion of B_1 -facets in [17] we introduce B_1 -pyramid facets of the Newton polyhedron $\Gamma_+(f)$ of f and show that for some configurations of them the candidate poles of $Z_{\text{top},f}(s)$ contributed only by them are fake i.e. not actual poles. Moreover in Sections 4, 5 and 6, following the strategy of [17] we prove some key results to show that the candidate poles of $Z_{\text{top},f}(s)$ contributed by some non- B_1 -pyramid facets of $\Gamma_+(f)$ yield monodromy eigenvalues at some points in a neighborhood of the origin $0 \in \mathbb{C}^n$. In particular we obtain Theorem 4.3. Its proof partially relies upon the new notion of a fully supermodular function, which is inspired by supermodular functions in game theory and may be of independent interest. See Section 5 for the details. In Section 6 we prove Proposition 6.1 which enables us to reduce the problem of non-convenient polynomials to that of convenient ones. In this way, we can confirm the monodromy conjecture of Denef-Loeser [6] for most of non-degenerate hypersurfaces in higher dimensions. Indeed in Section 7 we prove the conjecture for non-degenerate polynomials of four variables under some additional assumptions (see Theorem 7.10 for the details).

Acknowledgements: The second author is very grateful to the University of Nice for their hospitality.

2 The monodromy conjecture for topological zeta functions

In this section, we briefly recall the monodromy conjecture for local topological zeta functions and related results. Let $f : (\mathbb{C}^n, 0) \rightarrow (\mathbb{C}, 0)$ be a germ of a non-trivial analytic function. We assume that f is defined on an open neighborhood X of the origin $0 \in \mathbb{C}^n$. Let $\pi : Y \rightarrow X$ be an embedded resolution of the complex hypersurface $f^{-1}(0) \subset X$ and E_j ($j \in J$) the irreducible components of the normal crossing divisor $\pi^{-1}(f^{-1}(0)) \subset Y$. For $j \in J$ we denote by N_j (resp. $\nu_j - 1$) the multiplicity of the divisor associated to $f \circ \pi$ (resp. $\pi^*(dx_1 \wedge \cdots \wedge dx_n)$) along $E_j \subset Y$. For a non-empty subset $I \subset J$ we set

$$E_I = \bigcap_{i \in I} E_i, \quad E_I^\circ = E_I \setminus \left(\bigcup_{j \notin I} E_j \right). \quad (2.1)$$

In [6] Denef and Loeser defined the local topological zeta function $Z_{\text{top},f}(s) \in \mathbb{C}(s)$ associated to f (at the origin) by

$$Z_{\text{top},f}(s) = \sum_{I \neq \emptyset} \chi(E_I^\circ \cap \pi^{-1}(0)) \times \prod_{i \in I} \frac{1}{N_i s + \nu_i}, \quad (2.2)$$

where $\chi(\cdot)$ denotes the topological Euler characteristic. More precisely, they introduced $Z_{\text{top},f}(s)$ by p -adic integrals and showed that it does not depend on the choice of the embedded resolution $\pi : Y \rightarrow X$ by algebraic methods. Later in [7] and [8], they redefined $Z_{\text{top},f}(s)$ by using the motivic zeta function of f and reproved this independence

of π more elegantly. For a point $x \in f^{-1}(0)$ let $F_x \subset X \setminus f^{-1}(0)$ be the Milnor fiber of f at x and $\Phi_{j,x} : H^j(F_x; \mathbb{C}) \xrightarrow{\sim} H^j(F_x; \mathbb{C})$ ($j \in \mathbb{Z}$) the Milnor monodromies associated to it. Then the monodromy conjecture of Denef-Loeser for the local topological zeta function $Z_{\text{top},f}(s)$ is stated as follows.

Conjecture: (Denef-Loeser [6, Conjecture 3.3.2]) Assume that $s_0 \in \mathbb{C}$ is a pole of $Z_{\text{top},f}(s)$. Then $\exp(2\pi i s_0) \in \mathbb{C}$ is an eigenvalue of the monodromy $\Phi_{j,x} : H^j(F_x; \mathbb{C}) \xrightarrow{\sim} H^j(F_x; \mathbb{C})$ for some point $x \in f^{-1}(0)$ in a neighborhood of the origin $0 \in \mathbb{C}^n$ and $j \in \mathbb{Z}$.

In [6] the authors also formulated an even stronger conjecture concerning the Bernstein-Sato polynomial $b_f(s)$ of f . Namely they conjectured that the poles of $Z_{\text{top},f}(s)$ are roots of $b_f(s)$. From now on, we assume that f is a non-trivial polynomial on \mathbb{C}^n such that $f(0) = 0$ and recall the results of Denef-Loeser [6, Section 5] and Varchenko [29]. For $f(x) = \sum_{v \in \mathbb{Z}_+^n} c_v x^v \in \mathbb{C}[x_1, \dots, x_n]$ we define its support $\text{supp} f \subset \mathbb{Z}_+^n$ by

$$\text{supp} f = \{v \in \mathbb{Z}_+^n \mid c_v \neq 0\} \subset \mathbb{Z}_+^n. \quad (2.3)$$

We denote by $\Gamma_+(f) \subset \mathbb{R}_+^n$ the convex hull of $\cup_{v \in \text{supp} f} (v + \mathbb{R}_+^n)$ in \mathbb{R}_+^n and call it the Newton polyhedron of f at the origin $0 \in \mathbb{C}^n$. The polynomial f such that $f(0) = 0$ is called convenient if $\Gamma_+(f)$ intersects the positive part of any coordinate axis of \mathbb{R}^n .

Definition 2.1. (Kouchnirenko [15]) We say that f is non-degenerate (at the origin $0 \in \mathbb{C}^n$) if for any compact face $\tau \prec \Gamma_+(f)$ the complex hypersurface

$$\{x \in (\mathbb{C}^*)^n \mid f_\tau(x) = 0\} \subset (\mathbb{C}^*)^n \quad (2.4)$$

in $(\mathbb{C}^*)^n$ is smooth and reduced, where we set

$$f_\tau(x) = \sum_{v \in \tau \cap \mathbb{Z}_+^n} c_v x^v \in \mathbb{C}[x_1, \dots, x_n]. \quad (2.5)$$

It is well-known that generic polynomials having a fixed Newton polyhedron are non-degenerate (see for example [27, Chapter V, paragraph 2]). For $a = (a_1, \dots, a_n) \in \mathbb{R}_+^n$ we set

$$N(a) = \min_{v \in \Gamma_+(f)} \langle a, v \rangle, \quad \nu(a) = |a| = \sum_{i=1}^n a_i \quad (2.6)$$

and

$$F(a) = \{v \in \Gamma_+(f) \mid \langle a, v \rangle = N(a)\} \prec \Gamma_+(f). \quad (2.7)$$

We call $F(a)$ the supporting face of the vector $a \in \mathbb{R}_+^n$ on $\Gamma_+(f)$. For a face $\tau \prec \Gamma_+(f)$ we set

$$\tau^\circ = \overline{\{a \in \mathbb{R}_+^n \mid F(a) = \tau\}} \subset \mathbb{R}_+^n. \quad (2.8)$$

Note that τ° is an $(n - \dim \tau)$ -dimensional rational polyhedral convex cone in \mathbb{R}_+^n . We call it the dual cone of τ . Let $\Delta = \mathbb{R}_+ a(1) + \dots + \mathbb{R}_+ a(l)$ ($a(i) \in \mathbb{Z}_+^n$) be a rational simplicial cone in \mathbb{R}_+^n , where the $a(i)$ are linearly independent over \mathbb{R} and primitive. Let $\text{aff}(\Delta) \simeq \mathbb{R}^l$ be the affine span of Δ in \mathbb{R}^n and $s(\Delta) \subset \Delta$ the l -dimensional lattice simplex whose vertices are $a(1), \dots, a(l)$ and the origin $0 \in \Delta \subset \mathbb{R}_+^n$. We denote by $\text{mult}(\Delta) \in \mathbb{Z}_{>0}$ the l -dimensional normalized volume $\text{Vol}_{\mathbb{Z}}(s(\Delta))$ of $s(\Delta)$ i.e. $l!$ times the usual volume of

$s(\Delta)$ with respect to the affine lattice $\text{aff}(\Delta) \cap \mathbb{Z}^n \simeq \mathbb{Z}^l$ in $\text{aff}(\Delta)$. By using this integer $\text{mult}(\Delta)$ we set

$$J_\Delta(s) = \frac{\text{mult}(\Delta)}{\prod_{i=1}^l \{N(a(i))s + \nu(a(i))\}} \in \mathbb{C}(s). \quad (2.9)$$

For a face $\tau \prec \Gamma_+(f)$ we choose a decomposition $\tau^\circ = \cup_{1 \leq i \leq r} \Delta_i$ of its dual cone τ° into rational simplicial cones Δ_i of dimension $l := \dim \tau^\circ$ such that $\dim(\Delta_i \cap \Delta_j) < l$ ($i \neq j$) and set

$$J_\tau(s) = \sum_{i=1}^r J_{\Delta_i}(s) \in \mathbb{C}(s). \quad (2.10)$$

According to [6, Lemme 5.1.1], this rational function $J_\tau(s)$ does not depend on the choice of the decomposition of τ° . It is also well-known that we can decompose τ° into rational simplicial cones without adding new edges. Then we have the following formula for $Z_{\text{top},f}(s)$.

Theorem 2.2. (*Denef-Loeser [6, Théorème 5.3 (ii)]*) *Assume that $f(x) \in \mathbb{C}[x_1, \dots, x_n]$ is non-degenerate. Then we have*

$$Z_{\text{top},f}(s) = \sum_{\gamma} J_\gamma(s) + \frac{s}{s+1} \sum_{\tau} (-1)^{\dim \tau} \text{Vol}_{\mathbb{Z}}(\tau) \cdot J_\tau(s), \quad (2.11)$$

where in the sum \sum_{γ} (resp. \sum_{τ}) the face $\gamma \prec \Gamma_+(f)$ (resp. $\tau \prec \Gamma_+(f)$) ranges through the vertices of $\Gamma_+(f)$ (resp. the compact ones such that $\dim \tau \geq 1$) and $\text{Vol}_{\mathbb{Z}}(\tau) \in \mathbb{Z}_{>0}$ is the $(\dim \tau)$ -dimensional normalized volume of τ with respect to the affine lattice $\text{aff}(\tau) \cap \mathbb{Z}^n \simeq \mathbb{Z}^{\dim \tau}$ in $\text{aff}(\tau) \simeq \mathbb{R}^{\dim \tau}$.

Recall that a face τ of $\Gamma_+(f)$ is called a facet if $\dim \tau = n - 1$. For a facet $\tau \prec \Gamma_+(f)$ let $a(\tau) = (a(\tau)_1, \dots, a(\tau)_n) \in \tau^\circ \cap \mathbb{Z}_+^n$ be its primitive conormal vector and set

$$N(\tau) = \min_{v \in \Gamma_+(f)} \langle a(\tau), v \rangle, \quad (2.12)$$

$$\nu(\tau) = |a(\tau)| = \sum_{i=1}^n a(\tau)_i = \left\langle a(\tau), \begin{pmatrix} 1 \\ \vdots \\ 1 \end{pmatrix} \right\rangle. \quad (2.13)$$

We call $N(\tau)$ the lattice distance of τ from the origin $0 \in \mathbb{R}^n$. It follows from Theorem 2.2 that any pole $s_0 \neq -1$ of $Z_{\text{top},f}(s)$ is contained in the finite set

$$\left\{ -\frac{\nu(\tau)}{N(\tau)} \mid \tau \prec \Gamma_+(f) \text{ is a facet not lying in a coordinate hyperplane} \right\} \subset \mathbb{Q}. \quad (2.14)$$

Its elements are called candidate poles of $Z_{\text{top},f}(s)$. We say that a candidate pole $s_0 \in \mathbb{C}$ of $Z_{\text{top},f}(s)$ is contributed by a facet $\tau \prec \Gamma_+(f)$ if we have $s_0 = -\frac{\nu(\tau)}{N(\tau)}$.

Finally we recall the result of Varchenko [29]. For a polynomial $f(x) \in \mathbb{C}[x_1, \dots, x_n]$ such that $f(0) = 0$, we define its monodromy zeta function $\zeta_{f,0}(t) \in \mathbb{C}(t)$ at the origin $0 \in \mathbb{C}^n$ by

$$\zeta_{f,0}(t) = \prod_{j \in \mathbb{Z}} \left\{ \det(\text{id} - t\Phi_{j,0}) \right\}^{(-1)^j} \in \mathbb{C}(t). \quad (2.15)$$

Similarly one can define also $\zeta_{f,x}(t) \in \mathbb{C}(t)$ for any point $x \in f^{-1}(0)$. Then by considering the decomposition of the nearby cycle perverse sheaf $\psi_f(\mathbb{C}_{\mathbb{C}^n})[n-1]$ with respect to the monodromy eigenvalues of f and the concentrations of its components at generic points $x \in f^{-1}(0)$ (see e.g. [9] and [14]), in order to prove the monodromy conjecture, it suffices to show that for any pole $s_0 \in \mathbb{C}$ of $Z_{\text{top},f}(s)$ the complex number $\exp(2\pi i s_0) \in \mathbb{C}$ is a root or a pole of $\zeta_{f,x}(t)$ for some point $x \in f^{-1}(0)$ in a neighborhood of the origin $0 \in \mathbb{C}^n$ (see Denef [5, Lemma 4.6]). For a subset $S \subset \{1, 2, \dots, n\}$ we define a coordinate subspace $\mathbb{R}^S \simeq \mathbb{R}^{|S|}$ of \mathbb{R}^n by

$$\mathbb{R}^S = \{v = (v_1, \dots, v_n) \in \mathbb{R}^n \mid v_i = 0 \text{ for any } i \notin S\} \quad (2.16)$$

and set

$$\mathbb{R}_+^S = \mathbb{R}^S \cap \mathbb{R}_+^n \simeq \mathbb{R}_+^{|S|}. \quad (2.17)$$

For a compact face $\tau \prec \Gamma_+(f)$ we take the minimal coordinate subspace \mathbb{R}^S of \mathbb{R}^n containing τ and set $s_\tau = |S|$. If τ satisfies the condition $\dim \tau = s_\tau - 1$ we set

$$\zeta_\tau(t) = \left(1 - t^{N(\tau)}\right)^{\text{Vol}_{\mathbb{Z}}(\tau)} \in \mathbb{C}[t], \quad (2.18)$$

where $N(\tau) \in \mathbb{Z}_{>0}$ is the lattice distance of the affine hyperplane $\text{aff}(\tau) \simeq \mathbb{R}^{\dim \tau}$ in \mathbb{R}^S from the origin $0 \in \mathbb{R}^S$.

Theorem 2.3. (Varchenko [29]) *Assume that $f(x) \in \mathbb{C}[x_1, \dots, x_n]$ is non-degenerate. Then we have*

$$\zeta_{f,0}(t) = \prod_{\tau} \left\{ \zeta_\tau(t) \right\}^{(-1)^{\dim \tau}}, \quad (2.19)$$

where in the product \prod_{τ} the face $\tau \prec \Gamma_+(f)$ ranges through the compact ones satisfying the condition $\dim \tau = s_\tau - 1$.

Definition 2.4. *We say that a face τ of $\Gamma_+(f)$ is a V -face (or a Varchenko face) if it is compact and satisfies the condition $\dim \tau = s_\tau - 1$.*

To end this section, we note the following simple fact. Also for a general face $\tau \prec \Gamma_+(f)$ we define its lattice distance $N(\tau) \in \mathbb{Z}_{>0}$ to be that of the affine hyperplane $\text{aff}(\tau) \simeq \mathbb{R}^{\dim \tau}$ in $\text{aff}(\tau \cup \{0\}) \simeq \mathbb{R}^{\dim \tau + 1}$ from the origin $0 \in \text{aff}(\tau \cup \{0\}) \subset \mathbb{R}^n$.

Lemma 2.5. *For two faces $\tau, \gamma \prec \Gamma_+(f)$ such that $\gamma \prec \tau$, we have $N(\gamma) \mid N(\tau)$.*

Proof. We may assume that γ is a facet of τ . Let Σ_0 be the dual fan of $\Gamma_+(f)$ and Σ a smooth subdivision of Σ_0 . Then there exists an $(n - \dim \gamma)$ -dimensional smooth cone $\Delta \in \Sigma$ contained in γ° such that $\dim(\Delta \cap \tau^\circ) = \dim \tau^\circ = n - \dim \tau = (n - \dim \gamma) - 1$. Set $l = n - \dim \gamma$ and let $a(1), \dots, a(l) \in \mathbb{Z}_+^n$ be the primitive vectors on the edges of Δ . Then it is easy to see that for any $v \in \gamma$ we have

$$N(\tau) = \gcd\left(\langle a(1), v \rangle, \dots, \langle a(l), v \rangle\right). \quad (2.20)$$

We have also a similar description of $N(\tau)$ in terms of the primitive vectors on the edges of $\Delta \cap \tau^\circ \prec \Delta$. If we use a point $v \in \gamma \prec \tau$ to express $N(\gamma)$ and $N(\tau)$ simultaneously, we find $N(\gamma) \mid N(\tau)$. This completes the proof. \square

Remark 2.6. For two V -faces $\tau, \gamma \prec \Gamma_+(f)$ such that $\gamma \prec \tau$ we can prove Lemma 2.5 more easily as follows. Let \mathbb{R}^{S_τ} (resp. \mathbb{R}^{S_γ}) be the minimal coordinate subspace of \mathbb{R}^n containing τ (resp. γ) and

$$\text{aff}(\tau) : \langle a(\tau), v \rangle = N(\tau) \quad (2.21)$$

the equation of the affine hyperplane $\text{aff}(\tau) \subset \mathbb{R}^{S_\tau}$. Then by restricting it to $\mathbb{R}^{S_\gamma} \subset \mathbb{R}^{S_\tau}$ to find that of $\text{aff}(\gamma)$ we obtain the equality

$$\text{gcd}\{a(\tau)_i \ (i \in S_\tau)\} \cdot N(\gamma) = N(\tau). \quad (2.22)$$

3 Fake poles of topological zeta functions

In this section, we define B_1 -pyramid facets of the Newton polyhedron $\Gamma_+(f)$ and show that for some configurations of them the candidate poles of $Z_{\text{top},f}(s)$ contributed only by them are fake i.e. not actual poles. Our definition is a straightforward generalization of that of Lemahieu-Van Proeyen [17].

For a subset $S \subset \{1, 2, \dots, n\}$ let $\pi_S : \mathbb{R}_+^n \longrightarrow \mathbb{R}_+^{S^c} \simeq \mathbb{R}_+^{n-|S|}$ be the natural projection. We say that an $(n-1)$ -dimensional polyhedron τ in \mathbb{R}_+^n is non-compact for $S \subset \{1, 2, \dots, n\}$ if the Minkowski sum $\tau + \mathbb{R}_+^S$ is contained in τ .

Definition 3.1. (cf. Lemahieu-Van Proeyen [17]) Let τ be an $(n-1)$ -dimensional polyhedron in \mathbb{R}_+^n .

1. We say that τ is a B_1 -pyramid of compact type for the variable v_i if τ is a compact pyramid over the base $\gamma = \tau \cap \{v_i = 0\}$ and its unique vertex $P \prec \tau$ such that $P \notin \gamma$ has height one from the hyperplane $\{v_i = 0\} \subset \mathbb{R}_+^n$.
2. We say that τ is a B_1 -pyramid of non-compact type if there exists a non-empty subset $S \subset \{1, 2, \dots, n\}$ such that τ is non-compact for S and $\pi_S(\tau) \subset \mathbb{R}_+^{S^c} \simeq \mathbb{R}_+^{n-|S|}$ is a B_1 -pyramid of compact type for some variable v_i ($i \notin S$).
3. We say that τ is a B_1 -pyramid if it is a B_1 -pyramid of compact or non-compact type.

Let us recall the condition in Loeser [20]. For two distinct facets τ and τ' of $\Gamma_+(f)$ let $\beta(\tau, \tau') \in \mathbb{Z}$ be the greatest common divisor of the 2×2 minors of the matrix $(a(\tau), a(\tau')) \in M(n, 2; \mathbb{Z})$. If $N(\tau) \neq 0$ (e.g. if τ is compact) we set

$$\lambda(\tau, \tau') = \nu(\tau') - \frac{\nu(\tau)}{N(\tau)} N(\tau'), \quad \mu(\tau, \tau') = \frac{\lambda(\tau, \tau')}{\beta(\tau, \tau')} \in \mathbb{Q}. \quad (3.1)$$

In [20] the author considered only compact facets τ of $\Gamma_+(f)$ which satisfy the following technical condition:

“For any facet $\tau' \prec \Gamma_+(f)$ such that $\tau' \neq \tau$ and $\tau' \cap \tau \neq \emptyset$ we have $\mu(\tau, \tau') \notin \mathbb{Z}$.”

He showed that the candidate pole of $Z_{\text{top},f}(s)$ associated to such a compact facet τ is a root of the local Bernstein-Sato polynomial of f . Now let $\tau \prec \Gamma_+(f)$ be a facet containing a B_1 -pyramid of compact type for the variable v_i and set $\gamma = \tau \cap \{v_i = 0\}$. Let $\tau_0 \prec \Gamma_+(f)$ be the unique (non-compact) facet such that $\gamma \prec \tau_0$, $\tau_0 \neq \tau$ and $\tau_0 \subset \{v_i = 0\}$.

Then we can easily show that $\beta(\tau, \tau_0) = \mu(\tau, \tau_0) = 1$. Namely such a facet τ does not satisfy the above-mentioned condition of [20]. In this paper, we treat also candidate poles associated to non-compact facets of $\Gamma_+(f)$. Roughly speaking, non-isolated singularities of the hypersurface $f^{-1}(0) \subset \mathbb{C}^n$ correspond to non-convenient Newton polyhedra (see [16]).

Proposition 3.2. *Assume that f is non-degenerate and let the facet $\tau \prec \Gamma_+(f)$ be a B_1 -pyramid. Assume also that the candidate pole*

$$s_0 = -\frac{\nu(\tau)}{N(\tau)} \neq -1 \quad (3.2)$$

of $Z_{\text{top},f}(s)$ is contributed only by τ . Then s_0 is fake i.e. not an actual pole of $Z_{\text{top},f}(s)$.

Proof. Since the proof for B_1 -pyramids of non-compact type is similar, we prove the assertion only for B_1 -pyramids of compact type. Without loss of generality we may assume that τ is a compact pyramid over the base $\gamma = \tau \cap \{v_n = 0\}$ and its unique vertex $P \prec \tau$ such that $P \notin \gamma$ has height one from the hyperplane $\{v_n = 0\} \subset \mathbb{R}_+^n$. First let us consider the simplest case where γ and hence τ are simplices. Let A_1, A_2, \dots, A_{n-1} be the vertices of the $(n-2)$ -dimensional simplex γ . For $1 \leq i \leq n-1$ denote by σ_i the facet of τ whose vertices are P and the A_j ($j \neq i$). As in the proof of [17, Proposition 14] let $\tau_i \prec \Gamma_+(f)$ ($1 \leq i \leq n-1$) (resp. $\tau_0 \prec \Gamma_+(f)$) be the unique facet such that $\sigma_i \prec \tau_i$ and $\tau_i \neq \tau$ (resp. $\gamma \prec \tau_0$, $\tau_0 \neq \tau$ and $\tau_0 \subset \{v_n = 0\}$). Then it is easy to see that the primitive conormal vector $a(\tau) \in \mathbb{Z}_+^n$ of τ is in the interior of the cone generated by $a(\tau_1), \dots, a(\tau_{n-1})$ and $a(\tau_0) = (0, 0, \dots, 0, 1)$. For $1 \leq i \leq n-1$ denote by V_{A_i} the multiplicity of the (n) -dimensional cone generated by $a(\tau)$, $a(\tau_0)$ and $a(\tau_j)$ ($j \neq i$). Also we denote by V_P the multiplicity of the (n) -dimensional cone generated by $a(\tau)$ and $a(\tau_1), \dots, a(\tau_{n-1})$. Finally we set $l(\tau)(s) = N(\tau)s + \nu(\tau)$ and $l(\tau_i)(s) = N(\tau_i)s + \nu(\tau_i)$ ($1 \leq i \leq n-1$). Then by the argument in Case 1 of the proof of [17, Proposition 14] we obtain an equality

$$Z_{\text{top},f}(s) \equiv \frac{V_P}{l(\tau)(s) \prod_{j=1}^{n-1} l(\tau_j)(s)} + \frac{1}{s+1} \sum_{i=1}^{n-1} \frac{V_{A_i}}{l(\tau)(s) \prod_{j \neq i} l(\tau_j)(s)} \quad (3.3)$$

modulo holomorphic functions at $s = s_0 \in \mathbb{C}$. In order to prove the assertion, it suffices to show that the polynomial in the right hand side of the equality

$$(s+1)l(\tau)(s) \left(\prod_{j=1}^{n-1} l(\tau_j)(s) \right) Z_{\text{top},f}(s) \equiv V_P(s+1) + \sum_{i=1}^{n-1} V_{A_i} l(\tau_i)(s) \quad (3.4)$$

is divisible by the factor $l(\tau)(s) = N(\tau)s + \nu(\tau)$. Let $w = (*, *, \dots, *, 1) \in \mathbb{Z}_+^n$ be the coordinate of the point $P \in \tau \subset \mathbb{R}_+^n$ for which we have $\langle a(\tau_0), w \rangle = 1$. Let V be the multiplicity of the (n) -dimensional cone generated by $a(\tau_0), a(\tau_1), \dots, a(\tau_{n-1})$. Now observe that we have the geometric equality

$$Va(\tau) = V_P a(\tau_0) + \sum_{i=1}^{n-1} V_{A_i} a(\tau_i). \quad (3.5)$$

Then we obtain

$$V_P(s+1) + \sum_{i=1}^{n-1} V_{A_i} l(\tau_i)(s) \quad (3.6)$$

$$= \langle V_P a(\tau_0) + \sum_{i=1}^{n-1} V_{A_i} a(\tau_i), w \rangle s + \left\langle V_P a(\tau_0) + \sum_{i=1}^{n-1} V_{A_i} a(\tau_i), \begin{pmatrix} 1 \\ \vdots \\ 1 \end{pmatrix} \right\rangle \quad (3.7)$$

$$= \langle V a(\tau), w \rangle s + \left\langle V a(\tau), \begin{pmatrix} 1 \\ \vdots \\ 1 \end{pmatrix} \right\rangle = V l(\tau)(s) \quad (3.8)$$

and the proof for the case where γ and τ are simplices is now complete.

Now let us consider the general case where τ is a B_1 -pyramid of compact type over the base $\gamma \prec \tau$ such that $\gamma \subset \{v_n = 0\}$. Let $\tau_0 \prec \Gamma_+(f)$ be the unique facet such that $\gamma \prec \tau_0$, $\tau_0 \neq \tau$ and $\tau_0 \subset \{v_n = 0\}$. Let A_1, A_2, \dots, A_m ($m \geq n-1$) be the vertices of γ . For $1 \leq i \leq m$ we denote the dual cone A_i° of $A_i \prec \Gamma_+(f)$ by C_{A_i} . Similarly we set $C_P = P^\circ$. Then we have

$$a(\tau) \in \text{Int}(C_P \cup C_{A_1} \cup \dots \cup C_{A_m}). \quad (3.9)$$

In order to construct nice decompositions of $C_P, C_{A_1}, \dots, C_{A_m}$ into n -dimensional rational simplicial cones, we shall introduce a new dummy vector $b \in \text{Int}C_P \cap \mathbb{Z}_+^n$ satisfying the condition

$$-\frac{\nu(b)}{N(b)} \neq s_0 \quad (3.10)$$

as follows. First, by our assumption $s_0 \neq -1$ the coordinate vector $w = (*, *, \dots, *, 1) \in \mathbb{Z}_+^n$ of the summit P of the pyramid τ is not parallel to the vector

$$\begin{pmatrix} 1 \\ 1 \\ \vdots \\ 1 \end{pmatrix} \in \mathbb{Z}_+^n. \quad (3.11)$$

Then for a sufficiently large primitive vector $b \in \text{Int}C_P \cap \mathbb{Z}_+^n$ we can achieve the desired condition $N(b)s_0 + \nu(b) \neq 0$. For $1 \leq i \leq m$ let $e_i \prec \tau$ be the edge of τ connecting the two points P and A_i and $F_i \prec C_P$ the corresponding facet of the cone C_P containing its edge $\tau^\circ = \mathbb{R}_+ a(\tau) \prec C_P$. All the facets of C_P containing τ° are obtained in this way. Since the point $A_i \prec e_i$ is a vertex of τ_0 , its dual cone $C_{A_i} = A_i^\circ$ contains not only F_i but also the 1-dimensional cone $\tau_0^\circ = \mathbb{R}_+ a(\tau_0)$. For $1 \leq i \leq m$ set

$$F_i^\sharp = \mathbb{R}_+ a(\tau_0) + F_i, \quad F_i^\flat = \mathbb{R}_+ b + F_i. \quad (3.12)$$

In Figure 1 below we presented the transversal hyperplane sections of the cones F_i, F_i^\sharp and F_i^\flat .




Figure 1

Then by our construction $\cup_{i=1}^m (F_i^\# \cup F_i^b)$ is an n -dimensional cone in \mathbb{R}^n such that

$$a(\tau) \in \text{Int} \left\{ \bigcup_{i=1}^m (F_i^\# \cup F_i^b) \right\}. \quad (3.13)$$

Now we decompose F_1, F_2, \dots, F_m into rational simplicial cones. By using secondary polytopes (see [12, Chapter 7]), we can obtain such a subdivision of $F_1 \cup \dots \cup F_m \subset \partial C_P$ without adding new edges. We may assume also that it satisfies the axiom of fans. Let $\Delta_1, \dots, \Delta_r$ be the $(n-1)$ -dimensional simplicial cones thus obtained in $F_1 \cup \dots \cup F_m \subset \partial C_P$ and containing the edge $\tau^\circ = \mathbb{R}_+ a(\tau)$. For $1 \leq i \leq r$ set

$$\Delta_i^\# = \mathbb{R}_+ a(\tau_0) + \Delta_i, \quad \Delta_i^b = \mathbb{R}_+ b + \Delta_i. \quad (3.14)$$




Figure 2

Then $\cup_{i=1}^r (\Delta_i^\# \cup \Delta_i^b)$ is an n -dimensional cone in \mathbb{R}^n such that

$$a(\tau) \in \text{Int} \left\{ \bigcup_{i=1}^r (\Delta_i^\# \cup \Delta_i^b) \right\}. \quad (3.15)$$

By the argument in Case 1 of the proof of [17, Proposition 14] the contributions to $Z_{\text{top},f}(s)$ from the faces $\sigma \prec \tau$ of τ such that $\dim \sigma \geq 1$ cancel each other and it suffices to calculate

only the ones from the vertices $P, A_1, \dots, A_m \in \tau$ inside the cone $\cup_{i=1}^r (\Delta_i^\sharp \cup \Delta_i^b)$. For $1 \leq i \leq r$ let $\rho_{i,1}, \dots, \rho_{i,n-2} \in \partial C_P$ be the primitive vectors on the edges of Δ_i such that $\rho_{i,j} \notin \tau^\circ = \mathbb{R}_+ a(\tau)$ and set

$$l_{i,j}(s) = N(\rho_{i,j})s + \nu(\rho_{i,j}) \in \mathbb{C}[s] \quad (1 \leq j \leq n-2). \quad (3.16)$$

Moreover we set $l(b)(s) = N(b)s + \nu(b)$. For $1 \leq i \leq r$ we denote by V_i (resp. U_i) the multiplicity of the (n -dimensional) cone generated by $a(\tau), \rho_{i,1}, \dots, \rho_{i,n-2}$ and $a(\tau_0)$ (resp. b) in \mathbb{R}^n . Namely we set $\text{mult}(\Delta_i^\sharp) = V_i$ and $\text{mult}(\Delta_i^b) = U_i$. Then we obtain an equality

$$Z_{\text{top},f}(s) \equiv \sum_{i=1}^r \left\{ \frac{V_i}{(s+1)l(\tau)(s) \prod_{j=1}^{n-2} l_{i,j}(s)} + \frac{U_i}{l(b)(s)l(\tau)(s) \prod_{j=1}^{n-2} l_{i,j}(s)} \right\} \quad (3.17)$$

modulo holomorphic functions at $s = s_0 \in \mathbb{C}$. Let k be the cardinality of the set $\cup_{i=1}^r \{\rho_{i,1}, \rho_{i,2}, \dots, \rho_{i,n-2}\} \subset \mathbb{R}^n$ and set

$$\{\rho_1, \rho_2, \dots, \rho_k\} := \bigcup_{i=1}^r \{\rho_{i,1}, \rho_{i,2}, \dots, \rho_{i,n-2}\} \quad (3.18)$$

and

$$l_j(s) = N(\rho_j)s + \nu(\rho_j) \in \mathbb{C}[s] \quad (1 \leq j \leq k). \quad (3.19)$$

In order to prove the assertion, it suffices to show that the function

$$(s+1)l(\tau)(s)l(b)(s) \left(\prod_{j=1}^k l_j(s) \right) Z_{\text{top},f}(s) \quad (3.20)$$

$$\equiv \sum_{i=1}^r \left(\prod_{j: \rho_j \notin \Delta_i} l_j(s) \right) \cdot \{V_i l(b)(s) + U_i (s+1)\} \quad (3.21)$$

is divisible by the factor $l(\tau)(s) = N(\tau)s + \nu(\tau)$. For $1 \leq i \leq r$ we define an n -dimensional simplicial cone \square_i in \mathbb{R}^n by

$$\square_i = \mathbb{R}_+ a(\tau_0) + \mathbb{R}_+ \rho_{i,1} + \dots + \mathbb{R}_+ \rho_{i,n-2} + \mathbb{R}_+ b. \quad (3.22)$$




Figure 3

Then by applying the argument in the case where τ is a simplex to $\square_1, \dots, \square_r$, we can prove that (3.20) is divisible by $l(\tau)(s)$. Note that between two simplicial cones \square_i and \square_j ($i \neq j$) having a common facet there is a nice cancelling in our calculation. This completes the proof. \square

We can show also a similar result even in some cases where a candidate pole of $Z_{\text{top},f}(s)$ is contributed by several (adjacent) B_1 -pyramid facets of $\Gamma_+(f)$. Indeed we have the following higher-dimensional analogue of the result in the proof of [17, Proposition 14].

Proposition 3.3. *Assume that f is non-degenerate and let the facets $\tau_1, \dots, \tau_k \prec \Gamma_+(f)$ be B_1 -pyramids such that*

$$s_0 := -\frac{\nu(\tau_1)}{N(\tau_1)} = \dots = -\frac{\nu(\tau_k)}{N(\tau_k)} \neq -1 \quad (3.23)$$

and their common candidate pole $s_0 \in \mathbb{Q}$ of $Z_{\text{top},f}(s)$ is contributed only by τ_1, \dots, τ_k . Assume also that if τ_i and τ_j ($i \neq j$) have a common facet they are B_1 -pyramids for the same variable. Then s_0 is fake i.e. not an actual pole of $Z_{\text{top},f}(s)$.

Proof. If τ_i and τ_j ($i \neq j$) do not have a common facet, then by the proof of Proposition 3.2 after a suitable subdivision of the dual fan of $\Gamma_+(f)$ into rational simplicial cones we can calculate their contributions to $Z_{\text{top},f}(s)$ separately. So we may assume that the B_1 -pyramid facets τ_1, \dots, τ_k have the common summit $P \in \tau_1 \cap \dots \cap \tau_k$. For the sake of simplicity, here we shall treat only the case where $k = 2$, τ_1 (resp. τ_2) is a B_1 -pyramid over the base $\gamma_1 = \tau_1 \cap \{v_n = 0\}$ (resp. $\gamma_2 = \tau_2 \cap \{v_n = 0\}$) and $\tau_1 \cap \tau_2$ is the (unique) common facet of τ_1 and τ_2 . The proofs for the other cases are similar. Let $\tau_0 \prec \Gamma_+(f)$ be the unique facet of $\Gamma_+(f)$ such that $\gamma_1, \gamma_2 \prec \tau_0$, $\tau_0 \neq \tau_i$ ($i = 1, 2$) and $\tau_0 \subset \{v_n = 0\}$. We denote by P the common summit of τ_1 and τ_2 . As in the proof of Proposition 3.2, let F_1, F_2, \dots, F_m be the facets of the dual cone C_P of $P \prec \Gamma_+(f)$ containing $\tau_1^\circ = \mathbb{R}_+ a(\tau_1)$ or $\tau_2^\circ = \mathbb{R}_+ a(\tau_2)$ and subdivide $F_1 \cup \dots \cup F_m \subset \partial C_P$ into rational simplicial cones without adding new edges. We may assume that this subdivision satisfies the axiom of fans. Let $\Delta_1, \dots, \Delta_r$ be the $(n-1)$ -dimensional simplicial cones thus obtained in $F_1 \cup \dots \cup F_m \subset \partial C_P$ and containing τ_1° or τ_2° . As in the proof of Proposition 3.2, we take a new primitive vector $b \in \text{Int}C_P \cap \mathbb{Z}_+^n$ such that

$$-\frac{\nu(b)}{N(b)} \neq s_0. \quad (3.24)$$

For $1 \leq i \leq r$ set

$$\Delta_i^\sharp = \mathbb{R}_+ a(\tau_0) + \Delta_i, \quad \Delta_i^b = \mathbb{R}_+ b + \Delta_i. \quad (3.25)$$

Then $\square := \cup_{i=1}^r (\Delta_i^\sharp \cup \Delta_i^b)$ is an n -dimensional cone in \mathbb{R}^n such that

$$a(\tau_1), a(\tau_2) \in \text{Int}\square. \quad (3.26)$$

After a suitable change of the numbering of $\Delta_1, \dots, \Delta_r$, for some $0 < l < r$ we have also

$$\mathbb{R}_+ a(\tau_1) + \mathbb{R}_+ a(\tau_2) \prec \Delta_i \iff 1 \leq i \leq l. \quad (3.27)$$

Then we obtain an n -dimensional cone $\square' := \cup_{i=1}^l (\Delta_i^\sharp \cup \Delta_i^b)$ in \square such that $\mathbb{R}_+ a(\tau_1) + \mathbb{R}_+ a(\tau_2) \subset \square'$. Let us introduce a new primitive vector $b' \in \text{Int}\square' \cap \text{Int}C_P \cap \mathbb{Z}_+^n$ such that

$$-\frac{\nu(b')}{N(b')} \neq s_0 \quad (3.28)$$

and decompose $\square' \subset \square$ into rational simplicial cones by taking the convex hulls of $\mathbb{R}_+ b'$ with the facets of \square' .




Figure 4

Note that the resulting new decomposition of \square is different from the original one $\square = \cup_{i=1}^r (\Delta_i^\sharp \cup \Delta_i^b)$ inside \square' but still satisfies the axiom of fans. By (a slight modification of) [6, Lemme 5.1.1] and the condition $b' \in \text{Int}C_P$ we can use it for the calculation of $Z_{\text{top},f}(s)$. Then the contributions to $Z_{\text{top},f}(s)$ from τ_1 and τ_2 can be calculated separately, because the stars of the cones $\mathbb{R}_+ a(\tau_1)$ and $\mathbb{R}_+ a(\tau_2)$ are disjoint. Now the assertion follows immediately from the proof of Proposition 3.2. \square

As we see in the next proposition, in the case $n = 4$ there are also some other facets $\tau \prec \Gamma_+(f)$ whose candidate poles of $Z_{\text{top},f}(s)$ are fake.

Proposition 3.4. *In the case $n = 4$ assume that f is non-degenerate and let $\tau \prec \Gamma_+(f)$ be a facet with six vertices A, B, P, Q, X, Y of the form*

$$\begin{cases} A = (1, 0, *, *) \\ B = (1, 0, *, *) \\ P = (0, 1, *, *) \\ Q = (0, 1, *, *) \\ X = (0, 0, *, *) \\ Y = (0, 0, *, *) \end{cases} \quad (3.29)$$

as in Figure 5 below. It can be degenerated so that $P = Q$ etc. Assume also that the candidate pole

$$s_0 = -\frac{\nu(\tau)}{N(\tau)} \neq -1 \quad (3.30)$$

of $Z_{\text{top},f}(s)$ is contributed only by τ . Then s_0 is fake i.e. not an actual pole of $Z_{\text{top},f}(s)$.




Figure 5

Proof. Note that τ itself is not a B_1 -pyramid but it is a union of two B_1 -pyramids e.g. $AXPQ$ and $QXABY$. We define facets $\sigma_1, \sigma_2, \sigma_3$ of τ by $\sigma_1 = XAP$, $\sigma_2 = PABQ$, $\sigma_3 = YQB$ respectively. As in the proof of [17, Proposition 14] let $\tau_i \prec \Gamma_+(f)$ ($1 \leq i \leq 3$) be the unique facet such that $\sigma_i \prec \tau_i$ and $\tau_i \neq \tau$. Moreover for $i = 1, 2$ let $\rho_i \prec \Gamma_+(f)$ be the unique facet such that $\rho_i \subset \{v_i = 0\} \simeq \mathbb{R}^3$, $\tau \cap \{v_i = 0\} \prec \rho_i$ and $\rho_i \neq \tau$. As in the proof of Proposition 3.2 we define the multiplicities $V_A, V_B, V_P, V_Q, V_X, V_Y \in \mathbb{Z}$ of the (n -dimensional) cones associated to the points A, B, P, Q, X, Y and the polynomials $l(\tau)(s), l(\tau_1)(s), l(\tau_2)(s), l(\tau_3)(s)$ of degree one. Then by the argument in Case 1 of the proof of [17, Proposition 14] we obtain an equality

$$Z_{\text{top},f}(s) \equiv \frac{1}{(s+1)l(\tau)} \left\{ \frac{V_X}{l(\tau_1)} + \frac{V_A}{l(\tau_1)l(\tau_2)} + \frac{V_B}{l(\tau_2)l(\tau_3)} + \frac{V_Y}{l(\tau_3)} \right\} \quad (3.31)$$

$$+ \frac{V_P}{l(\tau)l(\tau_1)l(\tau_2)} + \frac{V_Q}{l(\tau)l(\tau_2)l(\tau_3)}. \quad (3.32)$$

modulo holomorphic functions at $s_0 \in \mathbb{C}$. In order to prove the assertion, it suffices to show that the polynomial in the right hand side of the equality

$$(s+1)l(\tau) \left(\prod_{i=1}^3 l(\tau_i) \right) Z_{\text{top},f}(s) \equiv l(\tau_3) \left\{ l(\tau_2)V_X + V_A + (s+1)V_P \right\} \quad (3.33)$$

$$+ l(\tau_1) \left\{ l(\tau_2)V_Y + V_B + (s+1)V_Q \right\} \quad (3.34)$$

is divisible by the factor $l(\tau)(s) = N(\tau)s + \nu(\tau)$. Let K_1 (resp. K'_1) be the multiplicity of the (4-dimensional) cone generated by $a(\rho_1), a(\rho_2), a(\tau_1), a(\tau_2)$ (resp. $a(\rho_1), a(\rho_2), a(\tau), a(\tau_2)$). Then we have the geometric equality

$$K_1 a(\tau) = V_P a(\rho_2) + V_X a(\tau_2) + V_A a(\rho_1) + K'_1 a(\tau_1). \quad (3.35)$$

From this we obtain

$$l(\tau_2)V_X + V_A + (s+1)V_P = l(\tau)K_1 - l(\tau_1)K'_1. \quad (3.36)$$

Moreover let K_2 be the multiplicity of the (4-dimensional) cone generated by $a(\rho_1), a(\rho_2), a(\tau_2), a(\tau_3)$. Then similarly we obtain

$$l(\tau_2)V_Y + V_B + (s+1)V_Q = l(\tau)K_2 + l(\tau_3)K'_1. \quad (3.37)$$

Now the assertion immediately follows. \square

Note that the facet in Proposition 3.4 splits into two B_1 -pyramids for different variables whose intersection does not contain any 1-dimensional V -face. Motivated by Proposition 3.4 we introduce the following definition.

Definition 3.5. *We call the face τ of $\Gamma_+(f)$ in Proposition 3.4 for $n = 4$ a B_2 -facet.*

Lemma 3.6. *For $n = 4$ if a compact facet $\tau \prec \Gamma_+(f)$ is not a B_1 -pyramid nor a B_2 -facet then it splits into lattice simplices so that one of them is not of type B_1 .*

Proof. The facet τ contains a face F not contained in a coordinate hyperplane. Note the following facts about F :

(1): F has at most 4 vertices. Otherwise it contains a triangle whose sides are not in coordinate hyperplanes, and the union of this triangle and any vertex of $\tau \setminus F$ gives a non- B_1 -simplex in τ .

(2): If F is a quadrilateral, then some pair of its opposite edges are contained in coordinate hyperplanes, say, $\{v_1 = 0\}$ and $\{v_2 = 0\}$, otherwise we get the same contradiction as in (1). In this case, if a vertex of F at the hyperplane $\{v_1 = 0\}$ has $v_2 > 1$, then this vertex, the two vertices of $F \cap \{v_2 = 0\}$ and any other vertex of $\tau \setminus F$ form a non- B_1 -simplex in τ . Thus both vertices of F in the hyperplane $\{v_1 = 0\}$ have $v_2 = 1$ and vice versa. Thus τ is a B_2 -facet.

(3): If F is a triangle, then at least one of its edges is contained in a coordinate hyperplane, otherwise we get the same contradiction as in (1).

(3.1): If the triangle F has exactly one edge in a coordinate hyperplane, say, $\{v_1 = 0\}$, then the coordinate v_1 of the other vertex of F equals 1, otherwise F together with any vertex from $\tau \setminus F$ form a non- B_1 -simplex in τ . Also in this case, all other vertices of τ should be in the hyperplane $\{v_1 = 0\}$, because otherwise such vertex together with F form a non- B_1 -simplex in τ . Thus τ is a B_1 -pyramid for v_1 .

(3.2): If the triangle F has all three edges in coordinate hyperplanes, then τ is a B_1 -tetrahedron.

(3.3): The only remaining case is that F is a triangle, exactly two of whose faces are in coordinate hyperplanes. Since its third edge is not in a coordinate hyperplane, then it should be an edge of another 2-dimensional face G of τ not contained in a coordinate hyperplane. If G is also a triangle, exactly two of whose faces are in coordinate hyperplanes, then τ is a B_1 -tetrahedron. If not, then G is of one of the types (2) or (3.1), and thus τ is B_1 or B_2 as shown in the corresponding paragraphs. \square

4 Candidate poles and eigenvalues of monodromies

In this section, we introduce some higher-dimensional analogues of the results in Lemahieu-Van Proeyen [17, Section 3]. Let $f(x)$ be a polynomial on \mathbb{C}^n such that $f(0) = 0$. Also for lattice simplices τ contained in compact facets of $\Gamma_+(f)$ we define a polynomial $\zeta_\tau(t) = (1 - t^{N(\tau)})^{\text{Vol}_\mathbb{Z}(\tau)} \in \mathbb{C}[t]$ similarly and use them freely in this section.

Proposition 4.1. *Let $\tau \prec \Gamma_+(f)$ be a compact facet such that $\gamma = \tau \cap \{v_i = 0\}$ is its facet. Then $\frac{\zeta_\tau}{\zeta_\gamma} \in \mathbb{C}(t)$ is a polynomial of t . If we assume moreover that τ is not a B_1 -pyramid for the variable v_i , then the complex number*

$$\lambda = \exp\left(-2\pi i \frac{\nu(\tau)}{N(\tau)}\right) \in \mathbb{C} \quad (4.1)$$

is a root of the polynomial.

Proof. By Lemma 2.5 we can easily prove that $\frac{\zeta_\tau}{\zeta_\gamma} \in \mathbb{C}(t)$ is a polynomial. Let us prove the remaining assertion. If τ is not a pyramid over $\gamma = \tau \cap \{v_i = 0\}$, then we have $\text{Vol}_{\mathbb{Z}}(\tau) > \text{Vol}_{\mathbb{Z}}(\gamma)$ and the assertion is obvious. So it suffices to consider the case where τ is a pyramid over $\gamma = \tau \cap \{v_i = 0\}$ but its unique vertex $P \prec \tau$ such that $P \notin \gamma$ has height $h \geq 2$ from the hyperplane $\{v_i = 0\} \subset \mathbb{R}_+^n$. In this case, we define two hyperplanes H_τ and L_τ in \mathbb{R}^n by

$$H_\tau = \{v \in \mathbb{R}^n \mid \langle a(\tau), v \rangle = N(\tau)\}, \quad (4.2)$$

$$L_\tau = \{v \in \mathbb{R}^n \mid \langle a(\tau), v \rangle = \left\langle a(\tau), \begin{pmatrix} 1 \\ \vdots \\ 1 \end{pmatrix} \right\rangle = \nu(\tau)\}. \quad (4.3)$$

Note that $P \in \tau \subset H_\tau$ and L_τ is the hyperplane passing through the point $(1, 1, \dots, 1) \in \mathbb{R}_+^n$ and parallel to H_τ . Namely H_τ is the affine span $\text{aff}(\tau) \simeq \mathbb{R}^{n-1}$ of τ . Moreover the affine subspace $L_\tau \cap \{v_i = 0\} \subset \mathbb{R}^n$ is parallel to the affine span $H_\tau \cap \{v_i = 0\} \subset \mathbb{R}^n$ of $\gamma = \tau \cap \{v_i = 0\}$. This implies that $\lambda = \exp(-2\pi i \frac{\nu(\tau)}{N(\tau)}) \in \mathbb{C}$ is a root of $\zeta_\gamma(t)$ if and only if $L_\tau \cap \{v_i = 0\}$ is rational i.e. $L_\tau \cap \{v_i = 0\} \cap \mathbb{Z}^n \neq \emptyset$. On the other hand, it is easy to see that the affine subspace $H_\tau \cap \{v_i = h - 1\} \subset \mathbb{R}^n$ is a parallel translation of $L_\tau \cap \{v_i = 0\}$ by a lattice vector. Hence if $L_\tau \cap \{v_i = 0\}$ is rational, then $H_\tau \cap \{v_i = h - 1\} \cap \mathbb{Z}^n \neq \emptyset$ and the lattice height of the pyramid τ from its base $\gamma = \tau \cap \{v_i = 0\}$ is $h \geq 2$ i.e. $\text{Vol}_{\mathbb{Z}}(\tau) \geq 2\text{Vol}_{\mathbb{Z}}(\gamma)$. It follows that the polynomial $\frac{\zeta_\tau}{\zeta_\gamma}$ is divided by the factor $t - \lambda$. This completes the proof. \square

Motivated by this proposition, we introduce the following definitions.

Definition 4.2. *Let τ be an $(n - 1)$ -dimensional lattice simplex contained in a compact facet of $\Gamma_+(f)$.*

1. *We say that τ has a (non-empty) corner of codimension r if there exist $1 \leq i_1 < i_2 < \dots < i_r \leq n$ such that $\tau \cap \{v_i = 0\}$ is a facet of τ if and only if $i \in \{i_1, i_2, \dots, i_r\}$ and $\tau \cap (\cap_{j=1}^r \{v_{i_j} = 0\}) \neq \emptyset$.*
2. *If τ has a corner of codimension r for $1 \leq i_1 < i_2 < \dots < i_r \leq n$, then we set*

$$F_\tau(t) = \prod_{I \subset \{i_1, i_2, \dots, i_r\}} \left\{ \zeta_{\tau \cap \{v_i = 0\} (i \in I)}(t) \right\}^{(-1)^{|I|}} \in \mathbb{C}(t). \quad (4.4)$$

In the next section we will prove the following result.

Theorem 4.3. *Let τ be an $(n - 1)$ -dimensional lattice simplex contained in a compact facet of $\Gamma_+(f)$. Assume that for some $r \geq 1$ it has a (non-empty) corner of codimension r . Then $F_\tau(t) \in \mathbb{C}(t)$ is a polynomial of t . If we assume moreover that τ is not a B_1 -pyramid, then the complex number*

$$\lambda = \exp\left(-2\pi i \frac{\nu(\tau)}{N(\tau)}\right) \in \mathbb{C} \quad (4.5)$$

is a root of the polynomial.

We can generalize Theorem 4.3 slightly to allow also simplices with empty corners as follows.

Proposition 4.4. *Let $\tau = A_1 A_2 \cdots A_n$ be an $(n - 1)$ -dimensional lattice simplex in \mathbb{R}_+^n . Assume that for any $1 \leq i \leq n$ its vertex A_i is on the positive part of the i -th coordinate axis of \mathbb{R}^n and set*

$$F_\tau(t) = \prod_{\sigma \prec \tau} \left\{ \zeta_\sigma(t) \right\}^{(-1)^{n-1-\dim\sigma}} \in \mathbb{C}(t), \quad (4.6)$$

where in the product $\prod_{\sigma \prec \tau}$ the face $\sigma \prec \tau$ of τ ranges through the non-empty ones. Then

$$F_\tau(t) \cdot (1 - t)^{(-1)^n} \in \mathbb{C}(t) \quad (4.7)$$

is a polynomial. If we assume moreover that τ is not a B_1 -simplex, then the complex number

$$\lambda = \exp\left(-2\pi i \frac{\nu(\tau)}{N(\tau)}\right) \in \mathbb{C} \quad (4.8)$$

is a root of the polynomial.

Proof. By the embedding $\mathbb{R}^n \hookrightarrow \mathbb{R}^n \times \mathbb{R}$, $v \mapsto (v, 0)$ we regard τ as a lattice simplex in $\mathbb{R}^n \times \mathbb{R}$ and set $Q(0, 1) \in \mathbb{R}^n \times \mathbb{R}$. Let τ' be the convex hull of $\{Q\} \cup \tau$ in $\mathbb{R}^n \times \mathbb{R}$. Then by considering the vertex $Q \prec \tau'$ as the corner of the simplex τ' we define a polynomial $F_{\tau'}(t) \in \mathbb{C}[t]$ by Theorem 4.3. In this situation, it is easy to see that we have an equality

$$F_{\tau'}(t) = F_\tau(t) \cdot (1 - t)^{(-1)^n} \quad (4.9)$$

from which the first assertion immediately follows. Since we have $N(\tau') = N(\tau)$ and $\nu(\tau') = \nu(\tau) + N(\tau)$, the second assertion also follows from Theorem 4.3. \square

Together with Proposition 3.2, following the strategy of Lemahieu-Van Proeyen [17] we can now confirm the monodromy conjecture for non-degenerate hypersurfaces in many cases also for $n \geq 4$. In the case $n = 4$ there are some B_2 -facets of $\Gamma_+(f)$ (along the intersections of two coordinate hyperplanes in \mathbb{R}_+^4) which are not B_1 -pyramids but are divided into two B_1 -pyramids. See Proposition 3.4 for the details. Their contributions to the monodromy zeta function $\zeta_{f,0}(t)$ are trivial.

5 The proof of Theorem 4.3

5.1 Preliminaries

For the proof of Theorem 4.3 we shall introduce some new notions and their basic properties. Let S be a finite set and denote its power set by 2^S . Namely elements of 2^S are subsets $I \subset S$ of S . Then for a function $\phi : 2^S \rightarrow \mathbb{Z}$ we define new ones $\phi^\downarrow, \phi^\uparrow : 2^S \rightarrow \mathbb{Z}$ by

$$\phi^\downarrow(I) = \sum_{J \subset I} \phi(J), \quad \phi^\uparrow(I) = \sum_{J \subset I} (-1)^{|I|-|J|} \phi(J). \quad (5.1)$$

We call ϕ^\downarrow (resp. ϕ^\uparrow) the antiderivative (resp. derivative) of ϕ . Then we can easily check that $\phi^{\uparrow\downarrow} = \phi^{\downarrow\uparrow} = \phi$.

Definition 5.1. (i) We say that the function ϕ is fully supermodular if $\phi^\uparrow(I) \geq 0$ for any subset $I \subset S$.

(ii) The function ϕ is called strictly fully supermodular if it is fully supermodular and $\phi^\uparrow(S) > 0$.

Lemma 5.2. The product of two fully supermodular functions $\phi, \psi : 2^S \rightarrow \mathbb{Z}$ is fully supermodular. Moreover it is strictly fully supermodular if and only if there exist subsets $I, J \subset S$ of S such that $I \cup J = S$ and $\phi^\uparrow(I), \psi^\uparrow(J) > 0$.

Proof. For any subset $R \subset S$ of S we have

$$(\phi\psi)^\uparrow(R) = (\phi^{\uparrow\downarrow}\psi^{\uparrow\downarrow})^\uparrow(R) = \sum_{I \cup J \subset R} (-1)^{|R|-|I|-|J|} \phi^\uparrow(I) \psi^\uparrow(J) \quad (5.2)$$

$$= \sum_{I \cup J = R} \phi^\uparrow(I) \psi^\uparrow(J). \quad (5.3)$$

Then the assertion immediately follows. \square

5.2 Reduction to the case $r = n - 1$

First we shall reduce the proof of Theorem 4.3 to the case $r = n - 1$. For simplicity assume that the corner $\gamma \prec \tau$ of the simplex $\tau \subset \partial\Gamma_+(f)$ is defined by $\gamma = \tau \cap \{v_1 = v_2 = \dots = v_r = 0\}$. We set

$$\lambda = \exp\left(-2\pi i \frac{\nu(\tau)}{N(\tau)}\right) \in \mathbb{C}. \quad (5.4)$$

As in the proof of Proposition 4.1 we define two parallel affine hyperplanes H_τ and L_τ in \mathbb{R}^n by

$$H_\tau = \{v \in \mathbb{R}^n \mid \langle a(\tau), v \rangle = N(\tau)\}, \quad (5.5)$$

$$L_\tau = \{v \in \mathbb{R}^n \mid \langle a(\tau), v \rangle = \left\langle a(\tau), \begin{pmatrix} 1 \\ \vdots \\ 1 \end{pmatrix} \right\rangle = \nu(\tau)\}. \quad (5.6)$$

Let $W = \{v_1 = v_2 = \dots = v_r = 0\} \simeq \mathbb{R}^{n-r} \subset \mathbb{R}^n$ be the linear subspace of \mathbb{R}^n spanned by γ . Similarly, for a face σ of τ containing γ let $W_\sigma \simeq \mathbb{R}^{\dim\sigma+1} \subset \mathbb{R}^n$ be the linear subspace

of \mathbb{R}^n spanned by σ . Then $\zeta_\sigma(\lambda) = 0$ if and only if the affine hyperplane $L_\tau \cap W_\sigma \subset W_\sigma$ of W_σ is rational i.e. $L_\tau \cap W_\sigma \cap \mathbb{Z}^n \neq \emptyset$. Let $\Phi_0 : W \xrightarrow{\sim} W$ be a unimodular transformation of W such that $\Phi_0(\gamma) \subset W \cap \{v_n = k\}$ for some $k \in \mathbb{Z}_{>0}$. Then we can easily extend it to a unimodular transformation $\Phi : \mathbb{R}^n \xrightarrow{\sim} \mathbb{R}^n$ of \mathbb{R}^n which preserves W_σ for any $\sigma \prec \tau$ containing γ and the point

$$\begin{pmatrix} 1 \\ 1 \\ \vdots \\ 1 \end{pmatrix} \in \mathbb{R}^n. \quad (5.7)$$

We can choose such Φ so that the heights of τ and $\Phi(\tau)$ from each coordinate hyperplane in \mathbb{R}^n containing W are the same. Indeed, for the invertible matrix $A_0 \in \mathrm{GL}_{n-r}(\mathbb{Z})$ representing $\Phi_0 : W \xrightarrow{\sim} W$ it suffices to define $\Phi : \mathbb{R}^n \xrightarrow{\sim} \mathbb{R}^n$ by taking an invertible matrix $A \in \mathrm{GL}_n(\mathbb{Z})$ of the form

$$A = \left(\begin{array}{c|c} I_r & 0 \\ * & A_0 \end{array} \right) \in \mathrm{GL}_n(\mathbb{Z}) \quad (5.8)$$

such that

$$A \begin{pmatrix} 1 \\ \vdots \\ 1 \end{pmatrix} = \begin{pmatrix} 1 \\ \vdots \\ 1 \end{pmatrix}, \quad (5.9)$$

where $I_r \in \mathrm{GL}_r(\mathbb{Z})$ stands for the identity matrix of size r . By this construction of Φ , τ is a B_1 -pyramid if and only if $\Phi(\tau)$ is so. Set $\tau' = \Phi(\tau)$ and define two parallel affine hyperplanes $H_{\tau'}$ and $L_{\tau'}$ in \mathbb{R}^n similarly to the case of τ so that we have $\Phi(H_\tau) = H_{\tau'}$. Since $\Phi(L_\tau)$ is parallel to $\Phi(H_\tau) = H_{\tau'}$ and passes through the point

$$\Phi \left(\begin{pmatrix} 1 \\ \vdots \\ 1 \end{pmatrix} \right) = \begin{pmatrix} 1 \\ \vdots \\ 1 \end{pmatrix} \in \mathbb{R}^n, \quad (5.10)$$

we have also $\Phi(L_\tau) = L_{\tau'}$. Now let us set

$$\lambda' = \exp \left(-2\pi i \frac{\nu(\tau')}{N(\tau')} \right) \in \mathbb{C}. \quad (5.11)$$

Then for any $\sigma \prec \tau$ containing γ we have $\zeta_\sigma(\lambda) = 0 \iff \zeta_{\Phi(\sigma)}(\lambda') = 0$. Since the lattice distance $N(\sigma) > 0$ of σ (resp. $N(\Phi(\sigma)) > 0$ of $\Phi(\sigma)$) from the origin $0 \in W_\sigma \simeq \mathbb{R}^{\dim \sigma + 1}$ is equal to the number of rational hyperplanes in $W_\sigma \simeq \mathbb{R}^{\dim \sigma + 1}$ parallel to σ (resp. $\Phi(\sigma)$) between $\mathrm{aff}(\sigma)$ (resp. $\mathrm{aff}(\Phi(\sigma))$) and the origin, we have $N(\sigma) = N(\Phi(\sigma))$ and hence $\zeta_\sigma(t) \equiv \zeta_{\Phi(\sigma)}(t)$. Then we obtain an equality $F_\tau(t) = F_{\tau'}(t)$, where we slightly generalized Definition 4.2 in an obvious way to define $F_{\tau'}(t)$. Hence, to prove Theorem 4.3 we may assume that the corner γ of τ is contained in $W \cap \{v_n = k\}$ for some $k \in \mathbb{Z}_{>0}$. Let $\pi : \mathbb{R}^n \rightarrow \mathbb{R}^{r+1}$, $v \mapsto (v_1, \dots, v_r, v_n)$ be the projection. Then by the definition of normalized volumes, for any face σ of τ containing its corner $\gamma \subset W \cap \{v_n = k\}$ we have $\mathrm{Vol}_{\mathbb{Z}}(\sigma) = \mathrm{Vol}_{\mathbb{Z}}(\pi(\sigma)) \cdot \mathrm{Vol}_{\mathbb{Z}}(\gamma)$ and hence $\zeta_\sigma(t) = \{\zeta_{\pi(\sigma)}(t)\}^{\mathrm{Vol}_{\mathbb{Z}}(\gamma)}$. We thus obtain an equality $F_\tau(t) = \{F_{\pi(\tau)}(t)\}^{\mathrm{Vol}_{\mathbb{Z}}(\gamma)}$. Moreover we have $N(\tau) = N(\pi(\tau))$ and $\nu(\tau) = \nu(\pi(\tau))$. This implies that we have only to consider the case $r = n - 1$.

5.3 The proof of the case $r = n - 1$

We have reduced our proof to the case where $r = n - 1$, a vertex Q of our simplex $\tau = QA_1A_2 \cdots A_{n-1}$ has the form $Q = (0, 0, \dots, 0, k)$ for some $k \in \mathbb{Z}_{>0}$ and its edges are given by

$$\overrightarrow{QA_1} = \begin{pmatrix} a_1 \\ 0 \\ \vdots \\ 0 \\ b_1 \end{pmatrix}, \quad \overrightarrow{QA_2} = \begin{pmatrix} 0 \\ a_2 \\ \vdots \\ 0 \\ b_2 \end{pmatrix}, \quad \dots, \quad \overrightarrow{QA_{n-1}} = \begin{pmatrix} 0 \\ 0 \\ \vdots \\ a_{n-1} \\ b_{n-1} \end{pmatrix}, \quad (5.12)$$

where $a_1, a_2, \dots, a_{n-1} \in \mathbb{Z}_{>0}$ and $b_1, b_2, \dots, b_{n-1} \in \mathbb{Z}$. We set

$$D = \prod_{i=1}^{n-1} a_i, \quad K_i = \frac{b_i}{a_i} \cdot D \quad (1 \leq i \leq n-1) \quad (5.13)$$

and $K = \sum_{i=1}^{n-1} K_i$. Moreover for a subset $I \subset S = \{1, 2, \dots, n-1\}$ we denote by $\tau_I \prec \tau$ the face of τ whose vertices are Q and A_i ($i \in I$) and set

$$D_I = \prod_{i \in I} a_i, \quad \gcd_I = \text{GCD}(D, K_i \ (i \in I)). \quad (5.14)$$

Then it is easy to see that the $|I|$ -dimensional normalized volume $\text{Vol}_{\mathbb{Z}}(\tau_I)$ of τ_I is given by the formula

$$\text{Vol}_{\mathbb{Z}}(\tau_I) = \gcd_I \cdot \frac{D_I}{D} \quad (5.15)$$

and hence we have

$$N(\tau_I) = \frac{D_I}{\text{Vol}_{\mathbb{Z}}(\tau_I)} \cdot k = \frac{D}{\gcd_I} \cdot k. \quad (5.16)$$

In particular, we obtain

$$N(\tau) = \frac{D}{\gcd_S} \cdot k. \quad (5.17)$$

For a subset $I \subset S = \{1, 2, \dots, n-1\}$ we set

$$\zeta_I(t) = \left\{ 1 - t^{N(\tau_I)} \right\}^{\text{Vol}_{\mathbb{Z}}(\tau_I)} \in \mathbb{C}[t] \quad (5.18)$$

so that we have the equality

$$F_{\tau}(t) = \prod_{I \subset S} \left\{ \zeta_I(t) \right\}^{(-1)^{n-1-|I|}}. \quad (5.19)$$

Lemma 5.3. *The complex number*

$$\lambda = \exp\left(-2\pi i \frac{\nu(\tau)}{N(\tau)}\right) \in \mathbb{C} \quad (5.20)$$

is a root of the polynomial $\zeta_I(t)$ if and only if $\gcd_I | K$.

Proof. Since the primitive conormal vector of the $(n - 1)$ -dimensional simplex τ is equal to

$$a(\tau) = \frac{1}{\gcd_S} \begin{pmatrix} -K_1 \\ -K_2 \\ \vdots \\ -K_{n-1} \\ D \end{pmatrix}, \quad (5.21)$$

we have

$$\nu(\tau) = \frac{D - K}{\gcd_S} \quad (5.22)$$

and

$$\frac{\nu(\tau)N(\tau_I)}{N(\tau)} = \frac{D - K}{\gcd_I}. \quad (5.23)$$

Note that λ is a root of $\zeta_I(t)$ if and only if $\frac{\nu(\tau)N(\tau_I)}{N(\tau)}$ is an integer. Then the assertion follows immediately from (5.23) and the fact $\gcd_I|D$. This completes the proof. \square

By this lemma the multiplicity of $t - \lambda$ in the rational function $Z_\tau(t)$ is equal to

$$\sum_{I: \gcd_I|K} (-1)^{n-1-|I|} \gcd_I \cdot \frac{D_I}{D}. \quad (5.24)$$

Similarly we obtain the following result.

Lemma 5.4. *For any $m \in \mathbb{Z}$ the complex number $\exp(2\pi i \frac{m}{N(\tau)}) \in \mathbb{C}$ is a root of the polynomial $\zeta_I(t)$ if and only if $\gcd_I|(m \cdot \gcd_S)$.*

Proposition 5.5. *The function $F_\tau(t)$ is a polynomial in t .*

Proof. By Lemma 5.4 it suffices to show that for any $m \in \mathbb{Z}$ the alternating sum

$$G_m = \sum_{I: \gcd_I|(m \cdot \gcd_S)} (-1)^{n-1-|I|} \gcd_I \cdot \frac{D_I}{D} \quad (5.25)$$

is non-negative. Fix $m \in \mathbb{Z}$ and for a prime number p denote its multiplicities in the prime decompositions of a_i, b_i and m by $\alpha(p)_i, \beta(p)_i$ and $\delta(p)$ respectively. We set

$$\gamma(p) = \delta(p) + \min_{1 \leq i \leq n-1} \{\beta(p)_i - \alpha(p)_i, 0\} \quad (5.26)$$

and define a function $\phi_p : 2^S \rightarrow \mathbb{Z}$ by

$$\phi_p(I) = \begin{cases} p^{\min_{i \in I} \{\beta(p)_i - \alpha(p)_i, 0\} + \sum_{i \in I} \alpha(p)_i} & \left(\min_{i \in I} \{\beta(p)_i - \alpha(p)_i, 0\} \leq \gamma(p) \right), \\ 0 & \text{(otherwise).} \end{cases} \quad (5.27)$$

Then it is easy to see that for the function $\phi = \prod_{p: \text{prime}} \phi_p : 2^S \rightarrow \mathbb{Z}$ we have

$$\phi^\uparrow(S) = G_m. \quad (5.28)$$

By Lemma 5.2 we have only to prove that for any prime number p the function $\phi_p : 2^S \rightarrow \mathbb{Z}$ is fully supermodular. For this purpose, we reorder the pairs (a_i, b_i) ($1 \leq i \leq n-1$) so that we have

$$\beta(p)_1 - \alpha(p)_1 \leq \beta(p)_2 - \alpha(p)_2 \leq \cdots \leq \beta(p)_{n-1} - \alpha(p)_{n-1}. \quad (5.29)$$

Fix a subset $I = \{i_1, i_2, i_3, \dots\} \subset S = \{1, 2, \dots, n-1\}$ ($i_1 < i_2 < i_3 < \dots$) of S . We will show the non-negativity of the alternating sum

$$\phi_p^\uparrow(I) = \sum_{J \subset I} (-1)^{|I|-|J|} \phi_p(J). \quad (5.30)$$

We define $q \geq 0$ to be the maximal number such that $\beta(p)_{i_q} - \alpha(p)_{i_q} < 0$ (resp. $\beta(p)_{i_q} - \alpha(p)_{i_q} \leq \gamma(p)$) in the case $\gamma(p) \geq 0$ (resp. $\gamma(p) < 0$). First let us consider the case $\gamma(p) \geq 0$. Then for $1 \leq l \leq q$ the part of the alternating sum (5.30) over the subsets $J \subset I$ such that $\min J = i_l$ is equal to

$$(-1)^{l-1} p^{\beta(p)_{i_l}} \prod_{j>l} (p^{\alpha(p)_{i_j}} - 1). \quad (5.31)$$

Moreover the remaining part of (5.30) is equal to

$$(-1)^q \prod_{j>q} (p^{\alpha(p)_{i_j}} - 1). \quad (5.32)$$

We thus obtain the equality

$$\phi_p^\uparrow(I) = p^{\beta(p)_{i_1}} \prod_{j>1} (p^{\alpha(p)_{i_j}} - 1) - p^{\beta(p)_{i_2}} \prod_{j>2} (p^{\alpha(p)_{i_j}} - 1) + \cdots \quad (5.33)$$

$$\cdots + (-1)^{q-1} p^{\beta(p)_{i_q}} \prod_{j>q} (p^{\alpha(p)_{i_j}} - 1) + (-1)^q \prod_{j>q} (p^{\alpha(p)_{i_j}} - 1). \quad (5.34)$$

Note that for any $1 \leq j \leq q$ we have $\beta(p)_{i_j} - \alpha(p)_{i_j} < 0$ and obtain an inequality

$$p^{\beta(p)_{i_{j-1}}} (p^{\alpha(p)_{i_j}} - 1) \geq p^{\alpha(p)_{i_j}} - 1 \geq p^{\beta(p)_{i_j}}. \quad (5.35)$$

Thus, subdividing the terms in the above expression of $\phi_p^\uparrow(I)$ into pairs, we get the desired non-negativity $\phi_p^\uparrow(I) \geq 0$. Finally let us consider the case $\gamma(p) < 0$. In this case, we have the following expression of $\phi_p^\uparrow(I)$:

$$\phi_p^\uparrow(I) = p^{\beta(p)_{i_1}} \prod_{j>1} (p^{\alpha(p)_{i_j}} - 1) - p^{\beta(p)_{i_2}} \prod_{j>2} (p^{\alpha(p)_{i_j}} - 1) + \cdots + (-1)^{q-1} p^{\beta(p)_{i_q}} \prod_{j>q} (p^{\alpha(p)_{i_j}} - 1). \quad (5.36)$$

Then by using the inequality (5.35) we can prove the non-negativity $\phi_p^\uparrow(I) \geq 0$ as in the previous case $\gamma(p) \geq 0$. This completes the proof. \square

Proposition 5.6. *Assume that τ is not a B_1 -simplex. Then the complex number*

$$\lambda = \exp\left(-2\pi i \frac{\nu(\tau)}{N(\tau)}\right) \in \mathbb{C} \quad (5.37)$$

is a root of the polynomial $F_\tau(t)$.

Proof. By Lemma 5.3 it suffices to show that the alternating sum

$$G = \sum_{I: \gcd_I | K} (-1)^{n-1-|I|} \gcd_I \cdot \frac{D_I}{D} \quad (5.38)$$

is positive. For a prime number p denote its multiplicities in the prime decompositions of a_i, b_i and K by $\alpha(p)_i, \beta(p)_i$ and $\kappa(p)$ respectively. We set

$$\mu(p) = \kappa(p) - \sum_{i=1}^{n-1} \alpha(p)_i \quad (5.39)$$

and define a function $\psi_p : 2^S \rightarrow \mathbb{Z}$ by

$$\psi_p(I) = \begin{cases} p^{\min_{i \in I} \{\beta(p)_i - \alpha(p)_i, 0\} + \sum_{i \in I} \alpha(p)_i} & \left(\min_{i \in I} \{\beta(p)_i - \alpha(p)_i, 0\} \leq \mu(p) \right), \\ 0 & \text{(otherwise).} \end{cases} \quad (5.40)$$

Then it is easy to see that for the function $\psi = \prod_{p: \text{prime}} \psi_p : 2^S \rightarrow \mathbb{Z}$ we have

$$\psi^\uparrow(S) = G. \quad (5.41)$$

Now let us set $S_p = \{1 \leq i \leq n-1 \mid \alpha(p)_i = 0\}$ and $I_p = S \setminus S_p = \{1 \leq i \leq n-1 \mid \alpha(p)_i > 0\}$. By our assumption we have $a_i > 1$ for any $1 \leq i \leq n-1$ and hence $\cup_{p: \text{prime}} I_p = S$. By Lemma 5.2, in order to show the positivity $\psi^\uparrow(S) > 0$ it suffices to prove that for any prime number p we have $\psi_p^\uparrow(I_p) > 0$. As in the proof of Proposition 5.5 we reorder the pairs (a_i, b_i) ($1 \leq i \leq n-1$) so that we have

$$\beta(p)_1 - \alpha(p)_1 \leq \beta(p)_2 - \alpha(p)_2 \leq \dots \leq \beta(p)_{n-1} - \alpha(p)_{n-1} \quad (5.42)$$

and $\alpha(p)_i \geq \alpha(p)_{i+1}$ whenever $\beta(p)_i - \alpha(p)_i = \beta(p)_{i+1} - \alpha(p)_{i+1}$. Moreover we set $I_p = \{i_1, i_2, i_3, \dots\}$ ($i_1 < i_2 < i_3 < \dots$). We define $q \geq 0$ to be the maximal number such that $\beta(p)_{i_q} - \alpha(p)_{i_q} < 0$ (resp. $\beta(p)_{i_q} - \alpha(p)_{i_q} \leq \mu(p)$) in the case $\mu(p) \geq 0$ (resp. $\mu(p) < 0$). Then we have the same expressions of $\psi_p^\uparrow(I_p) > 0$ as in the proof of Proposition 5.5. In the case $\mu(p) \geq 0$ we have

$$\psi_p^\uparrow(I_p) = p^{\beta(p)_{i_1}} \prod_{j>1} (p^{\alpha(p)_{i_j}} - 1) - p^{\beta(p)_{i_2}} \prod_{j>2} (p^{\alpha(p)_{i_j}} - 1) + \dots \quad (5.43)$$

$$\dots + (-1)^{q-1} p^{\beta(p)_{i_q}} \prod_{j>q} (p^{\alpha(p)_{i_j}} - 1) + (-1)^q \prod_{j>q} (p^{\alpha(p)_{i_j}} - 1). \quad (5.44)$$

In the case $\mu(p) < 0$ we have

$$\psi_p^\uparrow(I_p) = p^{\beta(p)_{i_1}} \prod_{j>1} (p^{\alpha(p)_{i_j}} - 1) - p^{\beta(p)_{i_2}} \prod_{j>2} (p^{\alpha(p)_{i_j}} - 1) + \dots + (-1)^{q-1} p^{\beta(p)_{i_q}} \prod_{j>q} (p^{\alpha(p)_{i_j}} - 1). \quad (5.45)$$

By the definitions of $I_p = \{i_1, i_2, i_3, \dots\} \subset S$ and $q \geq 0$ we have $i \in I_p$ for any $i \leq i_q$. Eventually we find that $i_j = j$ for any $j \leq q$. First let us consider the case $I_p = \emptyset$. Then we have

$$\psi_p(I_p) = \begin{cases} p^0 = 1 & (\mu(p) \geq 0), \\ 0 & (\mu(p) < 0). \end{cases} \quad (5.46)$$

In the case $\mu(p) \geq 0$ we thus obtain the positivity $\psi_p^\dagger(I_p) > 0$. But in the case $\mu(p) < 0$ the condition $I_p = \emptyset$ implies $q = 0$ and such a case cannot occur by the following lemma.

Lemma 5.7. *The case $q = 0$ and $\mu(p) < 0$ cannot occur.*

Proof. Assume that $q = 0$ and $\mu(p) < 0$. By the definition of $\mu(p)$ we have

$$\text{mult}_p(K) = \text{mult}_p(D \cdot p^{\mu(p)}). \quad (5.47)$$

On the other hand, it follows from the condition $\mu(p) < 0$ and $q = 0$ that

$$\mu(p) < \beta(p)_{i_1} - \alpha(p)_{i_1} \leq \beta(p)_{i_2} - \alpha(p)_{i_2} \leq \dots \dots \dots \quad (5.48)$$

Namely for any $i \in I_p$ we have $\mu(p) < \beta(p)_i - \alpha(p)_i$ and hence

$$\text{mult}_p(K_i) > \text{mult}_p(D \cdot p^{\mu(p)}). \quad (5.49)$$

Moreover for any $i \in S_p = S \setminus I_p$ we have

$$\text{mult}_p(K_i) \geq \text{mult}_p(D) > \text{mult}_p(D \cdot p^{\mu(p)}), \quad (5.50)$$

where we used the condition $\mu(p) < 0$ in the second inequality. We thus obtain the inequality

$$\text{mult}_p(K) = \text{mult}_p\left(\sum_{i \in S} K_i\right) > \text{mult}_p(D \cdot p^{\mu(p)}) \quad (5.51)$$

which contradicts (5.47). □

By this lemma, it remains for us to treat the case $I_p \neq \emptyset$. From now on, we assume that $I_p \neq \emptyset$. Note that the inequality (5.35) becomes an equality only in the case $p = 2$, $\beta(p)_{i_{j-1}} = \beta(p)_{i_j} = 0$ and $\alpha(p)_{i_j} = 1$. By Lemma 5.7 this means that the sums (5.43) and (5.45) may be zero only in the following two cases:

Case 1: $p = 2$, $\mu(p) \geq 0$, $q = 2m + 1$ for $m \geq 0$ and $(\alpha(p)_1, \beta(p)_1) = (a, 0)$ for $a > 0$, $(\alpha(p)_2, \beta(p)_2) = \dots \dots \dots = (\alpha(p)_q, \beta(p)_q) = (1, 0)$.

Case 2: $p = 2$, $\mu(p) < 0$, $q = 2m$ for $m \geq 1$ and $(\alpha(p)_1, \beta(p)_1) = (a, 0)$ for $a > 0$, $(\alpha(p)_2, \beta(p)_2) = \dots \dots \dots = (\alpha(p)_q, \beta(p)_q) = (1, 0)$.

Indeed, in the case $\mu(p) \geq 0$ and $q = 2m$ for $m \geq 0$, if $q < |I_p|$ the last term $(-1)^q \prod_{j>q} (p^{\alpha(p)_{i_j}} - 1)$ of the alternating sum (5.43) is positive. Even if $q = |I_p|$ we still have the positivity

$$(-1)^q \prod_{j>q} (p^{\alpha(p)_{i_j}} - 1) = \psi_p(\emptyset) = 1 > 0. \quad (5.52)$$

Let us show that none of the above two cases can occur.

Case 1: Set $\alpha(p) = \sum_{i \in S} \alpha(p)_i$. Then $2^{\alpha(p)} | D$ and for any $i \in S_2$ we have $2^{\alpha(p)} | K_i$. We thus obtain the equality

$$K \equiv 2^{\alpha(p)-a} \cdot \text{odd} + (q-1) \cdot 2^{\alpha(p)-1} \cdot \text{odd} + \sum_{j>q} 2^{\alpha(p)+\beta(p)_{i_j}-\alpha(p)_{i_j}} \quad (5.53)$$

$$\equiv 2^{\alpha(p)-a} \cdot \text{odd} + (q-1) \cdot 2^{\alpha(p)-1} \cdot \text{odd} \equiv 2^{\alpha(p)-a} \cdot \text{odd} \quad (5.54)$$

of mod $2^{\alpha(p)}$, where we used also the fact that $\beta(p)_{i_j} - \alpha(p)_{i_j} \geq 0$ for any $j > q$. We conclude that $2^{\alpha(p)}$ does not divide K , which contradicts our assumption $\mu(p) \geq 0$.

Case 2: By the condition $q \geq 2$ we have $-1 = \beta(p)_{i_2} - \alpha(p)_{i_2} \leq \mu(p)$. Then by $\mu(p) < 0$ we obtain $\mu(p) = -1$. As in **Case 1**, by using the fact that $q-1$ is odd and $\mu(p) = -1$, if $a = 1$ we obtain the equality

$$K \equiv \sum_{j>q} 2^{\alpha(p)+\beta(p)_{i_j}-\alpha(p)_{i_j}} \equiv 0 \quad (5.55)$$

of mod $2^{\alpha(p)}$. But this result $2^{\alpha(p)} | K$ contradicts our assumption $\mu(p) < 0$. If $a > 1$ we obtain the equality

$$K \equiv 2^{\alpha(p)-a} \cdot \text{odd} \quad (5.56)$$

of mod $2^{\alpha(p)-1}$. But it also contradicts $\mu(p) = -1$.

This completes the proof. \square

6 Some auxiliary results

In this section, to reduce the problem of non-convenient polynomials to that of convenient ones we prepare some technical results. Especially we have the following generalization of [17, Lemma 9].

Proposition 6.1. *Assume that f is non-degenerate at the origin $0 \in \mathbb{C}^n$. Then except for finitely many $c \in \mathbb{C}$ the polynomial $f(x_1, x_2, \dots, x_{n-1}, x_n + c)$ is non-degenerate at the origin $0 \in \mathbb{C}^n$.*

Proof. Let $\tau' \prec \Gamma_+(f)$ be a face of $\Gamma_+(f)$ which is non-compact for the variable v_n and denote its image by the projection $\mathbb{R}^n \rightarrow \mathbb{R}^{n-1}$ by $\sigma \subset \mathbb{R}^{n-1}$. Assume that σ is compact. Here we shall treat only the case where τ' is a facet and hence $\dim \sigma = n-2$. The other cases can be treated similarly. By a unimodular transformation of \mathbb{R}^n we regard τ' as a lattice polytope in its affine span $\text{aff}(\tau') \simeq \mathbb{R}^{n-1}$ and the τ' -part $f_{\tau'}$ of f as a Laurent polynomial on $T' = (\mathbb{C}^*)^{n-1}$. Then by our assumption for any compact face τ of τ' the hypersurface $\{f_{\tau} = 0\} \subset T'$ is smooth and reduced. Moreover the σ -part of the polynomial $f(x_1, x_2, \dots, x_{n-1}, x_n + c)$ is naturally identified with $f_{\tau'}(x_1, \dots, x_{n-2}, c)$. Therefore, in order to prove the assertion, it suffices to show that except for finitely many $c \in \mathbb{C}$ the hypersurface

$$W_c = \{(x_1, \dots, x_{n-2}) \mid f_{\tau'}(x_1, \dots, x_{n-2}, c) = 0\} \subset (\mathbb{C}^*)^{n-2} \quad (6.1)$$

in $T' \cap \{x_{n-1} = c\} \simeq (\mathbb{C}^*)^{n-2}$ is smooth and reduced. Let $h : T' = (\mathbb{C}^*)^{n-1} \rightarrow \mathbb{C}$ be the function defined by $h(x_1, \dots, x_{n-1}) = x_{n-1}$. Then the set of $c \in \mathbb{C}$ for which $W_c \subset (\mathbb{C}^*)^{n-2}$ is not smooth or not reduced is contained in the discriminant variety of the map $h|_{\{f_{\tau'}=0\}} : \{f_{\tau'} = 0\} \rightarrow \mathbb{C}$. For $\varepsilon > 0$ let $B(0; \varepsilon)^* = \{c \in \mathbb{C} \mid 0 < |c| < \varepsilon\}$ be the punctured disk centered at the origin $0 \in \mathbb{C}$. Then there exists a sufficiently small $0 < \varepsilon \ll 1$ such that the hypersurface $W_c \subset (\mathbb{C}^*)^{n-2}$ is smooth and reduced for any $c \in B(0; \varepsilon)^*$. Indeed, let $\Delta = \tau' \cap \{v_n \leq l\} \subset \tau'$ ($l \gg 0$) be the truncation of τ' . Let Σ_0 be the dual fan of the $(n-1)$ -dimensional polytope Δ in \mathbb{R}^{n-1} and Σ its smooth subdivision. We denote by X_Σ the toric variety associated to Σ (see [11] and [26] etc.). Then X_Σ is a smooth compactification of $T' = (\mathbb{C}^*)^{n-1}$. Recall that $T' = (\mathbb{C}^*)^{n-1}$ acts naturally on X_Σ and the T' -orbits in it are parametrized by the cones in the smooth fan Σ . For a cone $C \in \Sigma$ denote by $T_C \simeq (\mathbb{C}^*)^{n-1-\dim C} \subset X_\Sigma$ the T' -orbit associated to C . By our assumption above, if $C \in \Sigma$ corresponds to a compact face τ of τ' then the hypersurface $W = \overline{\{f_{\tau'} = 0\}} \subset X_\Sigma$ intersects $T_C \subset X_\Sigma$ transversally. We denote the meromorphic extension of $h : T' = (\mathbb{C}^*)^{n-1} \rightarrow \mathbb{C}$ to X_Σ by the same letter h . Note that h has no point of indeterminacy on the whole X_Σ (because it is a monomial). Then as $|c| \rightarrow 0$ the level set $h^{-1}(c) \subset X_\Sigma$ of h tends to the union of the T' -orbits which correspond to the compact faces of τ' . More precisely, if a cone $C \in \Sigma$ corresponds to a compact face of τ' then there exists an affine chart \mathbb{C}_y^{n-1} of X_Σ on which

$$T_C = \left\{ y = (y_1, \dots, y_{n-1}) \mid y_i = 0 \ (1 \leq i \leq \dim C), \ y_i \neq 0 \ (\dim C + 1 \leq i \leq n-1) \right\} \quad (6.2)$$

and $h(y) = y_1^{m_1} y_2^{m_2} \cdots y_k^{m_k}$ ($m_i \in \mathbb{Z}_{>0}$) for some $k \geq 1$. By this explicit description of h we see that for $0 < |c| \ll 1$ the hypersurface $h^{-1}(c)$ intersects W transversally. It follows that

$$W_c = W \cap h^{-1}(c) \cap T' \subset h^{-1}(c) \cap T' \simeq (\mathbb{C}^*)^{n-2} \quad (6.3)$$

is smooth and reduced for $0 < |c| \ll 1$. This completes the proof. \square

7 The monodromy conjecture for $n = 4$

In this section, we will prove the monodromy conjecture for non-degenerate polynomials of four variables under an additional assumption. From now on, let us consider the case $n = 4$. Assume that $f(x) \in \mathbb{C}[x_1, \dots, x_4]$ is non-degenerate at the origin $0 \in \mathbb{C}^4$. After subdividing the compact facets of $\Gamma_+(f)$ into 3-dimensional lattice simplices, we have the following lemmas. Also for such simplices τ we define their V -faces (see Definition 2.4) and the integers $N(\tau)$, $\nu(\tau)$ etc. in an obvious way.

Definition 7.1. *Let τ be a 3-dimensional lattice simplex in a compact facet of $\Gamma_+(f)$ and σ a V -face in $\Gamma_+(f)$. Then we say that σ contributes with respect to τ if the complex number*

$$\lambda = \exp\left(-2\pi i \frac{\nu(\tau)}{N(\tau)}\right) \in \mathbb{C} \quad (7.1)$$

is a root of the polynomial $\zeta_\sigma(t)$.

Lemma 7.2. *Let $\tau = APQR$ be a 3-dimensional lattice simplex in a compact facet of $\Gamma_+(f)$ such that the vertex $v = A \prec \tau$ and τ itself are its only contributing V -faces with*

respect to τ . Assume that $\text{Vol}_{\mathbb{Z}}(\tau) = 1$. Then there exists another 3-dimensional lattice simplex $\sigma \neq \tau$ in a compact facet of $\Gamma_+(f)$ containing $v = A$ which has no V -face γ containing $v = A$ such that $\gamma \neq \sigma$ and $\gamma \neq v = A$.

Proof. Let us denote $A(\alpha, 0, 0, 0)$, $P(p_0, p_1, p_2, p_3)$, $Q(q_0, q_1, q_2, q_3)$ and $R(r_0, r_1, r_2, r_3)$. We will show that $p_1 = q_2 = r_3 = 0$ contradicts our assumption $\text{Vol}_{\mathbb{Z}}(\tau) = 1$. This implies the existence of at least one more 3-dimensional lattice simplex $\sigma \neq \tau$ in the star of $v = A$ which has no V -face γ containing $v = A$ such that $\gamma \neq \sigma$ and $\gamma \neq v = A$. So suppose now that $p_1 = q_2 = r_3 = 0$. We define a matrix M by

$$M = \begin{pmatrix} 0 & p_2 & p_3 \\ q_1 & 0 & q_3 \\ r_1 & r_2 & 0 \end{pmatrix}. \quad (7.2)$$

Since the vertex $v = A \prec \tau$ and τ itself are the only V -faces of τ by our assumption, $p_2, p_3, q_1, q_3, r_1, r_2$ are positive integers. Let $\hat{\tau}$ be the convex hull of $\{0\} \cup \tau$ in \mathbb{R}^4 . Then it follows from $\text{Vol}_{\mathbb{Z}}(\tau) = 1$ that we have

$$\alpha \cdot \det(M) = \text{Vol}_{\mathbb{Z}}(\hat{\tau}) = \text{Vol}_{\mathbb{Z}}(\tau) \cdot N(\tau) = N(\tau). \quad (7.3)$$

Let

$$\text{aff}(\tau) : ax + by + cz + dw = N(\tau) = a \cdot \alpha \quad (7.4)$$

be the equation of $\text{aff}(\tau)$ with $\gcd(a, b, c, d) = 1$. Then we get $a = \det(M)$. Also, as $\text{Vol}_{\mathbb{Z}}(\tau) = 1$, the integers a, b, c and d are the 3×3 minors of the matrix

$$\begin{pmatrix} p_0 - \alpha & 0 & p_2 & p_3 \\ q_0 - \alpha & q_1 & 0 & q_3 \\ r_0 - \alpha & r_1 & r_2 & 0 \end{pmatrix}. \quad (7.5)$$

As we supposed that $v = A$ contributes with respect to τ , we have $a|(b + c + d)$. For the integer $k = (b + c + d)/a$ we thus obtain

$$\begin{vmatrix} p_0 - \alpha & 0 & p_2 & p_3 \\ q_0 - \alpha & q_1 & 0 & q_3 \\ r_0 - \alpha & r_1 & r_2 & 0 \\ -k & 1 & 1 & 1 \end{vmatrix} = 0. \quad (7.6)$$

Hence $(-k, 1, 1, 1)$ is a rational linear combination of the vectors \overrightarrow{AP} , \overrightarrow{AQ} and \overrightarrow{AR} . As $\text{Vol}_{\mathbb{Z}}(\tau) = 1$, the cone generated by \overrightarrow{AP} , \overrightarrow{AQ} and \overrightarrow{AR} is smooth and hence $(-k, 1, 1, 1)$ is an integer linear combination of these vectors. In particular there should be an integer solution $(x, y, z) \in \mathbb{Z}^3$ for the equation

$$\begin{pmatrix} 0 & q_1 & r_1 \\ p_2 & 0 & r_2 \\ p_3 & q_3 & 0 \end{pmatrix} \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}. \quad (7.7)$$

By Cramer's rule we then find in particular that

$$x = \frac{q_1 r_2 + q_3 (r_1 - r_2)}{\det(M)}. \quad (7.8)$$

Without loss of generality we may assume that $r_1 \geq r_2$. Then we have

$$\det(M) = p_3q_1r_2 + p_2q_3r_1 \leq q_1r_2 + q_3(r_1 - r_2), \quad (7.9)$$

which contradicts the positivity of $p_2, p_3, q_1, q_3, r_1, r_2$. This completes the proof. \square

Lemma 7.3. *Let $\tau = APQR$ be a 3-dimensional lattice simplex in a compact facet of $\Gamma_+(f)$ such that $v = A(\alpha, 0, 0, 0)$ and $\sigma = PQR$ are its V -faces. Then*

$$\frac{\zeta_\tau(t) \cdot (1-t)}{\zeta_v(t) \cdot \zeta_\sigma(t)} \quad (7.10)$$

is a polynomial.

Proof. If $\text{Vol}_{\mathbb{Z}}(\tau) > \text{Vol}_{\mathbb{Z}}(\sigma)$ then the assertion is obvious. So suppose that $\text{Vol}_{\mathbb{Z}}(\tau) = \text{Vol}_{\mathbb{Z}}(\sigma)$. Let

$$\text{aff}(\tau) : ax + by + cz + dw = N(\tau) \quad (7.11)$$

be the equation of $\text{aff}(\tau)$ with $\gcd(a, b, c, d) = 1$. Since we have

$$N(v) = \alpha = \frac{N(\tau)}{a}, \quad N(\sigma) = \frac{N(\tau)}{\gcd(b, c, d)} \quad (7.12)$$

and $\gcd(a, b, c, d) = 1$, the condition

$$\text{Vol}_{\mathbb{Z}}(\tau) = \text{Vol}_{\mathbb{Z}}(\sigma) \iff \gcd(N(v), N(\sigma)) = 1 \quad (7.13)$$

implies that $N(\tau) = a \cdot \gcd(b, c, d)$, $N(\sigma) = a$ and $N(v) = \alpha = \gcd(b, c, d)$. Hence we get

$$\frac{\zeta_\tau(t)}{\zeta_v(t) \cdot \zeta_\sigma(t)} = \frac{(1 - t^{a \cdot \alpha})^{\text{Vol}_{\mathbb{Z}}(\sigma)}}{(1 - t^a) \cdot (1 - t^a)^{\text{Vol}_{\mathbb{Z}}(\sigma)}}. \quad (7.14)$$

However, the only common zero of $\zeta_v(t)$ and $\zeta_\sigma(t)$ is equal to 1. \square

Lemma 7.4. *Let $\tau = APQR$ for $A(\alpha, 0, 0, 0), P(0, 0, p_2, p_3), Q(0, q_1, 0, q_3), R(0, r_1, r_2, 0)$ be a 3-dimensional lattice simplex in a compact facet of $\Gamma_+(f)$ that is not of type B_1 ($\iff \alpha \geq 2$). Assume that $v = A \prec \tau$ and $\sigma = PQR \prec \tau$ are its contributing V -faces with respect to τ and*

$$\lambda = \exp\left(-2\pi i \frac{\nu(\tau)}{N(\tau)}\right) \neq 1. \quad (7.15)$$

Then for the polynomial

$$F(t) = \frac{\zeta_\tau(t) \cdot (1-t)}{\zeta_v(t) \cdot \zeta_\sigma(t)} \in \mathbb{C}[t] \quad (7.16)$$

(see Lemma 7.3) we have $F(\lambda) = 0$.

Proof. Since the V -face $\sigma = PQR \prec \tau$ contributes with respect to τ , by the proof of Proposition 4.1 we have $\text{Vol}_{\mathbb{Z}}(\tau) = \alpha \cdot \text{Vol}_{\mathbb{Z}}(\sigma)$. Let

$$\text{aff}(\tau) : ax + by + cz + dw = N(\tau) = a \cdot \alpha \quad (7.17)$$

be the equation of $\text{aff}(\tau)$ with $\gcd(a, b, c, d) = 1$. Then by the proof of Lemma 7.3 the condition

$$\text{Vol}_{\mathbb{Z}}(\tau) = \alpha \cdot \text{Vol}_{\mathbb{Z}}(\sigma) \iff \gcd(N(v), N(\sigma)) = \alpha \quad (7.18)$$

implies that $N(\tau) = a \cdot \alpha \cdot \gcd(b, c, d)$, $N(\sigma) = a \cdot \alpha$ and $N(v) = \alpha = \alpha \cdot \gcd(b, c, d)$. Hence we get $\gcd(b, c, d) = 1$, $N(\tau) = N(\sigma) = a \cdot \alpha$ and

$$F(t) = \frac{(1 - t^{a \cdot \alpha})^{\alpha \cdot \text{Vol}_{\mathbb{Z}}(\sigma)} \cdot (1 - t)}{(1 - t^{\alpha}) \cdot (1 - t^{a \cdot \alpha})^{\text{Vol}_{\mathbb{Z}}(\sigma)}}. \quad (7.19)$$

It follows from our assumption $\alpha \geq 2$ that we have $F(\lambda) = 0$ unless $\alpha = 2$ and $\text{Vol}_{\mathbb{Z}}(\sigma) = 1$. Let us show that the case where $\alpha = 2$, $\text{Vol}_{\mathbb{Z}}(\sigma) = 1$ and $\text{Vol}_{\mathbb{Z}}(\tau) = 2$ cannot occur. Assume that $\text{Vol}_{\mathbb{Z}}(\tau) = 2$ and $\text{Vol}_{\mathbb{Z}}(\sigma) = 1$. Since another V -face $v = A \prec \tau$ contributes with respect to τ , we have also $a \mid (b+c+d)$. Then by $N(\tau) = 2a$ we obtain $e^{-2\pi i v(\tau)/N(\tau)} = 1$ or $e^{-2\pi i v(\tau)/N(\tau)} = -1$. It suffices to study what happens when $\lambda = e^{-2\pi i v(\tau)/N(\tau)} = -1 \iff 2a \mid (b+c+d)$. As $\text{Vol}_{\mathbb{Z}}(\tau) = 2$ and $\alpha = 2$, the even integers $2a, 2b, 2c$ and $2d$ are the 3×3 minors of the matrix

$$\begin{pmatrix} -2 & 0 & p_2 & p_3 \\ -2 & q_1 & 0 & q_3 \\ -2 & r_1 & r_2 & 0 \end{pmatrix},$$

and hence the expressions for a, b, c and d become

$$\begin{aligned} a &= \frac{q_1 r_2 p_3 + r_1 p_2 q_3}{2}, & b &= r_2 p_3 + p_2 q_3 - q_3 r_2, \\ c &= p_3 q_1 + q_3 r_1 - r_1 p_3, & d &= q_1 r_2 + r_1 p_2 - p_2 q_1. \end{aligned} \quad (7.20)$$

As in the proof of Lemma 7.2 for the integer $k = (b+c+d)/a$ we can show that the vector $(-k, 1, 1, 1)$ is a rational linear combination of \overrightarrow{AP} , \overrightarrow{AQ} and \overrightarrow{AR} . As $\text{Vol}_{\mathbb{Z}}(\tau) = 2$, the cone generated by \overrightarrow{AP} , \overrightarrow{AQ} and \overrightarrow{AR} has multiplicity two and so in particular there should be an integer solution $(x, y, z) \in \mathbb{Z}^3$ for the equation

$$\begin{pmatrix} 0 & q_1 & r_1 \\ p_2 & 0 & r_2 \\ p_3 & q_3 & 0 \end{pmatrix} \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} 2 \\ 2 \\ 2 \end{pmatrix}. \quad (7.21)$$

We define a matrix M by

$$M = \begin{pmatrix} 0 & q_1 & r_1 \\ p_2 & 0 & r_2 \\ p_3 & q_3 & 0 \end{pmatrix}. \quad (7.22)$$

Then by Cramer's rule we then find that

$$x = \frac{2(r_2 q_1 + r_1 q_3 - r_2 q_3)}{\det(M)}, \quad y = \frac{2(r_2 p_3 + r_1 p_2 - r_1 p_3)}{\det(M)}, \quad (7.23)$$

$$\text{and } z = \frac{2(q_1 p_3 + p_2 q_3 - p_2 q_1)}{\det(M)}. \quad (7.24)$$

Now we study the possible signs of x, y and z . If $p_2 \geq p_3$ and $q_1 \geq q_3$, then $y > 0$ and $x > 0$. If $p_2 \geq p_3$ and $q_1 \leq q_3$, then $y > 0$ and $z > 0$. If $p_2 \leq p_3$ and $r_1 \leq r_2$, then

$z > 0$ and $y > 0$ and so on. Thus we find that at least two of the integers x, y and z are always positive. By permuting them, we may assume that $x > 0$ and $y > 0$. As none of $p_2, p_3, q_1, q_3, r_1, r_2$ is equal to 0, the equation $p_3x + q_3y = 2$ obtained by (7.21) implies that $p_3 = q_3 = 1$ and $x = y = 1$. Consequently we get

$$a = \frac{q_1r_2 + r_1p_2}{2}, \quad b = p_2, \quad c = q_1, \quad d = q_1r_2 + r_1p_2 - p_2q_1 \quad (7.25)$$

and $\det(M) = q_1r_2 + r_1p_2$. As we supposed that $2a|(b + c + d)$, we have

$$(q_1r_2 + r_1p_2)|(p_2 + q_1 + q_1r_2 + r_1p_2 - p_2q_1) \iff \det(M)|(p_2 + q_1 - p_2q_1) \quad (7.26)$$

and z is an even integer. Hence, again by (7.21) and by using that $x = y = 1$, we find that p_2 and q_1 should be even. Then we have

$$\gcd(b, c, d) = \gcd(p_2, q_1, q_1r_2 + r_1p_2 - p_2q_1) \geq 2. \quad (7.27)$$

However, it contradicts our previous result $\gcd(b, c, d) = 1$. This completes the proof. \square

Definition 7.5. For a (not necessarily compact) facet τ of $\Gamma_+(f)$ we define its essential dimension $\text{ess.dim}\tau$ by

$$\text{ess.dim}\tau = \max_{\sigma \prec \tau: \text{compact}} \dim\sigma. \quad (7.28)$$

Note that a facet τ of $\Gamma_+(f)$ is compact if and only if $\text{ess.dim}\tau = 3$. Let \mathcal{F} be the set of V -faces of $\Gamma_+(f)$. First define a subset $\mathcal{F}_3 \subset \mathcal{F}$ of \mathcal{F} by $\sigma \in \mathcal{F}_3 \iff \sigma$ is contained in a compact facet τ of $\Gamma_+(f)$. Next define a subset $\mathcal{F}_2 \subset \mathcal{F} \setminus \mathcal{F}_3$ by $\sigma \in \mathcal{F}_2 \iff \sigma \notin \mathcal{F}_3$ and σ is contained in a facet τ of $\Gamma_+(f)$ such that $\text{ess.dim}\tau = 2$. Similarly we define a subset $\mathcal{F}_1 \subset \mathcal{F} \setminus (\mathcal{F}_2 \cup \mathcal{F}_3)$ by $\sigma \in \mathcal{F}_1 \iff \sigma \notin \mathcal{F}_2 \cup \mathcal{F}_3$ and σ is contained in a facet τ of $\Gamma_+(f)$ such that $\text{ess.dim}\tau = 1$. Finally we set $\mathcal{F}_0 = \mathcal{F} \setminus (\mathcal{F}_1 \cup \mathcal{F}_2 \cup \mathcal{F}_3)$. Namely the V -faces in \mathcal{F}_0 are those contained in facets τ of $\Gamma_+(f)$ such that $\text{ess.dim}\tau = 0$. Note that for any $0 \leq i \leq 3$ and $\sigma \in \mathcal{F}_i$ we have $\dim\sigma \leq i$. Thus \mathcal{F} is a disjoint union of $\mathcal{F}_0, \mathcal{F}_1, \mathcal{F}_2$ and \mathcal{F}_3 :

$$\mathcal{F} = \mathcal{F}_0 \sqcup \mathcal{F}_1 \sqcup \mathcal{F}_2 \sqcup \mathcal{F}_3. \quad (7.29)$$

For $0 \leq i \leq 3$ we define a rational function $R_i(t) \in \mathbb{C}(t)$ by

$$R_i(t) = \prod_{\sigma \in \mathcal{F}_i} \left\{ \zeta_\sigma(t) \right\}^{(-1)^{i-\dim\sigma}} \in \mathbb{C}(t) \quad (7.30)$$

so that we have the product decomposition

$$\zeta_{f,0}^{-1}(t) = R_3(t) \cdot R_2^{-1}(t) \cdot R_1(t) \cdot R_0^{-1}(t) \quad (7.31)$$

of $\zeta_{f,0}^{-1}(t)$ by Theorem 2.3.

Definition 7.6. We say that $\Gamma_+(f)$ is 0-convenient if for any 0-dimensional V -face $v \prec \Gamma_+(f)$ the set $\text{Int}\mathbb{R}_+^n \cap \partial\Gamma_+(f)$ is contained in the union of compact facets of $\Gamma_+(f)$ in a neighborhood of v .

Now we have the following result.

Theorem 7.7. *Assume that $f(x) \in \mathbb{C}[x_1, \dots, x_4]$ is non-degenerate at the origin $0 \in \mathbb{C}^4$ and $\Gamma_+(f)$ is 0-convenient. Let $\tau \prec \Gamma_+(f)$ be a (not necessarily compact) facet that is not a B_1 -pyramid nor a B_2 -facet. Then the complex number*

$$\lambda = \exp\left(-2\pi i \frac{\nu(\tau)}{N(\tau)}\right) \in \mathbb{C} \quad (7.32)$$

associated to it is an eigenvalue of the monodromy of f at some point in a neighborhood of the origin $0 \in \mathbb{C}^4$.

Proof. We follow the strategy in the proof of [17, Theorem 10]. Here we shall treat only the case where τ is compact. The other cases can be treated similarly as in the proof of [17, Theorem 10]. We subdivide compact facets of $\Gamma_+(f)$ into 3-dimensional lattice simplices τ_1, \dots, τ_k and for $1 \leq i \leq k$ define a polynomial $F_{\tau_i}(t) \in \mathbb{C}[t]$ by Theorem 4.3 and Proposition 4.4. Then we have

$$R_3(t) = \prod_{i=1}^k F_{\tau_i}(t) \cdot \prod_{\rho} \zeta_{\rho}(t) \cdot \prod_{\gamma} \zeta_{\gamma}^{-1}(t), \quad (7.33)$$

where in the product \prod_{ρ} (resp. \prod_{γ}) the V -face ρ (resp. γ) ranges through the 1-dimensional (resp. 0-dimensional) ones in \mathcal{F}_3 which were not used in the construction of $F_{\tau_1}(t), \dots, F_{\tau_k}(t)$. Since τ contains a non- B_1 -simplex by Lemma 3.6, as in the proof of [17, Theorem 10] by using Theorem 4.3, Proposition 4.4 and Lemmas 7.2, 7.3 and 7.4 we can show that $R_3(\lambda) = 0$. Indeed, if there exists a 0-dimensional V -face γ in (7.33) not contained in any 1-dimensional V -face ρ in (7.33) then by the 0-convenience of $\Gamma_+(f)$ we are in the situation of Lemmas 7.2, 7.3 and 7.4 and can use them to deduce $R_3(\lambda) = 0$. First assume that $R_2(\lambda) = 0$. In this case, we subdivide 2-dimensional V -faces in \mathcal{F}_2 into 2-dimensional lattice simplices $\sigma_1, \dots, \sigma_l$ and for $1 \leq i \leq l$ define a polynomial $F_{\sigma_i}(t) \in \mathbb{C}[t]$ by Theorem 4.3. Then we have

$$R_2(t) = \prod_{i=1}^l F_{\sigma_i}(t) \cdot \prod_{\gamma} \zeta_{\gamma}(t), \quad (7.34)$$

where in the product \prod_{γ} the V -face $\gamma \in \mathcal{F}_2$ ranges through the 0-dimensional ones which were not used in the construction of $F_{\sigma_1}(t), \dots, F_{\sigma_l}(t)$. By our assumption $R_2(\lambda) = 0$ there exists $1 \leq i \leq l$ such that λ is a root of the polynomial

$$Q(t) := F_{\sigma_i}(t) \cdot \prod_{\gamma \prec \sigma_i} \zeta_{\gamma}(t). \quad (7.35)$$

Let $\tilde{\sigma} \prec \Gamma_+(f)$ be the (unique) non-compact facet containing σ_i . For simplicity, assume that $\tilde{\sigma}$ is non-compact for the variable v_4 . For a sufficiently generic small complex number $c \in \mathbb{C}$ let us set

$$g(x_1, \dots, x_4) = f(x_1, x_2, x_3, x_4 + c) \in \mathbb{C}[x_1, \dots, x_4]. \quad (7.36)$$

Then by Proposition 6.1 the new polynomial $g(x)$ is also non-degenerate at the origin $0 \in \mathbb{C}^4$ and its Newton polyhedron $\Gamma_+(g)$ is the projection of $\Gamma_+(f)$ with respect to the

variable v_4 . By this construction, we can easily see that the polynomial $Q(t)$ shows up in the product decomposition of $\zeta_{g,0}(t)$ by Theorem 2.3. If $\zeta_{g,0}(\lambda) \neq 0$ then there exists a 1-dimensional V -face $\rho \prec \Gamma_+(g) \cap \{v_4 = 0\}$ of $\Gamma_+(g) \cap \{v_4 = 0\} \subset \{v_4 = 0\} \simeq \mathbb{R}^3$ satisfying the condition $F_\rho(\lambda) = 0$ and contained in a non-compact facet $\tilde{\rho}$ of $\Gamma_+(g) \cap \{v_4 = 0\}$ such that $\text{ess.dim} \tilde{\rho} = 1$. For simplicity assume that $\tilde{\rho}$ is non-compact for the variable v_3 and for a sufficiently generic small $c' \in \mathbb{C}$ set

$$h(x_1, \dots, x_4) = g(x_1, x_2, x_3 + c', x_4) \in \mathbb{C}[x_1, \dots, x_4]. \quad (7.37)$$

Then by Proposition 6.1 the polynomial $h(x)$ is also non-degenerate at the origin $0 \in \mathbb{C}^4$ and its Newton polyhedron $\Gamma_+(h)$ is the projection of $\Gamma_+(g)$ with respect to the variable v_3 . Now it follows from the condition $F_\rho(\lambda) = 0$ we have $\zeta_{h,0}^{-1}(\lambda) = 0$. Finally let us consider the remaining case $R_2(\lambda) \neq 0$. Since $R_1(t)$ is a polynomial, it suffices to treat only the case $R_0(\lambda) = 0$. However by the 0-convenience of $\Gamma_+(f)$ we have $\mathcal{F}_0 = \emptyset$ and hence $R_0(t) \equiv 1$. This completes the proof. \square

Definition 7.8. *A B -wall is a triangle in \mathbb{R}_+^4 with vertices of the form $(0, 0, a, b)$, $(1, 0, c, d)$, $(0, 1, e, f)$ up to reordering the coordinates.*

Definition 7.9. *We say that $\Gamma_+(f)$ is good if it is 0-convenient and no two of its B_1 -pyramid facets for different variables contributing to the same candidate pole of $Z_{\text{top},f}(s)$ have a common 1-dimensional V -face containing a 0-dimensional one.*

Finally we obtain the following result.

Theorem 7.10. *Assume that $f(x) \in \mathbb{C}[x_1, \dots, x_4]$ is non-degenerate at the origin $0 \in \mathbb{C}^4$, $\Gamma_+(f)$ is good and no two B_i -facets ($i = 1, 2$) contributing to the same candidate pole of $Z_{\text{top},f}(s)$ intersect by a B -wall. Let $s_0 \in \mathbb{C}$ be a pole of $Z_{\text{top},f}(s)$. Then the complex number $\exp(2\pi i s_0) \in \mathbb{C}$ is an eigenvalue of the monodromy of f at some point in a neighborhood of the origin $0 \in \mathbb{C}^4$.*

Proof. By the results in Section 3, Theorems 4.3, 7.7 and Proposition 4.4, we can prove the assertion following the strategy in proof of [17, Theorem 15] as follows. By Propositions 3.3, 4.4 and Theorems 4.3, 7.7 it remains to consider only the case where the pole s_0 of $Z_{\text{top},f}(s)$ is contributed by several B_1 -pyramid facets of $\Gamma_+(f)$ for different variables. Suppose that there exist two B_1 -pyramid facets $\tau, \sigma \prec \Gamma_+(f)$ for different variables such that

$$s_0 = -\frac{\nu(\tau)}{N(\tau)} = -\frac{\nu(\sigma)}{N(\sigma)}. \quad (7.38)$$

If $\tau \cap \sigma$ is their common facet, then by reducing the problem to the case where $\tau \cap \sigma$ is compact by Proposition 6.1 and using a V -face in it we can get the corresponding monodromy eigenvalue $\exp(2\pi i s_0) \in \mathbb{C}$ at some point in a neighborhood of the origin $0 \in \mathbb{C}^4$ by the proof of [17, Theorem 15]. Otherwise, their contributions to $Z_{\text{top},f}(s)$ can be calculated separately and by Proposition 3.2 they produce only a fake pole. \square

By our results obtained in this paper, we can prove also a similar assertion in higher dimensions $n \geq 5$ under some weak combinatorial assumption on $\Gamma_+(f)$.

References

- [1] A'Campo, N. A. “La fonction zêta d’une monodromie”, *Comment. Math. Helv.*, 50 (1975): 233-248.
- [2] Artal Bartolo, E., Cassou-Noguès, P., Luengo, I. and Melle Hernández, A. “Monodromy conjecture for some surface singularities”, *Ann. Scient. Ecole Norm. Sup.*, 35 (2002): 405-460.
- [3] Bories, B. and Veys, W. “Igusa’s p -adic local zeta function and the Monodromy Conjecture for non-degenerated surface singularities”, arXiv: arXiv:1306.6012v1.
- [4] Denef, J. “Report on Igusa’s local zeta function”, Séminaire Bourbaki, Vol. 1990/91. *Astérisque*, No. 201-203 (1991): 359-386.
- [5] Denef, J. “Degree of local zeta functions and monodromy”, *Compos. Math.*, No. 89 (1994): 207-216.
- [6] Denef, J. and Loeser, F. “Caractéristique d’Euler-Poincaré, fonctions zêta locales et modifications analytiques ”, *J. Amer. Math. Soc.*, 5 (1992): 705-720.
- [7] Denef, J. and Loeser, F. “Motivic Igusa zeta functions”, *J. Alg. Geom.*, 7 (1998): 505-537.
- [8] Denef, J. and Loeser, F. “Geometry on arc spaces of algebraic varieties”, *Progr. Math.*, 201 (2001): 327-348.
- [9] Dimca, A. *Sheaves in topology*, Universitext, Springer-Verlag, Berlin, 2004.
- [10] Esterov, A. and Takeuchi, K. “Motivic Milnor fibers over complete intersection varieties and their virtual Betti numbers”, *Int. Math. Res. Not.*, Vol. 2012, No. 15 (2012): 3567-3613.
- [11] Fulton, W. *Introduction to toric varieties*, Princeton University Press, 1993.
- [12] Gelfand, I.M., Kapranov, M. and Zelevinsky, A., *Discriminants, resultants and multidimensional determinants*, Birkhäuser, 1994.
- [13] González Villa, M. and Lemahieu, A., “The monodromy conjecture for plane meromorphic functions”, *Bull. of the London Math. Soc.*, 46, no. 3 (2014): 441-453 doi:10.1112/blms/bdt098.
- [14] Hotta, R., Takeuchi, K. and Tanisaki, T., *D-modules, perverse sheaves and representation theory*, Progress in Math., Birkhäuser, Boston, 2008.
- [15] Kouchnirenko, A. G. “Polyèdres de Newton et nombres de Milnor”, *Invent. Math.*, 32 (1976): 1-31.
- [16] Kouchnirenko, A. G. “A criterion for the existence of a nondegenerate quasihomogeneous function with given weights” (Russian), *Uspehi Mat. Nauk*, 32, no. 3 (195) (1977): 169-170.

- [17] Lemahieu, A. and Van Proeyen, L. “Monodromy conjecture for nondegenerate surface singularities”, *Transactions of AMS*, Vol. 363, No. 9 (2011): 4801-4829.
- [18] Lemahieu, A. and Veys, W., “Zeta functions and monodromy for surfaces that are general for a toric idealistic cluster”, *Int. Math. Res. Not.*, Vol. 2009, No. 1 (2009): 11-62.
- [19] Loeser, F. “Fonctions d’Igusa p -adiques et polynômes de Bernstein”, *Amer. J. Math.*, 110 (1988): 1-22.
- [20] Loeser, F. “Fonctions d’Igusa p -adiques, polynômes de Bernstein et polyèdres de Newton”, *J. Reine Angew. Math.*, 412 (1990): 75-96.
- [21] Matsui, Y. and Takeuchi, K. “Milnor fibers over singular toric varieties and nearby cycle sheaves”, *Tohoku Math. Journal*, 63 (2011): 113-136.
- [22] Matsui, Y. and Takeuchi, K. “Monodromy zeta functions at infinity, Newton polyhedra and constructible sheaves”, *Mathematische Zeitschrift*, 268 (2011): 409-439.
- [23] Matsui, Y. and Takeuchi, K. “Monodromy at infinity of polynomial maps and Newton polyhedra, with Appendix by C. Sabbah”, *Int. Math. Res. Not.*, Vol. 2013, No. 8 (2013): 1691-1746.
- [24] Milnor, J. *Singular points of complex hypersurfaces*, Princeton University Press, 1968.
- [25] Nicaise, J. “An introduction to p -adic and motivic zeta functions and the monodromy conjecture”, Algebraic and analytic aspects of zeta functions and L -functions, *MSJ Mem.*, Vol. 21, (2010): 141-166.
- [26] Oda, T. *Convex bodies and algebraic geometry. An introduction to the theory of toric varieties*, Springer-Verlag, 1988.
- [27] Oka, M. *Non-degenerate complete intersection singularity*, Hermann, Paris (1997).
- [28] Okada, T. Takeuchi, K. “Meromorphic continuations of local zeta functions and their applications to oscillating integrals”, *Tohoku Math. Journal*, 65 (2013): 159-178.
- [29] Varchenko, A. N. “Zeta-function of monodromy and Newton’s diagram”, *Invent. Math.*, 37 (1976): 253-262.
- [30] Veys, W. “Determination of the poles of the topological zeta functions for curves”, *Manuscr. Math.*, 87 (1995): 435-448.