



HAL
open science

Trapped Magnetic Field Experiments and Characterization of Large-Sized Bulk MgB₂ Samples

Kévin Berger, Bruno Douine, Jacques Noudem, Frédéric Trillaud, Michael
Rudolf Koblishka, Jean Lévêque

► **To cite this version:**

Kévin Berger, Bruno Douine, Jacques Noudem, Frédéric Trillaud, Michael Rudolf Koblishka, et al.. Trapped Magnetic Field Experiments and Characterization of Large-Sized Bulk MgB₂ Samples. 5th International Conference on Superconductivity and Magnetism - ICSM2016, Apr 2016, Fethiye, Turkey. p. 182, ID 539. hal-01281147

HAL Id: hal-01281147

<https://hal.science/hal-01281147v1>

Submitted on 2 Mar 2016

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Trapped Magnetic Field Experiments and Characterization of Large-Sized Bulk MgB₂ Samples

K. Berger¹, B. Douine¹, J. Noudem^{2,3},

F. Trillaud⁴, M.R. Koblischka⁵, and J. L  v  que¹.

¹ University of Lorraine, GREEN, 54506 Vandoeuvre-l  s-Nancy, France

² CRISMAT, CNRS/ENSICAEN, 14050 Caen, France

³ Caen University, LUSAC, 50130 Cherbourg-Octeville, France

⁴ National Autonomous University of Mexico, Institute of Engineering, 04510 Mexico

⁵ Saarland University, Institute of Experimental Physics, 66123 Saarbr  cken, Germany

Contact e-mail: kevin.berger@univ-lorraine.fr

From the applications point of view, the advantage of the very low density of MgB₂ bulk samples must be taken into consideration and, the generation of strong magnetic flux densities using MgB₂ should be investigated [1]. In this contribution, we have studied the trapped magnetic field properties of samples of 20 mm, 40 mm and 50 mm in diameter. Samples with a diameter of 20 mm and 40 mm were processed by *ex-situ* Spark Plasma Sintering, since the sample of 50 mm of diameter, from Edison Spa, has been produced using a Reactive Mg-Liquid Infiltration process. Both processes are able to produce dense and high quality MgB₂ samples. For the sample of 40 mm diameter, some Field Cooling experiments were carried out at temperatures from 10 K to 30 K both on a single pellet of 4 mm high and on a stack of several pellets of the same diameter. With the help of modeling tools, we have simulated the behavior of such samples in order to deduce their parameters as $J_c(B)$ and n -value, Fig. 1.




Fig. 1: Two models of critical current density (a), and the simulated resulting trapped magnetic field at the center of the surface of a bulk MgB₂ of 20 mm of diameter and 4 mm high (b).

With these simulations and experiments, we will show that the most relevant parameter for the applications of large-sized bulk MgB₂ samples as permanent magnet is the irreversibility field H_{irr} and not the critical current density at zero field $J_c(0)$.

References

1. J. G. Noudem, M. Aburras, P. Bernstein, X. Chaud, M. Muralidhar, and M. Murakami, *Development in processing of MgB₂ cryo-magnet superconductors*, J. Appl. Phys. **116**, 163916 (2014).