



HAL
open science

An algorithm which generates linear extensions for a non-simply-laced d-complete poset with uniform probability

Kento Nakada

► **To cite this version:**

Kento Nakada. An algorithm which generates linear extensions for a non-simply-laced d-complete poset with uniform probability. 24th International Conference on Formal Power Series and Algebraic Combinatorics (FPSAC 2012), 2012, Nagoya, Japan. pp.655-660, 10.46298/dmtcs.3071 . hal-01283096

HAL Id: hal-01283096

<https://hal.science/hal-01283096>

Submitted on 5 Mar 2016

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

An algorithm which generates linear extensions for a non-simply-laced d -complete poset with uniform probability

Kento Nakada[†]

Graduate School of Education Master's Program, Okayama University, Japan



Abstract. The purpose of this paper is to present an algorithm which generates linear extensions for a non-simply-laced d -complete poset with uniform probability.

Résumé. Le but de ce papier est présenter un algorithme qui produit des extensions linéaires pour une non-simply-laced d -complete poset avec probabilité constante.

Keywords: d -complete posets, algorithm, linear extension, uniform generation

1 Introduction

In [7](Theorem 4.2), J. Stembridge classified irreducible minuscule elements of Kac-Moody Weyl group over a root system Φ into three classes below:

- Φ is simply-laced,
- Φ has the form  (namely, of type B), or
- Φ has the form  (we say *type F_m* , for simplicity).

In [5][6], the author and S. Okamura constructed an algorithm which generates reduced decompositions for a given minuscule element of simply-laced Weyl group with uniform probability. The algorithm in [6] is described in terms of graphs. Simply-laced minuscule elements are described as certain simple acyclic di-graphs. The transitive-closure of the graph is called a d -complete poset. Then, the reduced decompositions are identified with linear extensions of the graph. This algorithm gives a proof of the hook formula [1] for the number of reduced decompositions of a minuscule element in simply-laced case.

In this paper, we present an algorithm (algorithm A) in terms of graphs (See Section 2 for details). This algorithm is a generalization of an algorithm in [5][6]. We define a certain acyclic multi-di-graph corresponding to a minuscule element of type B (resp. type F_m) in Section 3 (resp. Section 4). Our

[†]Email: nakada@okayama-u.ac.jp

main result (Theorem 5.1) is that the algorithm A generates linear extensions for a minuscule element of type B and F_m with uniform probability. More precisely, the probability the algorithm A generates linear extension L of a graph S is given by:

$$\frac{\prod_{v \in S} (1 + \#H_S(v)^+)}{\#S!}, \quad (1.1)$$

where $H_S(v)^+$ is a certain subset of S (See Section 2 for detail). This (1.1) is independent from the choice of L . Hence, we get the hook formula for the number of linear extensions of a given shape S of type B and F_m . Namely, the number of linear extensions of a shape S is given by:

$$\frac{\#S!}{\prod_{v \in S} (1 + \#H_S(v)^+)}.$$

In section 6, we give a Lie theoretical description of shape of type B and F_m .

2 An algorithm for a graph Γ

Let $\Gamma = (\Gamma; A, o, i)$ be a finite acyclic multi-di-graph, where A denotes the set of arrows of Γ , $i(a)$ the sink of $a \in A$, and $o(a)$ the source of $a \in A$.

Definition 2.1 Put $d := \#\Gamma$. A bijection $L : \{1, \dots, d\} \rightarrow \Gamma$ is said to be a linear extension of Γ if:

$$L(k) = o(a) \text{ and } i(a) = L(l) \text{ implies } k > l, \quad k, l \in \{1, \dots, d\}, \quad a \in A.$$

The set of linear extensions of Γ is denoted by $\mathcal{L}(\Gamma)$.

For a given $v \in \Gamma$, we define a set $H_\Gamma(v)^+$ by:

$$H_\Gamma(v)^+ := \{a \in A(\Gamma) \mid v = o(a)\}.$$

For a given Γ , we call the following algorithm the *algorithm A* for Γ :

GNW1. Set $i := 0$ and set $\Gamma_0 := \Gamma$.

GNW2. (Now Γ_i has $d - i$ nodes.) Set $j := 1$ and pick a node $v_1 \in \Gamma_i$ with the probability $1/(d - i)$.

GNW3. If $\#H_{\Gamma_i}(v_j)^+ \neq 0$, pick an arrow $a_{j+1} \in H_{\Gamma_i}(v_j)^+$ with the probability $1/\#H_{\Gamma_i}(v_j)^+$. If not, go to GNW5.

GNW4. Set $v_{j+1} := i(a_j)$. Set $j := j + 1$ and return to GNW3.

GNW5. (Now $\#H_{\Gamma_i}(v_j)^+ = 0$.) Set $L(i + 1) := v_j$ and set $\Gamma_{i+1} := \Gamma_i \setminus v_j$ (the graph deleted v_j from Γ_i).

GNW6. Set $i := i + 1$. If $i < d$, return to GNW2; if $i = d$, terminate.

We note that the algorithm A stops in finite time since Γ is acyclic. By the definition of the algorithm A for Γ , the map $L : i \mapsto L(i)$ generated above is a linear extension of Γ . We denote by $\text{Prob}_\Gamma(L)$ the probability we get $L \in \mathcal{L}(\Gamma)$ by the algorithm A.

3 Shapes of type B

We denote by \mathbb{N} the set of non-negative integers. We define a set \mathbb{B} by:

$$\mathbb{B} := \{ (i, j) \in \mathbb{N} \times \mathbb{N} \mid i \leq j \}.$$

The set \mathbb{B} is depicted in FIGURE 3.1. We equip the \mathbb{B} with the partial order:

$$(i, j) \leq (i', j') \iff i \geq i' \text{ and } j \geq j'.$$




Fig. 3.1: The set \mathbb{B}

Definition 3.1 Let S be a finite order filter of \mathbb{B} . We induce to S a graph structure by:

$$(i, j) \rightarrow (i', j') \text{ if and only if } \begin{cases} "i = j \text{ and } i' = i, j' > j", \\ "i < j \text{ and } i' = i, j' > j", \\ "i < j \text{ and } i' > i, j' = j", \\ \text{or } "i < j \text{ and } i' = j, j' > i", \end{cases}$$

$$(i, j) \rightrightarrows (i', j') \text{ if and only if } "i < j \text{ and } i' = j' = j",$$

and there exists no other adjacency relation. Here, $v \rightarrow v'$ means there exists exactly one arrow from v to v' , and $v \rightrightarrows v'$ there exists exactly two arrows from v to v' . A graph S is called a shape of type B . See FIGURE 3.2 for examples of $H_S(v)^+$.




Fig. 3.2: $H_S(u)^+$, $H_S(v)^+$, and $H_S(w)^+$.

Remark 3.2 A shape of type B as poset is order-isomorphic to a shifted shape. Shifted shapes are also realized as d -complete posets over a root system of type D . The graph-structure of shapes of type D is described in [6] and compatible with notion of hooks (or called bars) of shifted shapes. The algorithm A depends not only on poset-structure but on graph-structure. Hence, we do not consider shapes of type B as shifted shapes.

4 Shapes of type F_m ($m \geq 2$).

We denote by \mathbb{Z} the set of integers. Let m be an integer greater than or equal to 2. We define a set \mathbb{F}_m by:

$$\mathbb{F}_m := \left\{ (i, j) \in \mathbb{N} \times \mathbb{Z} \left| \begin{array}{l} i = 0 \text{ and } j \geq -m, \\ i = 1 \text{ and } j \geq 0, \text{ or} \\ 2 \leq i \leq m \text{ and } j = 0 \end{array} \right. \right\}$$

For example, the set \mathbb{F}_3 is depicted in FIGURE 4.1. We equip the \mathbb{F}_m with the partial order:

$$(i, j) \leq (i', j') \iff i \geq i' \text{ and } j \geq j'.$$




Fig. 4.1: The set \mathbb{F}_3

Definition 4.1 Let S be a finite order filter of \mathbb{F}_m . We induce to S a graph structure by:

$$(i, j) \rightarrow (i', j') \text{ if and only if } \begin{cases} "i = 0, j \leq -1 \text{ and } i' \neq -j, j' > j", \\ "i = 0, j = 0, \text{ and } j' > 0", \\ "i = 1, j = 0, \text{ and } i' = 1, j' > 0", \\ "i = 1, j = 0, \text{ and } i' > 1, j' = 0", \\ "i \geq 2, j = 0, \text{ and } i' > i, j' = 0", \\ "j \geq 1 \text{ and } i' = i, j' > j", \\ \text{or } "j \geq 1 \text{ and } i' > i, j' = j", \end{cases}$$

$$(i, j) \Rightarrow (i', j') \text{ if and only if } "i = 0, j = 0, \text{ and } 0 < i', j' = 0",$$

and there exists no other adjacency relation. A graph S is called a shape of type F_m . See FIGURE 4.2 for examples of $H_S(v)^+$.




Fig. 4.2: $H_S(u)^+$, $H_S(v)^+$, and $H_S(w)^+$.

5 Main result

Now, we can state the main theorem:

Theorem 5.1 *Let S be a shape of type B or type F_m for some $m \geq 2$. Let $L \in \mathcal{L}(S)$. Then the algorithm A for S generates L with the probability*

$$\text{Prob}_S(L) = \frac{\prod_{v \in S} (1 + \#H_S(v)^+)}{\#S!}. \quad (5.1)$$

Since the right hand side of (5.1) is independent from the choice of $L \in \mathcal{L}(S)$, we have:

Corollary 5.2 *Let S be a shape of type B or type F_m for some $m \geq 2$. Then we have:*

$$\#\mathcal{L}(S) = \frac{\#S!}{\prod_{v \in S} (1 + \#H_S(v)^+)}.$$

6 Lie theoretical description of main result and Remarks

In this section, we fix a (not necessary simply-laced) Kac-Moody Lie algebra \mathfrak{g} with a simple root system $\Pi = \{\alpha_i \mid i \in I\}$. For all undefined terminology in this section, we refer the reader to [2] [3].

Definition 6.1 *An integral weight λ is said to be pre-dominant if:*

$$\langle \lambda, \beta^\vee \rangle \geq -1 \quad \text{for each } \beta^\vee \in \Phi_+^\vee,$$

where Φ_+^\vee denotes the set of positive real coroots. The set of pre-dominant integral weights is denoted by $P_{\geq -1}$. For $\lambda \in P_{\geq -1}$, we define the set $D(\lambda)^\vee$ by:

$$D(\lambda)^\vee := \{ \beta^\vee \in \Phi_+^\vee \mid \langle \lambda, \beta^\vee \rangle = -1 \}.$$

The set $D(\lambda)^\vee$ is called the shape of λ . If $\#D(\lambda)^\vee < \infty$, then λ is called finite.

Proposition 6.2 (see [4]) *Let $\lambda \in P_{\geq -1}$ be finite and $\beta^\vee, \gamma^\vee \in D(\lambda)^\vee$ satisfy $\beta^\vee > \gamma^\vee$ in the ordinary order of coroots. Then we have:*

$$\langle \beta, \gamma^\vee \rangle = 0, 1, \text{ or } 2.$$

By proposition 6.2, we introduce graph-structure into $D(\lambda)^\vee$ by:

$$\begin{aligned} \beta^\vee \rightarrow \gamma^\vee &\Leftrightarrow \beta^\vee > \gamma^\vee \text{ and } \langle \beta, \gamma^\vee \rangle = 1. \\ \beta^\vee \rightrightarrows \gamma^\vee &\Leftrightarrow \beta^\vee > \gamma^\vee \text{ and } \langle \beta, \gamma^\vee \rangle = 2. \end{aligned}$$

If $\beta^\vee \not> \gamma^\vee$, or $\beta^\vee > \gamma^\vee$ and $\langle \beta, \gamma^\vee \rangle = 0$, then no arrows from β^\vee to γ^\vee exist.

Thus, we get a finite acyclic multi-di-graph $D(\lambda)^\vee$ for a finite $\lambda \in P_{\geq -1}$.


Remark 6.3 *The finite pre-dominant integral weights λ are identified with the minuscule elements w [4]. And, we have $D(\lambda)^\vee = \{ \beta^\vee \in \Phi_+^\vee \mid w^{-1}(\beta^\vee) < 0 \}$. Furthermore, the linear extensions of $D(\lambda)^\vee$ are identified with the reduced decompositions of w [4] by the following one-to-one correspondence:*

$$\text{Red}(w) \ni (s_{i_1}, s_{i_2}, \dots, s_{i_d}) \longleftrightarrow L \in \mathcal{L}(D(\lambda)^\vee), \quad L(k) = s_{i_1} \cdots s_{i_{k-1}}(\alpha_{i_k})^\vee \in D(\lambda)^\vee \quad (k = 1, \dots, d),$$

where $\text{Red}(w)$ denotes the set of reduced decompositions of w , $d = \ell(w)$ the length of w .

6.1 Case of type B

Suppose that the underlying Dynkin diagram is of type B :




Let $W = \langle s_0, s_1, s_2, \dots \rangle$ be the Weyl group. Let Λ_0 be the 0-th fundamental weight. Then each $\lambda \in W\Lambda_0$ is a finite pre-dominant integral weight. And, $D(\lambda)^\vee$ is graph-isomorphic with some shape of type B defined in section 3.

Remark 6.4 Let $W_0 := \langle s_1, s_2, \dots \rangle$ be a maximal parabolic subgroup of W , which is the Weyl group of type A . Then a minimal coset representative w in W/W_0 is called a Lagrangian Grassmannian element.

Let $\lambda \in W\Lambda_0$. Then the corresponding minuscule element w in remark 6.3 is a Lagrangian Grassmannian element. Our result gives the number of reduced decompositions of Lagrangian Grassmannian element w .

6.2 Case of type F_m ($m \geq 2$)

Let $m \in \mathbb{Z}$ be greater than or equal to 2. Suppose that the underlying Dynkin diagram is of type F_m :



Let $W = \langle s_{-m}, \dots, s_{-2}, s_{-1}, s_0, s_1, \dots \rangle$ be the Weyl group. Let Λ_{-m} be the $(-m)$ -th fundamental weight. Then each $\lambda \in P_{\geq -1} \cap W\Lambda_{-m}$ is a finite pre-dominant integral weight. And, $D(\lambda)^\vee$ is graph-isomorphic with some shape of type F_m defined in section 4.

References

- [1] J. B. Carrell, *Vector fields, flag varieties and Schubert calculus*, Proc. Hyderabad Conference on Algebraic Groups (ed. S.Ramanan), Manoj Prakashan, Madras, 1991.
- [2] V. G. Kac, "Infinite Dimensional Lie Algebras," Cambridge Univ. Press, Cambridge, UK, 1990.
- [3] R. V. Moody and A. Pianzola, "Lie Algebras With Triangular Decompositions," Canadian Mathematical Society Series of Monograph and Advanced Text, 1995.
- [4] K. Nakada, *Colored hook formula for a generalized Young diagram*, Osaka J. of Math. **Vol.45 No.4** (2008), 1085-1120.
- [5] K. Nakada and S. Okamura, *Uniform generation of standard tableaux of a generalized Young diagram*, submitted to Journal of Algebraic Combinatorics.
- [6] K. Nakada and S. Okamura, *An algorithm which generates linear extensions for a generalized Young diagram with uniform probability*, DMTCS (Discrete Mathematics and Theoretical Computer Science), <http://www.dmtcs.org/dmtcs-ojs/index.php/proceedings/article/view/dmAN0170/3216>
- [7] J. R. Stembridge, *Minuscule elements of Weyl groups*, J.Algebra **235** (2001), 722-743.