



HAL
open science

Polymorphisms in the feline TNFA and CD209 genes are associated with the outcome of feline coronavirus infection

Ying-Ting Wang, Li-En Hsieh, Yu-Rou Dai, Ling-Ling Chueh

► **To cite this version:**

Ying-Ting Wang, Li-En Hsieh, Yu-Rou Dai, Ling-Ling Chueh. Polymorphisms in the feline TNFA and CD209 genes are associated with the outcome of feline coronavirus infection. *Veterinary Research*, 2014, 45 (1), pp.123. <10.1186/s13567-014-0123-6>. <hal-01290623>

HAL Id: hal-01290623

<https://hal.science/hal-01290623v1>

Submitted on 18 Mar 2016

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire HAL, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

RESEARCH

Open Access

Polymorphisms in the feline TNFA and CD209 genes are associated with the outcome of feline coronavirus infection

Ying-Ting Wang^{1†}, Li-En Hsieh^{1†}, Yu-Rou Dai² and Ling-Ling Chueh^{1,2*}

Abstract

Feline infectious peritonitis (FIP), caused by feline coronavirus (FCoV) infection, is a highly lethal disease without effective therapy and prevention. With an immune-mediated disease entity, host genetic variant was suggested to influence the occurrence of FIP. This study aimed at evaluating cytokine-associated single nucleotide polymorphisms (SNPs), i.e., tumor necrosis factor alpha (TNF- α), receptor-associated SNPs, i.e., C-type lectin DC-SIGN (*CD209*), and the five FIP-associated SNPs identified from Birman cats of USA and Denmark origins and their associations with the outcome of FCoV infection in 71 FIP cats and 93 FCoV infected non-FIP cats in a genetically more diverse cat populations. A promoter variant, *fTNFA* - 421 T, was found to be a disease-resistance allele. One SNP was identified in the extracellular domain (ECD) of *fCD209* at position +1900, a G to A substitution, and the A allele was associated with FIP susceptibility. Three SNPs located in the introns of *fCD209*, at positions +2276, +2392, and +2713, were identified to be associated with the outcome of FCoV infection, with statistical relevance. In contrast, among the five Birman FIP cat-associated SNPs, no genotype or allele showed significant differences between our FIP and non-FIP groups. As disease resistance is multifactorial and several other host genes could involve in the development of FIP, the five genetic traits identified in this study should facilitate in the future breeding of the disease-resistant animal to reduce the occurrence of cats succumbing to FIP.

Introduction

Feline infectious peritonitis (FIP), a highly lethal disease with nearly 100% mortality among ill cats once clinical signs appear, is caused by feline coronavirus (FCoV) infection [1]. Despite the ubiquitous existence of FCoV around the world, the prevalence of FIP is less than 5% [2]. There is currently no therapy proven to be effective for the treatment of FIP, and once diagnosis is confirmed, euthanasia is generally inevitable. Although this disease has been described for over fifty years [3], studies attempting to develop vaccines with different approaches have all failed due to the immunopathogenic features of infection by this virus [4]. However, among many FCoV experimental inoculations studies, some cats survived challenge with the virulent strain of FCoV [2,5-10], whereas certain

pedigreed cats were reported to be more likely to succumb to FIP than mixed bred cats [2,11,12]. All these findings indicate that genetic polymorphisms between cats might affect their susceptibility to FIP.

FIP is an immunopathological consequence of the abnormal production of various cytokines. Imbalanced Th1/Th2 immune responses with scarce or absent interferon-gamma (IFN- γ) is consistently found in FIP cases [4,8,10,13-15] and association of genetic polymorphisms in the IFN- γ gene with FIP occurrence has recently been identified [16]. In addition to IFN- γ , the upregulation of tumor necrosis factor-alpha (TNF- α) during the development of FIP has been reported to result in lymphopenia [17]. Feline dendritic cell (DC)-specific intercellular adhesion molecule-grabbing non-integrin (fDC-SIGN, encoded by *fCD209*), a key coreceptor during the infection of both type I and II FCoV [18], was found to affect binding and infection of type I FCoV. fDC-SIGN is also involved in the infection of type II FCoV, albeit not through the initial binding [19]. Despite the close relationship to FCoV

* Correspondence: linglingchueh@ntu.edu.tw

†Equal contributors

¹Graduate institute of Veterinary Medicine, School of Veterinary Medicine, National Taiwan University, Taipei 10617, Taiwan

²Department of Veterinary Medicine, School of Veterinary Medicine, National Taiwan University, Taipei 10617, Taiwan

infection, polymorphisms in the *fCD209* and feline TNF- α (*fTNFA*) genes and their association with FIP occurrence have never been investigated.

Recently, the surveillance of FIP-associated single nucleotide polymorphisms (SNPs) in Birman cats from USA and Denmark was conducted using a commercialized feline SNP array [20], and five SNPs were found to be significantly associated with FIP occurrence. However, it is unclear whether these disease-associated SNPs are Birman cat specific or can also be applied to other purebred or mixed breed cat populations.

To elucidate the genetic traits that contribute to FIP susceptibility, the *fTNFA* and *fCD209* genes were screened to identify disease-associated SNPs. The five SNPs identified from Birman cats proposed to be genetically associated with the occurrence of FIP were further evaluated in populations with more variable genetic backgrounds. Among all the polymorphisms analyzed, SNPs located in the *fTNFA* and *fCD209* genes were found to be associated with the outcome of FCoV infection, with statistical relevance.

Materials and methods

Animals and specimens

Samples were collected from 71 FIP cats and 93 FCoV-infected asymptomatic cats from 2005 to 2014 at the National Taiwan University Animal Hospital for an association analysis. This study required no specific ethical approval, as the analysis was performed retrospectively from samples of diseased animals routinely submitted to our diagnostic laboratory.

Seventy-one FIP cats, including 35.2% (25/71) purebred and 64.8% (45/71) mixed breed, were confirmed by necropsy. Pedigree cats including Scottish Fold (6/25), American Shorthair (4/25), Chinchilla (3/25), Exotic Shorthair (3/25), Siamese (2/25), European Shorthair (1/25) and Russian Blue (1/25). Most of the FIP cats (46/71) were less than one years old, the other 19 FIP cats were between the ages of one and three. The rest six FIP cats elder than

three were from 3.5 to 10 years old. Also, FCoV detection by reverse transcription-nested polymerase chain reaction (RT-nPCR) [21] was confirmed in disease-associated tissue, including body effusions and/or internal organs with the typical lesions of FIP. Ninety-three asymptomatic healthy cats were included as a control group, including 75.3% (50/93) mixed breed and 24.7% (23/93) purebred, of an age of three years old or less and showing no FIP-related signs upon enrolment in this study. Pedigree cats including American Shorthair (8/23), Scottish Fold (7/23), Chinchilla (4/23), Persian (2/23), Abyssinian (1/23) and European Shorthair (1/23). All the asymptomatic cats were positive for FCoV detection in at least one sample collected, including whole blood, nasal/oral/conjunctival/rectal swabs, and feces. In addition, the detection of two feline retroviruses, i.e., feline leukemia virus and feline immunodeficiency virus, was performed [22,23]. FIP cats and FCoV infected non-FIP cats with a positive result for either of the feline retroviruses were excluded from the association study.

Identification of SNPs in target sequences

Genomic DNA was isolated from buccal swabs or whole blood samples from each cat using a genomic DNA mini kit (Geneaid Biotech, New Taipei City, Taiwan). Partial *fTNFA* and *fCD209* sequences and five FIP-associated SNPs identified in Birman cats, as reported by Golovko et al., namely, *A1.196617776*, *A1.206840008*, *Un.59861682*, *A2.191286425*, and *E2.65509996*, were amplified [20] using the polymerase chain reaction (PCR). The primers and conditions are listed in Tables 1 and 2. Briefly, each reaction contained 1 μ L of template DNA, 500 nM of each primer, 200 μ M dNTP, 1.5 mM MgCl₂, and 0.6 U Phusion DNA polymerase (Thermo Scientific, Waltham, USA) in a total volume of 30 μ L with 1 \times Phusion HF buffer. The amplified products were subsequently sequenced using an auto-sequencer ABI 3730XL (Applied Biosystems, San Mateo, USA), and the obtained sequences were aligned by Geneious 4.8.5 (Biomatters, Auckland, New Zealand). The

Table 1 Primers used for SNP identification of *fTNFA* and *fCD209* gene

Target gene/region	Position ^a	Orientation ^b	Sequence (5' - 3')	T _A ^c	Amplicon size
<i>fTNFA</i> /5'-PRR ^d	-847 to -827	F	GAATCCCAGGGTTGCTTCA	65 °C	1018 bp
	+171 to +153	R	GCCGATCACTCCAAAGTGC		
<i>fCD209</i> /5'-PRR	-1057 to -1038	F	GAAGCGGGCTTCTTGTGAC	65 °C	1076 bp
	+19 to +1	R	GCTCCTGGGGTCACACAT		
<i>fCD209</i> /ECD ^e	+1818 to +1838	F	CCAAGATCTGATGCATCTGCT	67 °C	1350 bp
	+3168 to +3149	R	ATGAGCTCGTTCCTGATCT		

^aThe nucleotide positions are numerated from the translation start point (+1).

^bF: forward; R: reverse.

^cAnnealing temperature.

^d5'-proximal regulatory region.

^eExtracellular domain.

Table 2 Primers used for SNP identification of five suspected FIP-associated SNPs in Birman cats

SNP name	Chr-SNP position ^a	Orientation ^b	Sequence (5' - 3')	T _A ^c	Amplicon size
A1.196617776	A1-154265118	F	GGCAGTCAGAGAATGAGACAC	61 °C	337 bp
		R	TTGCCAGTTCTGCAGATTG		
A1.206840008	A1-164728174	F	AGGTGAAGTGTGTGTGCAT	61 °C	388 bp
		R	ATGTTCTGCTAGATGAGCCG		
Un.59861682	A1-155715831	F	CTCATCCCAGTTGATCACAC	61 °C	230 bp
		R	TTCTCCTGGAAAACCT		
A2.191286425	A2-126618108	F	AGCGTATCAAGTGCCTGC	61 °C	299 bp
		R	CCTTCCTGTTAGGTGCTTG		
E2.65509996	E2-54165589	F	CGCTTCAGTTTCCTTCCAG	61 °C	417 bp
		R	TCTGAGCCTTGGTCTTCTG		

^aChr: chromosome.

^bF: forward; R: reverse.

^cAnnealing temperature.

polymorphisms were further identified; the nucleotide positions of the SNPs are numerated from the translation start point (+1).

Association analysis

The association between the targeted SNPs and the occurrence of FIP was analyzed using Fisher's exact test, and a *P* value < 0.05 was considered to be a significant association.

Results

Polymorphism at *fTNFA* - 421 was found to be significantly associated with resistance to FIP

Because the overproduction of TNF- α is widely reported in FIP animals and is considered to contribute to the pathogenesis of FIP, we first screened the polymorphisms at the 5' terminus of the *fTNFA* gene, including the proximal regulatory region (PRR), the 5'-untranslated region (UTR), and part of exon 1, in 71 FIP and 93 control cats. Eight SNPs and three repeat regions were identified in the analyzed 1018 bp (Figure 1). One SNP located in exon

1 at position +23 results in the substitution of CGG to CAG, causing an amino acid change from Arg to Gly (R8G); the remaining SNPs were found in the PRR. The mean allele frequencies of the minor alleles ranged from 4.3% to 39.9%. To examine the association between the identified SNPs in *fTNFA* and the outcome of FCoV infection, the frequency of each genotype and allele was analyzed (Additional file 1). Only one allele (T allele) at position -421 appeared to be significantly associated with resistance to FIP (*P* = 0.009, OR = 3.925), whereas the others showed no significance to the disease (Table 3 and Additional file 1).

Polymorphisms in the extracellular domain and introns 6 and 7 of *fCD209* were found to be significantly associated with the disease outcome

fCD209 is an important co-receptor for both type I and II FCoV infection. To demonstrate an association between polymorphisms and FIP, we sequenced polymorphisms of the PRR, 5'-UTR, and extracellular domain (ECD) of




Figure 1 A schematic of the 5' terminal of the *fTNFA* gene analyzed in this study. A partial *fTNFA* sequence of 1018 bp was sequenced in this study, including the PRR, the 5'-UTR, and part of exon 1. All the SNPs and the corresponding positions are indicated with lines. Gray box: exon 1. Black boxes: repeat regions. SNPs located in the exon were shaded.

Table 3 FIP-associated genetic polymorphisms identified in the *fifNG*, *ftnFA*, and *fCD209* gene and their associations with FIP

Target gene/position	Susceptible allele	Resistant allele	P value	OR ^a	Reference
<i>fifNG</i> +428	C	T	0.03	3.4 (1.1-10.3)	[16]
<i>ftnFA</i> - 421	C	T	0.009	3.9 (1.3-11.8)	This study
<i>fCD209</i> + 1900	A	G	0.014	3.7 (1.3-10.5)	This study
<i>fCD209</i> + 2276	C	T	0.038	NA ^b	This study
<i>fCD209</i> + 2392	G	A	0.016	2.6 (1.2-5.5)	This study
<i>fCD209</i> + 2713	T	C	0.039	1.75 (1.1-2.9)	This study

^aOdds Ratio.
^bNot available.

fCD209 in 71 FIP and 93 control cats. Twenty-four SNPs and one AG repeat region were identified in the PRR, and one SNP was found in the 5'-UTR (Figure 2). Furthermore, polymorphism screening of ECD, revealed 25 SNPs and a G repeat in the 1350-bp region analyzed (Figure 2). Four SNPs were located in exons, including two SNPs in exon 6 at positions +1900 (TGG > TAG, W128*) and +1952 (AAC > AAA, N145K), one in exon 7 at position +2498 (ACG > ACT, T178T), and one in exon 8 at position +3070 (TTC > TCC, F241S). The remaining 21 SNPs were located in introns 6 and 7 (Figure 2). The mean allele frequencies of the minor alleles ranged from 0.6% to 47.6%.

To further demonstrate the association between the identified SNPs in *fCD209* and the outcome of FCoV infection, the frequency of each genotype and allele was analyzed. Among the 24 SNPs analyzed in the PRR and the 5'-UTR of the *fCD209* gene, no genotype or allele showed a significant association with the outcome of the infection (Additional file 2). In contrast, four SNPs, +1900 in the ECD of exon 6, +2276 and +2392 in intron 6, and +2713 in intron 7, identified in *fCD209* were found to be significantly associated with FIP. *fCD209* + 1900, located in the ECD, which is characterized by a G-to-A substitution that

leads to a premature stop codon (TGG > TAG, W128*), had a significantly increased frequency in FIP animals (18.31%) compared with the control animals (5.38%) ($P = 0.011$, OR = 3.95), and the A allele was shown to be significantly associated with susceptibility to FIP ($P = 0.014$, OR = 3.65) (Table 3 and Additional file 2). Furthermore, three SNPs, +2276, +2392, and +2713, located in introns 6 and 7 were found to be associated with disease susceptibility. A higher frequency of the T allele at position +2276 in the control cats was significantly associated with resistance to FIP ($P = 0.038$). Similarly, the A allele at position +2392, with a higher frequency in FCoV-infected asymptomatic cats, showed a significant association with disease resistance ($P = 0.016$, OR = 2.57). Moreover, the T allele at position +2713 was identified with a higher frequency in FIP cats, showing a significant association with disease susceptibility ($P = 0.039$, OR = 1.75) (Table 3 and Additional file 2).

Evaluation of the association between the FIP-associated SNPs reported in Birman cats and disease susceptibility in a cat population with higher genetic variability

Five SNPs were reported to be associated with the occurrence of FIP in Birman cats in a recent study using




Figure 2 A schematic of the partial *fCD209* gene analyzed in this study. The PRR, the 5'-UTR, and the ECD of *fCD209* were sequenced in this study. All the SNPs and the corresponding positions were indicated with lines. Gray boxes: exons. Black boxes: repeat regions. White box: 5' UTR. SNPs located in the exon were shaded.

genome-wide association analysis [20]. Due to the lack of information concerning the nucleotide sequence of the SNPs, the sequences were identified. The nucleotide sequences of *A1.196617776*, *A1.206840008*, *Un.59861682*, *A2.191286425*, and *E2.65509996* consisted of C/A, G/A, G/A, C/T, and C/T, respectively. To test for an association of these SNPs with the occurrence of FIP also applies to other breeds of cats, the frequency of each genotype and allele of the targeted SNPs was analyzed in all the FIP and control cats enrolled in this study, i.e., 29.3% purebred and 70.7% mixed breed. Among the five SNPs analyzed, neither the genotype nor allele percentage showed a significant association with the outcome of the disease (Additional file 3).

Association study

Among all the FIP animals analyzed in this study, 64 (64/71, 90.1%) cats were found to be effusive form (wet), and the rest seven of them (7/71, 9.9%) were non-effusive (dry). The association between the identified FIP-related SNPs (*fTNFA* - 421, *fCD209* + 1900, + 2276, + 2392 and +2713) and the form of FIP were analyzed. None of the SNPs was found to be associated with the biotype of FIP.

The number of the FIP-associated SNPs harbored is correlated to the disease outcome

This study attempted to distinguish multiple genetic traits associated with FIP susceptibility. The FIP-associated genetic polymorphisms identified in the *fIFNG* [16], *fTNFA*, and *fCD209* genes are summarized in Table 3. Moreover, the association between the number of FIP-associated SNPs harbored, including resistant or susceptible genotype/alleles identified at *fIFNG* +428, *fTNFA* - 421, and *fCD209* + 1900, + 2276, + 2392, and +2713, and the occurrence of FIP, was analyzed in this study (Table 4). The number of FIP-resistant SNPs carried was found to be associated with the protection of cats from FIP ($P = 0.002$), and the odds ratios of FIP and non-FIP cats carrying one or more (≥ 2) resistance SNPs were 3.06 and 6.01, respectively; this result indicates that cats carrying more resistant SNPs appear to have a lower chance of developing FIP.

However, the FIP cats were identified with a higher frequency as carrying more than one FIP-susceptible SNP (50.7%) than the control cats (29.0%), showing a significant association with disease susceptibility ($P = 0.0059$, OR = 2.51) (Table 4).

Discussion

FIP is an important infectious disease in cats, with nearly 100% mortality. However, an understanding of the host determinants in the occurrence of FIP has been limited to date. An imbalance between cellular and humoral immunity - with excess antibodies contributing to disease progression [4,24] and a significant decrease in IFN- γ production [4,8,10,13-15] - has been consistently observed in FIP animals. We recently identified the first host gene - *fIFNG* - showing an association between host genetic polymorphisms and FIP [16]. A T allele at *fIFNG* +428 was identified as a resistant allele, and the heterozygous genotypes (CT) at positions +401 and +408 were identified as associated with susceptibility to type I FCoV-induced FIP. In this study, five additional SNPs from *fTNFA* and *fCD209* were identified.

During the development of FIP, the upregulation of TNF- α , an important pro-inflammatory cytokine, has been consistently found [4,17,24,25]. The overproduction of TNF- α induces apoptosis in CD8+ T cells and is associated with the upregulation of a type II FCoV receptor, i.e., feline aminopeptidase N, which accelerates macrophage infection by the virus [26]. Moreover, TNF- α together with granulocyte monocyte-colony stimulating factor (GM-CSF), G-CSF, and other neutrophil survival factors are suggested to prolong the survival of neutrophils, activate monocytes/macrophages, and contribute to the formation of the pyogranulomatous lesions of FIP [25]. In several human diseases, the most commonly identified genetic polymorphisms are located at position -238 and -308 of the promoter region of *TNFA*, suggesting an effect on the binding of transcription factors [27], with susceptibility for the development of several viral diseases, including severe acute respiratory syndrome (SARS) [28], dengue hemorrhagic fever (DHF) [29], and hepatitis

Table 4 Association of the number of the disease-associated SNPs harbored, including *fIFNG* +428, *fTNFA* - 421, and *fCD209* + 1900, + 2276, + 2392, and +2713, and the outcome of FIP

Genotypes/alleles harbored	FIP number (%)	Non-FIP number (%)	P value	OR (95% CI)
Resistant SNPs				
= 0	59 (83.1%)	54 (58.1%)	0.002	Reference
= 1	10 (14.1%)	28 (30.1%)		3.06 (1.36 - 6.88)
≥ 2	2 (3%)	11 (11.8%)		6.01 (1.27 - 28.35)
Susceptible SNPs				
= 0	35 (49.3%)	66 (71.0%)	0.0059	Reference
≥ 1	36 (50.7%)	27 (29.0%)		2.51 (1.32 - 4.80)

B virus (HBV) infection [30]. Compared to the human gene (*TNFA* - 308), SNP located in a slightly upstream region of the feline gene (*fTNFA* - 421) was found to be significantly associated with the occurrence of FIP. The variant *fTNFA* - 421 T allele was significantly associated with resistance to FIP. Through DNA transcriptional factor binding site prediction, we found that an *fTNFA* - 421 C to T mutation might affect the binding of some transcription factors, such as myeloid zinc finger 1 (MZF1) [31], a transcriptional regulator [32]. Through promoter binding, MZF1 was identified to function as a transcription activator of hematopoietic cells in vitro [32]. Because macrophages are hematopoietic cells, the loss of the binding site for such transcription factors might decrease TNF- α production in macrophages, which might prevent the immunopathogenesis and result in a resistant phenotype.

DC-SIGN, which recognizes high-mannose oligosaccharides as its ligand, is a co-receptor augmenting many viral infections, including human immunodeficiency virus [33], dengue virus [34], HBV [35], and SARS-coronavirus [36]. In addition, feline DC-SIGN also serves as a co-receptor and is involved in infection by FCoV [19]. A human SNP in the promoter region of *CD209* (-336 A/G) was identified as related to disease prognosis [36]. A variant of *CD209* - 336 was reported to affect the binding of SP1-like transcription factor and might modulate transcriptional activities [34], indicating that *CD209* - 336 plays a crucial role in disease pathogenesis. However, in our study, none of the SNPs in the promoter region showed an association with the outcome of FCoV infection. The identified FIP-associated SNPs, +1900 G/A, +2276 C/T, +2392 G/A, and +2713 C/T, were all located at the 3' end of *fCD209*. The polymorphism identified at *fCD209* + 1900 is located in the lectin binding domain of the ECD; a G to A substitution leads to a change from a tryptophan at amino acid 128 to a stop codon. This mutation might lead to an abortive mRNA [37] or a truncated protein if the mRNA is successfully translated. DC-SIGN serves as a pattern recognition receptor that interact with numerous pathogens, including FCoV, and mediates the clustering of DC with naive T cells [38]. Additionally, with a type II transmembrane domain, the truncated fDC-SIGN identified in this study (+1900) might still be expressed on the cell surface. However, it remains to be elucidated how the truncated protein, bearing only one half of the authentic protein, affect the normal function of DC-SIGN, i.e., pathogen recognition or T cell activation. The other three FIP-associated SNPs were located in introns 6 and 7. Although these polymorphisms are located in introns, which apparently would not affect the protein, a regulatory effect cannot be excluded. In humans, the SNP at *IFNG* +874, located in intron 1, was found to alter the binding activity of nuclear factor kappa-light-chain-enhancer of activated B cells and

influenced the production of IFN- γ [39]. In swine, an SNP (G3072A) in intron 3 of insulin-like growth factor 2 affected the binding of muscle growth regulator and was associated with the muscle content [40].

Several pedigreed cats, including Abyssinians, Himalayans, Birmans, Bengals, Ragdolls, and Rexes, were reported to have a higher incidence of FIP than other breed cats [2]. Recently, a genome-wide association study of Danish and American Birman cats populations identified five SNPs involved in FIP susceptibility [20]. However, none of these SNPs showed a similar correlation in the present study, concordant with the postulation by the authors that these associations might only be relevant to Birman breed or other breeds with similar genetic traits [24].

In addition to the three SNPs in *fIFNG*, we identified in this study five more SNPs from two genes - one in *fTNFA* and four in *fCD209* - that are associated with the occurrence of FIP. As the susceptibility or resistance to viral infections is a complex phenotype regulated by multiple interacting genes and gene networks, genes related to innate and adaptive immunity and other host genes remain to be pursued. The combination of all the FIP susceptibility genotypes into a single typing diagnosis assay should facilitate the screening of FIP-resistant cats in breeding and eventually decrease the loss of cats to this incurable disease.

Additional files

Additional file 1: Frequencies of the *fTNFA* genotypes and alleles and associations with the outcome of FCoV infection. The promoter variant, *fTNFA* - 421 T, was found to be a disease-resistance genotype.

Additional file 2: Frequencies of the *fCD209* genotypes and alleles and associations with the outcome of FCoV infection. Polymorphisms of *fCD209*, including *fCD209* + 1900, + 2276, + 2392 and +2713, were found to be significantly associated with the disease outcome.

Additional file 3: Frequencies of the genotypes and alleles of the proposed FIP-associated SNPs in Birman cats and associations with FIP. Neither the percentage of genotypes nor alleles was found to be associated with the outcome of FCoV infection in disease and non-disease group.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

YTW performed the sampling and preparation, DNA/RNA extraction, FCoV detection, the amplification and sequencing of *fIFNG*, *fTNFA*, *fCD209* and five Birman FIP cat-associated SNPs and further analysis and prepared the manuscript. LEH participated in primers design, DNA extraction, *fIFNG*, *fTNFA* and *fCD209* amplification, sequencing and further analysis and prepared the manuscript. YRD performed DNA extraction and *fIFNG*, *fTNFA* and *fCD209* amplification. LLC conceived of the study, participated in study design and coordination and contributed to the preparation of the manuscript. All authors read and approved the final manuscript.

Acknowledgements

We would like to thank Betty A. Wu-Hsieh from Graduate Institute of Immunology, National Taiwan University for the valuable suggestion in the selection of candidate genes. This work was supported by the grant MOST 103-2313-B-002-042 from the Ministry of Science and Technology, Taiwan.

Received: 28 August 2014 Accepted: 26 November 2014
Published online: 16 December 2014

References

- Hartmann K: Feline infectious peritonitis. *Vet Clin North Am Small Anim Pract* 2005, **35**:39–79.
- Pedersen NC: A review of feline infectious peritonitis virus infection: 1963–2008. *J Feline Med Surg* 2009, **11**:225–258.
- Holzworth J: Some important disorders of cats. *Cornell Vet* 1963, **53**:157–160.
- Kipar A, Meli ML: Feline infectious peritonitis: still an enigma? *Vet Pathol* 2014, **51**:505–526.
- Satoh R, Furukawa T, Kotake M, Takano T, Motokawa K, Gemma T, Watanabe R, Arai S, Hohdatsu T: Screening and identification of T helper 1 and linear immunodominant antibody-binding epitopes in the spike 2 domain and the nucleocapsid protein of feline infectious peritonitis virus. *Vaccine* 2011, **29**:1791–1800.
- de Groot-Mijnes JD, van Dun JM, van der Most RG, de Groot RJ: Natural history of a recurrent feline coronavirus infection and the role of cellular immunity in survival and disease. *J Virol* 2005, **79**:1036–1044.
- Hajjema BJ, Volders H, Rottier PJ: Live, attenuated coronavirus vaccines through the directed deletion of group-specific genes provide protection against feline infectious peritonitis. *J Virol* 2004, **78**:3863–3871.
- Kiss I, Poland AM, Pedersen NC: Disease outcome and cytokine responses in cats immunized with an avirulent feline infectious peritonitis virus (FIPV)-UCD1 and challenge-exposed with virulent FIPV-UCD8. *J Feline Med Surg* 2004, **6**:89–97.
- Hohdatsu T, Yamato H, Ohkawa T, Kaneko M, Motokawa K, Kusuhara H, Kaneshima T, Arai S, Koyama H: Vaccine efficacy of a cell lysate with recombinant baculovirus-expressed feline infectious peritonitis (FIP) virus nucleocapsid protein against progression of FIP. *Vet Microbiol* 2003, **97**:31–44.
- Dean GA, Olivry T, Stanton C, Pedersen NC: In vivo cytokine response to experimental feline infectious peritonitis virus infection. *Vet Microbiol* 2003, **97**:1–12.
- Pesteanu-Somogyi LD, Radzai C, Pressler BM: Prevalence of feline infectious peritonitis in specific cat breeds. *J Feline Med Surg* 2006, **8**:1–5.
- Worthing KA, Wigney DI, Dhand NK, Fawcett A, McDonagh P, Malik R, Norris JM: Risk factors for feline infectious peritonitis in Australian cats. *J Feline Med Surg* 2012, **14**:405–412.
- Giordano A, Paltrinieri S: Interferon-gamma in the serum and effusions of cats with feline coronavirus infection. *Vet J* 2009, **180**:396–398.
- Gelain ME, Meli M, Paltrinieri S: Whole blood cytokine profiles in cats infected by feline coronavirus and healthy non-FCoV infected specific pathogen-free cats. *J Feline Med Surg* 2006, **8**:389–399.
- Gunn-Moore DA, Caney SM, Gruffydd-Jones TJ, Helps CR, Harbour DA: Antibody and cytokine responses in kittens during the development of feline infectious peritonitis (FIP). *Vet Immunol Immunopathol* 1998, **65**:221–242.
- Hsieh LE, Chueh LL: Identification and genotyping of feline infectious peritonitis-associated single nucleotide polymorphisms in the feline interferon-gamma gene. *Vet Res* 2014, **45**:57.
- Takano T, Hohdatsu T, Hashida Y, Kaneko Y, Tanabe M, Koyama H: A “possible” involvement of TNF-alpha in apoptosis induction in peripheral blood lymphocytes of cats with feline infectious peritonitis. *Vet Microbiol* 2007, **119**:121–131.
- Regan AD, Ousterout DG, Whittaker GR: Feline lectin activity is critical for the cellular entry of feline infectious peritonitis virus. *J Virol* 2010, **84**:7917–7921.
- Van Hamme E, Desmarests L, Dewerchin HL, Nauwynck HJ: Intriguing interplay between feline infectious peritonitis virus and its receptors during entry in primary feline monocytes. *Virus Res* 2011, **160**:32–39.
- Golovko L, Lyons LA, Liu H, Sorensen A, Wehnert S, Pedersen NC: Genetic susceptibility to feline infectious peritonitis in Birman cats. *Virus Res* 2013, **175**:58–63.
- Herrewegh AA, de Groot RJ, Cepica A, Egberink HF, Horzinek MC, Rottier PJ: Detection of feline coronavirus RNA in feces, tissues, and body fluids of naturally infected cats by reverse transcriptase PCR. *J Clin Microbiol* 1995, **33**:684–689.
- Nishimura Y, Goto Y, Pang H, Endo Y, Mizuno T, Momoi Y, Watarai T, Tsujimoto H, Hasegawa A: Genetic heterogeneity of env gene of feline immunodeficiency virus obtained from multiple districts in Japan. *Virus Res* 1998, **57**:101–112.
- Stiles J, Bienzle D, Render JA, Buyukmihci NC, Johnson EC: Use of nested polymerase chain reaction (PCR) for detection of retroviruses from formalin-fixed, paraffin-embedded uveal melanomas in cats. *Vet Ophthalmol* 1999, **2**:113–116.
- Pedersen NC: An update on feline infectious peritonitis: virology and immunopathogenesis. *Vet J* 2014, **201**:123–132.
- Takano T, Azuma N, Satoh M, Toda A, Hashida Y, Satoh R, Hohdatsu T: Neutrophil survival factors (TNF-alpha, GM-CSF, and G-CSF) produced by macrophages in cats infected with feline infectious peritonitis virus contribute to the pathogenesis of granulomatous lesions. *Arch Virol* 2009, **154**:775–781.
- Takano T, Hohdatsu T, Toda A, Tanabe M, Koyama H: TNF-alpha, produced by feline infectious peritonitis virus (FIPV)-infected macrophages, upregulates expression of type II FIPV receptor feline aminopeptidase N in feline macrophages. *Virology* 2007, **364**:64–72.
- Smith AJ, Humphries SE: Cytokine and cytokine receptor gene polymorphisms and their functionality. *Cytokine Growth Factor Rev* 2009, **20**:43–59.
- Wang S, Wei M, Han Y, Zhang K, He L, Yang Z, Su B, Zhang Z, Hu Y, Hui W: Roles of TNF-alpha gene polymorphisms in the occurrence and progress of SARS-Cov infection: a case-control study. *BMC Infect Dis* 2008, **8**:27.
- Perez AB, Sierra B, Garcia G, Aguirre E, Babel N, Alvarez M, Sanchez L, Valdes L, Volk HD, Guzman MG: Tumor necrosis factor-alpha, transforming growth factor-beta1, and interleukin-10 gene polymorphisms: implication in protection or susceptibility to dengue hemorrhagic fever. *Hum Immunol* 2010, **71**:1135–1140.
- Xia Q, Zhou L, Liu D, Chen Z, Chen F: Relationship between TNF- < alpha > gene promoter polymorphisms and outcomes of hepatitis B virus infections: a meta-analysis. *PLoS One* 2011, **6**:e19606.
- Heinemeyer T, Wingender E, Reuter I, Hermjakob H, Kel AE, Kel OV, Ignatieva EV, Ananko EA, Podkolodnaya OA, Kolpakov FA, Podkolodny NL, Kolchanov NA: Databases on transcriptional regulation: TRANSFAC, TRRD and COMPEL. *Nucleic Acids Res* 1998, **26**:362–367.
- Morris JF, Rauscher FJ 3rd, Davis B, Klemsz M, Xu D, Tenen D, Hromas R: The myeloid zinc finger gene, MZF-1, regulates the CD34 promoter in vitro. *Blood* 1995, **86**:3640–3647.
- Boily-Larouche G, Milev MP, Zijenah LS, Labbe AC, Zannou DM, Humphrey JH, Ward BJ, Poudrier J, Moulard AJ, Cohen EA, Roger M: Naturally-occurring genetic variants in human DC-SIGN increase HIV-1 capture, cell-transfer and risk of mother-to-child transmission. *PLoS One* 2012, **7**:e40706.
- Sakuntabhai A, Turbpaiboon C, Casademont I, Chuansumrit A, Lowhnoo T, Kajaste-Rudnitski A, Kalayanarooj SM, Tangnaratchakik K, Tangthawornchaikul N, Vasanawathana S, Chaiyaratana W, Yenchitsomanus PT, Suriyaphol P, Avirutnan P, Chokephaibulkit K, Matsuda F, Yoksan S, Jacob Y, Lathrop GM, Malasit P, Despres P, Julier C: A variant in the CD209 promoter is associated with severity of dengue disease. *Nat Genet* 2005, **37**:507–513.
- Rebbani K, Ezzikouri S, Marchio A, Ababou M, Kitab B, Dejean A, Kandil M, Pineaou P, Benjelloun S: Common polymorphic effectors of immunity against hepatitis B and C modulate susceptibility to infection and spontaneous clearance in a Moroccan population. *Infect Genet Evol* 2014, **26**:1–7.
- Chan KY, Xu MS, Ching JC, So TM, Lai ST, Chu CM, Yam LY, Wong AT, Chung PH, Chan VS, Lin CL, Sham PC, Leung GM, Peiris JS, Khoo US: CD209 (DC-SIGN) -336A > G promoter polymorphism and severe acute respiratory syndrome in Hong Kong Chinese. *Hum Immunol* 2010, **71**:702–707.
- Waston JD, Baker TA, Bell SP, Gann A, Levine M, Losick R: Translation. In *Molecular Biology of the Gene*. Edited by W B. New York: Cold harbor laboratory press; 2008:457–519.
- den Dunnen J, Gringhuis SI, Geijtenbeek TB: Innate signaling by the C-type lectin DC-SIGN dictates immune responses. *Cancer Immunol Immunother* 2009, **58**:1149–1157.
- Pravica V, Perrey C, Stevens A, Lee JH, Hutchinson IV: A single nucleotide polymorphism in the first intron of the human IFN-gamma gene: absolute correlation with a polymorphic CA microsatellite marker of high IFN-gamma production. *Hum Immunol* 2000, **61**:863–866.
- Butter F, Kappei D, Buchholz F, Vermeulen M, Mann M: A domesticated transposon mediates the effects of a single-nucleotide polymorphism responsible for enhanced muscle growth. *EMBO Rep* 2010, **11**:305–311.

doi:10.1186/s13567-014-0123-6

Cite this article as: Wang *et al.*: Polymorphisms in the feline TNFA and CD209 genes are associated with the outcome of feline coronavirus infection. *Veterinary Research* 2014 **45**:123.