



# Strong modularity of reducible Galois representations

Nicolas Billerey, Ricardo Menares

## ► To cite this version:

Nicolas Billerey, Ricardo Menares. Strong modularity of reducible Galois representations. 2016. hal-01295749v1

**HAL Id: hal-01295749**

**<https://hal.science/hal-01295749v1>**

Preprint submitted on 31 Mar 2016 (v1), last revised 23 May 2016 (v2)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# STRONG MODULARITY OF REDUCIBLE GALOIS REPRESENTATIONS

NICOLAS BILLEREY <sup>¶</sup> AND RICARDO MENARES <sup>†</sup>

ABSTRACT. Let  $\rho: \text{Gal}(\overline{\mathbf{Q}}/\mathbf{Q}) \rightarrow \text{GL}_2(\overline{\mathbf{F}}_l)$  be an odd, semi-simple Galois representation. Here,  $l \geq 5$  is prime and  $\overline{\mathbf{F}}_l$  is an algebraic closure of the finite field  $\mathbf{Z}/l\mathbf{Z}$ . When the representation is irreducible, the strongest form of Serre's original modularity conjecture (now a theorem of Khare and Wintenberger) asserts that  $\rho$  arises from a cuspidal eigenform of type  $(N, k, \varepsilon)$  over  $\overline{\mathbf{F}}_l$ , where  $N$ ,  $k$  and  $\varepsilon$  are, respectively, the level, weight and character attached to  $\rho$  by Serre.

In this paper we characterize, under the assumption  $l > k + 1$ , reducible semi-simple representations, that we call strongly modular, such that the same result holds. This characterization generalizes a classical theorem of Ribet pertaining to the case  $N = 1$ . When the representation is not strongly modular, we give a necessary and sufficient condition on primes  $p$  not dividing  $Nl$  for which  $\rho$  arises in level  $Np$ , hence generalizing a classical theorem of Mazur concerning the case  $(N, k) = (1, 2)$ .

The proofs rely on the classical analytic theory of Eisenstein series and on local properties of automorphic representations attached to newforms.

## INTRODUCTION

Let  $l$  be a prime number. We denote by  $\overline{\mathbf{F}}_l$  and  $\overline{\mathbf{Q}}$  algebraic closures of  $\mathbf{F}_l = \mathbf{Z}/l\mathbf{Z}$  and the rational field  $\mathbf{Q}$  respectively. In this article we are interested in Galois representations of the form

$$(1) \quad \rho: \text{Gal}(\overline{\mathbf{Q}}/\mathbf{Q}) \longrightarrow \text{GL}_2(\overline{\mathbf{F}}_l),$$

where  $\rho$  is a continuous homomorphism. Let  $N \geq 1$  and  $k \geq 2$  be two integers with  $N$  coprime to  $l$  and let  $\varepsilon: (\mathbf{Z}/N\mathbf{Z})^\times \rightarrow \overline{\mathbf{F}}_l^\times$  be a character. Let  $f$  be a cusp form of type  $(N, k, \varepsilon)$  over  $\overline{\mathbf{F}}_l$  (in the sense of [Ser87, Déf. p. 193]) which is an eigenfunction for the  $p$ -th Hecke operator with eigenvalue  $a_p$  in  $\overline{\mathbf{F}}_l$  for each prime number  $p$ . By work of Deligne, to such a form  $f$ , one can attach a (unique up to isomorphism) semi-simple odd Galois representation  $\rho_f$  which is unramified outside  $Nl$  and satisfies the following property : If  $\text{Frob}_p$  denotes a Frobenius element at a prime  $p \nmid Nl$ , then the characteristic polynomial of  $\rho_f(\text{Frob}_p)$  is given by

$$X^2 - a_p X + \varepsilon(p)p^{k-1}.$$

---

*Date:* March 31, 2016.

2010 *Mathematics Subject Classification.* Primary 11F80, 11F33. Secondary 11F70.

<sup>¶</sup> Partially supported by CNRS and ANR-14-CE-25-0015 Gardio.

<sup>†</sup> Pontificia Universidad Católica de Valparaíso. Partially supported by PUCV grant 037.469/2015.

According to a standard terminology, a Galois representation  $\rho$  is called *modular* if it is isomorphic to  $\rho_f$  for some  $f$  as above. In that case, we also say that  $\rho$  arises from  $f$ .

Moreover, to any given Galois representation  $\rho$ , Serre attaches in [Ser87, §§1-2] a triple  $(N, k, \varepsilon)$ , which we refer to as the Serre type of  $\rho$ , consisting of an integer  $N \geq 1$  coprime to  $l$ , an integer  $k \geq 2$  and a group homomorphism  $\varepsilon: (\mathbf{Z}/N\mathbf{Z})^\times \rightarrow \overline{\mathbf{F}}_l^\times$  which are called the conductor, weight and character of  $\rho$  respectively.

In this paper, we shall say that a Galois representation  $\rho$  is *strongly modular* if it arises from a cuspidal eigenform  $f$  over  $\overline{\mathbf{F}}_l$  of type  $(N, k, \varepsilon)$  where  $(N, k, \varepsilon)$  is the Serre type of  $\rho$ .

With this terminology, the strong form ([Ser87, (3.2.4?)]) of Serre's modularity conjecture, proved by Khare and Wintenberger ([KW09a], [KW09b]), asserts that any odd, irreducible Galois representation  $\rho$  as in (1), with  $l \geq 5$ , is strongly modular.

In this article, we address the case where  $\rho$  is reducible. Let

$$\nu_1, \nu_2: \text{Gal}(\overline{\mathbf{Q}}/\mathbf{Q}) \longrightarrow \overline{\mathbf{F}}_l^\times$$

be continuous characters and assume that  $\rho = \nu_1 \oplus \nu_2$  defines an odd (semi-simple) Galois representation of Serre type  $(N, k, \varepsilon)$ . Then,  $\rho$  is modular (e.g. see [BM15, Thm. 2.1]) but need not be strongly modular. Our task is to provide a necessary and sufficient condition for such a reducible Galois representation to be strongly modular. Thanks to Ribet, such a characterization is known in the case  $N = 1$  under the assumption  $l > k + 1$  (see [Rib75, Lem. 5.2] or [BM15, Cor. 3.7] for a reformulation in this context). Under the same assumption, we prove in this paper a generalization of this result to arbitrary conductors.

Let  $\eta: \text{Gal}(\overline{\mathbf{Q}}/\mathbf{Q}) \rightarrow \overline{\mathbf{F}}_l^\times$  be a character unramified at  $l$ . For any integer  $k \geq 2$  satisfying  $l > k + 1$ , we define in paragraph 1.2 a mod  $l$  Bernoulli number  $B_{k,\eta} \in \overline{\mathbf{F}}_l$  associated with  $\eta$  (our  $B_{k,\eta}$  is essentially the reduction of a classical  $k$ -th Bernoulli number attached to a lift of  $\eta$ , but some care has to be taken due to denominators and the choice of place). For every prime number  $p$ , set

$$\eta(p) = \begin{cases} \eta(\text{Frob}_p) & \text{if } \eta \text{ is unramified at } p \\ 0 & \text{if } \eta \text{ is ramified at } p. \end{cases}$$

In this notation, the following is the main result of the paper.

**Theorem 1.** *Let  $\nu_1, \nu_2: \text{Gal}(\overline{\mathbf{Q}}/\mathbf{Q}) \rightarrow \overline{\mathbf{F}}_l^\times$  be characters defining an odd (semi-simple) Galois representation  $\rho = \nu_1 \oplus \nu_2$  of Serre type  $(N, k, \varepsilon)$  with  $l > k + 1$ . Then, there exist characters  $\varepsilon_1, \varepsilon_2: \text{Gal}(\overline{\mathbf{Q}}/\mathbf{Q}) \rightarrow \overline{\mathbf{F}}_l^\times$  unramified at  $l$  such that  $\rho = \varepsilon_1 \oplus \varepsilon_2 \chi_l^{k-1}$ , where  $\chi_l$  is the mod  $l$  cyclotomic character. Set  $\eta = \varepsilon_1^{-1} \varepsilon_2$ . The representation  $\rho$  is strongly modular if and only if*

$$\text{either } B_{k,\eta} = 0 \quad \text{or} \quad \eta(p)p^k = 1 \text{ for some prime } p \text{ dividing } N.$$

If the representation  $\rho$  alluded to above is not strongly modular, we give in Theorem 2 below a precise characterization (under the same assumption as before) of the primes  $M \nmid Nl$  for which  $\rho$  arises from a cusp form of type  $(NM, k, \varepsilon)$ . Such a theorem extends a result of Mazur ([Maz77,

Prop. 5.12]), that handles the case  $(N, k) = (1, 2)$ , to arbitrary weights and conductors.

**Theorem 2.** *In the same notation and under the same assumptions as in Theorem 1, assume moreover that  $\rho$  is not strongly modular. Let  $M$  be a prime number not dividing  $Nl$ . Then  $\rho$  arises from a modular form of type  $(NM, k, \varepsilon)$  if and only if*

$$\begin{cases} M \equiv 1 \pmod{l} & \text{if } (N, k) = (1, 2) \\ \eta(M)M^k = 1 & \text{if } (N, k) \neq (1, 2). \end{cases} \quad (\text{Mazur})$$

*In particular, there are infinitely many such primes.*

The paper is organized as follows. In Section 1, we define the Bernoulli numbers attached to mod  $l$  Galois characters that appear in the statement of Theorem 1 above and compute the constant term at the various cusps of a particular Eisenstein series which is of crucial use in the proofs of our results. After quickly recalling in Section 2 some background on cuspidal eigenforms and Hecke operators in the adelic setting, we prove in Section 3 our two main theorems. Although they are stated in positive characteristic, their proofs strongly rely on properties of characteristic zero eigenforms and their attached automorphic representations.

**Acknowledgements:** The authors wish to thank Vinayak Vatsal for inspiring discussions and the Pacific Institute for the Mathematical Sciences in Vancouver for providing ideal conditions to carry out part of this project.

## 1. BERNOULLI NUMBERS AND EISENSTEIN SERIES

In this section we recall some classical definitions and integrality results on Bernoulli numbers attached to Dirichlet characters. Also, we compute the constant term in the  $q$ -expansion at the cusps of the modular curve  $X_1(N)$  of some specific Eisenstein series that will be used in the sequel. The final computation is stated in Proposition 4 below.

**1.1. Notation and definitions.** Let  $\phi$  be a primitive Dirichlet character of conductor  $\mathfrak{f} \geq 1$ . The Gauss sum attached to  $\phi$  is defined by

$$W(\phi) = \sum_{n=1}^{\mathfrak{f}} \phi(n) e^{2i\pi n/\mathfrak{f}}.$$

It is a non-zero algebraic integer whose norm is a power of  $\mathfrak{f}$ . The (generalized) Bernoulli numbers  $(B_{m,\phi})_{m \geq 1}$  associated with  $\phi$  are defined by the following expansion

$$(2) \quad \sum_{n=1}^{\mathfrak{f}} \phi(n) \frac{te^{nt}}{e^{\mathfrak{f}t} - 1} = \sum_{m \geq 0} B_{m,\phi} \frac{t^m}{m!}.$$

Note that when  $\phi = \mathbf{1}$  is the trivial character (of conductor 1), then, for every integer  $m \geq 2$ , we have  $B_{m,\phi} = B_m$  where  $B_m$  denotes the classical  $m$ -th Bernoulli number.

**1.2. Bernoulli numbers of mod  $l$  characters.** Let  $\eta: \text{Gal}(\overline{\mathbf{Q}}/\mathbf{Q}) \rightarrow \overline{\mathbf{F}}_l^\times$  be a Galois character unramified at  $l$ . Denote by  $\mathfrak{c}_0$  the conductor of  $\eta$  (coprime to  $l$  by assumption) and identify  $\eta$  with a character

$$\eta: (\mathbf{Z}/\mathfrak{c}_0\mathbf{Z})^\times \longrightarrow \overline{\mathbf{F}}_l^\times.$$

The aim of this paragraph is to define the  $k$ -th Bernoulli number attached to  $\eta$  for any integer  $k \geq 2$  such that  $l > k + 1$ . This definition relies on integrality properties of Bernoulli numbers attached to Dirichlet characters which we now recall.

Let  $w$  be a place of  $\overline{\mathbf{Q}}$  above  $l$  and let  $\overline{\mathbf{Z}}_w$  be the local ring of  $w$ -integral algebraic numbers in  $\overline{\mathbf{Q}}$ . The residue field  $k_w$  of  $w$  identifies with an algebraic closure of  $\mathbf{F}_l$ . Fix an isomorphism  $\iota: k_w \rightarrow \overline{\mathbf{F}}_l$  and consider the composition map

$$\nu_w: \overline{\mathbf{Z}}_w \rightarrow k_w \xrightarrow{\iota} \overline{\mathbf{F}}_l.$$

We may then consider the multiplicative lift

$$\psi: (\mathbf{Z}/\mathfrak{c}_0\mathbf{Z})^\times \longrightarrow \overline{\mathbf{Z}}^\times$$

of  $\eta$  with respect to  $w$ . That is,  $\psi$  is the unique character with values in the roots of unity of prime-to- $l$  order such that

$$\nu_w(\psi(x)) = \eta(x), \quad \text{for all } x \in (\mathbf{Z}/\mathfrak{c}_0\mathbf{Z})^\times.$$

We now state the integrality result we need to define our Bernoulli numbers.

**Lemma 3.** *For any integer  $k \geq 2$  such that  $l > k + 1$ , the Bernoulli number  $B_{k,\psi}$  is  $w$ -integral, .*

*Proof.* Let  $k$  be an integer as in the statement of the lemma. We easily check on the definition (2) that if  $\psi(-1) \neq (-1)^k$ , then  $B_{k,\psi} = 0$ . Assume therefore that  $\psi(-1) = (-1)^k$ . If  $\psi$  is the trivial character, then  $k$  must be an even integer and the corresponding Bernoulli number  $B_{k,\psi}$  is nothing but the classical Bernoulli number  $B_k$ . The Van Staudt-Clausen theorem ensures that the prime divisors  $p$  of the denominator of  $B_k$  satisfy  $p - 1 \mid k$ . Since  $l > k + 1$ , the prime number  $l$  does not divide the denominator of  $B_k$ , as desired.

Assume therefore  $\psi$  is non trivial. Let

$$d = \begin{cases} 1 & \text{if } \mathfrak{c}_0 \text{ admits two different prime divisors} \\ 2 & \text{if } \mathfrak{c}_0 = 4 \\ 1 & \text{if } \mathfrak{c}_0 = 2^n, n > 2 \\ k\mathfrak{c}_0 & \text{if } \mathfrak{c}_0 > 2 \text{ is a prime number} \\ 1 - \psi(1+p) & \text{if } \mathfrak{c}_0 = p^n, p > 2, n > 1, p \text{ is a prime number} \end{cases}$$

By a theorem of Carlitz (see [Car59a] and [Car59b]),  $dk^{-1}B_{k,\psi}$  is an algebraic integer. Hence, we are reduced to verify that  $w$  does not divide  $d$ .

Assume that  $\mathfrak{c}_0 = p^n$ , where  $p$  is an odd prime number and  $n \geq 2$ . We assume by contradiction that  $w$  divides  $d = 1 - \psi(1+p)$ . Let  $H \subseteq (\mathbf{Z}/p^n\mathbf{Z})^\times$  be the subgroup spanned by  $1+p$ . Taking the reduction map attached to  $w$ , we conclude that  $\eta$  is trivial on  $H$ . Since  $H$  is the kernel of the natural map  $(\mathbf{Z}/p^n\mathbf{Z})^\times \rightarrow (\mathbf{Z}/p\mathbf{Z})^\times$ , we conclude that  $\eta$  can be factored through  $(\mathbf{Z}/p\mathbf{Z})^\times$ , contradicting the primitivity of  $\eta$ .

If  $\mathfrak{c}_0 \geq 3$  is not of the form discussed in the previous paragraph, the fact that  $l \nmid d$  clearly follows from the definition of  $d$  and the hypothesis on  $k, l$  and  $\mathfrak{c}_0$ .  $\square$

Using this result, we now set, for any integer  $k$  as above,

$$(3) \quad B_{k,\eta} = \nu_w(B_{k,\psi}) \in \overline{\mathbf{F}}_l.$$

Let  $w'$  be another place of  $\overline{\mathbf{Q}}$  over  $l$ . There exists  $\sigma \in \text{Gal}(\overline{\mathbf{Q}}/\mathbf{Q})$  such that  $w' = \sigma(w)$  and we identify the residue field  $k_{w'}$  (of the ring of  $w'$ -integral algebraic numbers in  $\overline{\mathbf{Q}}$ ) with  $\overline{\mathbf{F}}_l$  via  $\iota \circ \sigma^{-1}$ . Then  $\sigma(\psi)$  is the multiplicative lift of  $\eta$  with respect to the place  $w' = \sigma(w)$  and since we have

$$B_{k,\sigma(\psi)} = \sigma(B_{k,\psi})$$

the definition (3) is independent of the choice of the place  $w$ . We refer to  $B_{k,\eta} \in \overline{\mathbf{F}}_l$  as the  $k$ -th Bernoulli number associated with  $\eta$ .

**1.3. The setting.** In this paragraph we set some notation and definitions that will be used in the rest of this section. Let  $k \geq 2$  be an integer. We set

$$C_k = \frac{(-2i\pi)^k}{(k-1)!}.$$

Let

$$\chi_i: (\mathbf{Z}/\mathfrak{c}_i\mathbf{Z})^\times \longrightarrow \mathbf{C}^\times, \quad i = 1, 2$$

be primitive Dirichlet characters such that  $\chi_1(-1)\chi_2(-1) = (-1)^k$ . Denote by  $\overline{\chi}_i$  the complex conjugate of  $\chi_i$ ,  $i = 1, 2$ . Put  $N = \mathfrak{c}_1\mathfrak{c}_2$ . For  $k \geq 3$  and  $z$  in the complex upper-half plane  $\mathfrak{H}$ , let

$$G_k^{\chi_1, \chi_2}(z) = \sum_{(m,n) \in \mathbf{Z}^2 \setminus \{(0,0)\}} \frac{\chi_1(m)\overline{\chi}_2(n)}{(mz+n)^k}.$$

On the other hand, for any  $\varepsilon > 0$ , we consider

$$G_{2,\varepsilon}^{\chi_1, \chi_2}(z) = \sum_{(m,n) \in \mathbf{Z}^2 \setminus \{(0,0)\}} \frac{\chi_1(m)\overline{\chi}_2(n)}{(mz+n)^2 |mz+n|^{2\varepsilon}}, \quad (z \in \mathfrak{H}).$$

We remark that our functions  $G_k^{\chi_1, \chi_2}(z)$  ( $k \geq 3$ ) and  $G_{2,\varepsilon}^{\chi_1, \chi_2}(z)$  correspond to the functions  $E_k(z; \chi, \psi)$  and  $E_2(z, \varepsilon; \chi, \psi)$  respectively defined in Eq. (7.1.1) and (7.2.1) of [Miy06] with  $(\chi, \psi) = (\chi_1, \overline{\chi}_2)$ .

From now on, and until the end of this section, assume that either  $N > 1$  or  $k > 2$  and denote by  $E_k^{\chi_1, \chi_2}$  the function defined by

$$(4) \quad E_k^{\chi_1, \chi_2}(z) = -\delta(\chi_1) \frac{B_{k, \chi_2}}{2k} + \sum_{n \geq 1} \sigma_{k-1}^{\chi_1, \chi_2}(n) q^n, \quad (q = e^{2\pi iz}, z \in \mathfrak{H})$$

where

$$\sigma_{k-1}^{\chi_1, \chi_2}(n) = \sum_{m|n} \chi_1(n/m) \chi_2(m) m^{k-1}, \quad \delta(\chi_1) = \begin{cases} 1 & \text{if } \chi_1 \text{ is trivial} \\ 0 & \text{otherwise} \end{cases}$$

and  $B_{k, \chi_2}$  denotes the  $k$ -th Bernoulli number associated with  $\chi_2$  (see paragraph 1.1).

According to [Miy06], Thm. 7.1.3 and Eq. (7.1.13), we have

$$(5) \quad G_k^{\chi_1, \chi_2}(\mathfrak{c}_2 z) = \frac{2C_k W(\overline{\chi_2})}{\mathfrak{c}_2^k} E_k^{\chi_1, \chi_2}(z), \quad \text{for } k \geq 3,$$

and similarly using Thm. 7.2.12

$$(6) \quad \lim_{\varepsilon \rightarrow 0^+} G_{2, \varepsilon}^{\chi_1, \chi_2}(\mathfrak{c}_2 z) = \frac{2C_2 W(\overline{\chi_2})}{\mathfrak{c}_2^2} E_2^{\chi_1, \chi_2}(z).$$

According to *loc. cit.* §7.1 and §7.2 for  $k \geq 3$  and  $k = 2$  respectively, together with Thm. 4.7.1, we have that  $E_k^{\chi_1, \chi_2}$  is an Eisenstein series of weight  $k$ , level  $N$  and Nebentypus character  $\chi_1 \chi_2$ .

**1.4. Computation of the constant terms.** We keep the notation and assumptions of the previous paragraph and moreover denote by  $\mathfrak{c}_0$  the conductor of the primitive character  $(\overline{\chi_1} \chi_2)_0$  associated with  $\overline{\chi_1} \chi_2$ . For any integer  $M$  we denote by  $\alpha_M$  the usual degeneracy operator given by  $\alpha_M f(z) = f(Mz)$ .

For a given matrix  $\gamma \in \mathrm{SL}_2(\mathbf{Z})$ , we let

$$\Upsilon_k^{\chi_1, \chi_2}(\gamma, M) = \lim_{\mathrm{Im}(z) \rightarrow \infty} (\alpha_M E_k^{\chi_1, \chi_2}) |_k \gamma(z)$$

be the constant term of the Fourier expansion at  $\infty$  of  $\alpha_M E_k^{\chi_1, \chi_2} |_k \gamma$ . Here, the notation  $|_k$  refers to the classical slash operator acting on weight  $k$  modular forms.

The main goal of this section is the computation, embodied in Proposition 4 below, of the constant term  $\Upsilon_k^{\chi_1, \chi_2}(\gamma, M)$ .

**Proposition 4.** *Let  $\gamma = \begin{pmatrix} u & \beta \\ v & \delta \end{pmatrix} \in \mathrm{SL}_2(\mathbf{Z})$  and let  $M \geq 1$  be an integer. Put  $r = \gcd(v, M)$ ,  $v' = v/r$  and  $M' = M/r$ . If  $\mathfrak{c}_2 \nmid v'$ , then we have that  $\Upsilon_k^{\chi_1, \chi_2}(\gamma, M) = 0$ . Else, if  $\mathfrak{c}_2 \mid v'$ , then*

$$\Upsilon_k^{\chi_1, \chi_2}(\gamma, M) \neq 0 \iff \gcd\left(\frac{v'}{\mathfrak{c}_2}, \mathfrak{c}_1\right) = 1.$$

Moreover, in that case, we have that  $\Upsilon_k^{\chi_1, \chi_2}(\gamma, M)$  is given by the following non-zero algebraic number

$$\Upsilon_k^{\chi_1, \chi_2}(\gamma, M) = \xi \cdot \left(\frac{\mathfrak{c}_2}{M' \mathfrak{c}_0}\right)^k \cdot \frac{W((\chi_1 \overline{\chi_2})_0)}{W(\overline{\chi_2})} \cdot \frac{B_{k, (\overline{\chi_1} \chi_2)_0}}{2k} \prod_{p|N} \left(1 - (\chi_1 \overline{\chi_2})_0(p) p^{-k}\right)$$

where  $\xi = -\chi_2(\delta) \overline{\chi_2}(M') \chi_1(-v'/\mathfrak{c}_2)$  is a root of unity and  $p$  runs over the prime divisors of  $N$ .

*Remark 1.* The result above generalizes the special cases  $(\chi_1, \chi_2, M) = (\chi_1, \chi_1^{-1}, 1)$  and  $(\chi_1, k) = (1, \geq 3)$  stated in [BD14, Prop. 2.8] and [BM15, Prop. 1.2] respectively. In this paper, we not only need the above statement in its full generality and precision, but we also provide a unified and (slightly) simplified proof of these previous results.

The following result is easily deduced from the above proposition and will be of use in Section 3.

**Corollary 5.** *In the notation of Proposition 4, assume  $M$  and  $N$  are co-prime. Then, we have*

$$\Upsilon_k^{\chi_1, \chi_2}(\gamma, M) = \left(\frac{r}{M}\right)^k \overline{\chi_1}(r) \chi_2(r) \overline{\chi_2}(M) \Upsilon_k^{\chi_1, \chi_2}(\gamma, 1).$$

We break the proof of Proposition 4 in several steps. The proof is given at the end of this paragraph, except for the justification of an intermediary step in the case  $k = 2$ , which is dealt with in the next paragraph.

**Lemma 6.** *Under the same hypothesis as in Proposition 4, we have that*

$$\Upsilon_k^{\chi_1, \chi_2}(\gamma, M) = \frac{\mathfrak{c}_2^k}{2C_k W(\overline{\chi_2})} \cdot \sum_{(m,n) \in C} \frac{\chi_1(m) \overline{\chi_2}(n)}{(mM\mathfrak{c}_2\beta + n\delta)^k},$$

where  $C = \{(m, n) \in \mathbf{Z}^2 \setminus \{(0, 0)\} : mM\mathfrak{c}_2u + nv = 0\}$ .

*Proof of Lemma 6 in the case  $k > 2$ .* Using the relation (5), we have that

$$\frac{2C_k W(\overline{\chi_2})}{\mathfrak{c}_2^k} \Upsilon_k^{\chi_1, \chi_2}(\gamma, M) = \lim_{\text{Im}(z) \rightarrow \infty} (\alpha_{M\mathfrak{c}_2} G_k^{\chi_1, \chi_2})|_k \gamma.$$

Besides, we have

$$(7) \quad \alpha_{M\mathfrak{c}_2} G_k^{\chi_1, \chi_2}|_k \gamma(z) = \sum_{\substack{(m,n) \in \mathbf{Z}^2 \\ (m,n) \neq (0,0)}} \frac{\chi_1(m) \overline{\chi_2}(n)}{(z(mM\mathfrak{c}_2u + nv) + mM\mathfrak{c}_2\beta + n\delta)^k}$$

where the above sum is absolutely convergent since  $k \geq 3$ . We can therefore exchange limit and summation, yielding the result.  $\square$

*Remark 2.* When  $k = 2$ , the sum in (7) is not absolutely convergent and it becomes necessary to give additional considerations, that we present in paragraph 1.5, in order to justify the interchange of limit and summation. The full proof of Lemma 6 is thus achieved in Lemma 10 below.

We now prove the following key result assuming the validity of Lemma 6 for any  $k \geq 2$ .

**Lemma 7.** *Under the same hypothesis as in Proposition 4. If  $\mathfrak{c}_2 \nmid v'$ , then we have  $\Upsilon_k^{\chi_1, \chi_2}(\gamma, M) = 0$ . Else, if  $\mathfrak{c}_2 \mid v'$ , then we have*

$$\Upsilon_k^{\chi_1, \chi_2}(\gamma, M) = \frac{\overline{\chi_2}(M'u) \chi_1(-v'/\mathfrak{c}_2)}{M'^k} \cdot \frac{\mathfrak{c}_2^k}{C_k W(\overline{\chi_2})} \cdot L(k, \chi_1 \overline{\chi_2}),$$

where  $L(k, \chi_1 \overline{\chi_2}) = \sum_{n \geq 1} (\chi_1 \overline{\chi_2})(n) n^{-k}$ .

*Proof.* For simplicity, put  $\Upsilon = \Upsilon_k^{\chi_1, \chi_2}(\gamma, M)$ .

- (i) Assume  $u = 0$ . Then,  $-v\beta = 1$ , implying  $v \in \{\pm 1\}$ ,  $M' = M$  and  $v' = v$ . Also, the set  $C$  in Lemma 6 satisfies  $C = \mathbf{Z} \setminus \{0\} \times \{0\}$ . If  $\chi_2 \neq \mathbf{1}$  (that is, if  $\mathfrak{c}_2 \nmid v'$ ), we have that  $\chi_2(0) = 0$  and then  $\Upsilon = 0$  as claimed.

Assume now that  $\chi_2 = \mathbf{1}$ . Then,  $\mathfrak{c}_2 = W(\overline{\chi_2}) = 1$  and  $\chi_1(-1) = (-1)^k$ . These relations imply  $\chi_1(-v) = \beta^{-k}$ . On the other hand, Lemma 6 ensures that

$$2C_k \Upsilon = \sum_{\substack{m \in \mathbf{Z} \\ m \neq 0}} \frac{\chi_1(m)}{(mM\beta)^k} = \frac{2}{(\beta M)^k} L(k, \chi_1) = \frac{2\chi_1(-v)}{M^k} L(k, \chi_1),$$



concluding the proof in this case.

(ii) Assume  $u \neq 0$ . We have the following

**Claim.** *Let  $n \in \mathbf{Z} \setminus \{0\}$  with  $\gcd(n, \mathfrak{c}_2) = 1$ . Then, there exists  $m \in \mathbf{Z}$  such that  $(m, n) \in C$  if and only if  $M'u | n$  and  $\mathfrak{c}_2 | v'$ . Furthermore, in this case we have that*

$$(8) \quad m = -\frac{n}{M'u} \cdot \frac{v'}{\mathfrak{c}_2} \quad \text{and} \quad mM\mathfrak{c}_2\beta + n\delta = \frac{n}{u}.$$

*Proof of the claim.* If  $(m, n) \in C$ , then  $mM'\mathfrak{c}_2u + nv' = 0$ . We have that  $\gcd(M', v') = 1$  by definition. Moreover,  $\gamma \in \mathrm{SL}_2(\mathbf{Z})$  implies  $\gcd(u, v) = 1$ , hence  $M'u | n$ . On the other hand, since  $\gcd(\mathfrak{c}_2, n) = 1$ , we have that  $\mathfrak{c}_2 | v'$ .

Conversely, if  $M'u | n$  and  $\mathfrak{c}_2 | v'$ , then the integer  $m = -\frac{n}{M'u} \cdot \frac{v'}{\mathfrak{c}_2}$  satisfies  $(m, n) \in C$ .

Finally, if the equivalence is satisfied, we easily check using the relation  $u\delta - v\beta = 1$ , that the second relation in Eq. (8) holds.  $\square$

Using the claim and Lemma 6, we have that  $\Upsilon = 0$  if  $\mathfrak{c}_2 \nmid v'$ . Else, if  $\mathfrak{c}_2 | v'$ , then we have

$$\begin{aligned} \frac{2C_k W(\overline{\chi_2})}{\mathfrak{c}_2^k} \Upsilon &= \sum_{\substack{M'u | n \\ n \neq 0}} \frac{\chi_1(n/M'u) \chi_1(-v'/\mathfrak{c}_2) \overline{\chi_2}(n)}{\left(\frac{n}{u}\right)^k} \\ &= \chi_1(-v'/\mathfrak{c}_2) \sum_{\substack{n \in \mathbf{Z} \\ n \neq 0}} \frac{\chi_1(n) \overline{\chi_2}(nM'u)}{(M'n)^k} \\ &= \frac{\chi_1(-v'/\mathfrak{c}_2) \overline{\chi_2}(M'u)}{M'^k} \sum_{\substack{n \in \mathbf{Z} \\ n \neq 0}} \frac{\chi_1(n) \overline{\chi_2}(n)}{n^k} \\ &= \frac{\chi_1(-v'/\mathfrak{c}_2) \overline{\chi_2}(M'u)}{M'^k} 2L(k, \chi_1 \overline{\chi_2}), \end{aligned}$$

since  $\chi(-1) \overline{\chi_2}(-1) = (-1)^k$ . This finishes the proof of Lemma 7.  $\square$

*Proof of Proposition 4.* According to Lemma 7, it remains to deal with the case where  $\mathfrak{c}_2 | v'$ . In that case, by reducing the equality  $u\delta - v\beta = 1$  modulo  $\mathfrak{c}_2$ , we get  $u\delta \equiv 1 \pmod{\mathfrak{c}_2}$ . Besides, we have  $\gcd(M', \mathfrak{c}_2) | v'$  and hence  $\gcd(M', \mathfrak{c}_2) = 1$ . Therefore if we assume that  $\gcd(v'/\mathfrak{c}_2, \mathfrak{c}_1) = 1$ , it follows that

$$-\chi_1(-v'/\mathfrak{c}_2) \overline{\chi_2}(M'u) = -\chi_1(-v'/\mathfrak{c}_2) \chi_2(\delta) \overline{\chi_2}(M') = \xi$$

is a root of unity.

Besides, we have

$$L(k, \chi_1 \overline{\chi_2}) = L(k, (\chi_1 \overline{\chi_2})_0) \prod_{p|N} \left(1 - \frac{(\chi_1 \overline{\chi_2})_0(p)}{p^k}\right),$$

where  $(\chi_1 \overline{\chi_2})_0$  denotes the primitive character associated with  $\chi_1 \overline{\chi_2}$ . Moreover, it follows from the Euler product for  $L(k, (\chi_1 \overline{\chi_2})_0)$  that  $L(k, \chi_1 \overline{\chi_2}) \neq 0$ .

Now using the assumption  $(\chi_1 \overline{\chi_2})_0(-1) = (-1)^k$  and [Miy06, Thm. 3.3.4], we get that

$$L(k, (\chi_1 \overline{\chi_2})_0) = -W((\chi_1 \overline{\chi_2})_0) \cdot \frac{C_k}{c_0^k} \cdot \frac{B_{k, (\chi_1 \overline{\chi_2})_0}}{2k}.$$

Combining these facts together with Lemma 7 concludes the proof of Proposition 4.  $\square$

**1.5. The case of weight 2.** The goal of this paragraph is to prove Lemma 6 in the case  $k = 2$ . This is achieved in Lemma 10. For  $\varepsilon \geq 0$ , we use the notation

$$w^{2, \varepsilon} = w^2 |w|^{2\varepsilon}, \quad w \in \mathbf{C}.$$

Let  $y_0 > 0$  be a positive real number. The notation  $g_1 \ll_{y_0} g_2$  means that there exists a positive constant  $C$ , depending only on  $y_0$ , such that  $|g_1(r)| \leq C|g_2(r)|$  for all  $r$  in the common domain of  $g_1, g_2$ .

Let

$$S_\varepsilon(z) = \sum_{n \in \mathbf{Z}} \frac{1}{(z+n)^{2, \varepsilon}}, \quad z \in \mathbf{C} \setminus \mathbf{R}.$$

For  $z \in \mathfrak{H}$ , the function  $S_\varepsilon(z)$  corresponds to the function  $S(z; 2 + \varepsilon, \varepsilon)$  in the notation of [Miy06, (7.2.7)].

**Lemma 8.** *Fix  $y_0 > 0$ . Then, we have that*

$$S_\varepsilon(z) \ll_{y_0} \frac{1}{\Gamma(\varepsilon)|y|^{1+2\varepsilon}} + e^{-2\pi|y|}, \quad y = \text{Im}(z), \quad |y| \geq y_0, \quad 0 < \varepsilon \leq 1,$$

where for any real number  $s > 0$ ,  $\Gamma(s) = \int_0^\infty e^{-t} t^{s-1} dt$ .

*Proof.* Since we have  $S_\varepsilon(x - iy) = S_\varepsilon(-x + iy)$ , we can assume that  $y \geq y_0$ . For  $m \in \mathbf{Z}$ , let us denote by  $\xi_\varepsilon(y; m)$  the function  $\xi(y; 2 + \varepsilon, \varepsilon; m)$  of [Miy06, (7.2.11)]. According to Theorem 7.2.8 of *loc. cit.*, we then have

$$(9) \quad S_\varepsilon(z) = \xi_\varepsilon(y; 0) + \sum_{\substack{m \in \mathbf{Z} \\ m \neq 0}} e^{2\pi i m x} \xi_\varepsilon(y; m), \quad z = x + iy,$$

where the series converges absolutely. Besides, for  $m \in \mathbf{Z}$ , we have by *loc. cit.*, Theorem 7.2.5, that

$$\xi_\varepsilon(y; m) = \begin{cases} -\frac{(2\pi)^{2+\varepsilon}}{\Gamma(2+\varepsilon)} \frac{1}{(2y)^\varepsilon} m^{1+\varepsilon} e^{-2\pi y m} \omega(4\pi y m; 2 + \varepsilon, \varepsilon) & \text{if } m > 0 \\ -\frac{(2\pi)^{2+2\varepsilon}}{\Gamma(2+\varepsilon)\Gamma(\varepsilon)} \frac{1}{(4\pi y)^{1+2\varepsilon}} & \text{if } m = 0 \\ -\frac{(2\pi)^\varepsilon}{\Gamma(\varepsilon)} \frac{1}{(2y)^{2+\varepsilon}} \frac{1}{|m|^{1-\varepsilon}} e^{-2\pi y |m|} \omega(4\pi y |m|; \varepsilon, 2 + \varepsilon) & \text{if } m < 0. \end{cases}$$

The definition of the function  $\omega$  is stated in *loc. cit.* (7.2.31). It follows from Theorem 7.2.7 in *loc. cit.* that for all  $m \in \mathbf{Z} \setminus \{0\}$ ,  $y \geq y_0$  and  $0 < \varepsilon \leq 1$ , we have

$$\omega(4\pi y |m|; 2 + \varepsilon, \varepsilon) \ll_{y_0} 1 \quad \text{and} \quad \omega(4\pi y |m|; \varepsilon, 2 + \varepsilon) \ll_{y_0} 1.$$

Therefore, for all  $y \geq y_0$  and  $0 < \varepsilon \leq 1$  we have

$$\xi_\varepsilon(y; m) \ll_{y_0} \begin{cases} m^2 e^{-2\pi y m} & \text{if } m > 0 \\ \frac{1}{\Gamma(\varepsilon) y^{1+2\varepsilon}} & \text{if } m = 0 \\ e^{-2\pi y |m|} & \text{if } m < 0 \end{cases}$$

and equation (9) implies

$$S_\varepsilon(z) \ll_{y_0} \frac{1}{\Gamma(\varepsilon)y^{1+2\varepsilon}} + \sum_{m \geq 1} m^2 e^{-2\pi y m} + \sum_{m \geq 1} e^{-2\pi y m}.$$

On the other hand, for all  $y \geq y_0$ , we have

$$\sum_{m \geq 1} (m^2 + 1) e^{-2\pi y m} = \frac{e^{-2\pi y} (e^{-4\pi y} - e^{-2\pi y} + 2)}{(1 - e^{-2\pi y})^3} \ll_{y_0} e^{-2\pi y},$$

hence the result.  $\square$

**Lemma 9.** For any  $a_1, a_2, D \in \mathbf{Z}$  with  $D \neq 0$ , set

$$\sigma_\varepsilon(z; a_1, a_2, D) = \sum_{\substack{(m,n) \in \mathbf{Z}^2 \\ a_1 + Dm \neq 0}} \frac{1}{(z(a_1 + Dm) + a_2 + Dn)^{2,\varepsilon}}.$$

Then, we have that

$$\lim_{\text{Im}(z) \rightarrow \infty} \lim_{\varepsilon \rightarrow 0^+} \sigma_\varepsilon(z; a_1, a_2, D) = 0.$$

*Proof.* Assume  $y = \text{Im}(z) \geq 1$ . We have that

$$\begin{aligned} \sigma_\varepsilon(z; a_1, a_2, D) &= \frac{1}{D^{2,\varepsilon}} \sum_{\substack{m \in \mathbf{Z} \\ a_1 + Dm \neq 0}} \sum_{n \in \mathbf{Z}} \frac{1}{(z(\frac{a_1}{D} + m) + \frac{a_2}{D} + n)^{2,\varepsilon}} \\ &= \frac{1}{D^{2,\varepsilon}} \sum_{\substack{m \in \mathbf{Z} \\ a_1 + Dm \neq 0}} S_\varepsilon\left(z\left(\frac{a_1}{D} + m\right) + \frac{a_2}{D}\right). \end{aligned}$$

Define

$$y_0 = \min \left\{ \left| \text{Im}\left(z\left(\frac{a_1}{D} + m\right)\right) \right|; \text{Im}(z) \geq 1, m \in \mathbf{Z} : a_1 + Dm \neq 0 \right\}.$$

Since  $\mathbf{Z}$  is discrete, we have  $y_0 > 0$ . Using Lemma 8 with this choice of  $y_0$ , we find that for  $\varepsilon \leq 1 \leq y$  and  $m \in \mathbf{Z}$  such that  $a_1 + Dm \neq 0$ , we have

$$S_\varepsilon\left(z\left(\frac{a_1}{D} + m\right) + \frac{a_2}{D}\right) \ll_{y_0} \frac{1}{\Gamma(\varepsilon)y^{1+2\varepsilon}} \cdot \frac{1}{\left|\frac{a_1}{D} + m\right|^{1+2\varepsilon}} + e^{-2\pi y \left|\frac{a_1}{D} + m\right|}.$$

Therefore, we have

$$\sigma_\varepsilon(z; a_1, a_2, D) \ll_{y_0} \frac{1}{|D|^{2(1+\varepsilon)}} \left( \frac{1}{y^{1+2\varepsilon}} \cdot \frac{1}{\Gamma(\varepsilon)} \cdot \zeta(1+2\varepsilon) + \sum_{n \geq 1} e^{-\frac{2\pi y n}{|D|}} \right).$$

Since  $\sum_{n \geq 1} e^{-\frac{2\pi y n}{|D|}} \ll_{y_0} e^{-\frac{2\pi y}{|D|}}$ , we have that

$$\limsup_{\varepsilon \rightarrow 0^+} |\sigma_\varepsilon(z; a_1, a_2, D)| \ll_{y_0} \frac{1}{D^2} \left( \frac{1}{y} + e^{-\frac{2\pi y}{|D|}} \right).$$

This estimate justifies the claim.  $\square$

**Lemma 10.** Lemma 6 is true for  $k = 2$ .

*Proof.* Using relation (6), we have in particular that

$$(10) \quad \Upsilon_2^{\chi_1, \chi_2}(\gamma, M) = \frac{\mathfrak{c}_2^2}{2C_2 W(\overline{\chi_2})} \lim_{\operatorname{Im}(z) \rightarrow \infty} \lim_{\varepsilon \rightarrow 0^+} \left( \alpha_{M\mathfrak{c}_2} G_{2,\varepsilon}^{\chi_1, \chi_2} \right) |_2 \gamma.$$

For  $\varepsilon > 0$ , let

$$(11) \quad T_\varepsilon(z) = \sum_{(m,n) \in C} \frac{\chi_1(m) \overline{\chi_2}(n)}{(mM\mathfrak{c}_2\beta + n\delta)^{2,\varepsilon}}$$

and

$$R_\varepsilon(z) = \sum_{\substack{(m,n) \notin C \\ (m,n) \neq (0,0)}} \frac{\chi_1(m) \overline{\chi_2}(n)}{(z(mM\mathfrak{c}_2u + nv) + mM\mathfrak{c}_2\beta + n\delta)^{2,\varepsilon}},$$

where, as in Lemma 6,

$$C = \{(m, n) \in \mathbf{Z}^2 \setminus \{(0, 0)\} : mM\mathfrak{c}_2u + nv = 0\}.$$

Then, we have

$$\left( \alpha_{M\mathfrak{c}_2} G_{2,\varepsilon}^{\chi_1, \chi_2} \right) |_2 \gamma(z) = |vz + \delta|^\varepsilon \sum_{\substack{(m,n) \in \mathbf{Z}^2 \\ (m,n) \neq (0,0)}} \frac{\chi_1(m) \overline{\chi_2}(n)}{(mM\mathfrak{c}_2(uz + \beta) + n(vz + \delta))^{2,\varepsilon}}$$

and therefore

$$\lim_{\varepsilon \rightarrow 0^+} \left( \alpha_{M\mathfrak{c}_2} G_{2,\varepsilon}^{\chi_1, \chi_2} \right) |_2 \gamma(z) = \lim_{\varepsilon \rightarrow 0^+} (T_\varepsilon(z) + R_\varepsilon(z)).$$

Since the parameters appearing in the sum defining  $T_\varepsilon$  are linked by a linear relation, the series obtained by setting  $\varepsilon = 0$  in (11) is absolutely convergent. Hence, we have that

$$\lim_{\varepsilon \rightarrow 0^+} T_\varepsilon(z) = \sum_{(m,n) \in C} \frac{\chi_1(m) \overline{\chi_2}(n)}{(mM\mathfrak{c}_2\beta + n\delta)^2}.$$

In particular, this limit is independent of  $z$ . Hence, in light of Equation (10), in order to finish the proof we need to show that

$$(12) \quad \lim_{y \rightarrow \infty} \lim_{\varepsilon \rightarrow 0^+} R_\varepsilon(z) = 0.$$

We have that

$$(13) \quad R_\varepsilon(z) = \sum_{a=0}^{\mathfrak{c}_1-1} \sum_{b=0}^{\mathfrak{c}_2-1} \chi_1(a) \overline{\chi_2}(b) \sum_{\substack{(c,d) \in C_{a,b} \\ c \neq 0}} \frac{1}{(cz + d)^{2,\varepsilon}},$$

where

$$C_{a,b} = \{(mM\mathfrak{c}_2u + nv, mM\mathfrak{c}_2\beta + n\delta) : m \equiv a \pmod{\mathfrak{c}_1}, n \equiv b \pmod{\mathfrak{c}_2}\}.$$

Now we proceed to split each of the sums in (13) indexed by  $C_{a,b}$  in a finite number of sums of the type handled by Lemma 9. Let

$$\mathbb{M} = \begin{pmatrix} M\mathfrak{c}_1\mathfrak{c}_2u & M\mathfrak{c}_1\mathfrak{c}_2\beta \\ \mathfrak{c}_2v & \mathfrak{c}_2\delta \end{pmatrix}, \quad \theta^{a,b} = (aM\mathfrak{c}_2u + bv, aM\mathfrak{c}_2\beta + b\delta).$$

Then,  $C_{a,b} = \theta^{a,b} + \mathbf{Z}^2 \cdot \mathbb{M}$  (here, we represent the elements of  $\mathbf{Z}^2$  as row vectors). Let  $D := \det \mathbb{M} = M\mathfrak{c}_1\mathfrak{c}_2^2$ . By the elementary divisors theorem, we

have that  $D\mathbf{Z} \times D\mathbf{Z} \subset \mathbf{Z}^2 \cdot \mathbb{M}$  is a subgroup of index  $D$ . Let  $\{r_1, r_2, \dots, r_D\}$  be a system of representatives of the quotient  $\mathbf{Z}^2 \cdot \mathbb{M} / D\mathbf{Z} \times D\mathbf{Z}$ . Then, in the notation of Lemma 9, we have that

$$R_\varepsilon(z) = \sum_{a=0}^{\mathfrak{c}_1-1} \sum_{b=0}^{\mathfrak{c}_2-1} \chi_1(a) \overline{\chi_2}(b) \sum_{i=1}^D \sigma_\varepsilon \left( z; \theta_1^{a,b} + r_{i,1}, \theta_2^{a,b} + r_{i,2}, D \right),$$

where, for any vector  $w \in \mathbf{R}^2$  we write  $w = (w_1, w_2)$ . Then, using Lemma 9, we deduce the truth of Equation (12).  $\square$

## 2. ADELIZATION OF MODULAR FORMS AND HECKE OPERATORS

In this short section we briefly introduce some useful notation and make explicit our normalizations for modular forms and Hecke operators in the adelic setting.

Let  $G = \mathrm{GL}_2$  considered as an algebraic group over  $\mathbf{Q}$ . We denote by  $\mathbf{A}$  the ring of adèles of  $\mathbf{Q}$ . Let

$$G_{\mathbf{A}} = G(\mathbf{A}), \quad G_{\mathbf{Q}} = G(\mathbf{Q}) \quad \text{and} \quad G_{\mathbf{R}}^+ = \{\gamma \in G_{\mathbf{R}} = G(\mathbf{R}) : \det \gamma > 0\}.$$

For each prime number  $p$ , we denote by  $\iota_p: G_{\mathbf{Q}} \rightarrow G_{\mathbf{A}}$  the map induced by the ring homomorphism  $\mathbf{Q} \hookrightarrow \mathbf{Q}_p \rightarrow \mathbf{A}$ . We define similarly  $\iota_\infty: G_{\mathbf{R}} \rightarrow G_{\mathbf{A}}$  using the inclusion  $\mathbf{R} \hookrightarrow \mathbf{A}$ . We then embed  $G_{\mathbf{Q}}$  in  $G_{\mathbf{A}}$  diagonally (that is, using  $\prod_p \iota_p \times \iota_\infty$ ) and we embed  $G_{\mathbf{R}}^+$  at infinity (that is, using  $\iota_\infty$ ).

Let  $N \geq 1$  be a positive integer. For every prime number  $p$  set

$$K_p(N) = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathrm{GL}_2(\mathbf{Z}_p) : c \equiv 0 \pmod{p\mathbf{Z}_p} \right\}$$

and define  $K_0(N) = \prod_p K_p(N)$  as a subgroup of  $\mathrm{GL}_2(\mathbf{A}_f)$  where  $\mathbf{A}_f$  denotes the finite adèles of  $\mathbf{Q}$ . The strong approximation theorem ([Bum97, Thm. 3.3.1] for  $G$ ) then implies that

$$(14) \quad G_{\mathbf{A}} = G_{\mathbf{Q}} G_{\mathbf{R}}^+ K_0(N).$$

We denote by  $\omega$  the adelization (*loc. cit.* Prop. 3.1.2) of a given Dirichlet character  $\chi$  of modulus  $N$ , and define the group homomorphism

$$\lambda: \begin{matrix} K_0(N) & \longrightarrow & \mathbf{C}^\times \\ \left( \begin{pmatrix} a_p & b_p \\ c_p & d_p \end{pmatrix} \right)_p & \longmapsto & \prod_{p|N} \omega_p(d_p) \end{matrix}.$$

Let  $p$  be a prime divisor of  $N$ . For every integer  $n \in \{0, \dots, p-1\}$ , define

$$\xi_n = \begin{pmatrix} p & n \\ 0 & 1 \end{pmatrix}.$$

Let  $k_0 \in K_0(N)$ . Denote by  $\begin{pmatrix} a & b \\ c & d \end{pmatrix} \in K_p(N)$  the  $p$ -th component of  $k_0$ .

Let  $n \in \{0, \dots, p-1\}$  be an integer. Since  $p \mid N$ , we have that  $cn + d \in \mathbf{Z}_p^\times$  and we define  $m$  to be the unique integer in  $\{0, \dots, p-1\}$  such that

$$(cn + d)m \equiv an + b \pmod{p\mathbf{Z}_p}.$$

It then follows from the following matrix identity in  $\mathrm{GL}_2(\mathbf{Q}_p)$

$$\xi_m^{-1} \begin{pmatrix} a & b \\ c & d \end{pmatrix} \xi_n = \begin{pmatrix} a - mc & \frac{an+b-m(cn+d)}{p} \\ cp & cn+d \end{pmatrix}$$

that

$$(15) \quad k'_0 = \iota_p(\xi_m)^{-1} k_0 \iota_p(\xi_n) \in K_0(N) \quad \text{and} \quad \lambda(k'_0) = \lambda(k_0).$$

Let  $g \in \mathbf{G}_{\mathbf{A}}$ , that we decompose as

$$g = \gamma g_{\infty} k_0, \quad \gamma \in \mathbf{G}_{\mathbf{Q}}, \quad g_{\infty} \in \mathbf{G}_{\mathbf{R}}^+, \quad k_0 \in K_0(N)$$

using equation (14). We then check place by place that the following equality holds (see *loc. cit.*, p. 345)

$$(16) \quad g \iota_p(\xi_n) = (\gamma \xi_m) (\xi_{m,\infty}^{-1} g_{\infty}) (\xi_{m,f}^{-1} \iota_p(\xi_m) k'_0) \in \mathbf{G}_{\mathbf{Q}} \mathbf{G}_{\mathbf{R}}^+ K_0(N)$$

where  $n \in \{0, \dots, p-1\}$  and  $m \in \{0, \dots, p-1\}$ ,  $k'_0 \in K_0(N)$  are defined above. Here,  $\xi_{m,f}$  and  $\xi_{m,\infty}$  denote the finite and the infinite components of  $\xi_m \in \mathbf{G}_{\mathbf{A}}$  respectively.

Let  $k \geq 2$  be an integer. Denote by  $S_k(N, \chi)$  the space of cuspidal modular forms of weight  $k$ , level  $N$  and Nebentypus character  $\chi$ . To a modular form  $F \in S_k(N, \chi)$ , we attach

$$\phi_F: \mathbf{G}_{\mathbf{A}} \rightarrow \mathbf{C}, \quad \phi_F(g) = F(g_{\infty} \cdot i) j(g_{\infty}, i)^{-k} \lambda(k_0), \quad g = \gamma g_{\infty} k_0.$$

Here, for  $g_{\infty} = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathbf{G}_{\mathbf{R}}^+$ , we have  $j(g_{\infty}, z) = (cz + d) \det g_{\infty}^{-1/2}$ . Since

$$\mathbf{G}_{\mathbf{Q}} \cap \mathbf{G}_{\mathbf{R}}^+ K_0(N) = \Gamma_0(N) = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathrm{SL}_2(\mathbf{Z}) : c \equiv 0 \pmod{N} \right\}$$

and for every  $\gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma_0(N)$ , we have  $\lambda(\gamma) = \chi(d)^{-1}$  (as  $\omega$  is trivial on  $\mathbf{Q}^{\times}$ ), the function  $\phi_F$  is a well-defined automorphic form (*loc. cit.*, §3.6). Define  $\pi_F$  to be the linear span of right translates of  $\phi_F$  under  $\mathbf{G}_{\mathbf{A}}$  and assume that  $F$  is an eigenfunction for the Hecke operators away from  $N$ . Then  $\pi_F$  decomposes as a restricted tensor product  $\bigotimes' \pi_{F,v}$  where  $v$  runs over the places of  $\mathbf{Q}$  and  $\pi_{F,v}$  is an admissible irreducible representation of  $\mathbf{G}_{\mathbf{Q}_v}$  (*loc. cit.*, §3.3). We now define the  $p$ -th Hecke operator in this adelic setting as follows (note the factor  $1/\sqrt{p}$ )

$$(17) \quad \widetilde{U}_p = \frac{1}{\sqrt{p}} \sum_{n=0}^{p-1} \pi_{F,p}(\xi_n)$$

The following result will be used in the proof of Theorems 1 and 2.

**Lemma 11.** *Let  $U_p$  denote the  $p$ -th Hecke operator acting on  $S_k(N, \chi)$ . Then, we have*

$$p^{\frac{k-1}{2}} \widetilde{U}_p \phi_F = \phi_{U_p F}.$$

*Proof.* Let  $g = \gamma g_\infty k_0 \in \mathbf{G}_\mathbf{A}$ . Then, in the notation of Equation (16), we have (using the fact that the map  $n \mapsto m$  is a bijection of  $\{0, \dots, p-1\}$ )

$$\begin{aligned} \widetilde{U}_p \phi_F(g) &= \frac{1}{\sqrt{p}} \sum_{n=0}^{p-1} \phi_F(g \iota_p(\xi_n)) \\ &= \frac{1}{\sqrt{p}} \sum_{m=0}^{p-1} \phi_F\left((\gamma \xi_m) (\xi_{m,\infty}^{-1} g_\infty) (\xi_{m,f}^{-1} \iota_p(\xi_m) k'_0)\right) \\ &= \frac{1}{\sqrt{p}} \sum_{m=0}^{p-1} F\left((\xi_{m,\infty}^{-1} g_\infty) \cdot i\right) j\left(\xi_{m,\infty}^{-1} g_\infty, i\right)^{-k} \lambda\left(\xi_{m,f}^{-1} \iota_p(\xi_m) k'_0\right) \end{aligned}$$

Besides, from the definition of  $\xi_m$  and Equation (15), we have

$$\lambda\left(\xi_{m,f}^{-1} \iota_p(\xi_m) k'_0\right) = \lambda(k'_0) = \lambda(k_0),$$

and from the automorphy relation for  $F$ , we have

$$F\left((\xi_{m,\infty}^{-1} g_\infty) \cdot i\right) j\left(\xi_{m,\infty}^{-1} g_\infty, i\right)^{-k} = p^{-k/2} F\left(\frac{g_\infty \cdot i - m}{p}\right) j(g_\infty, i)^{-k}.$$

We conclude that

$$\widetilde{U}_p \phi_F(g) = \frac{1}{p^{(k+1)/2}} \sum_{m=0}^{p-1} F\left(\frac{g_\infty \cdot i - m}{p}\right) j(g_\infty, i)^{-k} \lambda(k_0).$$

Hence the desired formula since

$$U_p F(z) = \frac{1}{p} \sum_{m=0}^{p-1} F\left(\frac{z+m}{p}\right) = \frac{1}{p} \sum_{m=0}^{p-1} F\left(\frac{z-m}{p}\right), \quad z \in \mathfrak{H}.$$

□

### 3. PROOFS OF THE MAIN RESULTS

**3.1. Proof of Theorem 1.** Let  $\nu_1, \nu_2: \text{Gal}(\overline{\mathbf{Q}}/\mathbf{Q}) \rightarrow \overline{\mathbf{F}}_l^\times$  be characters such that  $\rho = \nu_1 \oplus \nu_2$  defines an odd (semi-simple) Galois representation of Serre type  $(N, k, \varepsilon)$ . Assume throughout that  $l > k+1$ . Each of the characters  $\nu_i$  ( $i = 1, 2$ ) can be decomposed as  $\nu_i = \varepsilon_i \chi_l^{a_i}$  where  $\varepsilon_i$  is unramified at  $l$ ,  $a_i$  is a non-negative integer and  $\chi_l$  denotes the mod  $l$  cyclotomic character. Without loss of generality, we may further assume that  $0 \leq a_1 \leq a_2 \leq l-2$ . According to Serre's definition of the weight  $k$  (see [Ser87, (2.3.2)]), we then have :

$$k = \begin{cases} 1 + la_1 + a_2 & \text{if } (a_1, a_2) \neq (0, 0) \\ l & \text{if } (a_1, a_2) = (0, 0) \end{cases}.$$

Since we have assumed  $l > k+1$ , it follows that  $(a_1, a_2) = (0, k-1)$ . This proves the first part of Theorem 1.

Let us then prove the equivalence. Denote by  $\mathbf{c}_1$  and  $\mathbf{c}_2$  the conductors of  $\varepsilon_1$  and  $\varepsilon_2$  respectively. We have the Serre parameters  $\varepsilon = \varepsilon_1 \varepsilon_2$  and  $N = \mathbf{c}_1 \mathbf{c}_2$ . If  $(N, k) = (1, 2)$ , then both  $\varepsilon_1$  and  $\varepsilon_2$  are trivial and therefore, in the notation of the theorem, we have

$$B_{2,\eta} = B_2 \pmod{l} \quad \text{and} \quad B_2 = \frac{1}{6} \not\equiv 0 \pmod{l}.$$

On the other hand, there is no non-zero cuspidal eigenform of weight 2 and level 1 over  $\overline{\mathbf{F}}_l$  for  $l \geq 5$ . Hence the desired equivalence in this case.

From now on, let us then assume that either  $N > 1$  or  $k > 2$ . Fix a place  $w$  of  $\overline{\mathbf{Q}}$  above  $l$  and denote by  $\chi_1$  and  $\chi_2$  the multiplicative lifts with respect to  $w$  (in the sense of paragraph 1.2) of  $\varepsilon_1$  and  $\varepsilon_2$  respectively. We view  $\chi = \chi_1 \chi_2$  as a Dirichlet character modulo  $N$ . The Eisenstein series  $E_k^{\chi_1, \chi_2}$  introduced in paragraph 1.3 (which is well-defined as we have  $(N, k) \neq (1, 2)$ ) has weight  $k$ , level  $N$  and Nebentypus character  $\chi$ . Moreover, it is a normalized eigenform for the full Hecke algebra at level  $N$ . In particular, if we write

$$E_k^{\chi_1, \chi_2}(z) = \sum_{n \geq 0} a_n (E_k^{\chi_1, \chi_2}) e^{2i\pi zn}, \quad (z \in \mathfrak{H})$$

then its eigenvalue for the action of the Hecke operator at an arbitrary prime  $p$  is given by

$$a_p (E_k^{\chi_1, \chi_2}) = \chi_1(p) + \chi_2(p)p^{k-1}.$$

Assume first that  $\rho = \varepsilon_1 \oplus \varepsilon_2 \chi_l^{k-1}$  is strongly modular. By assumption, there exists an eigenform  $f$  of type  $(N, k, \varepsilon)$  over  $\overline{\mathbf{F}}_l$  such that, in the notation of the Introduction, we have  $\rho_f \simeq \rho$ . Therefore, it exists  $F = \sum_{n \geq 1} A_n q^n$  a weight- $k$  cuspidal form of level  $N$  and Nebentypus character  $\chi$  such that  $A_n \in \overline{\mathbf{Z}}_w$  and

$$(18) \quad \nu_w(A_n) = a_n, \quad \text{for any integer } n \geq 1,$$

in the notation of paragraph 1.2. By Deligne-Serre lifting lemma ([DS74, Lem. 6.11]), one may further assume that  $F$  is a normalized eigenform for all the Hecke operators at level  $N$ . Denote by  $E$  the number field generated by the Hecke eigenvalues of  $F$  and by  $\lambda$  the prime ideal above  $l$  in  $E$  induced by  $w$ . Let  $E_\lambda$  be the completion of  $E$  at  $\lambda$ . Thanks to the isomorphism  $\rho \simeq \rho_f$  and (18), the  $\lambda$ -adic representation of  $F$

$$\rho_{F, \lambda}: \text{Gal}(\overline{\mathbf{Q}}/\mathbf{Q}) \longrightarrow \text{GL}_2(E_\lambda)$$

semi-simplifies to  $\rho$  after reduction modulo  $\lambda$ . Since  $F$  has level  $N$  and  $\rho$  conductor  $N$  away from  $l$ , the form  $F$  is actually a newform. For every prime  $p \nmid Nl$ , we have

$$\nu_w(A_p) = \varepsilon_1(p) + \varepsilon_2(p)p^{k-1},$$

where,  $\varepsilon_i(p) = \varepsilon_i(\text{Frob}_p)$  if  $\varepsilon_i$  is unramified at  $p$  and  $\varepsilon_i(p) = 0$  otherwise, for  $i = 1, 2$ . The next step is to extend these congruences to arbitrary primes  $p \neq l$ , as stated in the following key result. (Note that only the case  $N > 1$  requires a proof.)

**Proposition 12.** *In this notation, we have*

$$\nu_w(A_p) = \varepsilon_1(p) + \varepsilon_2(p)p^{k-1}, \quad \text{for every prime } p \neq l.$$

*Proof.* We have seen that the equality holds for primes not dividing  $Nl$ . Let  $p$  be a prime dividing  $N$  (note that, by definition,  $N$  is coprime to  $l$  and hence  $p \neq l$ ). We denote by  $\mathfrak{c}$  the conductor of  $\chi$ . We shall split the proof into three cases :

$$(1) \quad \text{ord}_p(N) = 1 \text{ and } \text{ord}_p(\mathfrak{c}) = 0;$$



- (2)  $\text{ord}_p(N) \geq 2$  and  $\text{ord}_p(\mathfrak{c}) < \text{ord}_p(N)$ ;
- (3)  $\text{ord}_p(N) = \text{ord}_p(\mathfrak{c})$ .

To deal with the first two cases, we first observe that if  $\text{ord}_p(\mathfrak{c}) < \text{ord}_p(N)$ , then both characters  $\chi_1$  and  $\chi_2$  are ramified at  $p$ . Indeed, since  $\text{ord}_p(N) > 0$  and  $N = \mathfrak{c}_1 \mathfrak{c}_2$ , at least one of the two characters  $\chi_1$  and  $\chi_2$  is ramified at  $p$ . On the other hand, if the other one is unramified at  $p$  then, we have

$$\text{ord}_p(\mathfrak{c}) = \text{ord}_p(\mathfrak{c}_1) + \text{ord}_p(\mathfrak{c}_2) = \text{ord}_p(N)$$

and a contradiction.

In the first case, using this observation, we obtain that  $1 = \text{ord}_p(N) = \text{ord}_p(\mathfrak{c}_1) + \text{ord}_p(\mathfrak{c}_2) \geq 2$  and hence a contradiction. Case (1) therefore does not occur.

In the second case, we have that  $A_p = 0$  ([Miy06, Thm. 4.6.17]) and by the above observation, both  $\chi_1, \chi_2$  (and hence  $\varepsilon_1$  and  $\varepsilon_2$ ) are ramified at  $p$ . We therefore have the desired equality as both sides are zero.

It therefore remains to deal with the last case. Let  $\phi_F$  be the adelization of  $F$  as defined in Section 2. Denote by  $\pi_F$  the corresponding automorphic representation. Since  $F$  is  $p$ -new, then  $\phi_F$  is a so-called new-vector for  $\pi_{F,p}$  (in the sense of [LW12, Thm. 2.2]). The endomorphism  $\widetilde{U}_p$  defined in Equation (17) acts on the (one-dimensional) vector space of new-vectors of  $\pi_{F,p}$  by multiplication by an eigenvalue that we denote by  $\lambda(\pi_{F,p})$ . It then follows from Lemma 11 that we have

$$\lambda(\pi_{F,p}) = A_p / p^{(k-1)/2}.$$

Using the assumption  $\text{ord}_p(N) = \text{ord}_p(\mathfrak{c})$ , we have that  $\lambda(\pi_{F,p})$  has absolute value 1 and therefore is  $\neq 0$  ([Miy06, Thm. 4.6.17]). On the other hand, we see from the classification of irreducible admissible infinite-dimensional smooth representations of  $\text{GL}_2(\mathbf{Q}_p)$  (as recalled in Table 1 of [LW12] for instance) that in this case  $\pi_{F,p}$  necessarily is a principal series  $\pi(\mu_1, \mu_2)$  associated with some characters  $\mu_1, \mu_2$  of  $\mathbf{Q}_p^\times$ . Equating the Hecke eigenvalues we find that

$$(19) \quad p^{(k-1)/2}(\mu_1^*(p) + \mu_2^*(p)) = A_p,$$

where

$$\mu_i^*(p) = \begin{cases} \mu_i(p) & \text{if } \mu_i \text{ is unramified at } p \\ 0 & \text{otherwise} \end{cases}, \quad \text{for } i = 1, 2.$$

Let  $\sigma^\lambda(\pi_{F,p})$  be the representation of the local Weil group  $W(\overline{\mathbf{Q}}_p/\mathbf{Q}_p)$  attached to  $\pi_{F,p}$  by the local Langlands correspondence. By a theorem of Carayol ([Car86, Thm. (A)]), it agrees with (the restriction to the Weil group of) the local representation  $\rho_{F,\lambda}|_{\text{Gal}(\overline{\mathbf{Q}}_p/\mathbf{Q}_p)}$ .

Let us denote by  $\overline{\mu}_1$  and  $\overline{\mu}_2$  the reductions modulo  $w$  of  $\mu_1$  and  $\mu_2$  respectively. According to §0.5 in *loc. cit.*, we therefore have the following equality of characters of  $\mathbf{Q}_p^\times$  with values in  $\overline{\mathbf{F}}_l^\times$ :

$$\left\{ \overline{\mu}_1 \chi_l^{(k-1)/2}, \overline{\mu}_2 \chi_l^{(k-1)/2} \right\} = \left\{ \varepsilon_1, \varepsilon_2 \chi_l^{k-1} \right\}.$$

The result now follows from Equation (19). □

Let us now consider the Eisenstein series  $E_k^{\chi_1, \chi_2}$ . Since both  $F$  and  $E_k^{\chi_1, \chi_2}$  are eigenfunctions for the full Hecke algebra at level  $N$ , it follows from the previous proposition and the multiplicativity of the Fourier coefficients that

$$\nu_w(A_n) = \nu_w(a_n(E_k^{\chi_1, \chi_2})), \quad \text{for all prime-to-}l \text{ integers } n.$$

Note that by Lemma 3 and Equation (4), the  $q$ -expansion of the Eisenstein series  $E_k^{\chi_1, \chi_2}$  lies in  $\overline{\mathbf{Z}}_w[[q]]$ . Let us denote by  $\overline{E}$  its reduction modulo  $w$ . Then, both  $f$  and  $\overline{E}$  have the same image under the  $\Theta$ -operator whose action on the  $q$ -expansions is  $q \frac{d}{dq}$  (see [Kat77, Ch. II]).

We remark that, since we are assuming that  $k \geq 2$  and  $l \nmid N$ , the space of modular forms for  $\Gamma_1(N)$  over  $\overline{\mathbf{F}}_l$  in the sense of Katz and in the sense of Serre are naturally isomorphic ([DI95], Theorem 12.3.7). Then, since  $l > k + 1$ , we can use [Kat77, Cor. 3] to assert that the  $\Theta$ -operator is injective. Hence,  $\overline{E}$  is a cuspidal form over  $\overline{\mathbf{F}}_l$ . This implies that  $w$  divides the constant term of  $E_k^{\chi_1, \chi_2}$  at each of the cusps.

In particular, it divides the constant term of the Fourier expansion at  $\infty$  of  $E_k^{\chi_1, \chi_2}|_k \gamma$  where  $\gamma = \begin{pmatrix} 1 & 0 \\ \mathfrak{c}_2 & 1 \end{pmatrix} \in \mathrm{SL}_2(\mathbf{Z})$ . According to Proposition 4 (applied to  $M = 1$  in its notation),  $w$  divides

$$\left( \frac{\mathfrak{c}_2}{\mathfrak{c}_0} \right)^k \frac{W((\chi_1 \overline{\chi_2})_0)}{W(\overline{\chi_2})} \frac{B_{k, (\overline{\chi_1} \chi_2)_0}}{2k} \prod_{p|N} \left( 1 - (\chi_1 \overline{\chi_2})_0(p) p^{-k} \right).$$

However  $\mathfrak{c}_0$ ,  $\mathfrak{c}_2$ ,  $W((\chi_1 \overline{\chi_2})_0)$ ,  $2k$  and  $W(\overline{\chi_2})$  are all coprime to  $l$ . Moreover,  $(\overline{\chi_1} \chi_2)_0$  is nothing but the multiplicative lift of  $\eta = \varepsilon_1^{-1} \varepsilon_2$  with respect to  $w$ . Hence, either  $B_{k, \eta} = 0$ , or there exists a prime  $p \mid N$  such that  $\eta(p) p^k = 1$ . This proves the direct implication in Theorem 1.

Conversely, assume that either condition of the theorem is satisfied. Then, by definition of the characters  $\chi_1$  and  $\chi_2$  and of the Bernoulli number  $B_{k, \eta}$ , the place  $w$  divides (the numerator of)

$$B_{k, (\overline{\chi_1} \chi_2)_0} \cdot \prod_{p|N} \left( (\overline{\chi_1} \chi_2)_0(p) p^k - 1 \right).$$

Then, according to Proposition 4 (with  $M = 1$ ), the constant term of the Eisenstein series  $E_k^{\chi_1, \chi_2}$  vanishes at each of the cusp of the modular curve  $X_1(N)$ . Let  $f$  be its reduction modulo  $w$ , which is an eigenform with coefficients in  $\overline{\mathbf{F}}_l$ . As we argued before,  $f$  can be seen both as a Katz or Serre modular form. Then, the  $q$ -expansion principle allows us to ensure that  $f$  is a cuspidal eigenform (cf. [DI95], Remark 12.3.5).

On the other hand, for every prime  $q \nmid Nl$ , we have

$$\rho_f(\mathrm{Frob}_q) = \nu_w(a_q(E_k^{\chi_1, \chi_2})) = \varepsilon_1(q) + \varepsilon_2(q) q^{k-1} = \rho(\mathrm{Frob}_q).$$

Therefore,  $\rho_f \simeq \rho$  and  $f$  is the desired eigenform. This finishes the proof of Theorem 1.

**3.2. Proof of Theorem 2.** In the case  $(N, k) = (1, 2)$  (where we necessarily have  $\rho \simeq \mathbf{1} \oplus \chi_l$  and hence  $\rho$  is not strongly modular), the result is due to Mazur ([Maz77, Prop. 5.12]).

We therefore assume throughout that  $(N, k) \neq (1, 2)$  and start by proving the direct implication.

Using the assumption that the representation  $\rho$  arises from a modular form of type  $(NM, k, \varepsilon)$ , we show as before that there exists  $F = \sum_{n \geq 1} A_n q^n$ , a weight- $k$  normalized cuspidal eigenform of level  $NM$  and Nebentypus character  $\chi$ , such that, if  $\lambda$  denotes the prime ideal of the coefficient field of  $F$  induced by  $w$ , then the  $\lambda$ -adic representation attached to  $F$  semi-simplifies to  $\rho$  after reduction modulo  $\lambda$ . Since  $\rho$  is not strongly modular,  $F$  necessarily is a newform. Therefore, considering its associated automorphic representation, we prove the following result using the same arguments as in Proposition 12.

**Proposition 13.** *In this notation, we have*

$$\nu_w(A_p) = \varepsilon_1(p) + \varepsilon_2(p)p^{k-1}, \quad \text{for every prime } p \neq l, M.$$

We now turn our attention to the local situation at  $M$  and prove the following statement.

**Proposition 14.** *We have*

- (1) *either  $\eta(M)M^k = 1$ ;*
- (2) *or,  $\eta(M)M^{k-2} = 1$  and  $\nu_w(A_M) = \varepsilon_1(M)$ .*

*Proof.* The form  $F$  is  $M$ -primitive in the sense of [LW12, Def. 2.7]. Therefore, according to Proposition 2.8 of *loc. cit.*, the local component at  $M$  of the automorphic representation of  $F$  corresponds to a Steinberg representation. Moreover, we have the following equality between sets of characters of a decomposition group at  $M$  in  $\text{Gal}(\overline{\mathbf{Q}}/\mathbf{Q})$  with values in  $\overline{\mathbf{F}}_l^\times$ :

$$\{\varepsilon_1, \varepsilon_2 \chi_l^{k-1}\} = \{\mu \chi_l^{k/2}, \mu \chi_l^{k/2-1}\},$$

where  $\mu$  is the unramified character that sends a Frobenius element at  $M$  to  $\mu(M) = \nu_w(A_M/M^{k/2-1})$ . We therefore have two cases to consider:

- Assume that, locally at  $M$ , we have  $\varepsilon_1 = \mu \chi_l^{k/2}$ . Then, in particular, we have  $\varepsilon_1(M)^2 = \mu(M)^2 M^k$ . On the other hand, according to [Miy06, Thm. 4.6.17], we have  $\mu(M)^2 = (\varepsilon_1 \varepsilon_2)(M)$ . Therefore, we get that  $\eta(M)M^k = 1$ . (Note that the other equality, namely  $\varepsilon_2 \chi_l^{k-1} = \mu \chi_l^{k/2-1}$ , does not provide any additional information.)
- Assume instead that, locally at  $M$ , we have  $\varepsilon_1 = \mu \chi_l^{k/2-1}$ . Then, on one hand, we have that  $\varepsilon_1(M) = \mu(M)M^{k/2-1}$  and hence  $\nu_w(A_M) = \varepsilon_1(M)$ . On the other hand, we have (using *loc. cit.*)  $M^{2k-2} \varepsilon_2(M)^2 = \mu(M)^2 M^k$ . Therefore we get that  $\eta(M)M^{k-2} = 1$ . Hence the result. (Once again, the other equality, namely  $\varepsilon_2 \chi_l^{k-1} = \mu \chi_l^{k/2}$ , does not give any other information.)

□

In order to finish the proof of Theorem 2, it therefore remains to show that, under the assumption that  $\rho$  is not strongly modular, condition (2) in Proposition 14 implies condition (1). For that purpose, let us assume that condition (2) is satisfied and consider the following Eisenstein series:

$$F_1 = E_k^{\chi_1, \chi_2} - \chi_2(M)M^{k-1} \alpha_M E_k^{\chi_1, \chi_2}.$$

It is a well-known fact that  $F_1$  is an eigenform for the full Hecke algebra at level  $NM$  with eigenvalues

$$a_p(F_1) = \chi_1(p) + \chi_2(p)p^{k-1}, \quad \text{for primes } p \neq M$$

and  $a_M(F_1) = \chi_1(M)$ . In particular, as a consequence of Proposition 13 and our assumption, we have

$$(20) \quad \nu_w(a_n(F_1)) = \nu_w(A_n), \quad \text{for every integer } n \text{ coprime to } l,$$

where  $\{a_n(F_1)\}_{n \geq 1}$  denote the coefficients of the Fourier expansion of  $F_1$  at  $\infty$ . By definition of  $F_1$ , Lemma 3 and Equation (4), this  $q$ -expansion lies in  $\overline{\mathbf{Z}}_w[[q]]$ . Let us thus denote by  $\overline{F}_1$  the reduction of  $F_1$  modulo  $w$ . According to (20),  $\overline{F}_1$  and the reduction of  $F$  modulo  $w$  have the same image under the  $\Theta$ -operator. Since  $l > k + 1$ , the injectivity of  $\Theta$  ([Kat77, Cor. 3]) implies that  $\overline{F}_1$  is cuspidal. Therefore, we have that  $w$  divides the numerator of the constant of the term of the Fourier expansion of  $F_1$  at each cusp of the modular curve at level  $NM$ . According to Corollary 5, such a constant term at the cusp  $1/(M\mathfrak{c}_2)$  is given (up to roots of unity) by

$$\Upsilon_k^{\chi_1, \chi_2}(\gamma, 1) \left( 1 - \overline{\chi}_1(M) \chi_2(M) M^{k-1} \right),$$

where  $\gamma \in \mathrm{SL}_2(\mathbf{Z})$  is such that  $\gamma \cdot \infty = 1/(M\mathfrak{c}_2)$ . On the other hand, for such a  $\gamma$ , thanks to Theorem 1 and Proposition 4, the assumption that  $\rho$  is not strongly modular guarantees that  $\Upsilon_k^{\chi_1, \chi_2}(\gamma, 1)$  is (non-zero and) not divisible by  $w$ . Therefore, it follows that  $\eta(M)M^{k-1} = 1$  and hence  $M \equiv 1 \pmod{l}$  (as we have assumed  $\eta(M)M^{k-2} = 1$ ). This implies the desired equality  $\eta(M)M^k = 1$  and concludes the proof of the direct implication.

In the other direction, assuming that  $\eta(M)M^k = 1$ , we now consider the Eisenstein series defined by

$$F_2 = E_k^{\chi_1, \chi_2} - \chi_1(M) \alpha_M E_k^{\chi_1, \chi_2}.$$

For any  $\gamma \in \mathrm{SL}_2(\mathbf{Z})$ , let us denote by  $a_0(F_2|_k\gamma)$  the constant term of the Fourier expansion at  $\infty$  of  $F_2|_k\gamma$ . According to Corollary 5, using its notation, we have that

$$a_0(F_2|_k\gamma) = \Upsilon_k^{\chi_1, \chi_2}(\gamma, 1) \left( 1 - \left( \frac{r}{M} \right)^k (\chi_1 \overline{\chi}_2)(M/r) \right),$$

where  $r = 1$  or  $M$ . In both cases, using the assumption  $\eta(M)M^k = 1$ , we have that  $\nu_w(a_0(F_2|_k\gamma)) = 0$ . Denote by  $f$  be the reduction of  $F_2$  modulo  $w$ . It is a well-defined cuspidal form of type  $(NM, k, \varepsilon)$  over  $\overline{\mathbf{F}}_l$  which is an eigenform for the full Hecke algebra at level  $NM$  with eigenvalue for the Hecke operator at  $p$  given by

$$\varepsilon_1(p) + \varepsilon_2(p)p^{k-1}, \quad \text{for all primes } p \neq M.$$

In particular, it proves that  $\rho$  arises from a form of type  $(NM, k, \varepsilon)$  as desired.

## REFERENCES

- [BD14] Nicolas Billerey and Luis V. Dieulefait. Explicit large image theorems for modular forms. *J. Lond. Math. Soc. (2)*, 89(2):499–523, 2014.
- [BM15] Nicolas Billerey and Ricardo Menares. On the modularity of reducible mod  $l$  Galois representations. *Math. Research Letters*, 2015?
- [Bum97] Daniel Bump. *Automorphic forms and representations*, volume 55 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 1997.
- [Car59a] Leonard Carlitz. Arithmetic properties of generalized Bernoulli numbers. *J. Reine Angew. Math.*, 202:174–182, 1959.
- [Car59b] Leonard Carlitz. Some arithmetic properties of generalized Bernoulli numbers. *Bull. Amer. Math. Soc.*, 65:68–69, 1959.
- [Car86] Henri Carayol. Sur les représentations  $l$ -adiques associées aux formes modulaires de Hilbert. *Ann. Sci. École Norm. Sup. (4)*, 19(3):409–468, 1986.
- [DI95] Fred Diamond and John Im. Modular forms and modular curves. In *Seminar on Fermat’s Last Theorem (Toronto, ON, 1993–1994)*, volume 17 of *CMS Conf. Proc.*, pages 39–133. Amer. Math. Soc., Providence, RI, 1995.
- [DS74] Pierre Deligne and Jean-Pierre Serre. Formes modulaires de poids 1. *Ann. Sci. École Norm. Sup. (4)*, 7:507–530 (1975), 1974.
- [Kat77] Nicholas M. Katz. A result on modular forms in characteristic  $p$ . In *Modular functions of one variable, V (Proc. Second Internat. Conf., Univ. Bonn, Bonn, 1976)*, pages 53–61. Lecture Notes in Math., Vol. 601. Springer, Berlin, 1977.
- [KW09a] Chandrashekhar Khare and Jean-Pierre Wintenberger. Serre’s modularity conjecture. I. *Invent. Math.*, 178(3):485–504, 2009.
- [KW09b] Chandrashekhar Khare and Jean-Pierre Wintenberger. Serre’s modularity conjecture. II. *Invent. Math.*, 178(3):505–586, 2009.
- [LW12] David Loeffler and Jared Weinstein. On the computation of local components of a newform. *Math. Comp.*, 81(278):1179–1200, 2012.
- [Maz77] Barry Mazur. Modular curves and the Eisenstein ideal. *Inst. Hautes Études Sci. Publ. Math.*, (47):33–186 (1978), 1977.
- [Miy06] Toshitsune Miyake. *Modular forms*. Springer Monographs in Mathematics. Springer-Verlag, Berlin, english edition, 2006. Translated from the 1976 Japanese original by Yoshitaka Maeda.
- [Rib75] Kenneth A. Ribet. On  $l$ -adic representations attached to modular forms. *Invent. Math.*, 28:245–275, 1975.
- [Ser87] Jean-Pierre Serre. Sur les représentations modulaires de degré 2 de  $\text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$ . *Duke Math. J.*, 54(1):179–230, 1987.

(1) UNIVERSITÉ CLERMONT AUVERGNE, UNIVERSITÉ BLAISE PASCAL, LABORATOIRE DE MATHÉMATIQUES, BP 10448, F-63000 CLERMONT-FERRAND, FRANCE. (2) CNRS, UMR 6620, LM, F-63171 AUBIÈRE, FRANCE

*E-mail address:* Nicolas.Billerey@math.univ-bpclermont.fr

INSTITUTO DE MATEMÁTICAS, PONTIFICIA UNIVERSIDAD CATÓLICA DE VALPARAÍSO, BLANCO VIEL 596, CERRO BARÓN, VALPARAÍSO, CHILE

*E-mail address:* ricardo.menares@pucv.cl