HAL
open science

Rational connectedness implies finiteness of quantum K-theory

Anders S. Buch, Pierre-Emmanuel Chaput, Leonardo C. Mihalcea, Nicolas Perrin

To cite this version:

Anders S. Buch, Pierre-Emmanuel Chaput, Leonardo C. Mihalcea, Nicolas Perrin. Rational connectedness implies finiteness of quantum K-theory. Asian Journal of Mathematics, 2016, 20 (1), pp.117-122. 10.4310/AJM.2016.v20.n1.a5 . hal-01297275

HAL Id: hal-01297275

https://hal.science/hal-01297275v1

Submitted on 3 Apr 2016

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

RATIONAL CONNECTEDNESS IMPLIES FINITENESS OF QUANTUM K-THEORY

ANDERS S. BUCH, PIERRE–EMMANUEL CHAPUT, LEONARDO C. MIHALCEA, AND NICOLAS PERRIN

Abstract. Let X be any generalized flag variety with Picard group of rank one. Given a degree d, consider the Gromov-Witten variety of rational curves of degree d in X that meet three general points. We prove that, if this Gromov-Witten variety is rationally connected for all large degrees d, then the structure constants of the small quantum K-theory ring of X vanish for large degrees.

1. Introduction

The (small) quantum K-theory ring $\mathrm{QK}(X)$ of a smooth complex projective variety X is a generalization of both the Grothendieck ring $K(X)$ of algebraic vector bundles on X and the small quantum cohomology ring of X. The ring $\mathrm{QK}(X)$ was defined by Givental [9] when X is a rational homogeneous space and by Lee [11] in general. In this paper we study this ring when X is a complex projective rational homogeneous space with $\mathrm{Pic}(X) = \mathbb{Z}$. Equivalently, we have $X = G/P$ where G is a complex semisimple algebraic group and $P \subset G$ is a maximal parabolic subgroup. The product in $\mathrm{QK}(X)$ of two arbitrary classes $\alpha, \beta \in K(X)$ is a power series

$$\alpha \star \beta = \sum_{d \geq 0} (\alpha \star \beta)_d \, q^d \, ,$$

where each coefficient $(\alpha \star \beta)_d \in K(X)$ is defined using the K-theory ring of the Kontsevich moduli space $\overline{\mathcal{M}}_{0,3}(X, d)$ of stable maps to X of degree d. For general homogeneous spaces it is an open problem if this power series can have infinitely many non-zero terms. The product $\alpha \star \beta$ is known to be finite if X is a Grassmann variety of type A [4]. More generally, when X is any cominuscule homogeneous space, it was proved by the authors in [3] that all products in $\mathrm{QK}(X)$ are finite. Let $d_X(2)$ denote the smallest possible degree of a rational curve connecting two general points in X. The main theorem of [3] states that $(\alpha \star \beta)_d = 0$ whenever X is cominuscule and $d > d_X(2)$, which is the best possible bound.

Given three general points $x, y, z \in X$, let $M_d(x, y, z) \subset \overline{\mathcal{M}}_{0,3}(X, d)$ denote the *Gromov-Witten variety* of stable maps that send the three marked points to x, y, and z. We will assume that this variety is rationally connected for all sufficiently large degrees d. Let d_{rc} be a positive integer such that $M_d(x, y, z)$ is rationally connected for $d \geq d_{\mathrm{rc}}$. We also let d_{cl} be the smallest length of a chain of lines connecting two general points in X. Our main result is the following theorem.

Date: October 18, 2014.

2000 *Mathematics Subject Classification.* Primary 14N35; Secondary 19E08, 14N15, 14M15, 14M20, 14M22.

The first author was supported in part by NSF grant DMS-1205351.

The third author was supported in part by NSA Young Investigator Award H98230-13-1-0208.

Theorem 1. *We have* $(\alpha \star \beta)_d = 0$ *for all* $d \geq d_{\mathrm{rc}} + d_{\mathrm{cl}}$.

The Gromov-Witten varieties $M_d(x, y, z)$ of large degrees are known to be rational when X is a cominuscule homogeneous space, an orthogonal Grassmannian $\mathrm{OG}(m, N)$ for $m \neq \frac{N}{2} - 1$, or any adjoint variety of type different from A or G_2. This was proved in [4] for Grassmannians of type A and in [5] in all other cases. Theorem 1 therefore establishes the finiteness of quantum K-theory for many new spaces. The *orthogonal Grassmannian* $\mathrm{OG}(m, N)$ is the variety of isotropic m-dimensional subspaces in the vector space $\mathbb{C}^N$ equipped with a non-degenerate symmetric bilinear form; these varieties account for all spaces G/P where G is a group of type B_n or D_n and P is a maximal parabolic subgroup. The variety $X = G/P$ is called *adjoint* if it is isomorphic to the closed orbit of the adjoint action of G on $\mathbb{P}(\mathrm{Lie}(G))$.

Remark 1.1. We thank Jason Starr for sending us an outline of an argument that uses the results of [6, 7] to prove that the Gromov-Witten varieties $M_d(x, y, z)$ of large degrees are rationally connected when X is any projective rational homogeneous space with $\mathrm{Pic}(X) = \mathbb{Z}$. As a consequence, Theorem 1 can be applied to all such spaces. We also thank Starr for making us aware of [6, Lemma 15.8].

2. Stable maps and Gromov-Witten varieties

We recall here some notation and results from [3]. Let $X = G/P$ be a homogeneous space defined by a semisimple complex linear algebraic group G and a parabolic subgroup $P \subset G$. Let $B \subset P$ be a Borel subgroup. Recall that a *Schubert variety* in X is an orbit closure of a Borel subgroup of G. Equivalently, it is a G-translate of the closure of a B-orbit in X; the latter orbit closure is a *B-stable Schubert variety*. Given an effective degree $d \in H_2(X; \mathbb{Z})$ and an integer $n \geq 0$, the Kontsevich moduli space $\overline{\mathcal{M}}_{0,n}(X, d)$ parametrizes the isomorphism classes of n-pointed stable (genus zero) maps $f : C \to X$ with $f_*[C] = d$, and comes with a total evaluation map $\mathrm{ev} = (\mathrm{ev}_1, \ldots, \mathrm{ev}_n) : \overline{\mathcal{M}}_{0,n}(X, d) \to X^n := X \times \cdots \times X$. Here a map is called *stable* if its automorphism group is finite, i.e. each of its contracted components has at least 3 special points. A detailed construction of this space can be found in the survey [8].

Let $\mathbf{d} = (d_0, d_1, \ldots, d_r)$ be a sequence of effective classes $d_i \in H_2(X; \mathbb{Z})$, let $\mathbf{e} = (e_0, \ldots, e_r) \in \mathbb{N}^{r+1}$, and set $|\mathbf{d}| = \sum d_i$ and $|\mathbf{e}| = \sum e_i$. Let $M_{\mathbf{d},\mathbf{e}} \subset \overline{\mathcal{M}}_{0,|\mathbf{e}|}(X, |\mathbf{d}|)$ be the closure of the locus of stable maps $f : C \to X$ defined on a chain C of $r + 1$ projective lines, such that the i-th projective line contains e_i marked points (numbered from $1 + \sum_{j<i} e_j$ to $\sum_{j \leq i} e_j$) and the restriction of f to this component has degree d_i. To ensure that these maps are indeed stable we assume that $e_i \geq 1 + \delta_{i,0} + \delta_{i,r}$ whenever $d_i = 0$. Moreover, we will assume that $e_0 > 0$ and $e_r > 0$. Set $\mathcal{Z}_{\mathbf{d},\mathbf{e}} = \mathrm{ev}(M_{\mathbf{d},\mathbf{e}}) \subset X^{|\mathbf{e}|}$. Given subvarieties $\Omega_1, \ldots, \Omega_m$ of X with $m \leq |\mathbf{e}|$, define a boundary Gromov-Witten variety by $M_{\mathbf{d},\mathbf{e}}(\Omega_1, \ldots, \Omega_m) = \bigcap_{i=1}^m \mathrm{ev}_i^{-1}(\Omega_i) \subset M_{\mathbf{d},\mathbf{e}}$. We also write $\Gamma_{\mathbf{d},\mathbf{e}}(\Omega_1, \ldots, \Omega_m) = \mathrm{ev}_{|\mathbf{e}|}(M_{\mathbf{d},\mathbf{e}}(\Omega_1, \ldots, \Omega_m)) \subset X$. If no sequence $\mathbf{e}$ is specified, we will use $\mathbf{e} = (3)$ when $r = 0$ and $\mathbf{e} = (2, 0, \ldots, 0, 1)$ when $r > 0$. This convention will be used only when $d_i \neq 0$ for $i > 0$. For this reason the sequence $\mathbf{d} = (d_0, \ldots, d_r)$ will be called a *stable sequence of degrees* if $d_i \neq 0$ for $i > 0$.

An irreducible variety Y has *rational singularities* if there exists a desingularization $\pi : \widetilde{Y} \to Y$ such that $\pi_* \mathcal{O}_{\widetilde{Y}} = \mathcal{O}_Y$ and $R^i \pi_* \mathcal{O}_{\widetilde{Y}} = 0$ for all $i > 0$. An arbitrary

variety has rational singularities if its irreducible components have rational singularities, are disjoint, and have the same dimension. We need the following result from [1, Lemma 3].

Lemma 2.1 (Brion)**.** *Let Z and S be varieties and let $\pi : Z \to S$ be a morphism. If Z has rational singularities, then the same holds for the general fibers of π.*

A morphism $f : Y \to Z$ of varieties is a *locally trivial fibration* if each point $z \in Z$ has an open neighborhood $U \subset Z$ such that $f^{-1}(U) \cong U \times f^{-1}(z)$ and f is the projection to the first factor. The following result is obtained by combining Propositions 2.2 and 2.3 in [3].

Proposition 2.2. *Let $B \subset G$ be a Borel subgroup, let Y be a B-variety, let $\Omega \subset X$ be a B-stable Schubert variety, and let $f : Y \to \Omega$ be a dominant B-equivariant map. Then f is a locally trivial fibration over the dense open B-orbit $\Omega^\circ \subset \Omega$.*

It was proved in [3, Prop. 3.7] that $M_{\mathbf{d},\mathbf{e}}$ is unirational and has rational singularities. Lemma 2.1 therefore implies that $M_{\mathbf{d},\mathbf{e}}(x_1, \dots, x_m)$ has rational singularities for all points $(x_1, \dots, x_m)$ in a dense open subset of $(\mathrm{ev}_1 \times \cdots \times \mathrm{ev}_m)(M_{\mathbf{d},\mathbf{e}}) \subset X^m$. Proposition 2.2 applied to the map $\mathrm{ev}_1 : M_{\mathbf{d},\mathbf{e}} \to X$ shows that $M_{\mathbf{d},\mathbf{e}}(x)$ is unirational for all points $x \in X$. Finally, [3, Lemma 3.9(a)] states that the variety $\mathcal{Z}_{d,2} = \mathrm{ev}(M_{d,2}) \subset X^2$ is rational and has rational singularities for any effective degree $d \in H_2(X; \mathbb{Z})$,

Proposition 2.3. *The variety $M_{\mathbf{d},\mathbf{e}}(x, y)$ is unirational for all points (x, y) in a dense open subset of the image $(\mathrm{ev}_1 \times \mathrm{ev}_2)(M_{\mathbf{d},\mathbf{e}}) \subset X^2$.*

Proof. Set $\Omega = \mathrm{ev}_2(M_{\mathbf{d},\mathbf{e}}(1.P)) \subset X$. Since $M_{\mathbf{d},\mathbf{e}}(1.P)$ is irreducible and P-stable, it follows that Ω is a P-stable Schubert variety. Let $U \subset \Omega$ be the dense open P-orbit. It follows from Proposition 2.2 that $\mathrm{ev}_2 : M_{\mathbf{d},\mathbf{e}}(1.P) \to \Omega$ is a locally trivial fibration over U. Since $M_{\mathbf{d},\mathbf{e}}(1.P)$ is unirational, this implies that $M_{\mathbf{d},\mathbf{e}}(1.P, x)$ is unirational for all $x \in U$. Finally notice that $(\mathrm{ev}_1 \times \mathrm{ev}_2)(M_{\mathbf{d},\mathbf{e}}) = G \times^P \Omega = (G \times \Omega)/P$, where P acts by $(g, x).p = (gp, p^{-1}.x)$, and $M_{\mathbf{d}}(x, y)$ is unirational for all points (x, y) in the dense open subset $G \times^P U \subset G \times^P \Omega$. □

Remark 2.4. It is proved in [6, Lemma 15.8] that, if $\mathbf{d} = (1^d) = (1, 1, \dots, 1)$ with d large, $\mathbf{e} = (1, 0^{d-2}, 1)$, and $\mathrm{Pic}(X) = \mathbb{Z}$, then the general fibers of $\mathrm{ev} : M_{\mathbf{d},\mathbf{e}} \to X^2$ are rationally connected. This also follows from Proposition 2.3. A more general statement is proved in [3, Prop. 3.2].

3. Rationally connected Gromov-Witten varieties

An algebraic variety Z is *rationally connected* if two general points $x, y \in Z$ can be joined by a rational curve, i.e. both x and y belong to the image of some morphism $\mathbb{P}^1 \to Z$. We need the following fundamental result from [10].

Theorem 3.1 (Graber, Harris, Starr)**.** *Let $f : Z \to Y$ be any dominant morphism of complete irreducible complex varieties. If Y and the general fibers of f are rationally connected, then Z is rationally connected.*

We assume from now on that $X = G/P$ is defined by a maximal parabolic subgroup $P \subset G$. Then we have $H_2(X; \mathbb{Z}) = \mathbb{Z}$, so the degree of a curve in X can be identified with an integer. We will further assume that the three-point Gromov-Witten varieties of X of sufficiently high degree are rationally connected. More

precisely, assume that there exists an integer d_{rc} such that $M_d(x,y,z)$ is rationally connected for all $d \geq d_{\mathrm{rc}}$ and all points (x,y,z) in a dense open subset $U_d \subset X^3$.

For $n \geq 2$ we set $d_X(n) = \min\{d \in \mathbb{N} \mid \mathcal{Z}_{d,n} = X^n\}$. This is the smallest integer such that, given n arbitrary points in X, there exists a curve of degree $d_X(n)$ through all n points. Finally we set $d_{\mathrm{cl}} = \min\{d \in \mathbb{N} \mid \mathcal{Z}_{(1^d),(1,0^{d-2},1)} = X^2\}$, where $(1^d) = (1,1,\ldots,1)$ denotes a sequence of d ones. This is the smallest length of a chain of lines connecting two general points in X. Notice that $d_X(3) \leq d_{\mathrm{rc}}$ and $d_X(2) \leq d_{\mathrm{cl}}$.

Theorem 3.2. *Let $\mathbf{d} = (d_0, d_1, \ldots, d_r)$ be a stable sequence of degrees such that $|\mathbf{d}| \geq d_{\mathrm{rc}} + d_{\mathrm{cl}} - 1$. Then we have $\mathcal{Z}_{\mathbf{d}} = \mathcal{Z}_{d_0,2} \times X$, and $M_{\mathbf{d}}(x,y,z)$ is rationally connected for all points (x,y,z) in a dense open subset of $\mathcal{Z}_{\mathbf{d}}$.*

Proof. Set $\mathbf{d}' = (d_1, \ldots, d_r)$ and $\mathbf{e}' = (1,0,\ldots,0,1) \in \mathbb{N}^r$. It follows from [3, Prop. 3.6] that $M_{\mathbf{d}}$ is the product over X of the maps $\mathrm{ev}_3 : M_{d_0,3} \to X$ and $\mathrm{ev}_1 : M_{\mathbf{d}',\mathbf{e}'} \to X$. The assumption implies that $d_0 \geq d_{\mathrm{rc}}$ or $|\mathbf{d}'| \geq d_{\mathrm{cl}}$.

Assume first that $|\mathbf{d}'| \geq d_{\mathrm{cl}}$. It then follows from the definition of d_{cl} that $\mathcal{Z}_{\mathbf{d}} = \mathcal{Z}_{d_0,2} \times X$. Let $X^\circ = Pw_0.P \subset X$ be the open P-orbit. By Proposition 2.3 and Lemma 2.1 we may choose a dense open subset $U \subset \mathcal{Z}_{d_0,2}$ such that, for all points $(x,y) \in U$ we have that $M_{d_0}(x,y)$ is unirational, $\Gamma_{d_0}(x,y) \cap X^\circ \neq \emptyset$, and $M_{\mathbf{d}}(x,y,1.P)$ has rational singularities. Let $(x,y) \in U$. We will show that $M_{\mathbf{d}}(x,y,1.P)$ is rationally connected. Let $p : M_{\mathbf{d}}(x,y,1.P) \to M_{d_0}(x,y)$ be the projection. Then the fibers of p are given by $p^{-1}(f) = M_{\mathbf{d}',\mathbf{e}'}(\mathrm{ev}_3(f), 1.P)$. Since the morphism $\mathrm{ev}_1 : M_{\mathbf{d}',\mathbf{e}'}(X, 1.P) \to X$ is surjective and P-equivariant, Proposition 2.2 implies that this map is locally trivial over X°. Since $M_{\mathbf{d}',\mathbf{e}'}(X,1.P)$ is unirational, we deduce that $M_{\mathbf{d}',\mathbf{e}'}(z',1.P)$ is unirational for all $z' \in X^\circ$. This implies that $p^{-1}(f)$ is unirational for all $f \in M_{d_0}(x,y,X^\circ)$, which is a dense open subset of $M_{d_0}(x,y)$ by choice of U. Since the general fibers of p are connected, it follows from Stein factorization that all fibers of p are connected. Therefore $M_{\mathbf{d}}(x,y,1.P)$ is connected. Since this variety also has rational singularities, we deduce that $M_{\mathbf{d}}(x,y,1.P)$ is irreducible. Finally, Theorem 3.1 applied to the map $p : M_{\mathbf{d}}(x,y,1.P) \to M_{d_0}(x,y)$ shows that $M_{\mathbf{d}}(x,y,1.P)$ is rationally connected.

Assume now that $d_0 \geq d_{\mathrm{rc}}$. In this case we have $\mathcal{Z}_{\mathbf{d}} = X^3$. Let $U \subset X^3$ be a dense open subset such that $M_{\mathbf{d}}(x,y,z)$ has rational singularities and $M_{d_0}(x,y,z)$ is rationally connected and has rational singularities for all $(x,y,z) \in U$. Using similar arguments, one can show that $M_{\mathbf{d}}(x,y,z)$ is rationally connected for all $(x,y,z) \in U$. This follows from Theorem 3.1 again, applied to the map $q : M_{\mathbf{d}}(x,y,z) \to M_{\mathbf{d}',\mathbf{e}'}(X,z)$. Details are left to the reader. □

4. Quantum K-theory

Let $K(X)$ denote the Grothendieck ring of algebraic vector bundles on X. An introduction to this ring can be found in e.g. [2, §3.3]. For each effective degree $d \in H_2(X;\mathbb{Z})$ we define a class $\Phi_d \in K(X^3)$ by

$$\Phi_d = \sum_{\mathbf{d}=(d_0,\ldots,d_r)} (-1)^r \, \mathrm{ev}_*[\mathcal{O}_{M_{\mathbf{d}}}] \,,$$

where the sum is over all stable sequences of degrees $\mathbf{d}$ such that $|\mathbf{d}| = d$, and $\mathrm{ev} : M_{\mathbf{d}} \to X^3$ is the evaluation map. Let $\pi_i : X^3 \to X$ be the projection to the i-th factor. For $\alpha, \beta \in K(X)$ we set $(\alpha \star \beta)_d = \pi_{3*}(\pi_1^*(\alpha) \cdot \pi_2^*(\beta) \cdot \Phi_d) \in K(X)$.

The quantum K-theory ring of X is an algebra over $\mathbb{Z}[\![q]\!]$, which as a $\mathbb{Z}[\![q]\!]$-module is given by $\mathrm{QK}(X) = K(X) \otimes_{\mathbb{Z}} \mathbb{Z}[\![q]\!]$. The multiplicative structure of $\mathrm{QK}(X)$ is defined by

$$\alpha \star \beta = \sum_d (\alpha \star \beta)_d \, q^d$$

for all classes $\alpha, \beta \in K(X)$, where the sum is over all effective degrees d. A theorem of Givental [9] states that $\mathrm{QK}(X)$ is an associative ring. We note that the definition of $\mathrm{QK}(X)$ given here is different from Givental's original construction; the equivalence of the two definitions follows from [3, Lemma 5.1].[1]

We need the following Gysin formula from [4, Thm. 3.1] (see also [3, Prop. 5.2] for the stated version.)

Proposition 4.1. *Let $f : X \to Y$ be a surjective morphism of projective varieties with rational singularities. If the general fibers of f are rationally connected, then $f_*[\mathcal{O}_X] = [\mathcal{O}_Y] \in K(Y)$.*

Corollary 4.2. *Let $\mathbf{d} = (d_0, \ldots, d_r)$ be a stable sequence of degrees such that $|\mathbf{d}| \geq d_{\mathrm{rc}} + d_{\mathrm{cl}} - 1$. Then we have $\mathrm{ev}_*[\mathcal{O}_{M_{\mathbf{d}}}] = [\mathcal{O}_{\mathcal{Z}_{\mathbf{d}}}] \in K(X^3)$.*

Proof. This holds because $\mathcal{Z}_{\mathbf{d}} = \mathcal{Z}_{d_0,2} \times X$ has rational singularities [3, Lemma 3.9], the general fibers of the map $\mathrm{ev} : M_{\mathbf{d}} \to \mathcal{Z}_{\mathbf{d}}$ are rationally connected by Theorem 3.2, and $M_{\mathbf{d}}$ has rational singularities by [3, Prop. 3.7]. □

Theorem 1 is equivalent to the following result.

Theorem 4.3. *We have $\Phi_d = 0$ for all $d \geq d_{\mathrm{rc}} + d_{\mathrm{cl}}$.*

Proof. It follows from Corollary 4.2 that, for $d \geq d_{\mathrm{rc}} + d_{\mathrm{cl}}$ we have

$$\Phi_d = \sum_{\mathbf{d}=(d_0,\ldots,d_r)} (-1)^r [\mathcal{O}_{\mathcal{Z}_{\mathbf{d}}}] \ \in K(X^3)\,,$$

where the sum is over all stable sequences of degrees $\mathbf{d}$ with $|\mathbf{d}| = d$. Since $\mathcal{Z}_{\mathbf{d}} = \mathcal{Z}_{d_0,2} \times X$, the terms of this sum depend only on d_0. Since $d \geq d_{\mathrm{cl}} + d_{\mathrm{rc}} > d_X(2)$, it follows that $\mathcal{Z}_{(d)} = \mathcal{Z}_{(d-1,1)} = X^3$, so the contributions from the sequences $\mathbf{d} = (d)$ and $\mathbf{d} = (d-1, 1)$ cancel each other out. Now let $0 \leq d' \leq d-2$. For each r with $1 \leq r \leq d - d'$, there are exactly $\binom{d-d'-1}{r-1}$ sequences $\mathbf{d}$ in the sum for which $d_0 = d'$ and the length of $\mathbf{d}$ is $r+1$. Since $\sum_{r=1}^{d-d'} (-1)^r \binom{d-d'-1}{r-1} = 0$, it follows that the corresponding terms cancel each other out. It follows that $\Phi_d = 0$, as claimed. □

Remark 4.4. Theorem 4.3 is true also for the equivariant K-theory ring $\mathrm{QK}_T(X)$ with the same proof.

Remark 4.5. If X is not the projective line, then the proof of Theorem 4.3 shows that $\Phi_d = 0$ for all $d \geq d_{\mathrm{rc}} + d_{\mathrm{cl}} - 1$. It would be interesting to determine the maximal value of d for which $\Phi_d \neq 0$. If X is a cominuscule variety, then this number is equal to $d_X(2)$, hence the maximal power of q that appears in products in the quantum K-theory ring of X is equal to the maximal power that appears in the quantum cohomology ring [3].

[1] Lemma 5.1 in [3] is stated only for cominuscule varieties, but its proof works verbatim for any projective rational homogeneous space X.

References

[1] M. Brion, *Positivity in the Grothendieck group of complex flag varieties*, J. Algebra **258** (2002), no. 1, 137–159, Special issue in celebration of Claudio Procesi's 60th birthday. MR 1958901 (2003m:14017)

[2] ———, *Lectures on the geometry of flag varieties*, Topics in cohomological studies of algebraic varieties, Trends Math., Birkhäuser, Basel, 2005, pp. 33–85. MR 2143072 (2006f:14058)

[3] A. S. Buch, P.-E. Chaput, L. Mihalcea, and N. Perrin, *Finiteness of cominuscule quantum K-theory*, Ann. Sci. Ec. Norm. Super. **46**, fascicule 3 (2013), pag. 477– 494.

[4] A. S. Buch and L. C. Mihalcea, *Quantum K-theory of Grassmannians*, Duke Math. J. **156** (2011), no. 3, 501–538. MR 2772069 (2011m:14092)

[5] P.-E. Chaput and N. Perrin, *Rationality of some Gromov-Witten varieties and application to quantum K-theory*, Commun. Contemp. Math. **13** (2011), no. 1, 67–90. MR 2772579 (2012k:14077)

[6] A. J. de Jong, X. He, and J. M. Starr, *Families of rationally simply connected varieties over surfaces and torsors for semisimple groups*, Publ. Math. Inst. Hautes Études Sci. (2011), no. 114, 1–85. MR 2854858

[7] A. J. de Jong and J. Starr, *Low degree complete intersections are rationally simply connected*, preprint http://www.math.sunysb.edu/∼jstarr/papers/nk1006g.pdf.

[8] W. Fulton and R. Pandharipande, *Notes on stable maps and quantum cohomology*, Algebraic geometry—Santa Cruz 1995, Proc. Sympos. Pure Math., vol. 62, Amer. Math. Soc., Providence, RI, 1997, pp. 45–96. MR 1492534 (98m:14025)

[9] A. Givental, *On the WDVV equation in quantum K-theory*, Michigan Math. J. **48** (2000), 295–304, Dedicated to William Fulton on the occasion of his 60th birthday. MR 1786492 (2001m:14078)

[10] T. Graber, J. Harris, and J. Starr, *Families of rationally connected varieties*, J. Amer. Math. Soc. **16** (2003), no. 1, 57–67 (electronic). MR 1937199 (2003m:14081)

[11] Y.-P. Lee, *Quantum K-theory. I. Foundations*, Duke Math. J. **121** (2004), no. 3, 389–424. MR 2040281 (2005f:14107)

Department of Mathematics, Rutgers University, 110 Frelinghuysen Road, Piscataway, NJ 08854, USA
E-mail address: asbuch@math.rutgers.edu

Domaine Scientifique Victor Grignard, 239, Boulevard des Aiguillettes, Université Henri Poincaré Nancy 1, B.P. 70239, F-54506 Vandoeuvre-lès-Nancy Cedex, France
E-mail address: pierre-emmanuel.chaput@univ-lorraine.fr

Department of Mathematics, Virginia Tech University, 460 McBryde, Blacksburg VA 24060, USA
E-mail address: lmihalce@math.vt.edu

Mathematisches Institut, Heinrich-Heine-Universität, D-40204 Düsseldorf, Germany
E-mail address: perrin@math.uni-duesseldorf.de