



HAL
open science

Positive linear functionals on BP^* -algebras

Mohammed El Azhari

► **To cite this version:**

Mohammed El Azhari. Positive linear functionals on BP^* -algebras. *Mathematica Slovaca*, 1995, 45 (3), pp.281-285. hal-01297816

HAL Id: hal-01297816

<https://hal.science/hal-01297816>

Submitted on 4 Apr 2016

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

POSITIVE LINEAR FUNCTIONALS ON BP*-ALGEBRAS

M. EL AZHARI

Abstract. Let A be a BP*-algebra with identity e , $P_1(A)$ be the set of all positive linear functionals f on A such that $f(e) = 1$, and let $M_s(A)$ be the set of all nonzero hermitian multiplicative linear functionals on A . We prove that $M_s(A)$ is the set of extreme points of $P_1(A)$. We also prove that, if $M_s(A)$ is equicontinuous, then every positive linear functional on A is continuous. Finally, we give an example of a BP*-algebra whose topological dual is not included in the vector space generated by $P_1(A)$, which gives a negative answer to a question posed by M. A. Hennings.

Mathematics Subject Classification 2010: 46K05, 46J05.

Keywords: positive linear functional, BP*-algebra, *-representation.

I. Preliminaries

Let A be a topological vector space, we denote by A^* the algebraic dual of A , and by A' the topological dual of A . Let A be an algebra and $x \in A$, we denote by $sp_A(x)$ the spectrum of x , by $\rho_A(x) = \sup\{|\lambda| : \lambda \in sp_A(x)\}$ the spectral radius of x , and by $R(A)$ the radical of A . A topological algebra is an algebra (over the complex field) which is also a Hausdorff topological vector space such that the multiplication is separately continuous. A locally convex algebra is a topological algebra whose topology is locally convex. Let A be a locally convex *-algebra. A linear functional f on A is called positive if $f(x^*x) \geq 0$ for all $x \in A$. A linear functional f on A is said to be hermitian if $f(x^*) = \overline{f(x)}$ for all $x \in A$. Let A be a commutative locally convex *-algebra with continuous involution and identity e . Let β be the family of all closed, bounded, absolutely convex, containing e , hermitian, idempotent subsets of A . For any $B \in \beta$, the linear span A_B of B , is a *-subalgebra of A which, when equipped with the Minkowski norm $\|\cdot\|_B$, becomes a normed *-algebra with identity and isometric involution. If $A = \cup\{A_B, B \in \beta\}$ and $(A_B, \|\cdot\|_B)$ is complete for all $B \in \beta$, we say that A is a BP*-algebra. We denote by $P(A)$ the set of all positive linear functionals on A , by $P_1(A)$ the subset of $P(A)$ consisting of those positive linear functionals f such that $f(e) = 1$, by $M(A)$ the set of all nonzero multiplicative linear functionals on A , and by $M_s(A)$ the set of all nonzero hermitian multiplicative linear functionals on A . A BP*-algebra A is called symmetric if $(e + x^*x)^{-1}$ exists for all $x \in A$.

II. Characterization of extreme points of $P_1(A)$

Let A be a BP*-algebra, $A = \cup\{A_B, B \in \beta\}$. In [5], T.Husain and S.A.Warsi tried to characterize the extreme points of $P_1(A)$, their method is essentially based on the fact that $P_1(A)$ is the projective limit of $P_1(A_B), B \in \beta$, this method gave no result in the general case. Using the known method of Banach algebras, we characterize the extreme points of $P_1(A)$.

Lemma 1. ([6; Theorem 4.4.12]) Let H be a Hilbert space and L any *-subalgebra of $B(H)$. Then L is irreducible on H if and only if the centralizer of L in $B(H)$ contains only scalar multiples of the identity operator.

Proof. See [6].

Theorem 2. Let A be a BP*-algebra, then the extreme points of $P_1(A)$ are exactly the nonzero hermitian multiplicative linear functionals on A .

Proof. Let $\chi \in M_s(A)$ and $x \in A, \chi(x^*x) = \overline{\chi(x)}\chi(x) = |\chi(x)|^2 \geq 0$, therefore $M_s(A) \subset P_1(A)$. Let $f \in P_1(A)$, f is hermitian and admissible [4; Theorem 5.2], so we can associate to f a cyclic representation T_f on a Hilbert space H_f [6; Theorem 4.5.4]. Let f be an extreme point of $P_1(A)$, T_f is an irreducible representation [5; Theorem 5.1]. By Lemma 1, the centralizer Z of $\{T_f x, x \in A\}$ in $B(H_f)$ contains only scalar multiples of the identity operator. Since A is commutative, $\{T_f x, x \in A\} \subset Z$, so for each $x \in A$ there exists $\chi(x) \in C$ such that $T_f x = \chi(x)I$. It is easily shown that the map $x \rightarrow \chi(x)$ from A to C is a nonzero hermitian multiplicative linear functional on A . Let h be a cyclic vector of H_f such that $f(x) = \langle T_f x(h), h \rangle$ for all $x \in A$, then $f(x) = \chi(x)\|h\|^2$, thus $\|h\|^2 = 1$ since $f(e) = \chi(e) = 1$, hence $f = \chi \in M_s(A)$. Conversely, let $\chi \in M_s(A)$ and let (T_χ, H_χ) be the cyclic representation associated with χ . We have $\dim H_\chi = \dim A/\text{Ker}\chi = 1$, therefore T_χ is irreducible, thus χ is an extreme point of $P_1(A)$ [5; Theorem 5.1].

III. On the symmetric spectral radius of a BP*-algebra

Let A be a BP*-algebra, we denote by ρ_s the symmetric spectral radius of A defined by $\rho_s(x) = \sup\{|\chi(x)| : \chi \in M_s(A)\}$ for all $x \in A$. We have $\rho_s(x) \leq \rho_A(x) < \infty$ for all $x \in A$.

Theorem 3. Let A be a BP*-algebra, then $\rho_s(x) = \sup\{f(x^*x) : f \in P_1(A)\}^{\frac{1}{2}}$ for all $x \in A$.

Proof. Put $|x| = \sup\{f(x^*x) : f \in P_1(A)\}^{\frac{1}{2}}$ for all $x \in A$. Let $\chi \in M_s(A) \subset P_1(A)$ and $x \in A, |\chi(x)|^2 = \chi(x^*x) \leq |x|^2$, thus $|\chi(x)| \leq |x|$, hence $\rho_s(x) \leq |x|$. Conversely, let $f \in P_1(A)$, by Theorem 2 and the Krein-Milman Theorem, there exists a net $(f_\alpha)_{\alpha \in \Lambda}$ in $\text{conv } M_s(A)$ such that $f_\alpha \rightarrow f$ in the weak topology $\sigma(A^*, A)$. For each $\alpha \in \Lambda$, there exist $\lambda_1, \dots, \lambda_n \in R^+, \lambda_1 + \dots + \lambda_n = 1$, and $\chi_1, \dots, \chi_n \in M_s(A)$ such that $f_\alpha = \lambda_1 \chi_1 + \dots + \lambda_n \chi_n$. Let $x \in A, f_\alpha(x^*x) \leq$

$\lambda_1|\chi_1(x)|^2 + \dots + \lambda_n|\chi_n(x)|^2 \leq \rho_s(x)^2$, since $f_\alpha(x^*x) \rightarrow f(x^*x)$, it follows that $f(x^*x) \leq \rho_s(x)^2$, therefore $\sup\{f(x^*x) : f \in P_1(A)\} \leq \rho_s(x)^2$, i.e. $|x| \leq \rho_s(x)$.

Corollary 4. Let A be a symmetric BP*-algebra, then $R(A) = \{x \in A : f(x^*x) = 0 \text{ for all } f \in P_1(A)\}$.

Proof. Each nonzero multiplicative linear functional on A is hermitian, therefore $R(A) = \{x \in A : \chi(x) = 0 \text{ for all } \chi \in M_s(A)\} = \{x \in A : \rho_s(x) = 0\}$.

Theorem 5. Let A be a BP*-algebra and $f \in P(A)$, then $|f(x)| \leq f(e)\rho_s(x)$ for all $x \in A$.

Proof. Let $f \in P(A)$. If $f(e) = 0$, by the Schwarz inequality, $|f(x)|^2 \leq f(e)f(x^*x)$ for all $x \in A$, then f vanishes on A . If $f(e) \neq 0$, put $f_1 = f(e)^{-1}f$, it is clear that $f_1 \in P_1(A)$. Let $x \in A$, $f_1(x^*x) \leq \rho_s(x)^2$ by Theorem 3, thus $|f_1(x)|^2 \leq f_1(x^*x) \leq \rho_s(x)^2$, therefore $|f_1(x)| \leq \rho_s(x)$, i.e. $|f(x)| \leq f(e)\rho_s(x)$.

Corollary 6. Let A be a BP*-algebra. If $M_s(A)$ is equicontinuous, then every positive linear functional on A is continuous.

Proof. If $M_s(A)$ is equicontinuous, then ρ_s is continuous.

Corollary 7. Let A be a barrelled BP*-algebra such that every nonzero hermitian multiplicative linear functional on A is continuous. Then every positive linear functional on A is continuous.

Proof $M_s(A)$ is bounded in the weak topology $\sigma(A', A)$. Since A is barrelled, $M_s(A)$ is equicontinuous by the Banach-Steinhaus Theorem.

Remarks.

1. Corollary 6 is an improvement of [4; Theorem 5.7].
2. Corollary 7 is a partial answer to the following problem [1]: is every positive linear functional on a barrelled BP*-algebra continuous ?

IV. On a question of M. A. Hennings

Let A be a BP*-algebra. In [3; question E], M. A. Hennings posed the following question: do the positive linear functionals on A spans the topological dual of A ?

M. A. Hennings [3, p.290 (lines 28-29)] stated that the general problem of whether this question is true or not is still open. Here we give a negative answer to this question, we first have the following proposition:

Proposition 8. Let A be a symmetric BP*-algebra. If $A' \subset \text{span}_{A^*}P(A)$, then A is semisimple.

Proof. We first remark that $\text{span}_{A^*}P(A) = \text{span}_{A^*}P_1(A)$. By Corollary 4, $R(A) = \{x \in A : f(x^*x) = 0 \text{ for all } f \in P_1(A)\}$. Let $x \in R(A)$, by the Schwarz inequality, $|f(x)|^2 \leq f(x^*x)$ for all $f \in P_1(A)$, thus $f(x) = 0$ for all $f \in P_1(A)$. Since $A' \subset \text{span}_{A^*}P_1(A)$, it follows that $g(x) = 0$ for all $g \in A'$, hence $x = 0$.

Remark. Proposition 8 is a generalization of the following Hennings result: If A is a symmetric unital MQ*-algebra such that $A' \subset \text{span}_{A^*}P(A)$, then A is semisimple.

Counter example Let E be the linear space of absolutely continuous functions on $[0,1]$, with the norm $\|f\| = \int_0^1 |f(t)|dt$, the convolution-multiplication $(f \times g)(x) = \int_0^x f(x-t)g(t)dt$, and the natural involution $f^*(x) = \overline{f(x)}$. $(E, \|\cdot\|)$ is a symmetric radical commutative Banach *-algebra [6; p.316]. Let $A = E \oplus Ce$ be the algebra obtained from E by adjunction of an identity e . A is a commutative Banach *-algebra with identity, A is symmetric [6; Theorem 4.7.9]. It is easily shown that $R(A) = E$, thus A is not semisimple, hence A' is not included in $\text{span}_{A^*}P(A)$ by Proposition 8.

V. On representations of BP*-algebras

Let A be a BP*-algebra and $f \in P(A)$. Let $L_f = \{x \in A : f(x^*x) = 0\}$, L_f is a linear space and the quotient space A/L_f is a pre-Hilbert space with the inner product $\langle x+L_f, y+L_f \rangle = f(y^*x)$, $x, y \in A$. Since f is hermitian and admissible, we can associate to f a cyclic representation on a Hilbert space H_f which is the completion of A/L_f . For each $x \in A$, $T_f x$ is defined by $T_f x(y+L_f) = xy+L_f$ for all $y \in A$.

Theorem 9. Let A be a BP*-algebra such that $M_s(A)$ is equicontinuous. Then for each $f \in P(A)$, the representation T_f is continuous.

Proof. Let $x, y \in A$, we have $f(y^*x^*xy) \leq f(y^*y)\rho_s(x^*x)$, thus $\langle xy+L_f, xy+L_f \rangle \leq \langle y+L_f, y+L_f \rangle \rho_s(x^*x)$, i.e. $\|T_f x(y+L_f)\| \leq \|y+L_f\| \rho_s(x^*x)^{\frac{1}{2}}$, hence $\|T_f x\| \leq \rho_s(x^*x)^{\frac{1}{2}} = \rho_s(x)$. Since $M_s(A)$ is equicontinuous, ρ_s is continuous, therefore T_f is continuous.

Remark. Theorem 9 is an improvement of [5; Theorem 4.2].

Acknowledgments. I would like to express my thanks to Professor T. Husain for help he has given me during the preparation of this work.

REFERENCES

[1] M. Akkar, M. El Azhari, M.Oudadess, On an expression of the spectrum in BP*-algebras, Period. Math. Hungar. 19 (1988), 65-67.

- [2] G. R. Allan, On a class of locally convex algebras, Proc. London Math. Soc. 17 (1967), 91-114.
- [3] M. A. Hennings, Topologies on BP*-algebras, J. Math. Anal. Appl. 140 (1989), 289-300.
- [4] T. Husain, S. A. Warsi, Positive functionals on BP*-algebras, Period. Math. Hungar. 8 (1977), 15-28.
- [5] T. Husain, S. A. Warsi, Representations of BP*-algebras, Math. Japon. 21 (1976), 237-247.
- [6] C. E. Richart, General Theory of Banach Algebras, Van Nostrand, Princeton N. J., 1960.
- [7] A. P. Robertson, W. Robertson, Topological Vector Spaces. Cambridge Tracts in Math. and Math. Phys. 53, Cambridge Univ. Press, London, 1973.

Ecole Normale Supérieure
Avenue Oued Akreuch
Takaddoum, BP 5118, Rabat
Morocco

E-mail: mohammed.elazhari@yahoo.fr