



Twenty Two Parameter Deformations of the Twelfth Peregrine Breather Solutions to the NLS Equation

Pierre Gaillard, Mickaël Gastineau

► To cite this version:

Pierre Gaillard, Mickaël Gastineau. Twenty Two Parameter Deformations of the Twelfth Peregrine Breather Solutions to the NLS Equation. *Advances in Research*, 2016, 7 (2), pp.1-11. 10.9734/AIR/2016/25636 . hal-01299469

HAL Id: hal-01299469

<https://hal.science/hal-01299469>

Submitted on 7 Apr 2016

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Families of deformations of the twelfth Peregrine breather solutions to the NLS equation with twenty two parameters.

⁺Pierre Gaillard, [×] Mickaël Gastineau

⁺ Université de Bourgogne,

9 Av. Alain Savary, Dijon, France : Dijon, France :

e-mail: Pierre.Gaillard@u-bourgogne.fr,

[×] IMCCE, Observatoire de Paris, PSL Research University,
CNRS, Sorbonne Universités, UPMC Univ Paris 06, Univ. Lille,

77 Av. Denfert-Rochereau, 75014 Paris, France :

e-mail: Mickael.Gastineau@obspm.fr

April 2, 2016

Abstract

The twelfth Peregrine breather (P_{12} breather) solution to the focusing one dimensional nonlinear Schrödinger equation (NLS) with its twenty two real parameters deformations solutions to the NLS equation are explicitly constructed here. New families of quasi-rational solutions of the NLS equation in terms of explicit quotients of polynomials of degree 156 in x and t by a product of an exponential depending on t are obtained. The patterns of the modulus of these solutions in the $(x; t)$ plane, in function of the different parameters are studied in details.

PACS : 35Q55, 37K10.

Keywords : NLS equation, wronskians, Peregrine breather, rogue waves.

1 Introduction

Since the first resolution of the one dimensional focusing nonlinear Schrödinger equation (NLS) in 1972 by Zakharov and Shabat using the inverse scattering method [1, 2], a lot of studies have been carried out. The first quasi-rational solution to NLS equation was constructed in 1983 by Peregrine [3]. Akhmediev, Eleonskii and Kulagin obtained the two-phase almost periodic solution to the NLS equation and obtained the first higher order analogue of the Peregrine breather [4, 5] in 1986; other families of higher order 3 and 4 were constructed in a series of articles by Akhmediev et al. [6, 7], using Darboux transformations.

Since the beginning of the years 2010, many works were published using various methods, in particular a formulation by means of wronskians was given in [8]. Recently, in 2013, it was found in [9], solutions expressed in terms of determinants of order $2N$ depending on $2N - 2$ real parameters. A new representation has been found as a ratio of a determinant of order $N + 1$ by another one of order N by Ling and Zhao in [10]. Very recently in 2014, another approach has been given in [11] using a dressing method where the solutions are expressed as the quotient of a determinant of order $N + 1$ by another one of order N .

In the present work, we construct new solutions to the focusing one dimensional nonlinear Schrödinger equation which appear as deformations of the (analogue) Peregrine breather of order 12 with 22 real parameters. These solutions are completely expressed as a quotient of two polynomials of degree 156 in x and t by an exponential de-

pending on t . Because of the length of the solutions in terms of polynomials of x and t , we cannot give them in this text; only plots in the (x, t) plane of the modulus of the solutions to analyze the evolution of the solutions in function of the different parameters are presented in details here.

2 Determinant representation of solutions to NLS equation

In [16, 32] the following result have been proved :

Theorem 2.1 *The function v defined by*

$$v(x, t) = \frac{\det((n_{jk})_{j,k \in [1, 2N]})}{\det((d_{jk})_{j,k \in [1, 2N]})} e^{(2it - i\varphi)}$$

is a quasi-rational solution to the NLS equation

$$iv_t + v_{xx} + 2|v|^2 v = 0,$$

where

$$\begin{aligned} n_{j1} &= f_{j,1}(x, t, 0), \\ n_{jk} &= \frac{\partial^{2k-2} f_{j,1}}{\partial \epsilon^{2k-2}}(x, t, 0), \\ n_{jN+1} &= f_{j,N+1}(x, t, 0), \\ n_{jN+k} &= \frac{\partial^{2k-2} f_{j,N+1}}{\partial \epsilon^{2k-2}}(x, t, 0), \\ d_{j1} &= g_{j,1}(x, t, 0), \\ d_{jk} &= \frac{\partial^{2k-2} g_{j,1}}{\partial \epsilon^{2k-2}}(x, t, 0), \\ d_{jN+1} &= g_{j,N+1}(x, t, 0), \\ d_{jN+k} &= \frac{\partial^{2k-2} g_{j,N+1}}{\partial \epsilon^{2k-2}}(x, t, 0), \\ 2 \leq k \leq N, 1 \leq j \leq 2N \end{aligned}$$

The functions f and g are defined for $1 \leq k \leq N$ by :

$$\begin{aligned}
f_{4j+1,k} &= \gamma_k^{4j-1} \sin A_k, \\
f_{4j+2,k} &= \gamma_k^{4j} \cos A_k, \\
f_{4j+3,k} &= -\gamma_k^{4j+1} \sin A_k, \\
f_{4j+4,k} &= -\gamma_k^{4j+2} \cos A_k, \\
f_{4j+1,N+k} &= \gamma_k^{2N-4j-2} \cos A_{N+k}, \\
f_{4j+2,N+k} &= -\gamma_k^{2N-4j-3} \sin A_{N+k}, \\
f_{4j+3,N+k} &= -\gamma_k^{2N-4j-4} \cos A_{N+k}, \\
f_{4j+4,k} &= \gamma_k^{2N-4j-5} \sin A_{N+k}, \\
g_{4j+1,k} &= \gamma_k^{4j-1} \sin B_k, \\
g_{4j+2,k} &= \gamma_k^{4j} \cos B_k, \\
g_{4j+3,k} &= -\gamma_k^{4j+1} \sin B_k, \\
g_{4j+4,k} &= -\gamma_k^{4j+2} \cos B_k, \\
g_{4j+1,N+k} &= \gamma_k^{2N-4j-2} \cos B_{N+k}, \\
g_{4j+2,N+k} &= -\gamma_k^{2N-4j-3} \sin B_{N+k}, \\
g_{4j+3,N+k} &= -\gamma_k^{2N-4j-4} \cos B_{N+k}, \\
g_{4j+4,N+k} &= \gamma_k^{2N-4j-5} \sin B_{N+k},
\end{aligned}$$

The arguments A_ν and B_ν of these functions are given for $1 \leq \nu \leq 2N$ by

$$\begin{aligned}
A_\nu &= \kappa_\nu x/2 + i\delta_\nu t - ix_{3,\nu}/2 - ie_\nu/2, \\
B_\nu &= \kappa_\nu x/2 + i\delta_\nu t - ix_{1,\nu}/2 - ie_\nu/2.
\end{aligned}$$

The terms κ_ν , δ_ν , γ_ν are defined by $1 \leq \nu \leq 2N$

$$\begin{aligned}
\kappa_j &= 2\sqrt{1-\lambda_j^2}, \quad \delta_j = \kappa_j \lambda_j, \\
\gamma_j &= \sqrt{\frac{1-\lambda_j}{1+\lambda_j}}, \quad \kappa_{N+j} = \kappa_j, \\
\delta_{N+j} &= -\delta_j, \quad \gamma_{N+j} = 1/\gamma_j, \\
1 &\leq j \leq N,
\end{aligned} \tag{2}$$

where λ_j are given for $1 \leq j \leq N$ by :

$$\lambda_j = 1 - 2j^2\epsilon^2, \quad \lambda_{N+j} = -\lambda_j. \tag{3}$$

The terms $x_{r,\nu}$ ($r = 3, 1$) are defined for $1 \leq \nu \leq 2N$ by :

$$x_{r,\nu} = (r-1) \ln \frac{\gamma_\nu - i}{\gamma_\nu + i}. \tag{4}$$

The parameters e_ν are given by

$$\begin{aligned}
e_j &= i \sum_{k=1}^{N-1} \tilde{a}_j \epsilon^{2k+1} j^{2k+1} \\
&\quad - \sum_{k=1}^{N-1} \tilde{b}_j \epsilon^{2k+1} j^{2k+1}, \\
e_{N+j} &= i \sum_{k=1}^{N-1} \tilde{a}_j \epsilon^{2k+1} j^{2k+1} \\
&\quad + \sum_{k=1}^{N-1} \tilde{b}_j \epsilon^{2k+1} j^{2k+1}, \\
1 &\leq j \leq N,
\end{aligned} \tag{5}$$

3 Quasi-rational solutions of order 12 with twenty two parameters

We have already explicitly constructed quasi rational solutions to NLS equation from order 2 until 11. We construct here deformations of the Peregrine breather P_{12} of order 12 depending on 22 parameters. We do not give the analytic expression of the solution to NLS equation of order 12 with twenty two parameters. The computations were done using the computer algebra systems Maple and TRIP [38]. For simplicity, we denote

$$\begin{aligned}
d_3 &:= \det((n_{jk})_{j,k \in [1, 2N]}), \\
d_1 &:= \det((d_{jk})_{j,k \in [1, 2N]}).
\end{aligned}$$

The number of terms of the polynomials of the numerator d_3 and denominator d_1 of the solutions are shown in the table below (Table 1) when other a_i and b_i are set to 0.

N=11	number of terms
$d3(a_1, b_1, x, t)$	1535989
$d1(a_1, b_1, x, t)$	777834
$d3(a_2, b_2, x, t)$	579793
$d1(a_2, b_2, x, t)$	293604
$d3(a_3, b_3, x, t)$	310973
$d1(a_3, b_3, x, t)$	157500
$d3(a_4, b_4, x, t)$	196817
$d1(a_4, b_4, x, t)$	99668
$d3(a_5, b_5, x, t)$	196817
$d1(a_5, b_5, x, t)$	70052
$d3(a_6, b_6, x, t)$	138317
$d1(a_6, b_6, x, t)$	52513
$d3(a_7, b_7, x, t)$	81416
$d1(a_7, b_7, x, t)$	41240
$d3(a_8, b_8, x, t)$	65909
$d1(a_8, b_8, x, t)$	33334
$d3(a_9, b_9, x, t)$	52787
$d1(a_9, b_9, x, t)$	26728
$d3(a_{10}, b_{10}, x, t)$	38627
$d1(a_{10}, b_{10}, x, t)$	19547
$d3(a_{11}, b_{10}, x, t)$	21558
$d1(a_{11}, b_{10}, x, t)$	10908

Table 1: Number of terms for the polynomials d_3 and d_1 of the solutions of the NLS equation.

We construct figures to show deformations of the Peregrine breather of order 12. We get different types of symmetries in the plots in the (x, t) plane. We give some examples of this fact in the following discussion.

3.1 Peregrine breather of order 12

If we choose $\tilde{a}_i = \tilde{b}_i = 0$ for $1 \leq i \leq 10$, we obtain the classical eleventh Peregrine breather




Figure 1: Solution of NLS, $N=12$, all parameters equal to 0, Peregrine breather P_{12} .

3.2 Variation of parameters

With other choices of parameters, we obtain all types of configurations : triangles and multiple concentric rings configurations with a maximum of 78 peaks.




Figure 2: Solution of NLS, $N=12$, $\tilde{a}_1 = 10^3$: triangle with 78 peaks; on the right, sight from top.





Figure 3: Solution of NLS, $N=12$, $\tilde{b}_1 = 10^3$: triangle with 78 peaks; on the right, sight from top.





Figure 5: Solution of NLS, $N=12$, $\tilde{b}_2 = 10^5$: 7 rings with 5, 10, 10, 10, 20, 10, 10, peaks on each of them with in the center P_2 ; on the right, sight from top.





Figure 4: Solution of NLS, $N=12$, $\tilde{a}_2 = 10^5$: 7 rings with 5, 10, 10, 10, 20, 10, 10, peaks on each of them with in the center P_2 ; on the right, sight from top.





Figure 6: Solution of NLS, $N=12$, $\tilde{a}_3 = 10^6$: 7 rings with 7, 14, 7, 21, 14, 7, 7 peaks with in the center P_1 ; on the right, sight from top.





Figure 7: Solution of NLS, $N=12$, $\tilde{b}_3 = 10^6$: 7 rings with 7, 14, 7, 21, 14, 7, 7 peaks with in the center P_1 ; on the right, sight from top.





Figure 9: Solution of NLS, $N=12$, $\tilde{b}_4 = 10^8$: 6 rings with 9, 18, 9, 9, 19, 9 peaks, in the center P_3 ; on the right, sight from top.





Figure 8: Solution of NLS, $N=12$, $\tilde{a}_4 = 10^8$: 6 rings with 9, 18, 9, 9, 19, 9 peaks, in the center P_3 ; on the right, sight from top.





Figure 10: Solution of NLS, $N=11$, $\tilde{a}_5 = 10^{11}$: 6 rings of 11, 11, 22, 11, 11, 11 peaks one peak in the center; on the right, sight from top.





Figure 11: Solution of NLS, $N=11$, $\tilde{b}_5 = 10^{11}$: 6 rings of 11, 11, 22, 11, 11, 11 peaks one peak in the center; on the right, sight from top.





Figure 13: Solution of NLS, $N=12$, $\tilde{b}_6 = 10^{13}$: 6 rings with 13 peaks without peak in the center; on the right, sight of top.





Figure 12: Solution of NLS, $N=12$, $\tilde{a}_6 = 10^{13}$: 6 rings with 13 peaks without peak in the center; on the right, sight of top.





Figure 14: Solution of NLS, $N=12$, $\tilde{a}_7 = 10^{15}$: 5 rings with 15 peaks and in the center the Peregrine breather of order 2; on the right, sight from top.




Figure 15: Solution of NLS, $N=12$, $\tilde{a}_7 = 10^{15}$: 5 rings with 15 peaks and in the center the Peregrine breather of order 2; on the right, sight from top.




Figure 17: Solution of NLS, $N=12$, $\tilde{b}_8 = 10^{18}$: 4 rings with 17 peaks and in the center the Peregrine breather of order 4; on the right, sight from top.




Figure 16: Solution of NLS, $N=12$, $\tilde{a}_8 = 10^{18}$: 4 rings with 17 peaks and in the center the Peregrine breather of order 4; on the right, sight from top.




Figure 18: Solution of NLS, $N=10$, $\tilde{a}_9 = 10^{20}$: 3 rings with 19 peaks and in the center the Peregrine breather of order 6; on the right, sight from top.




Figure 19: Solution of NLS, $N=10$, $\tilde{b}_9 = 10^{20}$: 3 rings with 19 peaks and in the center the Peregrine breather of order 6; on the right, sight from top.




Figure 21: Solution of NLS, $N=12$, $\tilde{b}_{10} = 10^{20}$: two rings with 21 peaks and in the center the Peregrine breather of order 8; on the right, sight from top.




Figure 20: Solution of NLS, $N=12$, $\tilde{a}_{10} = 10^{20}$: two rings with 21 peaks and in the center the Peregrine breather of order 8; on the right, sight from top.




Figure 22: Solution of NLS, $N=12$, $\tilde{a}_{11} = 10^{20}$: one ring with 23 peaks and in the center the Peregrine breather of order 10; on the right, sight from top.




Figure 23: Solution of NLS, $N=12$, $\tilde{b}_{11} = 10^{20}$: one ring with 23 peaks and in the center the Peregrine breather of order 10; on the right, sight from top.

4 Conclusion

Among the various methods currently built to determine explicit solutions of the nonlinear Schrödinger equation, by using the transformations of Darboux [6, 7, 39], the bilinear method of Hirota [40], the ratio of special determinants [10, 11] or the wronskians [41, 16, 32], the method presented in this text is the most effective.

We have constructed explicitly multi-parametric solutions to the NLS equation of order 12 with 22 real parameters. The explicit representation in terms of polynomials of degree 156 in x and t is obtained. We cannot give its expression because of its length. It confirms the property about the shape of the breather in the (x, t) coordinates, the maximum of amplitude equal to $2N + 1$ and the degree of polynomials

in x and t here equal to $N(N + 1)$.

We obtained different patterns in the $(x; t)$ plane by different choices of these parameters. So we obtain a classification of the rogue waves at order 12.

We obtain two types of patterns : the triangular shape and the concentric rings for the same index i for a_i or b_i non equals to 0.

In the cases $a_1 \neq 0$ or $b_1 \neq 0$ we obtain triangles with a maximum of 78 peaks; for $a_2 \neq 0$ or $b_2 \neq 0$, we have 7 rings with 5, 10, 10, 10, 20, 10, 10 peaks with in the center the Peregrine P_2 . For $a_3 \neq 0$ or $b_3 \neq 0$, we obtain 7 rings with 7, 14, 7, 21, 14, 7, 7 peaks with in the center one peak. For $a_4 \neq 0$ or $b_4 \neq 0$, we have 6 rings with 9, 18, 9, 9, 18, 9 peaks with in the center the Peregrine P_3 . For $a_5 \neq 0$ or $b_5 \neq 0$, we have 6 rings of 11, 11, 22, 11, 11, 11 peaks with in the center, one peak. For $a_6 \neq 0$ or $b_6 \neq 0$, we have 6 rings with 13 peaks on each of them without peak in the center. For $a_7 \neq 0$ or $b_7 \neq 0$, we have 5 rings with 15 peaks on each of them and in the center the Peregrine breather of order 2. For $a_8 \neq 0$ or $b_8 \neq 0$, we have 4 rings with 17 peaks on each of them and in the center the Peregrine breather of order 4. For $a_9 \neq 0$ or $b_9 \neq 0$, we have 3 rings with 19 peaks and in the center the Peregrine breather of order 6. For $a_{10} \neq 0$ or $b_{10} \neq 0$, we have 2 rings with 21 peaks and in the center the Peregrine breather of order 8. At least, for $a_{11} \neq 0$ or $b_{11} \neq 0$, we have only one ring with 23 peaks and in the center the Peregrine breather of order 10.

With the previous study, we can formulate the following conjectures about the structure of solutions to the NLS equation based on the parameters a_j and b_j ; we chose the convention that P_0 represent 0 peak. These conjectures

are given under the hypothesis that only one parameter is not equal to 0 :

at order N , for $a_1 \neq 0$ or $b_1 \neq 0$, the modulus of the solution to the NLS equation presents the configuration of a triangle with $N(N+1)/2$ peaks;

at order N , in the case $1 \leq i \leq [\frac{N}{2}]$, for $a_{N-i} \neq 0$ or $b_{N-i} \neq 0$, the modulus of the solution to the NLS equation presents i concentric rings with $2N - 2i + 1$ peaks and in the center the P_{N-2i} breather;

at order N , in the case $[\frac{N}{2}] < i \leq N - 2$, for $a_{N-i} \neq 0$ or $b_{N-i} \neq 0$, the modulus of the solution to the NLS equation presents n_k rings of $k(2N - 2i + 1)$ peaks, for $1 \leq k \leq r$, $rn_r(2N - 2i + 1) < N(N+1)/2$ with in the center the P_{N-2i} breather, verifying

$$\sum_{n_k=1}^r n_k k(2N - 2i + 1) = 2iN - 2i^2 + i.$$

It currently has many applications in various fields like nonlinear optics [42] or hydrodynamics [43]. We hope that this work will be able to lead to physical applications and to bring a better comprehension of the phenomenon of the rogue waves.

It would be relevant to continue this study to try to classify them in the general case of order N ($N > 11$) and to prove the preceding conjectures.

References

- [1] V. E. Zakharov, Stability of periodic waves of finite amplitude on

a surface of a deep fluid, J. Appl. Tech. Phys, V. **9**, 86-94, (1968)

- [2] V. E. Zakharov, A.B. Shabat Exact theory of two dimensional self focusing and one dimensional self modulation of waves in nonlinear media, Sov. Phys. JETP, V. **34**, 62-69, (1972)
- [3] D. Peregrine, Water waves, nonlinear Schrödinger equations and their solutions, J. Austral. Math. Soc. Ser. B, V. **25**, 16-43, (1983)
- [4] N. Akhmediev, V. Eleonski, N. Kulagin, Generation of periodic trains of picosecond pulses in an optical fiber : exact solutions, Sov. Phys. J.E.T.P., V. **62**, 894-899, (1985)
- [5] N. Akhmediev, V. Eleonski, N. Kulagin, Exact first order solutions of the nonlinear Schrödinger equation, Th. Math. Phys., V. **72**, N. 2, 183-196, (1987)
- [6] N. Akhmediev, A. Ankiewicz, J.M. Soto-Crespo, Rogue waves and rational solutions of nonlinear Schrödinger equation, Physical Review E, V. **80**, N. 026601, (2009)
- [7] N. Akhmediev, A. Ankiewicz, P.A. Clarkson, Rogue waves, rational solutions, the patterns of their zeros and integral relations, J. Phys. A : Math. Theor., V. **43**, 122002-1-9, (2010)
- [8] P. Gaillard, Families of quasi-rational solutions of the NLS equation and multi-rogue waves, J. Phys. A : Meth. Theor., V. **44**, 1-15, (2011)

- [9] P. Gaillard, Degenerate determinant representation of solution of the NLS equation, higher Peregrine breathers and multi-rogue waves, *Jour. Of Math. Phys.*, V. **54**, 013504-1-32, (2013)
- [10] L. Ling, L.C. Zhao, Trajectory Characters of rogue waves, *arXiv* : 1305.5599v1, 24 May, (2013)
- [11] A.A. Gelash, V.E. Zakharov, Superregular solitonic solutions : a novel scenario for the nonlinear stage of modulation instability, *Non linearity*, V. **27**, 1-39, (2014)
- [12] P. Gaillard, V.B. Matveev, Wronskian addition formula and its applications, *Max-Planck-Institut für Mathematik, MPI 02-31*, V. **161**, (2002)
- [13] P. Gaillard, A new family of deformations of Darboux-Pöschl-Teller potentials, *Lett. Math. Phys.*, V. **68**, 77-90, (2004)
- [14] P. Gaillard, V.B. Matveev, New formulas for the eigenfunctions of the two-particle Calogero-Moser system, *Lett. Math. Phys.*, V. **89**, 1-12, (2009)
- [15] P. Gaillard, V.B. Matveev, Wronskian and Casorai determinant representations for Darboux-Pöschl-Teller potentials and their difference extensions, *J. Phys A : Math. Theor.*, V. **42**, 1-16, (2009)
- [16] P. Gaillard, Wronskian representation of solutions of the NLS equation and higher Peregrine breathers, *J. Math. Sciences : Adv. Appl.*, V. **13**, N. 2, 71-153, (2012)
- [17] P. Gaillard, Wronskian representation of solutions of NLS equation and seventh order rogue waves, *J. Mod. Phys.*, V. **4**, N. 4, 246-266, (2013)
- [18] P. Gaillard, V.B. Matveev, Wronskian addition formula and Darboux-Pöschl-Teller potentials, *J. Math.*, V. **2013**, ID 645752, 1-10, (2013)
- [19] P. Gaillard, Two parameters deformations of ninth Peregrine breather solution of the NLS equation and multi rogue waves, *J. Math.*, V. **2013**, 1-111, (2013)
- [20] P. Gaillard, Two-parameters determinant representation of seventh order rogue waves solutions of the NLS equation, *J. Theor. Appl. Phys.*, V. **7**, N. **45**, 1-6, (2013)
- [21] P. Gaillard, Six-parameters deformations of fourth order Peregrine breather solutions of the NLS equation, *J. Math. Phys.*, V. **54**, 073519-1-22, (2013)
- [22] P. Gaillard, Deformations of third order Peregrine breather solutions of the NLS equation with four parameters, *Phys. Rev. E*, V. **88**, 042903-1-9, (2013)
- [23] P. Gaillard, Ten parameters deformations of the sixth order Peregrine breather solutions of the NLS equation, *Phys. Scripta*, V. **89**, 015004-1-7, (2014)
- [24] P. Gaillard, The fifth order Peregrine breather and its eight-parameters deformations solutions of the NLS equation, *Commun. Theor. Phys.*, V. **61**, 365-369, (2014)

- [25] P. Gaillard, Higher order Peregrine breathers, their deformations and multi-rogue waves, *J. Of Phys. : Conf. Ser.*, V. **482**, 012016-1-7, (2014)
- [26] P. Gaillard, M. Gastineau, Eighteen parameter deformations of the Peregrine breather of order ten solutions of the NLS equation, *Int. J. Mod. Phys. C*, V. **26**, N. 2, 1550016-1-14, (2014)
- [27] P. Gaillard, Two parameters wronskian representation of solutions of nonlinear Schrödinger equation, eight Peregrine breather and multi-rogue waves, *J. Math. Phys.*, V. **5**, 093506-1-12, (2014)
- [28] P. Gaillard, Hierarchy of solutions to the NLS equation and multi-rogue waves, *J. Phys. : Conf. Ser.*, V. **574**, 012031-1-5, (2015)
- [29] P. Gaillard, Tenth Peregrine breather solution of the NLS, *Ann. Phys.*, V. **355**, 293-298, (2015)
- [30] P. Gaillard, M. Gastineau, The Peregrine breather of order nine and its deformations with sixteen parameters solutions of the NLS equation *Phys. Lett. A*, V. **379**, 1309-1313, (2015)
- [31] P. Gaillard, Other 2N-2 parameters solutions to the NLS equation and 2N+1 highest amplitude of the modulus of the N-th order AP breather, *J. Phys. A: Math. Theor.*, V. **48**, 145203-1-23, (2015)
- [32] P. Gaillard, Multi-parametric deformations of the Peregrine breather of order N solutions to the NLS equation and multi-rogue waves, *Adv. Res.*, V. **4**, 346-364, (2015)
- [33] P. Gaillard, Higher order Peregrine breathers solutions to the NLS equation, *Jour. Phys. : Conf. Ser.*, V. **633**, 012106-1-6, (2015)
- [34] P. Gaillard, M. Gastineau Patterns of deformations of Peregrine breather of order 3 and 4, solutions to the NLS equation with multi-parameters, *Journal of Theoretical and Applied Physics*, (2016)
- [35] P. Gaillard, M. Gastineau Twenty parameters families of solutions to the NLS equation and the eleventh Peregrine breather, *Commun. Theor. Phys*, V. **65**, 136-144, (2016)
- [36] P. Gaillard, Rational solutions to the KPI equation and multi rogue waves, *Annals Of Physics*, V. **355**, 293-298, (2016)
- [37] P. Gaillard, Towards a classification of the quasi rational solutions to the NLS equation, *Teoreticheskaya Y Matematicheskaya Fyzyka*, (2015)
- [38] M. Gastineau, J. Laskar, TRIP 1.3.8 : TRIP Reference manual, IMCCE, Paris Observatory, (2014), <http://www.imcce.fr/trip/>
- [39] B. Guo, L. Ling, Q.P. Liu, Non-linear Schrodinger equation: Generalized Darboux transformation and rogue wave solutions, *Phys. Rev. E*, V. **85**, 026607, (2012)

- [40] Y. Ohta, J. Yang, General high-order rogue waves and their dynamics in the nonlinear Schrödinger equation, *Pro. R. Soc. A*, V. 468, 1716-1740, (2012)
- [41] P. Dubard, P. Gaillard, C. Klein, V.B. Matveev, On multi-rogue waves solutions of the NLS equation and positon solutions of the KdV equation, *Eur. Phys. J. Special Topics*, V. **185**, 247-258, (2010)
- [42] B. Kibler, J. Fatome, C. Finot, G. Millot, F. Dias, G. Genty, N. Akhmediev, J.M. Dudley, The Peregrine soliton in nonlinear fibre optics, *Nat. Phys.*, V. 6, 790-795, (2010)
- [43] A. Chabchoub, H. Hoffmann, M. Onorato, N. Akhmediev, Super rogue waves : observation of a higher-order breather in water waves, *Phys. Review X*, V. **2**, 1-6, (2012).