



HAL
open science

Adaptive Horizontal Gene Transfers between Multiple Cheese-Associated Fungi

Jeanne Ropars, Ricardo Rodriguez de La Vega, Manuela Lopez-Villavicencio, Jerome Gouzy, Erika Sallet, Emilie E. Dumas, Sandrine Lacoste, Robert Debuchy, Joelle Dupont, Antoine Branca, et al.

► **To cite this version:**

Jeanne Ropars, Ricardo Rodriguez de La Vega, Manuela Lopez-Villavicencio, Jerome Gouzy, Erika Sallet, et al.. Adaptive Horizontal Gene Transfers between Multiple Cheese-Associated Fungi. *Current Biology - CB*, 2015, 25 (19), pp.2562-2569. 10.1016/j.cub.2015.08.025 . hal-01302701

HAL Id: hal-01302701

<https://hal.science/hal-01302701>

Submitted on 14 Apr 2016

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License

Current Biology

Adaptive Horizontal Gene Transfers between Multiple Cheese-Associated Fungi

Highlights

- New HTRs are found in cheese fungi
- HTRs are flanked by specific transposable elements
- HTRs have spread in cheese-associated fungi through recent selective sweeps
- Experiments link two HTRs to growth and competitive advantages on cheese

Authors

Jeanne Ropars,
Ricardo C. Rodríguez de la Vega,
Manuela López-Villavicencio, ...,
Joëlle Dupont, Antoine Branca,
Tatiana Giraud

Correspondence

antoine.branca@u-psud.fr (A.B.),
tatiana.giraud@u-psud.fr (T.G.)

In Brief

Ropars et al. report newly discovered horizontally transferred regions, flanked by specific transposable elements that allow cheese-making fungi and food spoilers to grow faster and be better competitors on cheese. These findings have industrial and food safety implications and also improve our understanding of adaptation processes.

Accession Numbers

HG813308–HG813531
HG813601–HG814182
HG814183–HG815135
HG815136–HG815288
HG815290–HG816004
HG816029–HG818118



Adaptive Horizontal Gene Transfers between Multiple Cheese-Associated Fungi

Jeanne Ropars,^{1,2,8} Ricardo C. Rodríguez de la Vega,^{1,2,8} Manuela López-Villavicencio,³ Jérôme Gouzy,^{4,5} Erika Sallet,^{4,5} Émilie Dumas,^{1,2} Sandrine Lacoste,³ Robert Debuchy,^{6,7} Joëlle Dupont,³ Antoine Branca,^{1,2,9,*} and Tatiana Giraud^{1,2,9,*}

¹Ecologie, Systématique et Evolution, UMR8079, Univ. Paris-Sud, 91405 Orsay, France

²Ecologie, Systématique et Evolution, UMR8079, CNRS, 91405 Orsay, France

³Institut de Systématique, Evolution, Biodiversité, UMR 7205 CNRS-MNHN-UPMC-EPHE, Muséum national d'Histoire naturelle, Sorbonne Université, CP39, 57 Rue Cuvier, 75231 Paris Cedex 05, France

⁴Laboratoire des Interactions Plantes-Microorganismes (LIPM), UMR441, INRA, Castanet-Tolosan 31326, France

⁵Laboratoire des Interactions Plantes-Microorganismes (LIPM), UMR2594, CNRS, Castanet-Tolosan 31326, France

⁶Institut de Génétique et Microbiologie, UMR8621, Univ. Paris-Sud, 91405 Orsay, France

⁷Institut de Génétique et Microbiologie, UMR8621, CNRS, 91405 Orsay, France

⁸Co-first author

⁹Co-senior author

*Correspondence: antoine.branca@u-psud.fr (A.B.), tatiana.giraud@u-psud.fr (T.G.)

<http://dx.doi.org/10.1016/j.cub.2015.08.025>

This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

SUMMARY

Domestication is an excellent model for studies of adaptation because it involves recent and strong selection on a few, identified traits [1–5]. Few studies have focused on the domestication of fungi, with notable exceptions [6–11], despite their importance to bioindustry [12] and to a general understanding of adaptation in eukaryotes [5]. *Penicillium* fungi are ubiquitous molds among which two distantly related species have been independently selected for cheese making—*P. roqueforti* for blue cheeses like Roquefort and *P. camemberti* for soft cheeses like Camembert. The selected traits include morphology, aromatic profile, lipolytic and proteolytic activities, and ability to grow at low temperatures, in a matrix containing bacterial and fungal competitors [13–15]. By comparing the genomes of ten *Penicillium* species, we show that adaptation to cheese was associated with multiple recent horizontal transfers of large genomic regions carrying crucial metabolic genes. We identified seven horizontally transferred regions (HTRs) spanning more than 10 kb each, flanked by specific transposable elements, and displaying nearly 100% identity between distant *Penicillium* species. Two HTRs carried genes with functions involved in the utilization of cheese nutrients or competition and were found nearly identical in multiple strains and species of cheese-associated *Penicillium* fungi, indicating recent selective sweeps; they were experimentally associated with faster growth and greater competitiveness on cheese and contained genes highly expressed in the early stage of cheese maturation. These findings have industrial and food safety implications and improve our under-




standing of the processes of adaptation to rapid environmental changes.

RESULTS AND DISCUSSION

Multiple Recent Horizontal Gene Transfers between Distant *Penicillium* Species, Flanked by Specific Retrotransposons

We report here five newly sequenced and assembled *Penicillium* genomes, which we compared with previously available data [16–19]. The full dataset included the genome sequences of ten *Penicillium* species, six of which are either used as industrial inocula for cheese making (*Penicillium roqueforti* and *Penicillium camemberti*) or occur as contaminants in cheeses (Figure 1; Table S1). *P. camemberti* is only found in cheese and is thought to include a single clonal lineage originating from selection programs at the end of the 19th century from the blue-gray cheese molds used at that time, i.e., *Penicillium bifforme* and *Penicillium fuscoglaucum* [20, 21]. By contrast, *P. roqueforti* also occurs in habitats other than cheese, such as silage or wood, and displays substantial genetic diversity [22, 23]. For reconstructing a rooted phylogeny of these ten *Penicillium* species, we used four *Aspergillus* species as an outgroup. We concatenated alignments of 3,089 single-copy genes shared by at least ten species and reconstructed a fully resolved and well-supported maximum-likelihood phylogeny (Figures 1A and 1B).

We used this rooted phylogeny to investigate the occurrence of horizontal gene transfers (HGTs) between *Penicillium* species. As HGTs (also known as xenology [24]) result in incongruences between gene genealogies and species trees, we compared all individual gene genealogies with the species tree. For this goal, we used the Notung software [25–27] to infer the number of duplication, loss, and HGT events that reconciled the gene genealogies with the species tree. Notung is conservative regarding the inference of HGTs because it tests their temporal feasibility, assumes that HGTs occur with a low probability, and forces the poorly supported nodes to follow the species



(legend on next page)

tree. Only orthologous groups with at least one homolog in at least eight genomes were analyzed, further rendering our estimates of HGTs a lower bound. We found overall 104 HGTs between *Penicillium* species, distributed among 77 orthologous groups. Notung inferred the highest number of HGTs relative to branch length in the clade encompassing *P. camemberti*, *P. bifforme*, and their common ancestor (Figure 1B). *P. roqueforti* also acquired many xenologs relative to its branch length. 8 of the 21 horizontally acquired genes detected in *P. roqueforti* were inferred to come from *P. camemberti*, *P. bifforme*, or their common ancestor, indicating recent transfers from species sharing the same ecological niche (Figure S1). Only five of these eight genes could be assigned putative functions, i.e., a protein kinase, two transcription factors, a cation transporter, and an integrase-like protein. Cation transport seems particularly relevant for growth in cheeses as several ions are limiting in this medium (e.g., iron ions), and pH drastically drops during cheese maturation [28].

Another line of evidence for the abundance of HGTs in *Penicillium* fungi came from the finding of multiple large genomic islands that were almost 100% identical at the nucleotidic level between distant species, while being absent from closely related species (Figure 1A). The only substitutions detected in these genomic islands corresponded to repeat-induced point mutations, i.e., C-to-T transitions induced by a specific fungal defense mechanism against transposable elements (TEs) that can substitute multiple base pairs in a single meiosis [29]. In *P. roqueforti*, for example, seven genomic islands larger than 10 kb and displaying above 97% nucleotide identity with multiple other species were found. Only the largest region had previously been identified and was called *Wallaby* [16]. Such a high level of identity suggests that these genomic islands correspond to recent horizontally transferred regions (HTRs), although they could alternatively be recent introgressions. Two lines of evidence, however, support the HTR hypothesis rather than introgression: (1) the presence of several of these regions at non-homologous locations in the different *Penicillium* genomes (Figure S2; [16]) and (2) the low mean genome sequence identity between the *Penicillium* species sharing these regions, being less than 90%, an identity level at which no successful interspecific crosses have ever been reported in fungi [30]. Interestingly, these HTRs were flanked in *P. roqueforti* by copies of TEs from a particular family that were rare elsewhere in the genomes (Figures 1A and S3), the *i* non-LTR retrotransposons [6]. The other

abundant TEs (e.g., *mariner* DNA transposons and *copia* retrotransposons) were in contrast scattered in the genomes (Figure 1A). The genes present in these genomic islands of high sequence similarity partially overlapped with the HGTs detected by Notung. In *P. roqueforti* for example, 17% of the HGTs inferred by Notung were located in the seven large HTRs (Figure 1A). The fact that not all genes in HTRs were detected by Notung mainly results from the filter of this analysis where we used only orthologous groups with homologs in at least eight species. Further, 11% of the inferred HGTs in *P. roqueforti* clustered within 50 kb of the HTRs (Figure 1A), suggesting that these genomic regions may be prone to integrate foreign DNA. This is consistent with the previous finding that the genomic region where *Wallaby* is inserted in *P. roqueforti* carries other species-specific genes in each of *P. camemberti*, *P. rubens*, and *P. roqueforti* [16].

The identification of multiple very recent HTRs, with almost 100% identity in multiple species (Figure 1A), together with Notung inferences of horizontally transferred genes occurring also elsewhere in the genomes (at least 104 in total among *Penicillium* fungi), indicates that HGTs occur frequently among *Penicillium*. The clustering of *i* elements in the flanking regions of HTRs suggests that they may be involved in the mechanism of horizontal transfers. It has been shown that TEs can pass across species boundaries [31] and that they promote genomic rearrangements and recombination [32–34]. The capacity for mycelia fusions may also facilitate the exchange of genetic material in fungi [35].

Two Horizontally Transferred Genomic Regions Are Likely Involved in Cheese Adaptation and Have Spread in Cheese-Associated *Penicillium* through Recent Selective Sweeps



Five of the seven large HTRs detected in *P. roqueforti* were shared between *Penicillium* strains isolated from cheese, *P. camemberti* carrying four of them (Figure 1A). Two HTRs appeared of special relevance for cheese adaptation. *Wallaby* [16] carries a gene encoding an antifungal protein, known for inhibiting the growth of competitors. The second largest HTR was found at terminal edges of scaffolds and was therefore named *CheesyTer*. This 80-kb region, found as a single block in all the genomes studied, carried 37 putative genes, among which two had relevant putative functions for adaptation to cheese, i.e., lactose permease and beta-galactosidase (Figure 2A). Lactose

Figure 1. Horizontal Gene Transfers between *Penicillium* Fungi

(A) The syntenic blocks in *Penicillium* genomes larger than 10,000 bp are shown as heat-colored circles, aligned against the 23 *P. roqueforti* scaffolds that are larger than 10,000 bp (outer blue circle). The percentage of identity to the *P. roqueforti* genome is indicated by heat colors, from yellow for low identity level to red for high identity level. The seven large regions represented in red on the *P. roqueforti* outer circle (blue otherwise) display levels of identity above 97% between several distantly related species, while being absent from others, and are indicated as horizontally transferred regions (HTRs) numbers 1 to 5, *Wallaby* and *CheesyTer*, respectively. These regions are also characterized by the clustering of *i* LINE retrotransposons at their edges; the four most abundant transposable element families are shown on the four outermost gray circles. The red bars in the inner circle indicate the location of the horizontal gene transfer (HGT) events inferred by Notung. The topology of the species phylogeny obtained with 100% bootstrap support based on 3,089 single-copy genes is represented. Asterisks indicate the strains collected from cheese.

(B) Phylogenetic tree of *Penicillium* fungi based on the 3,089 single-copy genes, with branch lengths as estimated by RAxML. The asterisks are as in (A). Branch widths in the tree are proportional to HGT acquisition rate, i.e., number of genes acquired by HGT as inferred by Notung divided by the number of substitutions along the branch.

(C) Presence of *Wallaby* and *CheesyTer* in 416 strains from 65 fungal species. Each column represents a strain; positive (for at least one primer pair) and negative (for all three primer pairs) PCR amplifications are indicated in yellow and black, respectively. Species are ordered by origin (i.e., dairy environment above the blue line versus other environments above the green line) and by taxon (i.e., terverticillate *Penicillium* below the red lines versus other genera below gray lines).



(legend on next page)

is present during the first few days of cheese maturation, and it acts as a primary carbon source, being rapidly consumed by lactic acid bacteria [37]. The two lactose metabolism genes present in *CheesyTer* were among the most strongly expressed in *P. camemberti* during the first step of cheese rind maturation, during which lactose is available (Figure 2B; Table S2; Supplemental Experimental Procedures), indicating a role in the use of the cheese substrate.

We investigated the presence of *CheesyTer* and *Wallaby* by PCR in 416 strains from 65 fungal species from various environments (Figure 1C; Table S3; Supplemental Experimental Procedures). The presence of the transfers was found highly significantly associated with dairy environment both among species ($\chi^2 = 55.7$; degrees of freedom [df] = 1; p value = $8.571e-14$) and among strains within species ($\chi^2 = 45.2$; df = 1, p value = $1.774e-11$). Amplicons were actually obtained only for *Penicillium* species that are frequently isolated in the dairy environment, with the only exception of *P. rubens*, the penicillin-producer fungus, in which 16 strains out of 20 carried either one of the two HTRs. The *CheesyTer* and *Wallaby* fragments obtained by PCR showed zero substitution among all strains from all species, including synonymous sites and non-coding regions, as previously found for *Wallaby* [16]. This result confirms that the presence of *CheesyTer* and *Wallaby* is not ancestral in *Penicillium* species and that these genomic islands have instead been acquired very recently. This also indicates that the two HTRs have spread in several species through recent selective sweeps. *P. roqueforti* was found polymorphic for the presence of *Wallaby* and *CheesyTer*, with all tested strains carrying either both of these regions or neither of them (Table S3). Within *P. roqueforti*, these regions were present only in strains isolated from the cheese environment, suggesting a role in adaptation to the cheese environment.

The *Wallaby* and *CheesyTer* Horizontally Transferred Regions Are Experimentally Associated with Faster Growth and Greater Competitiveness on Cheese

We therefore investigated whether strains carrying *Wallaby* and *CheesyTer* showed higher fitness in terms of growth on cheese substrate or for competitiveness. We set up three experiments, focusing on *P. roqueforti*, because a large collection of strains was available, isolated from various environments, and including strains carrying both *Wallaby* and *CheesyTer* (hereafter named W+C+) and strains lacking them (hereafter named W-C-).

We first compared the growth of 50 *P. roqueforti* strains on a cheese medium and on a minimal medium (26 W+C+ and 24 W-C-; Table S4, tab a). Neither the presence of *Wallaby* and *CheesyTer*, as a main effect independent of the medium, nor the origin of the strain (i.e., cheese versus other environments)

significantly influenced the growth of *P. roqueforti* (Table S1). By contrast, the effect on growth of the medium and its interaction with the presence of the two genomic islands were significant (Figure 3; Table S1): W+C+ strains had a growth advantage on cheese medium but a slower growth on minimal medium.

Second, we investigated whether *P. roqueforti* strains carrying *Wallaby* and *CheesyTer* had a higher ability to exclude competitors. We measured the growth of three fungal strains belonging to species commonly found in cheese but lacking *Wallaby* and *CheesyTer* (*P. nalgiovense* FM193, *P. bifforme* LCP05529, and *Geotrichum candidum* FM074) on plates covered with lawns of *P. roqueforti* either W+C+ (n = 11) or W-C- (n = 12) strains (Table S4, tab b). These experiments were carried out on minimal, cheese, and malt agar media. No difference in growth was detected for the yeast *G. candidum* between lawns of W+C+ or W-C- *P. roqueforti* strains (Table S1). By contrast, W+C+ *P. roqueforti* strains significantly impaired the growth of the two *Penicillium* challengers on the cheese and malt media (Figure 3B). This was not the case on the minimal medium: the interaction between the presence of the transfers and the medium was significant (Table S1). Using the same experimental design, we then investigated the effect of the two genomic islands when present in the challengers. For this goal, we inoculated on *P. roqueforti* lawns (W+C+, n = 2, or W-C-, n = 2), on cheese medium, different strains of species displaying a polymorphism in the *Wallaby* and/or *CheesyTer* presence (Table S4, tab c). We used as challengers different strains of *P. camemberti* (W+C-, n = 1, or W+C+, n = 3), *P. bifforme* (W-C-, n = 2, or W+C+, n = 2), and *P. rubens* (W-C-, n = 1, or W+C-, n = 3). For all three species, we found that the *P. roqueforti* lawns significantly inhibited the growth of challengers and significantly more so when the *P. roqueforti* lawn carried *Wallaby* and *CheesyTer*. Interestingly, the presence of either *CheesyTer* or *Wallaby* in the challengers allowed better growth on W-C- *P. roqueforti* lawns while neither had significant effect on the growth on W+C+ *P. roqueforti* lawns (Table S1).

Third, we investigated competition among *P. roqueforti* strains carrying (W+C+, n = 8) or lacking (W-C-, n = 11) *Wallaby* and *CheesyTer*. We grew *P. roqueforti* strains on cheese medium as pairwise face-to-face confrontations, and we measured the deviations from symmetrical growth (Table S4, tab d; Figure 3C; Supplemental Experimental Procedures). For the W+C+ versus W+C+ confrontations, the mean growth deviation from a boundary in the exact middle of the Petri dish was not significantly different from zero (t test, t = 1.5; df = 36; p value = 0.14). Similar results were obtained for the W-C- versus W-C- confrontations (t test, t = 0.25; df = 70; p value = 0.80). For the W+C+ versus W-C- confrontations, deviations were measured by

Figure 2. Structure of *CheesyTer* and Gene Expression of *Wallaby* and *CheesyTer* in *P. camemberti*

(A) Structure of the *CheesyTer* island in *P. roqueforti* (scaffold PROQFM164_S02 from 30,000 bp to 120,000 bp). *CheesyTer* is entirely shared and syntenic between *P. roqueforti*, *P. bifforme*, and *P. camemberti*. The region in *P. fuscoglaucum* lacks some fragments (shown in yellow) but is syntenic otherwise (fragments shown in red). The putative functions of the genes are shown. *CheesyTer* is flanked by *i* transposable elements, represented in black, showing a high level of identity (e value < $1e-5$). This suggests that they are recently duplicated copies.

(B) Expression of *Wallaby* and *CheesyTer* genes in *P. camemberti* during the first 77 days of cheese rind maturation in industrial Camembert, represented as a heatmap of $\log_2(\text{FPKM}+1)$, a measure of transcript abundance (fragments per kilobase of exon per million fragments mapped). These data were generated in a previous study [36]. The putative functions of the genes are shown; the two genes of *CheesyTer* whose functions are likely involved in lactose metabolism, i.e., the putative lactose permease and beta-galactosidase, are highlighted in gray.




Figure 3. Fitness Advantages of *P. roqueforti* Carrying *Wallaby* and *CheesyTer*, for the Use of the Cheese Substrate and for Outgrowing Competitors

(A) Left: pictures of two *P. roqueforti* strains, with (LCP06166, top) and without (LCP06040, bottom) *Wallaby* and *CheesyTer* on minimal medium and cheese medium. Right: mean growth \pm SE (in mm) of *P. roqueforti* strains with and without *Wallaby* and *CheesyTer* on the two media.

(B) Left: pictures of a *P. biforme* challenger (LCP05529, without *Wallaby* and *CheesyTer*) on two different *P. roqueforti* lawns, on cheese, malt agar, and minimal media (bottom: strain LCP06149 with *Wallaby* and *CheesyTer*; top: strain LCP05885 without the genomic islands; the first line is a control, i.e., with no *P. roqueforti* lawn). Right: mean growth \pm SE (in mm) of a *P. biforme* (top) or a *P. camemberti* (bottom) challenger on *P. roqueforti* strain lawns with or without *Wallaby* and *CheesyTer*.

(C) Growth asymmetry (mean \pm SE in mm of deviations from the middle of the Petri dish) in pairwise confrontations of *P. roqueforti* strains with (W+C+) or without (W-C-) *Wallaby* and *CheesyTer*, on cheese medium, for the three types of possible pairs. The A and B letters correspond to significantly different means according to a Tukey-Kramer test. The picture shows examples of confrontations, at left LCP06271 (W+C+) against LCP06157 (W+C+) and at right LCP00148 (W+C+) against LCP06157 (W-C-).

taking the W+C+ strain as the focal strain; the mean growth deviation was significantly different from zero and positive, the W+C+ strains thus growing farther than the W-C- strain (t test, $t = 12.32$; $df = 90$; p value < 0.0001). The mean deviations were significantly higher in the W-C- versus W+C+ confrontations than in the W+C+ versus W+C+ or W-C- versus W-C- confrontations (Tukey-Kramer test, p value < 0.0001), while the means between these two latter were not significantly different. This experiment shows that the competitive advantage of W+C+ strains against W-C- strains also holds within the species *P. roqueforti*. Altogether, these experimental results strongly support the existence of fitness advantages for the *Penicillium* strains carrying the horizontally transferred genomic islands, both in the use of cheese substrate and in competition with fungal competitors.

Conclusions

Our present study on domesticated fungi shows how adaptation can occur rapidly in eukaryotes. The two cheese species studied here underwent parallel adaptation to the cheese medium, and this involved the transfers of identical regions across species boundaries. HGT events have been reported in fungi

particularly striking. Furthermore, we provide experimental evidence of fitness advantages for strains carrying these HTRs on a human-made medium. These findings altogether are potentially useful for guiding modern strain improvement programs. Indeed, together with the protocol for inducing sex in *P. roqueforti* [22, 42], the identification here of several key candidate genes important for cheese metabolism and competition may allow further selecting interesting traits for cheese industry using the great genetic variability present in *P. roqueforti* strains without *Wallaby* or *CheesyTer* [22]. In addition, our results suggest that caution is required concerning the introduction of genes into microorganisms, as these genes could readily be transferred to other species in the food environment. Indeed, the rapid spread of *Wallaby* and *CheesyTer* into many species of the dairy environment, even when occurring only as contaminants, indicates that transgenes may readily cross species boundaries in the food chain. Finally, the findings here of rapid adaptation through frequent horizontal gene transfers among distant species under selection in novel, human-made media contribute to our understanding of the evolutionary genomic mechanisms allowing rapid adaptation to environmental changes in eukaryotes.

ACCESSION NUMBERS

The accession numbers for the *Penicillium* genome sequences reported in this paper are GenBank: HG813601–HG814182 for *P. bifforme* FM169; HG816029–HG818118 for *P. carneum* LCP05634; HG814183–HG815135 for *P. fuscoglaucum* FM041; HG815136–HG815288 and HG815290–HG816004 for *P. nalgiense* FM193; and HG813308–HG813531 for *P. paneum* FM227.

SUPPLEMENTAL INFORMATION

Supplemental Information includes Supplemental Experimental Procedures, three figures, and four tables and can be found with this article online at <http://dx.doi.org/10.1016/j.cub.2015.08.025>.

AUTHOR CONTRIBUTIONS

Conceptualization, T.G., J.D., and J.R.; Methodology, A.B., T.G., M.L.-V., R.C.R.d.I.V., and J.R.; Software, A.B. and R.C.R.d.I.V.; Validation, É.D. and S.L.; Formal Analysis, A.B., T.G., M.L.-V., R.C.R.d.I.V., and J.R.; Investigation, A.B., M.L.-V., R.C.R.d.I.V., and J.R.; Resources, R.D., J.D., and J.G.; Data Curation, A.B., J.G., R.C.R.d.I.V., and E.S.; Writing – Original Draft, A.B., T.G., R.C.R.d.I.V., and J.R.; Writing – Review & Editing, A.B., T.G., M.L.-V., R.C.R.d.I.V., and J.R.; Visualization, A.B., M.L.-V., R.C.R.d.I.V., and J.R.; Supervision, A.B. and T.G.; Project Administration, A.B. and T.G.; Funding Acquisition, A.B., J.D., T.G., J.G., and R.C.R.d.I.V.

ACKNOWLEDGMENTS

This work was supported by the ANR FROMA-GEN grant (ANR-12-PDOC-0030) awarded to A.B., an “attractivite” grant from Paris-Sud University to A.B., the ERC starting grant GenomeFun 309403 awarded to T.G., a Marie Curie postdoctoral fellowship to R.C.R.d.I.V. (FP7 COFUND PRES-SUD No. 246556), and the ANR grant “Food Microbiomes” (ANR-08-ALIA-007-02) awarded to J.D. We thank Marco van den Berg for sharing raw data for the *Penicillium chrysogenum* Wisconsin 54-1255 strain (recently renamed *P. rubens*).

Received: January 29, 2015

Revised: July 9, 2015

Accepted: August 11, 2015

Published: September 24, 2015

REFERENCES

- Lin, T., Zhu, G., Zhang, J., Xu, X., Yu, Q., Zheng, Z., Zhang, Z., Lun, Y., Li, S., Wang, X., et al. (2014). Genomic analyses provide insights into the history of tomato breeding. *Nat. Genet.* **46**, 1220–1226.
- Bosse, M., Megens, H.J., Frantz, L.A.F., Madsen, O., Larson, G., Paudel, Y., Duijvesteijn, N., Harlizius, B., Hagemeijer, Y., Crooijmans, R.P., and Groenen, M.A. (2014). Genomic analysis reveals selection for Asian genes in European pigs following human-mediated introgression. *Nat. Commun.* **5**, 4392.
- Wu, G.A., Prochnik, S., Jenkins, J., Salse, J., Hellsten, U., Murat, F., Perrier, X., Ruiz, M., Scalabrini, S., Terol, J., et al. (2014). Sequencing of diverse mandarin, pummelo and orange genomes reveals complex history of admixture during citrus domestication. *Nat. Biotechnol.* **32**, 656–662.
- Tian, F., Stevens, N.M., and Buckler, E.S., 4th. (2009). Tracking footprints of maize domestication and evidence for a massive selective sweep on chromosome 10. *Proc. Natl. Acad. Sci. USA* **106** (Suppl 1), 9979–9986.
- Gladieux, P., Ropars, J., Badouin, H., Branca, A., Aguilera, G., de Vienne, D.M., Rodríguez de la Vega, R.C., Branco, S., and Giraud, T. (2014). Fungal evolutionary genomics provides insight into the mechanisms of adaptive divergence in eukaryotes. *Mol. Ecol.* **23**, 753–773.
- Galagan, J.E., Calvo, S.E., Cuomo, C., Ma, L.J., Wortman, J.R., Batzoglou, S., Lee, S.I., Baştürkmen, M., Spevak, C.C., Clutterbuck, J., et al. (2005). Sequencing of *Aspergillus nidulans* and comparative analysis with *A. fumigatus* and *A. oryzae*. *Nature* **438**, 1105–1115.
- Borneman, A.R., Desany, B.A., Riches, D., Affourtit, J.P., Forgan, A.H., Pretorius, I.S., Egholm, M., and Chambers, P.J. (2011). Whole-genome comparison reveals novel genetic elements that characterize the genome of industrial strains of *Saccharomyces cerevisiae*. *PLoS Genet.* **7**, e1001287.
- Gibbons, J.G., Salichos, L., Slot, J.C., Rinker, D.C., McGary, K.L., King, J.G., Klich, M.A., Tabb, D.L., McDonald, W.H., and Rokas, A. (2012). The evolutionary imprint of domestication on genome variation and function of the filamentous fungus *Aspergillus oryzae*. *Curr. Biol.* **22**, 1403–1409.
- Fay, J.C., and Benavides, J.A. (2005). Evidence for domesticated and wild populations of *Saccharomyces cerevisiae*. *PLoS Genet.* **1**, 66–71.
- Novo, M., Bigey, F., Beyne, E., Galeote, V., Gavory, F., Mallet, S., Cambon, B., Legras, J.L., Wincker, P., Casaregola, S., and Dequin, S. (2009). Eukaryote-to-eukaryote gene transfer events revealed by the genome sequence of the wine yeast *Saccharomyces cerevisiae* EC1118. *Proc. Natl. Acad. Sci. USA* **106**, 16333–16338.
- Marsit, S., Mena, A., Bigey, F., Sauvage, F.X., Couloux, A., Guy, J., Legras, J.L., Barrio, E., Dequin, S., and Galeote, V. (2015). Evolutionary advantage conferred by an eukaryote-to-eukaryote gene transfer event in wine yeasts. *Mol. Biol. Evol.* **32**, 1695–1707.
- Stajich, J.E., Berbee, M.L., Blackwell, M., Hibbett, D.S., James, T.Y., Spatafora, J.W., and Taylor, J.W. (2009). The fungi. *Curr. Biol.* **19**, R840–R845.
- McSweeney, P.L.H. (2004). Biochemistry of cheese ripening. *Int. J. Dairy Technol.* **57**, 127–144.
- Wolfe, B.E., Button, J.E., Santarelli, M., and Dutton, R.J. (2014). Cheese rind communities provide tractable systems for *in situ* and *in vitro* studies of microbial diversity. *Cell* **158**, 422–433.
- Irlinger, F., and Mounier, J. (2009). Microbial interactions in cheese: implications for cheese quality and safety. *Curr. Opin. Biotechnol.* **20**, 142–148.
- Cheeseman, K., Ropars, J., Renault, P., Dupont, J., Gouzy, J., Branca, A., Abraham, A.L., Ceppi, M., Conseiller, E., Debuchy, R., et al. (2014). Multiple recent horizontal transfers of a large genomic region in cheese making fungi. *Nat. Commun.* **5**, 2876.
- Marcet-Houben, M., Ballester, A.R., de la Fuente, B., Harries, E., Marcos, J.F., González-Candelas, L., and Gabaldón, T. (2012). Genome sequence of the necrotrophic fungus *Penicillium digitatum*, the main postharvest pathogen of citrus. *BMC Genomics* **13**, 646.
- Cerqueira, G.C., Arnaud, M.B., Inglis, D.O., Skrzypek, M.S., Binkley, G., Simion, M., Miyasato, S.R., Binkley, J., Orvis, J., Shah, P., et al. (2014). The Aspergillus Genome Database: multispecies curation and incorporation of RNA-Seq data to improve structural gene annotations. *Nucleic Acids Res.* **42**, D705–D710.
- van den Berg, M.A., Albang, R., Albermann, K., Badger, J.H., Daran, J.M., Driessen, A.J.M., Garcia-Estrada, C., Fedorova, N.D., Harris, D.M., Heijne, W.H.M., et al. (2008). Genome sequencing and analysis of the filamentous fungus *Penicillium chrysogenum*. *Nat. Biotechnol.* **26**, 1161–1168.
- Giraud, F., Giraud, T., Aguilera, G., Fournier, E., Samson, R., Cruaud, C., Lacoche, S., Ropars, J., Tellier, A., and Dupont, J. (2010). Microsatellite loci to recognize species for the cheese starter and contaminating strains associated with cheese manufacturing. *Int. J. Food Microbiol.* **137**, 204–213.
- Pitt, J.I., Cruickshank, R.H., and Leistner, L. (1986). *Penicillium commune*, *P. camemberti*, the origin of white cheese moulds, and the production of cyclopiazonic acid. *Food Microbiol.* **3**, 363–371.
- Ropars, J., López-Villavicencio, M., Dupont, J., Snirc, A., Gillot, G., Coton, M., Jany, J.L., Coton, E., and Giraud, T. (2014). Induction of sexual reproduction and genetic diversity in the cheese fungus *Penicillium roqueforti*. *Evol. Appl.* **7**, 433–441.
- Gillot, G., Jany, J.-L., Coton, M., Le Floch, G., Debaets, S., Ropars, J., López-Villavicencio, M., Dupont, J., Branca, A., Giraud, T., and Coton, E. (2015). Insights into *Penicillium roqueforti* morphological and genetic diversity. *PLoS ONE* **10**, e0129849.

24. Koonin, E.V. (2005). Orthologs, paralogs, and evolutionary genomics. *Annu. Rev. Genet.* **39**, 309–338.
25. Durand, D., Halldórsson, B.V., and Vernot, B. (2006). A hybrid micro-macroevolutionary approach to gene tree reconstruction. *J. Comput. Biol.* **13**, 320–335.
26. Vernot, B., Stolzer, M., Goldman, A., and Durand, D. (2008). Reconciliation with non-binary species trees. *J. Comput. Biol.* **15**, 981–1006.
27. Stolzer, M., Lai, H., Xu, M., Sathaye, D., Vernot, B., and Durand, D. (2012). Inferring duplications, losses, transfers and incomplete lineage sorting with nonbinary species trees. *Bioinformatics* **28**, i409–i415.
28. Monnet, C., Landaud, S., Bonnarne, P., and Swennen, D. (2015). Growth and adaptation of microorganisms on the cheese surface. *FEMS Microbiol. Lett.* **362**, 1–9.
29. Galagan, J.E., and Selker, E.U. (2004). RIP: the evolutionary cost of genome defense. *Trends Genet.* **20**, 417–423.
30. Le Gac, M., and Giraud, T. (2008). Existence of a pattern of reproductive character displacement in Homobasidiomycota but not in Ascomycota. *J. Evol. Biol.* **21**, 761–772.
31. Schaack, S., Gilbert, C., and Feschotte, C. (2010). Promiscuous DNA: horizontal transfer of transposable elements and why it matters for eukaryotic evolution. *Trends Ecol. Evol.* **25**, 537–546.
32. Baudry, C., Malinsky, S., Restituto, M., Kapusta, A., Rosa, S., Meyer, E., and Bétermier, M. (2009). PiggyMac, a domesticated piggyBac transposase involved in programmed genome rearrangements in the ciliate *Paramecium tetraurelia*. *Genes Dev.* **23**, 2478–2483.
33. Zhou, L., Mitra, R., Atkinson, P.W., Hickman, A.B., Dyda, F., and Craig, N.L. (2004). Transposition of hAT elements links transposable elements and V(D)J recombination. *Nature* **432**, 995–1001.
34. Rajaei, N., Chiruvella, K.K., Lin, F., and Aström, S.U. (2014). Domesticated transposase Kat1 and its fossil imprints induce sexual differentiation in yeast. *Proc. Natl. Acad. Sci. USA* **111**, 15491–15496.
35. Wisecaver, J.H., and Rokas, A. (2015). Fungal metabolic gene clusters-caravans traveling across genomes and environments. *Front Microbiol* **6**, 161.
36. Lessard, M.H., Viel, C., Boyle, B., St-Gelais, D., and Labrie, S. (2014). Metatranscriptome analysis of fungal strains *Penicillium camemberti* and *Geotrichum candidum* reveal cheese matrix breakdown and potential development of sensory properties of ripened Camembert-type cheese. *BMC Genomics* **15**, 235.
37. Leclercq-Perlat, M.N., Buono, F., Lambert, D., Latrille, E., Spinnler, H.E., and Corrieu, G. (2004). Controlled production of Camembert-type cheeses. Part I: Microbiological and physicochemical evolutions. *J. Dairy Res.* **71**, 346–354.
38. Marcet-Houben, M., and Gabaldón, T. (2010). Acquisition of prokaryotic genes by fungal genomes. *Trends Genet.* **26**, 5–8.
39. Wisecaver, J.H., Slot, J.C., and Rokas, A. (2014). The evolution of fungal metabolic pathways. *PLoS Genet.* **10**, e1004816.
40. Friesen, T.L., Stukenbrock, E.H., Liu, Z., Meinhardt, S., Ling, H., Faris, J.D., Rasmussen, J.B., Solomon, P.S., McDonald, B.A., and Oliver, R.P. (2006). Emergence of a new disease as a result of interspecific virulence gene transfer. *Nat. Genet.* **38**, 953–956.
41. Coelho, M.A., Gonçalves, C., Sampaio, J.P., and Gonçalves, P. (2013). Extensive intra-kingdom horizontal gene transfer converging on a fungal fructose transporter gene. *PLoS Genet.* **9**, e1003587.
42. Goarin, A., Silar, P., and Malagnac, F. (2015). Gene replacement in *Penicillium roqueforti*. *Curr. Genet.* **61**, 203–210.