

# **Non-contiguous finished genome sequence and description of Streptococcus varani sp. nov.**

Sofiane Bakour, Jaishriram Rathored, Cheikh Ibrahima Lo, Oleg Mediannikov, Mamadou Beye, Cyrille Bilé Ehounoud, Philippe Biagini, Didier Raoult, Pierre-Edouard Fournier, Florence Fenollar

## **To cite this version:**

Sofiane Bakour, Jaishriram Rathored, Cheikh Ibrahima Lo, Oleg Mediannikov, Mamadou Beye, et al.. Non-contiguous finished genome sequence and description of Streptococcus varani sp. nov.. New Microbes and New Infections, 2016, 11, pp.93-102. 10.1016/j.nmni.2016.03.004. hal-01303332

# **HAL Id: hal-01303332 <https://hal.science/hal-01303332>**

Submitted on 18 Apr 2018

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



[Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0](http://creativecommons.org/licenses/by-nc-nd/4.0/) [International License](http://creativecommons.org/licenses/by-nc-nd/4.0/)

## Non-contiguous finished genome sequence and description of Streptococcus varani sp. nov.

S. Bakour<sup>i</sup>, J. Rathored<sup>i</sup>, C. I. Lo<sup>i,2</sup>, O. Mediannikov<sup>i,2</sup>, M. Beye<sup>l</sup>, C. B. Ehounoud<sup>i</sup>, P. Biagini<sup>3</sup>, D. Raoult<sup>i,2,4</sup>, P.-E. Fournier<sup>i</sup> and F. Fenollar $1,2$ 

1) Aix-Marseille Université, URMITE, UM63, CNRS 7278, IRD 198, Inserm U1095, Faculté de médecine, Marseille cedex 05, France, 2) Campus international UCAD-IRD, Dakar, Senegal, 3) Equipe Emergence et Co-évolution Virale, UMR 7268 – Aix-Marseille Université/ EFS / CNRS, IHU Méditerranée Infection, Marseille, France and 4) Special Infectious Agents Unit, King Fahd Medical Research Center, King Abdul Aziz University, Jeddah, Saudi Arabia

#### Abstract

Strain FF10<sup>T</sup> (= CSUR P1489 = DSM 100884) was isolated from the oral cavity of a lizard (Varanus niloticus) in Dakar, Senegal. Here we used a polyphasic study including phenotypic and genomic analyses to describe the strain FF10 $^\top$ . Results support strain FF10 $^{\top}$  being a Gram-positive coccus, facultative anaerobic bacterium, catalase-negative, non-motile and non-spore forming. The sequenced genome counts 2.46 Mb with one chromosome but no plasmid. It exhibits a G+C content of 40.4% and contains 2471 protein-coding and 45 RNA genes. On the basis of these data, we propose the creation of Streptococcus varani sp. nov.

New Microbes and New Infections © 2016 The Authors. Published by Elsevier Ltd on behalf of European Society of Clinical Microbiology and Infectious Diseases.

Keywords: Culturomics, genome, Streptococcus varani, taxonogenomics, Varanus niloticus Original Submission: 5 February 2016; Accepted: 14 March 2016 Article published online: 19 March 2016

Corresponding author: P.-E. Fournier, URMITE UM63, CNRS7278, IRD198, INSERM1095, Faculté de Médecine, Aix Marseille Université, 27 boulevard Jean Moulin, 13385 Marseille cedex 5, France

E-mail: [pierre-edouard.fournier@univ-amu.fr](mailto:pierre-edouard.fournier@univ-amu.fr)

#### Introduction

The genus Streptococcus contains 113 recognized species and 22 subspecies [\(http://www.bacterio.net/streptococcus.html\)](http://www.bacterio.net/streptococcus.html) as of 5 February 2016. These species are Gram-positive cocci, chainforming, facultative anaerobes and catalase negative [1]. Many of them are associated with human or animal hosts. Previous studies have reported that a large number of Streptococcus species colonize the oral cavities of humans and animals [2]. Among oral streptococci isolated from animals, we note: Streptococcus ursoris isolated from the oral cavities of bears [1], Streptococcus orisratti isolated from the surface of the lower molars of Sprague-Dawley rats [3], Streptococcus oriloxodontae isolated from the oral cavities of elephants [4], and Streptococcus mutans isolated from dental plaque [5].

Recently, with next-generation sequencing technology able to sequence whole genomes in a short time, and mass spectrometric analysis of bacteria, we have had easy access to genetic and proteomic information  $[6]$ . Therefore, we propose a polyphasic approach combining genomic properties in combination with matrix-assisted laser-desorption/ ionization time-of-flight (MALDI-TOF) spectra and phenotypic characteristics to describe new bacterial species. The strain FF10<sup>T</sup> (= CSUR P1489 = DSM 100884) was isolated from the oral cavity of the Nile monitor (Varanus niloticus) in Dakar, Senegal.

Here, we present a summary classification and a set of features for Streptococcus varani sp. nov. strain FF10 $<sup>T</sup>$  (= CSUR</sup> P1489 = DSM 100884), including the description of its complete genome and annotation. These characteristics support the circumscription of the species S. varani.

New Microbe and New Infect 2016; 11: 93–102

New Microbes and New Infections © 2016 The Authors. Published by Elsevier Ltd on behalf of European Society of Clinical Microbiology and Infectious Diseases This is an open access article under the CC BY-NC-ND license ([http://creativecommons.org/licenses/by-nc-nd/4.0/\)](http://creativecommons.org/licenses/by-nc-nd/4.0/)

#### Classification and Features of the Strain

#### Strain identification

In 2014, a sample was collected from the oral cavity of a lizard reptile named 'Nile monitor' (Varanus niloticus) in Dakar, Senegal, and stored at –80°C. In October 2014, the strain FF10<sup>T</sup> (Table 1) was isolated from this sample by cultivation on 5% sheep blood-enriched Columbia agar (BioMérieux, Marcy l'Etoile, France) at 37°C with 5% CO<sub>2</sub>.

To identify the strain, MALDI-TOF MS protein analysis was performed as previously described [7] using a Microflex spectrometer (Bruker Daltonics, Leipzig, Germany). The 12  $FF10<sup>T</sup>$ spectra were imported into the MALDI BIOTYPER software (version 2.0, Bruker) and analysed by standard pattern matching (with default parameter settings) against the main spectra of 6252 bacteria. From the resulting scores, the tested species may or may not be identified compared with the instrument's database; a score  $\geq 2$  with a validly published species enables identification at the species level; a score  $> 1.7$  and  $< 2$  allows identification at the genus level; and a score < 1.7 does not enable any identification. For strain  $\mathsf{FFI0}^\mathsf{T}$ , the scores obtained were lower than 1.306, suggesting that our strain was not a member of any known species. The reference mass spectrum of S. varani strain FF10<sup>T</sup> was added to our database (Fig. 1). A gel view comparing the spectrum of strain  $FF10<sup>T</sup>$  with those of other Streptococcaceae species is shown in Fig. 2. The bacterium was identified using 16S rRNA PCR coupled with sequencing, as previously described  $[8]$ . Strain FF10<sup>T</sup> exhibited 96% 16S rRNA sequence similarity with Streptococcus minor strain DSM 17118





(GenBank accession number: AY232832), the phylogenetically closest bacterial species with standing in nomenclaturevalidated Streptococcus species (Fig. 3). This value was lower than the 98.7% 16S rRNA gene sequence threshold recommended by Stackebrandt and Ebers to delineate a new species without carrying out DNA–DNA hybridization [9]. Strepto $coccus$  varani strain  $FF10<sup>T</sup>$  16S rRNA accession number from GenBank Sequence Database is LN810501.

#### Growth characterization

Different growth temperatures (25°C, 30°C, 37°C, 45°C and 56°C) were tested. Growth was obtained between 25 and 37° C, with optimal growth at 37°C. Growth of the strain was tested also under anaerobic and microaerophilic conditions using GENbag anaer and GENbag microaer systems, respectively (bioMérieux), and under aerobic conditions, with or without  $5\%$  CO<sub>2</sub>. Strain growth was observed under anaerobic and microaerophilic conditions but optimal growth was observed under aerobic conditions. Colonies were translucent and yellow with a regular surface, haemolytic on 5% sheep blood-enriched Columbia agar (bioMérieux), and approximately 1 mm in diameter. A motility test was negative. Cells were Gram-positive cocci, unable to form spores (Fig. 4), and with mean diameter of 0.6 μm (range 0.4–0.8 μm) and mean length of 1.3 μm (range 0.7–1.9 μm) (Fig. 5).

### Biochemical characterization and antibiotic susceptibility

This bacterium,  $FF10<sup>T</sup>$ , exhibits neither catalase nor oxidase activities. Using an API ZYM strip (bioMérieux), positive reactions were observed for alkaline phosphatase, esterase, esterase-lipase, lipase, leucine arylamidase, acid phosphatase, β-glucosidase, β-galactosidase, α-mannosidase, and α-fucosidase. Negative reactions were noted for cystine arylamidase, valine arylamidase, trypsin, α-chymotrypsin, α-glucosidase, α-galactosidase, β-glucuronidase, naphthol-AS-BI-phosphohydrolase, and N-acetyl-β-glucosaminidase. Using an API 20 NE strip (bio-Mérieux), positive reactions were observed only for the esculin hydrolysis test whereas negative reactions were observed for nitrate reduction, urease, indole production, arginine dihydrolase, glucose fermentation, arabinose, mannose, mannitol, Nacetyl-glucosamine, maltose, gluconate, caprate, adipate, malate, citrate, phenyl-acetate assimilation, and gelatine hydrolysis.

Using API 50 CH strip (bioMérieux), negative reactions were observed for the fermentation of glycerol, erythritol, D-arabinose, L-arabinose, D-ribose, D-xylose, L-xylose, D-adonitol, methyl-β-D-xylopyranoside, D-galactose, D-glucose, D-fructose, D-mannose, L-sorbose, L-rhamnose, dulcitol, inositol, Dmannitol, D-sorbitol, methyl-αD-xylopyranoside, methyl-αDglucopyranoside, N-acetylglucosamine, amygdalin, arbutin,



FIG. 1. Reference mass spectrum from Streptococcus varani sp. nov. strain FF10<sup>T</sup>. Spectra from 12 individual colonies were compared and a reference spectrum was generated.



FIG. 2. Gel view comparing Streptococcus varani sp. nov. strain FF10<sup>T</sup> spectra with other members of the Streptococcus genus. The gel view displays the raw spectra of loaded spectrum files arranged in a pseudo-gel like look. The x-axis records the m/z value. The left y-axis displays the running spectrum number originating from subsequent spectra loading. The peak intensity is expressed by a Grey-scale scheme code. The colour bar and the right y-axis indicate the relation between the colour in which a peak is displayed and the peak intensity in arbitrary units. Displayed species are indicated on the left.



FIG. 3. Phylogenetic tree showing the position of Streptococcus varani sp. nov. strain  $FF10<sup>T</sup>$  relative to other type strains within the genus Streptococcus. The strains and their corresponding GenBank accession numbers for 16S rRNA genes are in parenthesis. The tree was constructed by the neighbour joining method with Kimura two-parameter model and 1000 bootstrap replications using the MEGA6 software and rooted by using Lactococcus lactis (EU723831) strain as the out-group.

salicin, D-cellobiose, D-maltose, D-lactose, D-mellibiose, D-saccharose, D-trehalose, inulin, D-melezitose, D-raffinose, amidon, glycogen, xylitol, gentiobiose, D-turanose, D-lyxose, D-tagatose, D-fucose, L-fucose, D-arabitol, L-arabitol, potassium gluconate, potassium 2-ketogluconate and potassium-5-ketogluconate.

Streptococcus varani strain  $FF10<sup>T</sup>$  was susceptible to penicillin, amoxicillin, amoxicillin/clavulanic acid, ticarcillin, ceftriaxone, cefalotin, imipenem, gentamicin, kanamycin, trimethoprim/sulfamethoxazole, erythromycin, doxycycline, metronidazole,



FIG. 4. Gram staining of Streptococcus varani sp. nov. strain FF10<sup>T</sup>.

vancomycin and rifampicin, but resistant to ciprofloxacin, nitrofurantoin and colistin.

The differential phenotypic characteristics with other Streptococcus species (S. minor [10], Streptococcus plurextorum [11], S. ursoris [1] and S. orisratti [3]) are summarized in Table 2.

#### Genome Description

#### Genome sequencing and assembly

The DNA was extracted using the phenol/chloroform method as previously described [12]. Genomic DNA (gDNA) of S. varani FF10 $<sup>T</sup>$  was sequenced on the MiSeq Technology (Illu-</sup> mina Inc., San Diego, CA, USA) using the mate-pair strategy. The gDNA was bar-coded in order to be mixed with 11 other projects with the Nextera Mate Pair sample prep kit (Illumina). The gDNA was quantified by a Qubit assay with the high sensitivity kit (Thermo Fisher Scientific, Waltham, MA, USA) to 62.4 mg/L. The mate-pair library was prepared with 1.5 μg of genomic DNA using the Nextera mate-pair Illumina guide. The gDNA sample was simultaneously fragmented and tagged with a mate-pair junction adapter. The fragmentation pattern was validated on an Agilent 2100 BioAnalyzer (Agilent Technologies Inc., Santa Clara, CA, USA) with a DNA 7500 LabChip. The



FIG. 5. Transmission electron microscopy of Streptococcus varani sp. nov. strain FF10 $^{\mathsf{T}}$ , using a TECNAI G20 (FEI) at an operating voltage of 200 keV. The scale bar represents 500 nm.

DNA fragments ranged from 1.5 kb up to 11 kb with optimal size of 5.8 kb. No size selection was performed and 591 ng of tagmented fragments were circularized. The circularized DNA was mechanically sheared to small fragments, optimally at 697 bp, on the Covaris device S2 in T6 tubes (Covaris, Woburn, MA, USA). The library profile was visualized on a High Sensitivity Bioanalyzer LabChip (Agilent Technologies Inc.) and the final concentration library was measured at 75.52 nmol/ L. The libraries were normalized at 2 nM and pooled. After a denaturation step and dilution at 15 pM, the pool of libraries was loaded onto the reagent cartridge and then onto the instrument along with the flow cell. Automated cluster generation and sequencing were performed in a single 39-h run in a  $2 \times 251$ -bp.

Total information of 5.9 Gb was obtained from a 624 K/mm<sup>2</sup> cluster density with a cluster passing quality control filters of 96.33% (12 040 000 clusters). Within this run, the index representation for S. varani  $FF10<sup>T</sup>$  was determined to 5.26%. The 610 160 paired reads were filtered according to the read qualities. These reads were trimmed then assembled.

#### Genome Annotation and Genome Analysis

Open reading frames (ORFs) were predicted using PRODIGAL ([http://prodigal.ornl.gov/\)](http://prodigal.ornl.gov/) with default parameters. However, the predicted ORFs were excluded if they spanned a sequencing gap region. The predicted bacterial protein sequences were searched against GenBank [13] and Clusters of Orthologous Groups (COG) databases using BLASTP. The tRNAs and rRNAs were predicted using the tRNAScan-SE [14] and RNAmmer [15] tools, respectively. Signal peptides and numbers of transmembrane helices were predicted using

<b>Characteristics</b>	S. varani	S. minor	S. plurextorum	S. ursoris	S. orisratti
Cell diameter (µm)	$0.4 - 0.8$	$\leq$	na	$0.5 - 0.7$	na
Oxygen requirement	Facultative anaerobic	Microaerobic	Facultative anaerobic	Facultative anaerobic	Facultative anaerobic
Gram stain	$+$	$+$	$+$	$+$	$+$
Motility		na	-	na	۰
Endospore formation		na	na	-	na
Production of:					
Alkaline phosphatase	$+$				
Acid phosphatase	$\ddot{}$	$+$	$\ddot{}$	$\ddot{}$	$+$
Catalase		na		$\ddot{}$	
Oxidase		na		na	na
Nitrate reductase		na	na	na	na
Urease		$\qquad \qquad -$			na
$\alpha$ -galactosidase		W		$\ddot{}$	
$\beta$ -galactosidase	$\div$				
β-glucuronidase					
$\alpha$ -glucosidase		na			na
β-glucosidase	$\div$	na		$\ddot{}$	na
Esterase		na		na	na
Esterase lipase	$\ddot{}$	na		na	na
Naphthol-AS-BI-phosphohydrolase		na		$\ddot{}$	$+$
N-acetyl-β-glucosaminidase	$\qquad \qquad -$	na			na
Utilization of:					
Gluconate			<b>DP</b>	na	na
D-ribose				$\ddot{}$	$\overline{\phantom{0}}$
D-xylose					na
D-fructose			$\ddot{}$		$\ddot{}$
D-glucose			$\ddot{}$		$\ddot{}$
D-mannose					
D-lactose			$\ddot{}$		
D-maltose			$\ddot{}$		÷
DL-arabinose					
Habitat	Lizard	Dog, cat and calf	Pigs	<b>Bears</b>	Rats

TABLE 2. Differential characteristics of Streptococcus varani sp. nov. strain FF10<sup>T</sup> with other strains (Streptococcus minor [8], Streptococcus plurextorum [9], Streptococcus ursoris [1] and Streptococcus orisratti [3])

+, positive; –, negative; DP, delayed production; W, weak reaction; na, not available.

TABLE 3. Nucleotide content and gene count levels of the genome

<b>Attribute</b>	Value	% of total <sup>a</sup>
Genome size (bp)	2 460 376	100
DNA coding (bp)	2 170 614	88.2
DNA G+C (bp)	993 991	40.4
Total genes	2516	100
Protein coding genes	2471	98.21
RNA genes	45	1.78
Genes with function prediction	1852	73.60
Genes assigned to COGs	1746	69.39
Genes with signal peptides	118	4.68
Genes with transmembrane helices	565	22.45
Genes associated with resistant genes	0	O
Genes associated with PKS or NRPS	5	0.19
<b>ORFans</b> genes	0	0
<b>CRISPRs</b> repeats	O	O
Genes associated with Pfam-A domains	2264	89.98

<sup>a</sup>The total is based on either the size of the genome in base pairs or the total number of protein coding genes in the annotated genome.

SignalP [16], and TMHMM [17], respectively. Mobile genetic elements were predicted using PHAST [14] and RAST [18]. ORFans were identified if their BLASTP E-value was lower than 1e-03 for alignment length >80 amino acids. If alignment lengths were <80 amino acids, we used an E-value of 1e-05. Such parameter thresholds have already been used in previous work to define ORFans. ARTEMIS [19] and DNA PLOTTER [20] were used for data management and visualization of genomic features, respectively. The MAUVE alignment tool (version 2.3.1) was used for multiple genomic sequence alignment [21].

The mean level of nucleotide sequence similarity at the genome level between S. varani and other bacteria (S. orisratti [\(ARCG01000001.1\)](ncbi-wgs:ARCG01000001.1), S. tigurinus (AORU01000001.1), Streptococcus parasanguinis (CP002843.1), Streptococcus agalactiae [\(AEQQ01000001.1\)](ncbi-wgs:AEQQ01000001.1), S. plurextorum ([AUIO01000001.1\)](ncbi-wgs:AUIO01000001.1), S. minor [\(AQYB01000001.1](ncbi-wgs:AQYB01000001.1)), Streptococcus oralis [\(ADMV01000001.1\)](ncbi-wgs:ADMV01000001.1), Streptococcus pyogenes (AE004092.2), Streptococcus vestibularis ([AEVI01000001.1](ncbi-wgs:AEVI01000001.1)), Streptococcus peroris [\(AEVF01000001.1\)](ncbi-wgs:AEVF01000001.1) and Lactococcus lactis (NC\_002662.1)) was estimated using the Average Genomic Identity of gene Sequences (AGIOS) home-made software [22]. Overall, this software combines the functionality of other software programs: PROTEINORTHO [23] (detects orthologous proteins between genomes compared two by two, then retrieves the corresponding genes) and the Needleman–Wunsch global



FIG. 6. Graphical circular map of the chromosome. From outside to the centre. Genes on the forward strand coloured by COG categories (only genes assigned to COG), genes on the reverse strand coloured by COG categories (only gene assigned to COG), RNA genes (tRNAs green, rRNAs red), GC content, and GC skew.





<sup>a</sup>The total is based on the total number of protein-coding genes in the annotated genome.

alignment algorithm (determines the mean percentage of nucleotide sequence identity among orthologous ORFs).

#### Genome properties

The genome is 2 460 376 bp long with 40.4% GC content (Table 3). It is composed of 16 contigs (Fig. 6). Of the 2516 predicted genes, 2471 were protein-coding genes, and 45 were RNAs (one gene is 5S rRNA, one gene is 16S rRNA, one gene is 23S rRNA, 42 genes are tRNA genes). No genes were identified as ORFans (0.0%). The remaining genes were annotated as hypothetical proteins (293 genes  $> 11.64%$ ). Table 4 distributes the genes into COG functional categories. The genome sequence has been deposited in GenBank under accession number [GCA\\_001375655](ncbi-wgs:GCA_001375655).

#### Comparison with other genomes

The draft genome of S. varani (2.46 Mb) is larger than those of S. agalactiae (2.11 Mb), S. minor (1.93 Mb), S. oralis (1.90 Mb), S. orisratti (2.41 Mb), S. parasanguinis (2.15 Mb), S. peroris (1.63Mb), S. plurextorum (2.10 Mb), S. pyogenes (1.85 Mb), S. tigurinus (2.18 Mb), S. vestibularis (1.84 Mb). The G+C content of S. varani (40.40%) is lower than those of S. minor (41.10%), S. oralis (44.40%), S. parasanguinis (41.70%), S. plurextorum (41.10) but higher than those of S. agalactiae (35.30%), S. orisratti (38.50%), S. peroris (39.10%), S. pyogenes (38.50%), S. tigurinus (40.30%) and S. vestibularis (39.60%).

Similarly, the protein-coding genes of S. varani (2471) are larger than those of S. agalactiae, S. minor, S. oralis, S. orisratti, S. parasanguinis, S. peroris, S. plurextorum, S. pyogenes, S. tigurinus and S. vestibularis (2139, 1903, 1795, 2319, 1982, 1608, 2081, 1776, 2146 and 1812, respectively). The gene content of S. varani (2516) is larger than that of S. agalactiae, S. minor, S. oralis, S. orisratti, S. parasanguinis, S. peroris, S. plurextorum, S. pyogenes, S. tigurinus and S. vestibularis (2164, 1948, 1830, 2348, 2055, 1641, 2116, 1854, 2197 and 1838, respectively). However, the COG category gene distribution is similar in all genomes compared (Fig. 7). Among species with standing in nomenclature, AGIOS values ranged from 81.54 between S. peroris and S. oralis to 65.60 between L. lactis and S. varani (Table 5). To evaluate the genomic similarity among the strains, we determined two parameters, dDDH, which exhibits high correlation with DDH [24], and AGIOS [22], which was designed to be independent of DDH (Table 6).



FIG. 7. Distribution of predicted genes of Streptococcus varani strain FF10<sup>T</sup> and 11 other closely related species into COG categories.

	S. varani	S. minor	S. agalactiae	S. oralis	S. orisratti	S. parasanguinis	S. peroris	S. plurextorum	S. pyogenes	S. tigurinus	S. vestibularis	lactis
Strebtococcus varani	2471	476	1079	1016	1133	1067	975	1044	972	1089	983	952
Streptococcus minor	76.99	1903	997	1000	1089	1045	962	1024	930	1023	989	902
Streptococcus agalactiae	69.62	69.74	2139	982	1082	1030	936	1020	1022	1036	1008	915
Streptococcus oralis	70.91	70.91	69.62	1795	1035	149	1171	994	911	1261	1031	887
Streptococcus orisratti	71.00	70.83	73.03	70.46	2319	1110	1001	1059	976	1092	1066	930
Streptococcus parasanguinis	70.27	70.37	69.26	74.95	69.97	1982	1101	1061	959	1156	1068	915
Streptococcus peroris	70.79	70.69	70.31	81.54	70.73	75.43	1608	950	870	1170	978	845
Streptococcus plurextorum	70.19	70.31	70.95	70.04	72.20	69.49	70.12	2081	951	1031	966	856
Streptococcus pyogenes	70.19	70.00	74.23	70.10	73.04	69.97	70.59	71.26	1176	932	903	840
Streptococcus tigurinus	70.58	70.52	69.82	93.01	70.27	74.91	81.42	69.96	69.92	2146	1051	892
Streptococcus vestibularis	70.47	70.45	71.67	71.82	72.94	72.11	72.00	71.56	72.05	71.69	1812	892
Lactococcus lactis	65.60	65.36	66.90	66.33	66.46	66.03	66.70	65.63	66.68	66.09	66.63	2650

TABLE 5. The numbers of orthologous protein shared between genomes (upper right)<sup>a</sup>

Average percentage similarity of nucleotides corresponding to orthologous protein shared between genomes (lower left) and numbers of proteins per genome (bold).

TABLE 6. Pairwise comparison of Streptococcus varani with eight other species using GGDC, formula 2 (DDH estimates based on

identities /  $HSP$  length)<sup>a</sup>



<sup>a</sup>The confidence intervals indicate the inherent uncertainty in estimating DDH values from intergenomic distances based on models derived from empirical test data sets (which are<br>always limited in size). These results are

#### Conclusion

On the basis of phenotypic, phylogenetic and genomic analyses, we formally propose the creation of Streptococcus varani sp. nov. that contains the strain  $FF10<sup>T</sup>$ . This bacterial strain has been isolated from the oral cavities of a lizard (Varanus niloticus) in Dakar, Senegal.

### Description of Streptococcus varani sp. nov. strain FF10<sup>T</sup>

Streptococcus varani (va.ra'ni. NL. gen. n. varani, from Varanus, the Latin name from the lizard which the type strain was isolated) is a non-motile, non-spore forming, facultative anaerobe, and Gram-positive coccus. Growth is obtained between 25°C and 37°C, with optimal growth at 37°C. The colonies are translucent and yellow with a regular surface,

haemolytic on 5% sheep blood-enriched Columbia agar, and approximately 1 mm in diameter. Cells have mean diameter of 0.6 μm and mean length of 1.3 μm. Streptococcus varani strain  $FF10<sup>T</sup>$  exhibits positive reactions for alkaline phosphatase, esterase, esterase-lipase, lipase, leucine arylamidase, acid phosphatase, β-glucosidase, β-galactosidase, α-mannosidase, α-fucosidase, and for the esculin hydrolysis test. Streptococcus varani strain  $FF10<sup>T</sup>$  is sensitive to penicillin, amoxicillin, amoxicillin/clavulanic acid, ticarcillin, ceftriaxone, cefalotin, imipenem, gentamicin, kanamycin, trimethoprim/sulfamethoxazole, erythromycin, doxycycline, metronidazole, vancomycin and rifampicin, but resistant to ciprofloxacin, nitrofurantoin and colistin. The G+C content of the genome is 40.4%. The 16S rRNA gene sequence and whole-genome shotgun sequence of strain  $FF10<sup>T</sup>$  are deposited in GenBank under accession numbers LN810501 and [GCA\\_001375655,](ncbi-wgs:GCA_001375655) respectively. The type strain FF10<sup>T</sup> (= CSUR P1489 = DSM 100884) was isolated from the oral cavity of a lizard (Varanus niloticus) in Dakar, Senegal.

#### Authors' Contributions

SB, JR, and MB performed the genomic analyses and drafted the manuscript. CIL and CBE performed the phenotypic characterization of the bacterium and drafted the manuscript. OM participated in its design and helped to draft the manuscript. PB helped to draft the manuscript. DR conceived the study and helped to draft the manuscript. PEF and FF conceived the study, participated in its design and coordination and helped to draft the manuscript. All authors read and approved the final manuscript.

#### Transparency Declaration

The authors declare no conflict of interest.

#### Acknowledgements

The authors thank the Xegen Company ([www.xegen.fr\)](http://www.xegen.fr) for automating the genomic annotation process. This study was funded by the Fondation Méditerranée Infection. We also thank Claudia Andrieu for administrative assistance.

#### **References**

- [1] [Shinozaki-Kuwahara N, Takada K, Hirasawa M.](http://refhub.elsevier.com/S2052-2975(16)30003-8/sref1) Streptococcus ursoris sp. [nov., isolated from the oral cavities of bears. Int J Syst Evol Microbiol](http://refhub.elsevier.com/S2052-2975(16)30003-8/sref1) [2015;61:40](http://refhub.elsevier.com/S2052-2975(16)30003-8/sref1)–4.
- [2] [Facklam R. What happened to the streptococci: overview of taxo](http://refhub.elsevier.com/S2052-2975(16)30003-8/sref2)[nomic and nomenclature changes. Clin Microbiol Rev 2002;15:613](http://refhub.elsevier.com/S2052-2975(16)30003-8/sref2)–30.
- [3] [Zhu H, Willcox MDP, Knox KW. A new species of oral](http://refhub.elsevier.com/S2052-2975(16)30003-8/sref3) Streptococcus [isolated from Sprague-Dawley rats,](http://refhub.elsevier.com/S2052-2975(16)30003-8/sref3) Streptococcus orisratti sp. nov. Int J [Syst Evol Microbiol 2000;50:55](http://refhub.elsevier.com/S2052-2975(16)30003-8/sref3)–61.
- [4] [Shinozaki-Kuwahara N, Saito M, Hirasawa M, Takada K.](http://refhub.elsevier.com/S2052-2975(16)30003-8/sref4) Streptococcus oriloxodontae [sp. nov., isolated from the oral cavities of elephants. Int J](http://refhub.elsevier.com/S2052-2975(16)30003-8/sref4) [Syst Evol Microbiol 2014;64:3755](http://refhub.elsevier.com/S2052-2975(16)30003-8/sref4)–9.
- [5] [Hamada S, Slade H. Biology, immunology, and cariogenicity of](http://refhub.elsevier.com/S2052-2975(16)30003-8/sref5) Streptococcus mutans[. Microbiol Rev 1980;44:331](http://refhub.elsevier.com/S2052-2975(16)30003-8/sref5)–84.
- [6] [Welker M, Moore ER. Applications of whole-cell matrix-assisted laser](http://refhub.elsevier.com/S2052-2975(16)30003-8/sref6) desorption/ionization time-of-fl[ight mass spectrometry in systematic](http://refhub.elsevier.com/S2052-2975(16)30003-8/sref6) [microbiology. Syst Appl Microbiol 2011;34:2](http://refhub.elsevier.com/S2052-2975(16)30003-8/sref6)–11.
- [7] [Seng P, Drancourt M, Gouriet F, La Scola B, Fournier PE, Rolain JM,](http://refhub.elsevier.com/S2052-2975(16)30003-8/sref7) [et al. Ongoing revolution in bacteriology: routine identi](http://refhub.elsevier.com/S2052-2975(16)30003-8/sref7)fication of [bacteria by matrix-assisted laser desorption ionization time-of-](http://refhub.elsevier.com/S2052-2975(16)30003-8/sref7)flight [mass spectrometry. Clin In-fect Dis 2009;49:543](http://refhub.elsevier.com/S2052-2975(16)30003-8/sref7)–51.
- [8] [Mourembou G, Rathored J, Lekana-Douki JB, Ndjoyi-Mbiguino A,](http://refhub.elsevier.com/S2052-2975(16)30003-8/sref8) [Fenollar F, Michelle C, et al. Non-contiguous](http://refhub.elsevier.com/S2052-2975(16)30003-8/sref8) finished genome [sequence and description of](http://refhub.elsevier.com/S2052-2975(16)30003-8/sref8) Kallipyga gabonensis sp. nov. New Mi[crobes New Infect 2015;9:15](http://refhub.elsevier.com/S2052-2975(16)30003-8/sref8)–23.
- [9] [Stackebrandt E, Ebers J. Taxonomic parameters revisited: tarnished](http://refhub.elsevier.com/S2052-2975(16)30003-8/sref9) [gold standards. Microbiol Today 2006;33:152](http://refhub.elsevier.com/S2052-2975(16)30003-8/sref9)–5.
- [10] [Vancanneyt M, Devriese L, Haesebrouck F, Baele M, Lefebvre K,](http://refhub.elsevier.com/S2052-2975(16)30003-8/sref10) Snauwaert C, et al. Streptococcus minor [sp. nov., from faecal samples](http://refhub.elsevier.com/S2052-2975(16)30003-8/sref10) and tonsils of domestic animals. Int | Syst Evol Microbiol 2004;54: [449](http://refhub.elsevier.com/S2052-2975(16)30003-8/sref10)–52.
- [11] [Vela A, Casamayor A, Sanchez Del Rey V, Dominguez L, Fernandez](http://refhub.elsevier.com/S2052-2975(16)30003-8/sref11)Garayzabal J. Streptococcus plurextorum [sp. nov., isolated from pigs. Int J](http://refhub.elsevier.com/S2052-2975(16)30003-8/sref11) [Syst Evol Microbiol 2009;59:504](http://refhub.elsevier.com/S2052-2975(16)30003-8/sref11)–8.
- [12] [Lagier JC, Bibi F, Ramasamy D, Azhar EI, Robert C, Yasir M, et al. Non](http://refhub.elsevier.com/S2052-2975(16)30003-8/sref12) contiguous-fi[nished genome sequence and description of](http://refhub.elsevier.com/S2052-2975(16)30003-8/sref12) Clostridium jeddahense [sp. nov.. BASE, EBSCOhost; 2014 \(accessed November 24,](http://refhub.elsevier.com/S2052-2975(16)30003-8/sref12) [2015\).](http://refhub.elsevier.com/S2052-2975(16)30003-8/sref12)
- [13] [Benson DA, Karsch-Mizrachi I, Clark K, Lipman DJ, Ostell J,](http://refhub.elsevier.com/S2052-2975(16)30003-8/sref13) [Sayers EW. GenBank. Nucl Acids Res 2012;40:D48](http://refhub.elsevier.com/S2052-2975(16)30003-8/sref13)–53.
- [14] [Lowe TM, Eddy SR. tRNAscan-SE: a program for improved detection](http://refhub.elsevier.com/S2052-2975(16)30003-8/sref14) [of transfer RNA genes in genomic sequence. Nucl Acids Res 1997;25:](http://refhub.elsevier.com/S2052-2975(16)30003-8/sref14) [955](http://refhub.elsevier.com/S2052-2975(16)30003-8/sref14)–64.
- [15] [Lagesen K, Hallin P, Rodland EA, Staerfeldt HH, Rognes T, Ussery DW.](http://refhub.elsevier.com/S2052-2975(16)30003-8/sref15) [RNAmmer: consistent and rapid annotation of ribosomal RNA genes.](http://refhub.elsevier.com/S2052-2975(16)30003-8/sref15) [Nucl Acids Res 2007;35:3100](http://refhub.elsevier.com/S2052-2975(16)30003-8/sref15)–8.
- [16] [Bendtsen JD, Nielsen H, von Heijne G, Brunak S. Improved prediction](http://refhub.elsevier.com/S2052-2975(16)30003-8/sref16) [of signal peptides: SignalP 3.0. J Mol Biol 2004;340:783](http://refhub.elsevier.com/S2052-2975(16)30003-8/sref16)–95.
- [17] [Krogh A, Larsson B, von Heijne G, Sonnhammer EL. Predicting](http://refhub.elsevier.com/S2052-2975(16)30003-8/sref17) [transmembrane protein topology with a hidden Markov model:](http://refhub.elsevier.com/S2052-2975(16)30003-8/sref17) [application to complete genomes. J Mol Biol 2001;305:567](http://refhub.elsevier.com/S2052-2975(16)30003-8/sref17)–80.
- [18] [Zhou Y, Liang Y, Lynch KH, Dennis JJ, Wishart DS. PHAST: a fast](http://refhub.elsevier.com/S2052-2975(16)30003-8/sref18) [phage search tool. Nucl Acids Res 2011;39:W347](http://refhub.elsevier.com/S2052-2975(16)30003-8/sref18)–52.
- [19] [Rutherford K, Parkhill J, Crook J, Horsnell T, Rice P, Rajandream MA,](http://refhub.elsevier.com/S2052-2975(16)30003-8/sref19) [et al. Artemis: sequence visualization and annotation. Bioinformatics](http://refhub.elsevier.com/S2052-2975(16)30003-8/sref19) [2000;16:944](http://refhub.elsevier.com/S2052-2975(16)30003-8/sref19)–5.
- [20] [Carver T, Thomson N, Bleasby A, Berriman M, Parkhill J. DNAPlotter:](http://refhub.elsevier.com/S2052-2975(16)30003-8/sref20) [circular and linear interactive genome visualization. Bioinformatics](http://refhub.elsevier.com/S2052-2975(16)30003-8/sref20) [2009;25:119](http://refhub.elsevier.com/S2052-2975(16)30003-8/sref20)–20.
- [21] [Darling AC, Mau B, Blattner FR, Perna NT. Mauve: multiple alignment](http://refhub.elsevier.com/S2052-2975(16)30003-8/sref21) [of conserved genomic sequence with rearrangements. Genome Res](http://refhub.elsevier.com/S2052-2975(16)30003-8/sref21) [2004;14:1394](http://refhub.elsevier.com/S2052-2975(16)30003-8/sref21)–403.
- [22] [Ramasamy D, Mishra AK, Lagier JC, Padhmanabhan R, Rossi-](http://refhub.elsevier.com/S2052-2975(16)30003-8/sref22)[Tamisier M, Sentausa E, et al. A polyphasic strategy incorporating](http://refhub.elsevier.com/S2052-2975(16)30003-8/sref22) [genomic data for the taxonomic description of new bacterial species.](http://refhub.elsevier.com/S2052-2975(16)30003-8/sref22) [Int J Syst Evol Microbiol 2014;64:384](http://refhub.elsevier.com/S2052-2975(16)30003-8/sref22)–91.
- [23] [Lechner M, Findeiss S, Steiner L, Marz M, Stadler PF, Prohaska SJ.](http://refhub.elsevier.com/S2052-2975(16)30003-8/sref23) [Proteinortho: detection of \(co\)orthologs in large-scale analysis. BMC](http://refhub.elsevier.com/S2052-2975(16)30003-8/sref23) [Bioinformatics 2011;12:124.](http://refhub.elsevier.com/S2052-2975(16)30003-8/sref23)
- [24] [Auch AF, Von Jan M, Klenk HP, Göker M. Digital DNA-DNA hy](http://refhub.elsevier.com/S2052-2975(16)30003-8/sref24)[bridization for microbial species delineation by means of genome-to](http://refhub.elsevier.com/S2052-2975(16)30003-8/sref24)[genome sequence comparison. Stand Genomic Sci 2010;2:117](http://refhub.elsevier.com/S2052-2975(16)30003-8/sref24)–34.
- [25] [Woese CR, Kandler O, Wheelis ML. Towards a natural system of](http://refhub.elsevier.com/S2052-2975(16)30003-8/sref25) [organisms: proposal for the domains](http://refhub.elsevier.com/S2052-2975(16)30003-8/sref25) Archaea, Bacteria, and Eucarya. [Proc Natl Acad Sci USA 1990;87:4576](http://refhub.elsevier.com/S2052-2975(16)30003-8/sref25)–9.
- [26] [Gibbons NE, Murray RGE. Proposals Concerning the Higher Taxa of](http://refhub.elsevier.com/S2052-2975(16)30003-8/sref26) Bacteria. Int | Syst Bacteriol 1978;28:1-6.
- [27] [Murray RGE. The Higher Taxa, or, a Place for Everything...? In: Holt JG,](http://refhub.elsevier.com/S2052-2975(16)30003-8/sref27) editor. Bergey'[s Manual of Systematic Bacteriology. First EditionVo](http://refhub.elsevier.com/S2052-2975(16)30003-8/sref27)[lume 1. Baltimore, MD: The Williams and Wilkins Co.; 1984. p. 31](http://refhub.elsevier.com/S2052-2975(16)30003-8/sref27)–4.
- [28] [Euzéby J. List of new names and new combinations previously effec](http://refhub.elsevier.com/S2052-2975(16)30003-8/sref28)[tively, but not validly, published. List no. 132. Int J Syst Evol Microbiol](http://refhub.elsevier.com/S2052-2975(16)30003-8/sref28) [2010;60:469](http://refhub.elsevier.com/S2052-2975(16)30003-8/sref28)–72.
- [29] [Ludwig W, Schleifer KH, Whitman WB. Class I.](http://refhub.elsevier.com/S2052-2975(16)30003-8/sref29) Bacilli class nov. In: De [Vos P, Garrity G, Jones D, Krieg NR, Ludwig W, Rainey FA,](http://refhub.elsevier.com/S2052-2975(16)30003-8/sref29) [Schleifer KH, Whitman WB, editors. Bergey](http://refhub.elsevier.com/S2052-2975(16)30003-8/sref29)'s Manual of Systematic [Bacteriology. Second EditionVolume 3. New York: Springer-Verlag;](http://refhub.elsevier.com/S2052-2975(16)30003-8/sref29) [2009. p. 19](http://refhub.elsevier.com/S2052-2975(16)30003-8/sref29)–20.
- [30] [Ludwig W, Schleifer KH, Whitman WB. Order II.](http://refhub.elsevier.com/S2052-2975(16)30003-8/sref30) Lactobacillales ord. [nov. In: De Vos P, Garrity G, Jones D, Krieg NR, Ludwig W, Rainey FA,](http://refhub.elsevier.com/S2052-2975(16)30003-8/sref30) [Schleifer KH, Whitman WB, editors. Bergey](http://refhub.elsevier.com/S2052-2975(16)30003-8/sref30)'s Manual of Systematic

New Microbes and New Infections @ 2016 The Authors. Published by Elsevier Ltd on behalf of European Society of Clinical Microbiology and Infectious Diseases, NMNI, 11, 93-102 This is an open access article under the CC BY-NC-ND license ([http://creativecommons.org/licenses/by-nc-nd/4.0/\)](http://creativecommons.org/licenses/by-nc-nd/4.0/)

[Bacteriology. Second EditionVolume 3. New York: Springer-Verlag;](http://refhub.elsevier.com/S2052-2975(16)30003-8/sref30) [2009. p. 464](http://refhub.elsevier.com/S2052-2975(16)30003-8/sref30).

- [31] [Skerman VBD, McGowan V, Sneath PHA. Approved Lists of Bacterial](http://refhub.elsevier.com/S2052-2975(16)30003-8/sref31) [Names. Int J Syst Bacteriol 1980;30:225](http://refhub.elsevier.com/S2052-2975(16)30003-8/sref31)–420.
- [32] [Deibel RH, Seeley HW. Family II.](http://refhub.elsevier.com/S2052-2975(16)30003-8/sref32) Streptococcaceae. In: Buchanan RE, Gibbons NE, editors. Bergey'[s Manual of Determinative Bacteri-ology.](http://refhub.elsevier.com/S2052-2975(16)30003-8/sref32) [Eighth Edition. Baltimore, MD: The Williams and Wilkins Co.; 1974. p.](http://refhub.elsevier.com/S2052-2975(16)30003-8/sref32) 490–[515.](http://refhub.elsevier.com/S2052-2975(16)30003-8/sref32)
- [33] Rosenbach FJ. In: Bergmann JF (ed), Microorganismen bei den Wund-Infections-Krankheiten des Menschen., Wiesbaden; 1884, p. 1–122.
- [34] [Deibel RH, Seeley HW. Genus I.](http://refhub.elsevier.com/S2052-2975(16)30003-8/sref34) Streptococcus Rosenbach 1884, 22. In: [Buchanan RE, Gibbons NE, editors. Bergey](http://refhub.elsevier.com/S2052-2975(16)30003-8/sref34)'s Manual of Determinative [Bacteriology. Eighth Edition. Baltimore: The Williams and Wilkins Co.;](http://refhub.elsevier.com/S2052-2975(16)30003-8/sref34) [1974. p. 490](http://refhub.elsevier.com/S2052-2975(16)30003-8/sref34)–509.