



HAL
open science

Error estimates for phaseless inverse scattering in the Born approximation at high energies

Alexey Agaltsov, Roman Novikov

► **To cite this version:**

Alexey Agaltsov, Roman Novikov. Error estimates for phaseless inverse scattering in the Born approximation at high energies. 2016. hal-01303885v1

HAL Id: hal-01303885

<https://hal.science/hal-01303885v1>

Preprint submitted on 18 Apr 2016 (v1), last revised 8 Dec 2016 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Error estimates for phaseless inverse scattering in the Born approximation at high energies

A. D. Agaltsov¹ and R. G. Novikov²

April 18, 2016

Abstract. We study explicit formulas for phaseless inverse scattering in the Born approximation at high energies for the Schrödinger equation with compactly supported potential in dimension $d \geq 2$. We obtain error estimates for these formulas in the configuration space.

1 Introduction

We consider the time-independent Schrödinger equation

$$-\Delta\psi + v(x)\psi = E\psi, \quad x \in \mathbb{R}^d, \quad d \geq 2, \quad E > 0, \quad (1)$$

where

$$v \in L^\infty(\mathbb{R}^d), \quad \text{supp } v \subset D, \quad (2)$$

where D is some fixed open bounded domain in \mathbb{R}^d .

In quantum mechanics equation (1) describes an elementary particle interacting with a macroscopic object contained in D at fixed energy E .

For equation (1) we consider the classical scattering solutions $\psi^+ = \psi^+(x, k)$, $x \in \mathbb{R}^d$, $k \in \mathbb{R}^d$, $k^2 = E$. These solutions ψ^+ can be specified by the following asymptotics as $|x| \rightarrow \infty$:

$$\begin{aligned} \psi^+(x, k) &= e^{ikx} + c(d, |k|) \frac{e^{i|k||x|}}{|x|^{(d-1)/2}} f(k, |k| \frac{x}{|x|}) + O(|x|^{-(d+1)/2}), \\ x \in \mathbb{R}^d, \quad k \in \mathbb{R}^d, \quad k^2 = E, \quad c(d, |k|) &= -\pi i (-2\pi i)^{(d-1)/2} |k|^{(d-3)/2}, \end{aligned} \quad (3)$$

for some a priori unknown f . The function f arising in (3) is defined on

$$\mathcal{M}_E = \{(k, l) \in \mathbb{R}^d \times \mathbb{R}^d : k^2 = l^2 = E\}, \quad (4)$$

and is known as the classical scattering amplitude for equation (1).

In quantum mechanics $|f(k, l)|^2$ describes the probability density of scattering of particle with initial impulse k into direction $l/|l| \neq k/|k|$, and is known as differential scattering cross section for equation (1); see, e.g., [11, Chapter 1, Section 6].

¹Centre de Mathématiques Appliquées, Ecole Polytechnique, 91128 Palaiseau, France; and Moscow State University, 119991 Moscow, Russia; email: agaltsov@cmap.polytechnique.fr

²Centre de Mathématiques Appliquées, Ecole Polytechnique, 91128 Palaiseau, France; IEPT RAS, 117997 Moscow, Russia; and Moscow Institute of Physics and Technology, 141700 Dolgoprudny, Russia; email: novikov@cmap.polytechnique.fr

The problem of finding ψ^+ and f from v is known as the direct scattering problem for equation (1). For solving this problem, one can use, in particular, the Lippmann-Schwinger integral equation for ψ^+ and an explicit integral formula for f , see, e.g., [5, 10, 26].

In turn, the problem of finding v from f is known as the inverse scattering problem (with phase information) and the problem of finding v from $|f|^2$ is known as the phaseless inverse scattering problem for equation (1).

There is a vast literature on the former inverse scattering problem with phase information; see [3, 4, 6, 7, 8, 9, 12, 13, 14, 15, 19, 20, 21, 22, 25] and references therein. In particular, it is well known that the scattering amplitude f uniquely determines v via the Born approximation formulas at high energies:

$$\widehat{v}(k-l) = f(k, l) + O(E^{-\frac{1}{2}}), \quad E \rightarrow +\infty, \quad (k, l) \in \mathcal{M}_E, \quad (5)$$

$$\widehat{v}(p) = (2\pi)^{-d} \int_{\mathbb{R}^d} e^{ipx} v(x) dx, \quad p \in \mathbb{R}^d, \quad (6)$$

and the inverse Fourier transform; see, e.g., [9, 25].

On the other hand, the literature for the phaseless case is much more limited; see [7, 26] and references therein for the case of the aforementioned phaseless inverse problem and see [16, 17, 18, 23, 24, 26] and references therein for the case of some similar inverse problems without phase information. In addition, it is well known that the phaseless scattering data $|f|^2$ does not determine v uniquely, even if $|f|^2$ is given completely for all positive energies. In particular, it is known that

$$\begin{aligned} f_y(k, l) &= e^{i(k-l)y} f(k, l), \\ |f_y(k, l)|^2 &= |f(k, l)|^2, \quad k, l \in \mathbb{R}^d, \quad k^2 = l^2 > 0, \end{aligned} \quad (7)$$

where f is the scattering amplitude for v and f_y is the scattering amplitude for $v_y = v(\cdot - y)$, where $y \in \mathbb{R}^d$; see [26] and references therein.

In the present work, in view of the aforementioned non-uniqueness for the problem of finding v from $|f|^2$, we consider the modified phaseless inverse scattering problem formulated below as Problem 1. Let

$$S = \{|f|^2, |f_1|^2, \dots, |f_m|^2\}, \quad (8)$$

where f is the scattering amplitude for v and f_1, \dots, f_m are the scattering amplitudes for v_1, \dots, v_m , where

$$v_j = v + w_j, \quad j = 1, \dots, m, \quad (9)$$

where w_1, \dots, w_m are additional a priori known background scatterers such that

$$\begin{aligned} w_j &\in L^\infty(\mathbb{R}^d), \quad \text{supp } w_j \subset \Omega_j, \\ \Omega_j &\text{ is an open bounded domain in } \mathbb{R}^d, \quad \Omega_j \cap D = \emptyset, \\ w_j &\neq 0, \quad w_{j_1} \neq w_{j_2} \text{ if } j_1 \neq j_2 \text{ (in } L^\infty(\mathbb{R}^d)), \end{aligned} \quad (10)$$

where $j, j_1, j_2 \in \{1, \dots, m\}$. Thus, S consists of the phaseless scattering data $|f|^2, |f_1|^2, \dots, |f_m|^2$ measured sequentially, first, for the unknown scatterer v and then for v in the presence of known scatterer w_j disjoint from v for $j = 1, \dots, m$.

Actually, in the present work we continue studies of [26] on the following inverse scattering problem for equation (1):

Problem 1. *Reconstruct potential v from the phaseless scattering data S for some appropriate background scatterers w_1, \dots, w_m .*

Studies of Problem 1 in dimension $d \geq 2$ were started in [26]. In dimension $d = 1$ for $m = 1$ studies of Problem 1 were started earlier in [2], where phaseless scattering data was considered for all $E > 0$.

Actually, the key result of [26] consists in a proper extension of formula (5) for the Fourier transform \widehat{v} of v to the phaseless case of Problem 1; see Section 2.

In the present work we proceed from the aforementioned result of [26] and study related approximate reconstruction of v in the configuration space. In this connection our results consist in obtaining related error estimates in the configuration space at high energies E ; see Section 4.

In addition, results of the present work are necessary for extending the iterative algorithm of [25] to the phaseless case of Problem 1. The latter extension will be given in [1].

2 Extension of formula (5) to the phaseless case

Actually, the key result of [26] consists in the following formulas for solving Problem 1 in dimension $d \geq 2$ for $m = 2$ at high energies E :

$$\begin{pmatrix} \operatorname{Re} \widehat{v} \\ \operatorname{Im} \widehat{v} \end{pmatrix} = \frac{1}{2} \begin{pmatrix} \operatorname{Re} \widehat{w}_1 & \operatorname{Im} \widehat{w}_1 \\ \operatorname{Re} \widehat{w}_2 & \operatorname{Im} \widehat{w}_2 \end{pmatrix}^{-1} \begin{pmatrix} |\widehat{v}_1|^2 - |\widehat{v}|^2 - |\widehat{w}_1|^2 \\ |\widehat{v}_2|^2 - |\widehat{v}|^2 - |\widehat{w}_2|^2 \end{pmatrix}, \quad (11)$$

$$\begin{aligned} |\widehat{v}_j(p)|^2 &= |f_j(k, l)|^2 + O(E^{-\frac{1}{2}}), \quad E \rightarrow +\infty, \\ p \in \mathbb{R}^d, \quad (k, l) \in \mathcal{M}_E, \quad k - l = p, \quad j = 0, 1, 2, \end{aligned} \quad (12)$$

where:

- $v_0 = v$, v_j is defined by (9), $j = 1, 2$, and $f_0 = f$, f_1, f_2 are the scattering amplitudes for v_0, v_1, v_2 , respectively;
- $\widehat{v} = \widehat{v}(p)$, $\widehat{v}_j = \widehat{v}_j(p)$, $\widehat{w}_j = \widehat{w}_j(p)$, $p \in \mathbb{R}^d$, are the Fourier transforms of v, v_j, w_j (defined as in (6));
- formula (11) is considered for all $p \in \mathbb{R}^d$ such that the determinant

$$\zeta_{\widehat{w}_1, \widehat{w}_2}(p) \stackrel{def}{=} \operatorname{Re} \widehat{w}_1(p) \operatorname{Im} \widehat{w}_2(p) - \operatorname{Im} \widehat{w}_1(p) \operatorname{Re} \widehat{w}_2(p) \neq 0. \quad (13)$$

The point is that using formulas (12) for $d \geq 2$ with

$$\begin{aligned} k &= k_E(p) = \frac{p}{2} + \left(E - \frac{p^2}{4}\right)^{1/2} \gamma(p), \\ l &= l_E(p) = -\frac{p}{2} + \left(E - \frac{p^2}{4}\right)^{1/2} \gamma(p), \\ |\gamma(p)| &= 1, \quad \gamma(p)p = 0, \end{aligned} \tag{14}$$

where $p \in \mathbb{R}^d$, $|p| \leq 2\sqrt{E}$, one can reconstruct $|\widehat{v}|^2$, $|\widehat{v}_1|^2$, $|\widehat{v}_2|^2$ from S at high energies for any $p \in \mathbb{R}^d$. And then using formula (11) one can reconstruct \widehat{v} completely, provided that condition (13) is fulfilled for almost all $p \in \mathbb{R}^d$.

Remark 1. Formulas (12) can be precised as formula (2.15) of [26]:

$$\begin{aligned} \left| |\widehat{v}_j(p)|^2 - |f_j(k, l)|^2 \right| &\leq c(D_j) N_j^3 E^{-\frac{1}{2}}, \\ p &= k - l, \quad (k, l) \in \mathcal{M}_E, \quad E^{\frac{1}{2}} \geq \rho(D_j, N_j), \quad j = 0, 1, 2, \end{aligned} \tag{15}$$

where $\|v_j\|_{L^\infty(D_j)} \leq N_j$, $j = 0, 1, 2$, and $D_0 = D$, $D_j = D \cup \Omega_j$, $j = 1, 2$, and formulas for the constants c, ρ are given in [26].

In addition, from the experimental point of view it seems to be, in particular, convenient to consider Problem 1 with $m = 2$ for the case when w_2 is just a translation of w_1 :

$$w_2(x) = w_1(x - y), \quad x \in \mathbb{R}^d, \quad y \in \mathbb{R}^d. \tag{16}$$

In this case

$$\widehat{w}_2(p) = e^{ipy} \widehat{w}_1(p), \quad \zeta_{\widehat{w}_1, \widehat{w}_2}(p) = \sin(py) |\widehat{w}_1(p)|^2, \quad p \in \mathbb{R}^d. \tag{17}$$

On the level of analysis, the principal complication of (11), (12) in comparison with (5) consists in possible zeros of the determinant $\zeta_{\widehat{w}_1, \widehat{w}_2}$ of (13). For some simplest cases, we study these zeros in the next section.

3 Zeros of the determinant $\zeta_{\widehat{w}_1, \widehat{w}_2}$

Let

$$\begin{aligned} Z_{\widehat{w}_1, \widehat{w}_2} &= \{p \in \mathbb{R}^d : \zeta_{\widehat{w}_1, \widehat{w}_2}(p) = 0\}, \\ Z_{\widehat{w}_j} &= \{p \in \mathbb{R}^d : \widehat{w}_j(p) = 0\}, \quad j = 1, 2, \end{aligned} \tag{18}$$

where ζ is defined by (13). From (13), (18) it follows that

$$Z_{\widehat{w}_1} \cup Z_{\widehat{w}_2} \subseteq Z_{\widehat{w}_1, \widehat{w}_2}. \tag{19}$$

In view of (19), in order to construct examples of w_1, w_2 such that the set $Z_{\widehat{w}_1, \widehat{w}_2}$ is as simple as possible, we use the following lemma:

Lemma 1. *Let*

$$w(x) = |x|^\nu K_\nu(|x|) \int_{\mathbb{R}^d} q(x-y)q(y) dy, \quad x \in \mathbb{R}^d, \nu > 0, \quad (20)$$

$$K_\nu(s) = \frac{\Gamma(\frac{1}{2} + \nu)}{\sqrt{\pi}} \left(\frac{2}{s}\right)^\nu \int_0^\infty \frac{\cos(st) dt}{(1+t^2)^{\frac{1}{2}+\nu}}, \quad s > 0, \quad (21)$$

$$\begin{aligned} q &\in L^\infty(\mathbb{R}^d), \quad q = \bar{q}, \quad q \neq 0 \text{ in } L^\infty(\mathbb{R}^d), \\ q(x) &= 0 \text{ if } |x| > r, \quad q(x) = q(-x), \quad x \in \mathbb{R}^d. \end{aligned} \quad (22)$$

Then

$$\begin{aligned} w &\in C(\mathbb{R}^d), \quad w = \bar{w}, \quad w(x) = 0 \text{ if } |x| > 2r, \quad x \in \mathbb{R}^d, \\ \widehat{w}(p) &= \overline{\widehat{w}(p)} \geq c_1(1+|p|)^{-\beta}, \quad p \in \mathbb{R}^d, \end{aligned} \quad (23)$$

for $\beta = d + 2\nu$ and some positive constant $c_1 = c_1(q, \nu)$, where \widehat{w} is the Fourier transform of w . In addition, if $q \geq 0$, then $w \geq 0$.

We recall that K_ν defined by (21) is the modified Bessel function of the second kind and order ν . In addition, Γ denotes the gamma function.

Lemma 1 is proved in Section 8.

As a corollary of Lemma 1, functions

$$w_j(x) = w(x - T_j), \quad x \in \mathbb{R}^d, \quad T_j \in \mathbb{R}^d, \quad (24)$$

where w is constructed in Lemma 1, give us examples of w_j satisfying (10) for fixed D , Ω_j and for appropriate radius r of Lemma 1 and translations T_j of (24), and such that

$$\begin{aligned} Z_{\widehat{w}_j} &= \emptyset, \\ |\widehat{w}_j(p)| &= \widehat{w}(p) \geq c_1(1+|p|)^{-\beta}, \quad p \in \mathbb{R}^d, \end{aligned} \quad (25)$$

where c_1, β are the same as in (23). In addition,

$$\begin{aligned} \zeta_{\widehat{w}_1, \widehat{w}_2}(p) &= \sin(py) |\widehat{w}(p)|^2, \quad y = T_2 - T_1 \neq 0, \quad p \in \mathbb{R}^d, \\ Z_{\widehat{w}_1, \widehat{w}_2} &= \{p \in \mathbb{R}^d : \sin(py) = 0\} = \{p \in \mathbb{R}^d : py \in \pi\mathbb{Z}\}, \end{aligned} \quad (26)$$

for w_1, w_2 of (24).

As another corollary of Lemma 1, we have that

$$\begin{aligned} \zeta_{\widehat{w}_1, \widehat{w}_2}(p) &= |\widehat{w}(p)|^2 \geq c_1^2(1+|p|)^{-2\beta}, \quad p \in \mathbb{R}^d, \\ Z_{\widehat{w}_1, \widehat{w}_2} &= \emptyset, \end{aligned} \quad (27)$$

if w_1 is defined as in (24) and $w_2 = iw_1$.

Note that complex-valued v and w_j naturally arise if we interpret equation (1) as a time-harmonic Helmholtz equation of acoustics or electrodynamics.

Finally, note that

$$\begin{aligned} Z_{\widehat{w}_1, \dots, \widehat{w}_{d+1}} &= \frac{\pi}{s} \mathbb{Z}^d, \quad \text{where} \\ Z_{\widehat{w}_1, \dots, \widehat{w}_{d+1}} &= Z_{\widehat{w}_1, \widehat{w}_2} \cap Z_{\widehat{w}_1, \widehat{w}_3} \cap \dots \cap Z_{\widehat{w}_1, \widehat{w}_{d+1}}, \end{aligned} \quad (28)$$

if w_1 is defined as in (24), and

$$w_2(x) = w_1(x - se_1), \dots, w_{d+1}(x) = w_1(x - se_d), \quad (29)$$

where (e_1, \dots, e_d) is the standard basis of \mathbb{R}^d and $s > 0$.

Thus, in principle, for Problem 1 with background scatterers w_1, \dots, w_{d+1} as in (29), for each $p \in \mathbb{R}^d \setminus \frac{\pi}{s}\mathbb{Z}^d$ formulas (11), (12) can be used with appropriate w_j in place of w_2 , where $j = 2, \dots, d+1$.

4 Error estimates in the configuration space

We recall that for inverse scattering with phase information the scattering amplitude f on \mathcal{M}_E processed by (5) and the inverse Fourier transform yield the approximate reconstruction

$$u(\cdot, E) = v + O(E^{-\alpha}) \quad \text{in } L^\infty(\mathbb{R}^d) \text{ as } E \rightarrow +\infty, \quad \alpha = \frac{n-d}{2n}, \quad (30)$$

if $v \in W^{n,1}(\mathbb{R}^d)$, $n > d$ (in addition to the initial assumption (2)), where $W^{n,1}(\mathbb{R}^d)$ denotes the standard Sobolev space of n -times differentiable functions in $L^1(\mathbb{R}^d)$:

$$\begin{aligned} W^{n,1}(\mathbb{R}^d) &= \{u \in L^1(\mathbb{R}^d) : \|u\|_{n,1} < \infty\}, \\ \|u\|_{n,1} &= \max_{|J| \leq n} \left\| \frac{\partial^{|J|} u}{\partial x^J} \right\|_{L^1(\mathbb{R}^d)}, \quad n \in \mathbb{N} \cup \{0\}. \end{aligned} \quad (31)$$

More precisely, the approximation $u(\cdot, E)$ in (30) is defined by

$$\begin{aligned} u(x, E) &= \int_{\mathcal{B}_{r(E)}} e^{-ipx} f(k_E(p), l_E(p)) dp, \quad x \in \mathbb{R}^d, \\ r(E) &= 2\tau E^{\frac{\alpha}{n-d}} \quad \text{for some fixed } \tau \in (0, 1], \end{aligned} \quad (32)$$

where

$$\mathcal{B}_r = \{p \in \mathbb{R}^d : |p| \leq r\}, \quad (33)$$

α is defined in (30), and $k_E(p), l_E(p)$ are defined as in (14) with some piecewise continuous vector-function γ on \mathbb{R}^d ; see, e.g., [25].

Analogues of $u(\cdot, E)$ for the phaseless case are given below in this section. In particular, related formulas depend on the zeros of determinant $\zeta_{\hat{w}_1, \hat{w}_2}$ of (13).

We consider

$$\begin{aligned} U_{\widehat{w}_1, \widehat{w}_2} &= \operatorname{Re} U_{\widehat{w}_1, \widehat{w}_2} + i \operatorname{Im} U_{\widehat{w}_1, \widehat{w}_2}, \\ \begin{pmatrix} \operatorname{Re} U_{\widehat{w}_1, \widehat{w}_2} \\ \operatorname{Im} U_{\widehat{w}_1, \widehat{w}_2} \end{pmatrix} &= \frac{1}{2} M_{\widehat{w}_1, \widehat{w}_2}^{-1}(p) b_{\widehat{w}_1, \widehat{w}_2}(p, E), \end{aligned} \quad (34)$$

$$M_{\widehat{w}_1, \widehat{w}_2}(p) = \begin{pmatrix} \operatorname{Re} \widehat{w}_1(p) & \operatorname{Im} \widehat{w}_1(p) \\ \operatorname{Re} \widehat{w}_2(p) & \operatorname{Im} \widehat{w}_2(p) \end{pmatrix}, \quad (35)$$

$$M_{\widehat{w}_1, \widehat{w}_2}^{-1}(p) = \frac{1}{\zeta_{\widehat{w}_1, \widehat{w}_2}(p)} \begin{pmatrix} \operatorname{Im} \widehat{w}_2(p) & -\operatorname{Im} \widehat{w}_1(p) \\ -\operatorname{Re} \widehat{w}_2(p) & \operatorname{Re} \widehat{w}_1(p) \end{pmatrix}, \quad (36)$$

$$b_{\widehat{w}_1, \widehat{w}_2}(p, E) = \begin{pmatrix} |f_1(p, E)|^2 - |f(p, E)|^2 - |\widehat{w}_1(p)|^2 \\ |f_2(p, E)|^2 - |f(p, E)|^2 - |\widehat{w}_2(p)|^2 \end{pmatrix}, \quad (37)$$

$$f(p, E) = f(k_E(p), l_E(p)), \quad f_j(p, E) = f_j(k_E(p), l_E(p)), \quad j = 1, 2, \quad (38)$$

where $\widehat{w}_1, \widehat{w}_2, f, f_1, f_2$ are the same as in (11), (12), $\zeta_{\widehat{w}_1, \widehat{w}_2}$ is defined by (13), $k_E(p), l_E(p)$ are the same as in (14), (32), and $p \in \mathcal{B}_{2\sqrt{E}}, d \geq 2$.

For Problem 1 for $d \geq 2, m = 2$, and for the case when $\zeta_{\widehat{w}_1, \widehat{w}_2}$ has no zeros (the case of (27) in Section 3) we have the following result:

Theorem 1. *Let v satisfy (2) and $v \in W^{n,1}(\mathbb{R}^d)$ for some $n > d$. Let w_1, w_2 be the same as in (27). Let*

$$\begin{aligned} u(x, E) &= \int_{\mathcal{B}_{r_1(E)}} e^{-ipx} U_{\widehat{w}_1, \widehat{w}_2}(p, E) dp, \quad x \in \mathbb{R}^d, \\ r_1(E) &= 2\tau E^{\frac{\alpha_1}{n-d}}, \quad \alpha_1 = \frac{n-d}{2(n+\beta)}, \quad \text{for some fixed } \tau \in (0, 1], \end{aligned} \quad (39)$$

where $U_{\widehat{w}_1, \widehat{w}_2}$ is defined by (34), \mathcal{B}_r is defined by (33), β is the number of (23), (27). Then

$$u(\cdot, E) = v + O(E^{-\alpha_1}) \quad \text{in } L^\infty(\mathbb{R}^d), \quad E \rightarrow +\infty. \quad (40)$$

Theorem 1 is proved in Section 5.

Next, we set

$$Z_{\widehat{w}_1, \widehat{w}_2}^\varepsilon = \{p \in \mathbb{R}^d : py \in (-\varepsilon, \varepsilon) + \pi\mathbb{Z}\}, \quad y \in \mathbb{R}^d \setminus 0, \quad 0 < \varepsilon < 1, \quad (41)$$

where $\widehat{w}_1, \widehat{w}_2$ and y are the same as in (24)–(26). One can see that $Z_{\widehat{w}_1, \widehat{w}_2}^\varepsilon$ is the open $\frac{\varepsilon}{|y|}$ -neighborhood of $Z_{\widehat{w}_1, \widehat{w}_2}$ defined in (26).

Note that

$$\begin{aligned} &\text{for any } p \in Z_{\widehat{w}_1, \widehat{w}_2}^\varepsilon \text{ there exists} \\ &\text{the unique } z(p) \in \mathbb{Z} \text{ such that } |py - \pi z(p)| < \varepsilon. \end{aligned} \quad (42)$$

In addition to $U_{\widehat{w}_1, \widehat{w}_2}$ of (34), we define

$$\begin{aligned} U_{\widehat{w}_1, \widehat{w}_2}^\varepsilon(p, E) &= \frac{1}{2} (U_{\widehat{w}_1, \widehat{w}_2}(p_-^\varepsilon, E) + U_{\widehat{w}_1, \widehat{w}_2}(p_+^\varepsilon, E)), \\ p_\pm^\varepsilon &= p_\perp + \pi z(p) \frac{y}{|y|^2} \pm \varepsilon \frac{y}{|y|^2}, \quad p_\perp = p - (py) \frac{y}{|y|^2}, \quad p \in \mathcal{B}_{2\sqrt{E}} \cap Z_{\widehat{w}_1, \widehat{w}_2}^\varepsilon, \end{aligned} \quad (43)$$

where $z(p)$ is the integer number of (42).

For Problem 1 for $d \geq 2$, $m = 2$, and for the case when $\zeta_{\widehat{w}_1, \widehat{w}_2}$ has zeros on hyperplanes (the case of (26) in Section 3) we have the following result:

Theorem 2. *Let v satisfy (2) and $v \in W^{n,1}(\mathbb{R}^d)$ for some $n > d$. Let w_1, w_2 be the same as in (24)–(26). Let*

$$\begin{aligned}
u(x, E) &= u_1(x, E) + u_2(x, E), \quad x \in \mathbb{R}^d, \\
u_1(x, E) &= \int_{\mathcal{B}_{r_2(E)} \setminus Z_{\widehat{w}_1, \widehat{w}_2}^{\varepsilon_2(E)}} e^{-ipx} U_{\widehat{w}_1, \widehat{w}_2}(p, E) dp, \\
u_2(x, E) &= \int_{\mathcal{B}_{r_2(E)} \cap Z_{\widehat{w}_1, \widehat{w}_2}^{\varepsilon_2(E)}} e^{-ipx} U_{\widehat{w}_1, \widehat{w}_2}^\varepsilon(p, E) dp, \\
r_2(E) &= 2\tau E^{\frac{\alpha_2}{n-d}}, \quad \varepsilon_2(E) = E^{-\frac{\alpha_2}{2}}, \\
\alpha_2 &= \frac{n-d}{2(n+\beta+\frac{n-d}{2})}, \quad \text{for some fixed } \tau \in (0, 1],
\end{aligned} \tag{44}$$

where $U_{\widehat{w}_1, \widehat{w}_2}$ and $U_{\widehat{w}_1, \widehat{w}_2}^\varepsilon$ are defined by (34), (43), \mathcal{B}_r and $Z_{\widehat{w}_1, \widehat{w}_2}^\varepsilon$ are defined by (33), (41), and β is the number of (23). Then

$$u(\cdot, E) = v + O(E^{-\alpha_2}) \quad \text{in } L^\infty(\mathbb{R}^d), \quad E \rightarrow +\infty. \tag{45}$$

Theorem 2 is proved in Section 6.

Next, we set

$$\begin{aligned}
Z_{\widehat{w}_1, \dots, \widehat{w}_{d+1}}^\varepsilon &= \mathcal{B}'_{\varepsilon/s} + \frac{\pi}{s} \mathbb{Z}^d, \quad 0 < \varepsilon < 1, \\
\mathcal{B}'_r &= \mathcal{B}_r \setminus \partial \mathcal{B}_r, \quad r > 0,
\end{aligned} \tag{46}$$

where $\widehat{w}_1, \dots, \widehat{w}_{d+1}$ are the same as in (28), (29), and \mathcal{B}_r is defined by (33). One can see that $Z_{\widehat{w}_1, \dots, \widehat{w}_{d+1}}^\varepsilon$ is the open $\frac{\varepsilon}{s}$ -neighborhood of $Z_{\widehat{w}_1, \dots, \widehat{w}_{d+1}}$ defined in (28).

Note that

$$\begin{aligned}
&\text{for any } p \in Z_{\widehat{w}_1, \dots, \widehat{w}_{d+1}}^\varepsilon \text{ there exists} \\
&\text{the unique } z(p) \in \mathbb{Z}^d \text{ such that } |sp - \pi z(p)| < \varepsilon.
\end{aligned} \tag{47}$$

In addition, we consider i' such that

$$\begin{aligned}
i' &= i'(p, s), \quad p = (p_1, \dots, p_d) \in \mathbb{R}^d \setminus \frac{\pi}{s} \mathbb{Z}^d, \quad s > 0, \\
i' &\text{ take values in } \{2, \dots, d+1\}, \\
|\sin(sp_{i'} - 1)| &\geq |\sin(sp_{i-1})| \quad \text{for all } i \in \{2, \dots, d+1\}.
\end{aligned} \tag{48}$$

Let

$$U_{\widehat{w}_1, \dots, \widehat{w}_{d+1}}(p, E) = U_{\widehat{w}_1, \widehat{w}_{i'}}(p, E), \quad p \in \mathbb{R}^d \setminus \frac{\pi}{s}Z^d, \quad (49)$$

$$U_{\widehat{w}_1, \dots, \widehat{w}_{d+1}}^\varepsilon(p, E) = \frac{1}{|\mathbb{S}^{d-1}|} \int_{\mathbb{S}^{d-1}} U_{\widehat{w}_1, \dots, \widehat{w}_{d+1}}\left(\frac{\varepsilon}{s}\vartheta + \frac{\pi}{s}z(p), E\right) d\vartheta, \quad p \in Z_{\widehat{w}_1, \dots, \widehat{w}_{d+1}}^\varepsilon, \quad (50)$$

$$\mathbb{S}^{d-1} = \{p \in \mathbb{R}^d : |p| = 1\}, \quad (51)$$

where $|\mathbb{S}^{d-1}|$ denotes the standard Euclidean volume of \mathbb{S}^{d-1} , $U_{\widehat{w}_1, \widehat{w}_{i'}}$ is defined in a similar way with $U_{\widehat{w}_1, \widehat{w}_2}$ of (34), $\widehat{w}_1, \dots, \widehat{w}_{d+1}$ are the same as in (28), (29), and i' is the same as in (48).

For Problem 1 for $d \geq 2$, $m = d + 1$, we also have the following result:

Theorem 3. *Let v satisfy (2) and $v \in W^{n,1}(\mathbb{R}^d)$ for some $n > d$. Let w_1, \dots, w_{d+1} be the same as in (29). Let*

$$\begin{aligned} u(x, E) &= u_1(x, E) + u_2(x, E), \quad x \in \mathbb{R}^d, \\ u_1(x, E) &= \int_{\mathcal{B}_{r_3(E)} \setminus Z_{\widehat{w}_1, \dots, \widehat{w}_{d+1}}^{\varepsilon_3(E)}} e^{-ipx} U_{\widehat{w}_1, \dots, \widehat{w}_{d+1}}(p, E) dp, \\ u_2(x, E) &= \int_{\mathcal{B}_{r_3(E)} \cap Z_{\widehat{w}_1, \dots, \widehat{w}_{d+1}}^{\varepsilon_3(E)}} e^{-ipx} U_{\widehat{w}_1, \dots, \widehat{w}_{d+1}}^\varepsilon(p, E) dp, \\ r_3(E) &= 2\tau E^{\frac{\alpha_3}{n-d}}, \quad \varepsilon_2(E) = E^{-\frac{\alpha_3}{d+1}}, \\ \alpha_3 &= \frac{n-d}{2(n+\beta+\frac{n-d}{d+1})}, \quad \text{for some fixed } \tau \in (0, 1], \end{aligned} \quad (52)$$

where $U_{\widehat{w}_1, \dots, \widehat{w}_{d+1}}$ and $U_{\widehat{w}_1, \dots, \widehat{w}_{d+1}}^\varepsilon$ are defined by (49), (50), \mathcal{B}_r and $Z_{\widehat{w}_1, \dots, \widehat{w}_{d+1}}^\varepsilon$ are defined by (33), (46), and β is the number of (23). Then

$$u(\cdot, E) = v + O(E^{-\alpha_3}) \quad \text{in } L^\infty(\mathbb{R}^d), \quad E \rightarrow +\infty. \quad (53)$$

Theorem 3 is proved in Section 7.

5 Proof of Theorem 1

Proposition 1. *Let v satisfy (2) and w_1, w_2 be the same as in (27), $d \geq 2$. Then:*

$$\begin{aligned} |\widehat{v}(p) - U_{\widehat{w}_1, \widehat{w}_2}(p, E)| &\leq c_2 |\widehat{w}(p)|^{-1} E^{-\frac{1}{2}} \quad \text{for } p \in \mathcal{B}_{2\sqrt{E}}, \quad E \geq \rho_1, \\ c_2 &= 2c(D_0)N_0^3 + 2c(D_1)N_1^3, \\ \rho_1 &= \max_{j=0,1} \rho(D_j, N_j), \end{aligned} \quad (54)$$

where w is the function of (23), (24), and $c, \rho, N_j, D_j, j = 0, 1, 2$, are the same as in estimates (15).

Proposition 1 follows from formulas (11), (14), estimates (15), definitions (34)–(38), and the properties that

$$\Omega_2 = \Omega_1, D_2 = D_1, N_2 = N_1. \quad (55)$$

In turn, properties (55) follow from (9) for $j = 1, 2$, (10) for $j = 1$, and from the equality $w_2 = iw_1$ assumed in (27).

Next, we represent v as follows:

$$\begin{aligned} v(x) &= v^+(x, r) + v^-(x, r), \quad x \in \mathbb{R}^d, r > 0, \\ v^+(x, r) &= \int_{\mathcal{B}_r} e^{-ipx} \widehat{v}(p) dp, \\ v^-(x, r) &= \int_{\mathbb{R}^d \setminus \mathcal{B}_r} e^{-ipx} \widehat{v}(p) dp. \end{aligned} \quad (56)$$

Since $v \in W^{n,1}(\mathbb{R}^d)$, $n > d$, we have

$$\begin{aligned} |v^-(x, r)| &\leq c_3 \|v\|_{n,1} r^{d-n}, \quad x \in \mathbb{R}^d, r > 0, \\ c_3 &= |\mathbb{S}^{d-1}| \frac{(2\pi)^{-d} d^n}{n-d}, \end{aligned} \quad (57)$$

where $\|\cdot\|_{n,1}$ is defined in (31), and $|\mathbb{S}^{d-1}|$ is the standard Euclidean volume of \mathbb{S}^{d-1} . Indeed,

$$\begin{aligned} |p_1^{k_1} \cdots p_d^{k_d} \widehat{v}(p)| &\leq (2\pi)^{-d} \|v\|_{n,1}, \quad p = (p_1, \dots, p_d) \in \mathbb{R}^d, \\ &\text{for any } k_1, \dots, k_d \in \mathbb{N} \cup \{0\}, k_1 + \cdots + k_d \leq n, \end{aligned} \quad (58)$$

assuming also that $p_j^0 = 1$. Taking an appropriate sum in (58) over all such k_1, \dots, k_d with $k_1 + \cdots + k_d = m \leq n$, we get

$$|p|^m |\widehat{v}(p)| \leq (|p_1| + \cdots + |p_d|)^m |\widehat{v}(p)| \leq (2\pi)^{-d} d^m \|v\|_{n,1}, \quad p \in \mathbb{R}^d. \quad (59)$$

The definition of v^- of (56) and inequalities (59) for $m = n$ imply (57).

In addition, using Proposition 1 and the estimate on \widehat{w} of (23), we obtain:

$$\begin{aligned} \left| v^+(x, r) - \int_{\mathcal{B}_r} e^{-ipx} U_{\widehat{w}_1, \widehat{w}_2}(p, E) dp \right| &\leq c_2 E^{-\frac{1}{2}} \int_{\mathcal{B}_r} |\widehat{w}(p)|^{-1} dp \\ &\leq c_1^{-1} c_2 E^{-\frac{1}{2}} \int_{\mathcal{B}_r} (1 + |p|)^\beta dp \leq c_4 E^{-\frac{1}{2}} r^{d+\beta}, \quad c_4 = |\mathbb{S}^{d-1}| \frac{2^{d+\beta}}{d+\beta} c_1^{-1} c_2, \\ &x \in \mathbb{R}^d, 1 \leq r \leq 2\sqrt{E}, E \geq \rho_1. \end{aligned} \quad (60)$$

As a corollary of (56), (57), (60), we have

$$\left| v(x) - \int_{\mathcal{B}_r} e^{-ipx} U_{\widehat{w}_1, \widehat{w}_2}(p, E) dp \right| \leq c_3 \|v\|_{n,1} r^{d-n} + c_4 E^{-\frac{1}{2}} r^{d+\beta}, \quad (61)$$

where $x \in \mathbb{R}^d$, $1 \leq r \leq 2\sqrt{E}$, $E \geq \rho_1$. In addition, if $r = r_1(E)$, where $r_1(E)$ is defined in (39), then

$$\begin{aligned} r^{d-n} &= (2\tau)^{d-n} E^{-\alpha_1}, \\ E^{-\frac{1}{2}} r^{d+\beta} &= (2\tau)^{d+\beta} E^{-\alpha_1}. \end{aligned} \quad (62)$$

Formula (40) follows from formulas (61) and (62).

Theorem 1 is proved.

6 Proof of Theorem 2

Proposition 2. *Let v satisfy (2) and w_1, w_2 be the same as in (24)–(26), $d \geq 2$. Then:*

$$\begin{aligned} |\widehat{v}(p) - U_{\widehat{w}_1, \widehat{w}_2}(p, E)| &\leq c_5 \varepsilon^{-1} (1 + |p|)^\beta E^{-\frac{1}{2}}, \\ p &\in \mathcal{B}_{2\sqrt{E}} \setminus Z_{\widehat{w}_1, \widehat{w}_2}^\varepsilon, \quad E \geq \rho_2, \quad 0 < \varepsilon < 1, \\ c_5 &= \frac{\pi}{2} (2c(D_0)N_0^3 + c(D_1)N_1^3 + c(D_2)N_2^3) c_1, \\ \rho_2 &= \max_{j=0,1,2} \rho(D_j, N_j), \end{aligned} \quad (63)$$

in addition, if $v \in W^{n,1}(\mathbb{R}^d)$, $n \geq 0$, then:

$$\begin{aligned} &|\widehat{v}(p) - U_{\widehat{w}_1, \widehat{w}_2}^\varepsilon(p, E)| \\ &\leq 2^\beta c_5 \varepsilon^{-1} (1 + |p|)^\beta E^{-\frac{1}{2}} + c_6 \varepsilon (1 + \frac{\pi}{|y|} |z(p)| + |p_\perp|)^{-n}, \\ &p \in \mathcal{B}_{2\sqrt{E}} \cap Z_{\widehat{w}_1, \widehat{w}_2}^\varepsilon, \quad E \geq \rho_2, \quad 0 < \varepsilon < \min\{1, \frac{1}{2}|y|\}; \\ &c_6 = 2^n \frac{(d+1)^{n+1}}{(2\pi)^d |y|} \max_{j=1, \dots, d} \|x_j v\|_{n,1}, \end{aligned} \quad (64)$$

where $c, \rho, N_j, D_j, j = 0, 1, 2$, are the same as in estimates (15), c_1, β are the same as in Lemma 1; $z(p), p_\perp$ are defined in (42), (43), $x_j v = x_j v(x)$, and $\|\cdot\|_{n,1}$ is defined in (31).

Proof of Proposition 2. It follows from formulas (24), (26) and (35), (36) that

$$\begin{aligned} M_{\widehat{w}_1, \widehat{w}_2}(p) &= \widehat{w}(p) \begin{pmatrix} \cos(T_1 p) & \sin(T_1 p) \\ \cos(T_2 p) & \sin(T_2 p) \end{pmatrix}, \quad p \in \mathbb{R}^d, \\ M_{\widehat{w}_1, \widehat{w}_2}^{-1}(p) &= \frac{1}{\sin(py) \widehat{w}(p)} \begin{pmatrix} \sin(T_2 p) & -\sin(T_1 p) \\ -\cos(T_2 p) & \cos(T_1 p) \end{pmatrix}, \quad p \in \mathbb{R}^d \setminus Z_{\widehat{w}_1, \widehat{w}_2}^\varepsilon. \end{aligned} \quad (65)$$

Also note that

$$|\sin(py)| \geq \frac{2\varepsilon}{\pi}, \quad p \in \mathbb{R}^d \setminus Z_{\widehat{w}_1, \widehat{w}_2}^\varepsilon, \quad 0 < \varepsilon < 1. \quad (66)$$

The estimate (63) follows from (11), (15), (23), (34), (37), (38) and from (65), (66).

It remains to prove (64). Using definition (43), one can write

$$\begin{aligned}
U_{\widehat{w}_1, \widehat{w}_2}^\varepsilon(p, E) - \widehat{v}(p) &= \varphi_1^\varepsilon(p, E) + \varphi_2^\varepsilon(p), \quad p \in \mathcal{B}_{2\sqrt{E}} \cap Z_{\widehat{w}_1, \widehat{w}_2}^\varepsilon, \\
\varphi_1^\varepsilon(p, E) &= \frac{1}{2}(U_{\widehat{w}_1, \widehat{w}_2}^\varepsilon(p_-^\varepsilon, E) - \widehat{v}(p_-^\varepsilon)) \\
&\quad + \frac{1}{2}(U_{\widehat{w}_1, \widehat{w}_2}^\varepsilon(p_+^\varepsilon, E) - \widehat{v}(p_+^\varepsilon)), \quad p \in \mathcal{B}_{2\sqrt{E}} \cap Z_{\widehat{w}_1, \widehat{w}_2}^\varepsilon, \\
\varphi_2^\varepsilon(p) &= \frac{1}{2}(\widehat{v}(p_-^\varepsilon) + \widehat{v}(p_+^\varepsilon)) - \widehat{v}(p), \quad p \in Z_{\widehat{w}_1, \widehat{w}_2}^\varepsilon.
\end{aligned} \tag{67}$$

Using estimate (63), formula (67) and the definitions of p_\pm^ε in (43), we get

$$\begin{aligned}
|\varphi_1^\varepsilon(p, E)| &\leq \frac{1}{2}c_5\varepsilon^{-1}E^{-\frac{1}{2}}((1 + |p_-^\varepsilon|)^\beta + (1 + |p_+^\varepsilon|)^\beta) \\
&\leq c_5\varepsilon^{-1}(1 + |p| + 2\frac{\varepsilon}{|y|})^\beta E^{-\frac{1}{2}} \leq 2^\beta c_5\varepsilon^{-1}(1 + |p|)^\beta E^{-\frac{1}{2}}, \\
&\text{for } \varepsilon \text{ as in (64), } p \in \mathcal{B}_{2\sqrt{E}} \cap Z_{\widehat{w}_1, \widehat{w}_2}^\varepsilon.
\end{aligned} \tag{68}$$

Next, using the definition of φ_2^ε in (67) and the mean value theorem, we obtain

$$|\varphi_2^\varepsilon(p)| \leq \frac{\varepsilon}{|y|} \max\{|\frac{y}{|y|}\nabla\widehat{v}(\xi)|: \xi \in [p_-^\varepsilon, p_+^\varepsilon]\}, \quad p \in Z_{\widehat{w}_1, \widehat{w}_2}^\varepsilon, \tag{69}$$

where $[p_-^\varepsilon, p_+^\varepsilon]$ denotes the segment joining p_-^ε to p_+^ε . Here, the mean value theorem was used for $\widehat{v}(\xi)$ on $[p_-^\varepsilon, p]$ and on $[p, p_+^\varepsilon]$.

Note also that

$$|\nabla\widehat{v}(\xi)| \leq d \max_{j=1, \dots, d} \left| \frac{\partial\widehat{v}}{\partial\xi_j}(\xi) \right|, \quad \xi = (\xi_1, \dots, \xi_d) \in [p_-^\varepsilon, p_+^\varepsilon]. \tag{70}$$

In addition, the following estimates hold:

$$\left| \frac{\partial\widehat{v}}{\partial\xi_j}(\xi) \right| \leq \frac{(1+d)^n}{(2\pi)^d(1+|\xi|)^n} \|x_j v\|_{n,1}, \quad \xi \in [p_-^\varepsilon, p_+^\varepsilon], \quad j = 1, \dots, d. \tag{71}$$

Indeed, taking the sum in (59) over all $m = 0, \dots, n$ with the binomial coefficients, we get

$$\begin{aligned}
(1 + |p|)^n |\widehat{v}(p)| &\leq (1 + |p_1| + \dots + |p_d|)^n |\widehat{v}(p)| \\
&\leq (2\pi)^{-d} (1+d)^n \|v\|_{n,1}, \quad p \in \mathbb{R}^d.
\end{aligned} \tag{72}$$

Estimates (71) follow from (72), where we replace v by $x_j v$ and use that v belongs to $W^{n,1}(\mathbb{R}^d)$ and is compactly supported.

Estimates (69)–(71) imply

$$|\varphi_2^\varepsilon(p)| \leq 2^{-n} c_6 \varepsilon \max\{(1 + |\xi|)^{-n}: \xi \in [p_-^\varepsilon, p_+^\varepsilon]\}, \quad p \in Z_{\widehat{w}_1, \widehat{w}_2}^\varepsilon. \tag{73}$$

Using also that

$$\xi = \tau \frac{y}{|y|} + p_\perp, \text{ where } |\tau - \frac{\pi}{|y|} z(p)| \leq \frac{\varepsilon}{|y|}, \text{ if } \xi \in [p_-^\varepsilon, p_+^\varepsilon], \tag{74}$$

and that $\varepsilon < |y|$, we obtain

$$\begin{aligned}
|\varphi_2^\varepsilon(p)| &\leq 2^{-n} c_6 \varepsilon \left(1 + \frac{1}{2} \left(\frac{\pi}{|y|} |z(p)| - \frac{\varepsilon}{|y|} + |p_\perp|\right)\right)^{-n} \\
&\leq c_6 \left(1 + \frac{\pi}{|y|} |z(p)| + |p_\perp|\right)^{-n}, \quad p \in Z_{\widehat{w}_1, \widehat{w}_2}^\varepsilon.
\end{aligned} \tag{75}$$

Estimate (64) follows from (68) and (75).

Proposition 2 is proved. \square

The final part of the proof of Theorem 2 is as follows. In a similar way with (56), we represent v as follows:

$$\begin{aligned}
v(x) &= v_1^+(x, r) + v_2^+(x, r) + v^-(x, r), \quad x \in \mathbb{R}^d, r > 0, \\
v_1^+(x, r) &= \int_{\mathcal{B}_r \setminus Z_{\widehat{w}_1, \widehat{w}_2}^\varepsilon} e^{-ipx} \widehat{v}(p) dp, \\
v_2^+(x, r) &= \int_{\mathcal{B}_r \cap Z_{\widehat{w}_1, \widehat{w}_2}^\varepsilon} e^{-ipx} \widehat{v}(p) dp, \\
v^-(x, r) &= \int_{\mathbb{R}^d \setminus \mathcal{B}_r} e^{-ipx} \widehat{v}(p) dp.
\end{aligned} \tag{76}$$

Since $v \in W^{n,1}(\mathbb{R}^d)$, estimate (57) holds.

Using estimates (63), (64), we get:

$$\begin{aligned}
& \left| v_1^+(x, r) - \int_{\mathcal{B}_r \setminus Z_{\widehat{w}_1, \widehat{w}_2}^\varepsilon} e^{-ipx} U_{\widehat{w}_1, \widehat{w}_2}(p, E) dp \right. \\
& \left. + v_2^+(x, r) - \int_{\mathcal{B}_r \cap Z_{\widehat{w}_1, \widehat{w}_2}^\varepsilon} e^{-ipx} U_{\widehat{w}_1, \widehat{w}_2}^\varepsilon(p, E) dp \right| \leq I_1 + I_2,
\end{aligned} \tag{77}$$

$$I_1 = 2^\beta c_5 \varepsilon^{-1} E^{-\frac{1}{2}} \int_{\mathcal{B}_r} (1 + |p|)^\beta dp, \tag{78}$$

$$I_2 = c_6 \varepsilon \int_{\mathcal{B}_r \cap Z_{\widehat{w}_1, \widehat{w}_2}^\varepsilon} \left(1 + \frac{\pi}{|y|} |z(p)| + |p_\perp|\right)^{-n} dp, \tag{79}$$

$$x \in \mathbb{R}^d, 1 \leq r \leq 2\sqrt{E}, E \geq \rho_2,$$

where ρ_2 is the same as in Proposition 2. In addition:

$$I_1 \leq c_7 \varepsilon^{-1} E^{\frac{1}{2}} r^{d+\beta}, \quad c_7 = |\mathbb{S}^{d-1}|^{\frac{2d+2\beta}{d+\beta}} c_5; \tag{80}$$

$$I_2 = c_6 \varepsilon \sum_{z \in \mathbb{Z}} \int_{\left\{ \begin{array}{l} \tau^2 + p_\perp^2 \leq r^2 \\ |\tau - \frac{\pi}{|y|} z| \leq \frac{\varepsilon}{|y|} \end{array} \right\}} \left(1 + \frac{\pi}{|y|} |z| + |p_\perp|\right)^{-n} d\tau dp_\perp,$$

where $\tau \in \mathbb{R}, p_\perp \in \mathbb{R}^d, p_\perp y = 0$,

$$\begin{aligned}
I_2 &\leq 2c_6 \frac{\varepsilon^2}{|y|} \sum_{z \in \mathbb{Z}} \int_{\xi \in \mathbb{R}^{d-1}, |\xi| \leq r} \left(1 + \frac{\pi}{|y|} |z| + |\xi|\right)^{-n} d\xi \leq c_6 c_8 \varepsilon^2, \\
c_8 &= \frac{2}{|y|} \frac{|\mathbb{S}^{d-2}|}{n-d+1} \sum_{z \in \mathbb{Z}} \left(1 + \frac{\pi}{|y|} |z|\right)^{d-n-1}.
\end{aligned} \tag{81}$$

In addition, if $r = r_2(E), \varepsilon = \varepsilon_2(E)$, where $r_2(E), \varepsilon_2(E)$ are defined in (44), then

$$\begin{aligned}
r^{d-n} &= (2\tau)^{d-n} E^{-\alpha_2}, \\
\varepsilon^{-1} E^{-\frac{1}{2}} r^{d+\beta} &= (2\tau)^{d+\beta} E^{-\alpha_2}, \\
\varepsilon^2 &= E^{-\alpha_2}.
\end{aligned} \tag{82}$$

Formula (45) follows from representation (76), estimates (57), (77), (80), (81) and formulas (82).

Theorem 2 is proved.

7 Proof of Theorem 3

Proposition 3. *Let v satisfy (2) and w_1, \dots, w_{d+1} be the same as in (24), (29), $d \geq 2$. Then:*

$$\begin{aligned} |\widehat{v}(p) - U_{\widehat{w}_1, \dots, \widehat{w}_{d+1}}(p, E)| &\leq c_9 \varepsilon^{-1} (1 + |p|)^\beta E^{-\frac{1}{2}}, \\ p &\in \mathcal{B}_{2\sqrt{E}} \setminus Z_{\widehat{w}_1, \dots, \widehat{w}_{d+1}}^\varepsilon, \quad E \geq \rho_3, \quad 0 < \varepsilon < 1, \\ c_9 &= \frac{\pi\sqrt{d}}{2} (2c(D_0)N_0^3 + c(D_1)N_1^3 + \max_{j=2, \dots, d+1} c(D_j)N_j^3) c_1, \\ \rho_3 &= \max_{j=0, \dots, d+1} \rho(D_j, N_j), \end{aligned} \quad (83)$$

in addition, if $v \in W^{n,1}(\mathbb{R}^d)$, $n \geq 0$, then:

$$\begin{aligned} &|\widehat{v}(p) - U_{\widehat{w}_1, \dots, \widehat{w}_{d+1}}^\varepsilon(p, E)| \\ &\leq 2^\beta c_9 \varepsilon^{-1} (1 + |p|)^\beta E^{-\frac{1}{2}} + 2c_6 \varepsilon (1 + 2\frac{\pi}{s} \|z(p)\|_2)^{-n}, \\ &p \in \mathcal{B}_{2\sqrt{E}} \cap Z_{\widehat{w}_1, \dots, \widehat{w}_{d+1}}^\varepsilon, \quad E \geq \rho_3, \quad 0 < \varepsilon < \min\{1, \frac{1}{2}s\}, \end{aligned} \quad (84)$$

where $c, \rho, D_j, N_j, j = 0, \dots, d+1$, are defined as in (15); c_1, β are the same as in Lemma 1, c_6 is the same as in Proposition 2, $z(p)$ is defined in (47) and $\|z(p)\|_2$ is the standard Euclidean norm of $z(p)$.

Proof of Proposition 3. In a similar way with formulas (65), one can write

$$\begin{aligned} M_{\widehat{w}_1, \widehat{w}_{i'}}(p) &= \widehat{w}_1(p) \begin{pmatrix} 1 & 0 \\ \cos(sp_{i'-1}) & \sin(sp_{i'-1}) \end{pmatrix}, \quad p \in \mathbb{R}^d \setminus \frac{\pi}{s} Z^d, \\ M_{\widehat{w}_1, \widehat{w}_{i'}}^{-1}(p) &= \frac{1}{\sin(sp_{i'-1}) \widehat{w}_1(p)} \begin{pmatrix} \sin(sp_{i'-1}) & 0 \\ -\cos(sp_{i'-1}) & 1 \end{pmatrix}, \quad p \in \mathbb{R}^d \setminus Z_{\widehat{w}_1, \dots, \widehat{w}_{d+1}}^\varepsilon, \end{aligned} \quad (85)$$

where $i' = i'(p, s)$ is defined in (48). Also note that

$$|\sin(sp_{i'-1})| \geq \frac{2\varepsilon}{\pi\sqrt{d}}, \quad p \in \mathbb{R}^d \setminus Z_{\widehat{w}_1, \dots, \widehat{w}_{d+1}}^\varepsilon, \quad 0 < \varepsilon < 1. \quad (86)$$

Estimate (83) follows from (11), (15), (23), (34), (37), (38), (48), (49) and from (85), (86).

It remains to prove (84). Using definition (50), we represent

$$\begin{aligned} U_{\widehat{w}_1, \dots, \widehat{w}_{d+1}}^\varepsilon(p, E) - \widehat{v}(p) &= \varphi_1^\varepsilon(p, E) + \varphi_2^\varepsilon(p), \quad p \in \mathcal{B}_{2\sqrt{E}} \cap Z_{\widehat{w}_1, \dots, \widehat{w}_{d+1}}^\varepsilon \\ \varphi_1^\varepsilon(p, E) &= \frac{1}{|\mathbb{S}^{d-1}|} \int_{\mathbb{S}^{d-1}} (U_{\widehat{w}_1, \dots, \widehat{w}_{d+1}}(\eta, E) - \widehat{v}(\eta)) \Big|_{\eta = \frac{\varepsilon}{s} \vartheta + \frac{\pi}{s} z(p)} d\vartheta, \\ \varphi_2^\varepsilon(p) &= \frac{1}{|\mathbb{S}^{d-1}|} \int_{\mathbb{S}^{d-1}} (\widehat{v}(\frac{\varepsilon}{s} \vartheta + \frac{\pi}{s} z(p)) - \widehat{v}(p)) d\vartheta, \end{aligned} \quad (87)$$

where $z(p)$ is defined in (47).

Using formulas (83), (87), we obtain

$$\begin{aligned} |\varphi_1^\varepsilon(p, E)| &\leq c_9 \varepsilon^{-1} E^{-\frac{1}{2}} \frac{1}{|\mathbb{S}^{d-1}|} \int_{\mathbb{S}^{d-1}} (1 + |\frac{\varepsilon}{s} \vartheta + \frac{\pi}{s} z(p)|)^\beta d\vartheta \\ &\leq c_9 \varepsilon^{-1} E^{-\frac{1}{2}} \frac{1}{|\mathbb{S}^{d-1}|} \int_{\mathbb{S}^{d-1}} (1 + |p| + 2\frac{\varepsilon}{s})^\beta d\vartheta \leq 2^\beta c_9 \varepsilon^{-1} (1 + |p|)^\beta E^{-\frac{1}{2}}, \end{aligned} \quad (88)$$

for ε as in (84).

Next, using the definition of φ_2^ε in formula (87) and the mean value theorem, we get the following estimate:

$$|\varphi_2^\varepsilon(p)| \leq 2\frac{\varepsilon}{s} \max\{|\nabla \widehat{v}(\xi)| : \xi \in \mathbb{R}^d, |\xi - \frac{\pi}{s} z(p)| \leq \frac{\varepsilon}{s}\}, p \in Z_{\widehat{w}_1, \dots, \widehat{w}_{d+1}}^\varepsilon. \quad (89)$$

Here, the mean value theorem was used for $\widehat{v}(\xi)$ on $[p, \frac{\varepsilon}{s} \vartheta + \frac{\pi}{s} z(p)]$, $\vartheta \in \mathbb{S}^{d-1}$. One can see that

$$\begin{aligned} &\text{estimates (70) and (71) hold for all } \xi \in \mathbb{R}^d \\ &\text{such that } |\xi - \frac{\pi}{s} z(p)| \leq \frac{\varepsilon}{s}, \text{ where } p \in Z_{\widehat{w}_1, \dots, \widehat{w}_{d+1}}^\varepsilon. \end{aligned} \quad (90)$$

It follows from (89), (90) and from the upper estimate on ε of (84), that

$$\begin{aligned} |\varphi_2^\varepsilon(p)| &\leq 2^{1-n} c_6 \varepsilon \max\{(1 + |\xi|)^{-n} : |\xi - \frac{\pi}{s} z(p)| \leq \frac{\varepsilon}{s}\} \\ &\leq 2^{1-n} c_6 \varepsilon (1 + \frac{\pi}{s} \|z(p)\|_2 - \frac{\varepsilon}{s})^{-n} \\ &\leq 2c_6 \varepsilon (1 + 2\frac{\pi}{s} \|z(p)\|_2)^{-n}, p \in Z_{\widehat{w}_1, \dots, \widehat{w}_{d+1}}^\varepsilon. \end{aligned} \quad (91)$$

Estimate (84) follows from estimates (88) and (91).

Proposition 3 is proved. \square

The final part of the proof of Theorem 3 is as follows. In a similar way with (76), we represent v as follows:

$$\begin{aligned} v(x) &= v_1^+(x, r) + v_2^+(x, r) + v^-(x, r), \quad x \in \mathbb{R}^d, r > 0, \\ v_1^+(x, r) &= \int_{\mathcal{B}_r \setminus Z_{\widehat{w}_1, \dots, \widehat{w}_{d+1}}^\varepsilon} e^{-ipx} \widehat{v}(p) dp, \\ v_2^+(x, r) &= \int_{\mathcal{B}_r \cap Z_{\widehat{w}_1, \dots, \widehat{w}_{d+1}}^\varepsilon} e^{-ipx} \widehat{v}(p) dp, \\ v^-(x, r) &= \int_{\mathbb{R}^d \setminus \mathcal{B}_r} e^{-ipx} \widehat{v}(p) dp. \end{aligned} \quad (92)$$

Since v belongs to $W^{n,1}(\mathbb{R}^d)$, estimate (57) is valid.

Using estimates (83), (84) we obtain

$$\begin{aligned}
& \left| v_1^+(x, r) - \int_{\mathcal{B}_r \setminus Z_{\widehat{w}_1, \dots, \widehat{w}_{d+1}}^\varepsilon} e^{-ipx} U_{\widehat{w}_1, \dots, \widehat{w}_{d+1}}(p, E) dp \right. \\
& \left. + v_2^+(x, r) - \int_{\mathcal{B}_r \cap Z_{\widehat{w}_1, \dots, \widehat{w}_{d+1}}^\varepsilon} e^{-ipx} U_{\widehat{w}_1, \dots, \widehat{w}_{d+1}}^\varepsilon(p, E) dp \right| \leq J_1 + J_2, \\
& J_1 = 2^\beta c_9 \varepsilon^{-1} E^{-\frac{1}{2}} \int_{\mathcal{B}_r} (1 + |p|)^\beta dp, \\
& J_2 = 2c_6 \varepsilon \int_{\mathcal{B}_r \cap Z_{\widehat{w}_1, \dots, \widehat{w}_{d+1}}^\varepsilon} \left(1 + 2\frac{\pi}{s} \|z(p)\|_2\right)^{-n} dp, \\
& x \in \mathbb{R}^d, \quad 1 \leq r \leq 2\sqrt{E}, \quad E \geq \rho_3,
\end{aligned} \tag{93}$$

where ρ_3 is the same as in Proposition 3. In addition,

$$\begin{aligned}
& J_1 \leq c_{10} \varepsilon^{-1} E^{-\frac{1}{2}} r^{d+\beta}, \quad c_{10} = |\mathbb{S}^{d-1}| \frac{2^{d+\beta}}{d+\beta} c_9, \\
& J_2 \leq c_{11} \varepsilon^{d+1}, \quad c_{11} = \left(\frac{1}{s}\right)^d |\mathcal{B}_1| \sum_{z \in \mathbb{Z}^d} \left(1 + 2\frac{\pi}{s} \|z\|_2\right)^{-n},
\end{aligned} \tag{94}$$

where $|\mathcal{B}_1|$ is the standard Euclidean volume of \mathcal{B}_1 . Finally, if $r = r_3(E)$, $\varepsilon = \varepsilon_3(E)$, where $r_3(E)$, $\varepsilon_3(E)$ are defined in (52), then

$$\begin{aligned}
& r^{d-n} = (2\tau)^{d-n} E^{-\alpha_3}, \\
& \varepsilon^{-1} E^{-\frac{1}{2}} r^{d+\beta} = (2\tau)^{d+\beta} E^{-\alpha_3}, \\
& \varepsilon^{d+1} = E^{-\alpha_3}.
\end{aligned} \tag{95}$$

Formula (53) follows from representation (92), estimates (57), (93), (94) and formulas (95).

Theorem 3 is proved.

8 Proof of Lemma 1

Note that

$$\widehat{w}(p) = \int_{\mathbb{R}^d} |\widehat{q}(\xi)|^2 \widehat{\omega}_\nu(p - \xi) d\xi, \quad p \in \mathbb{R}^d, \tag{96}$$

$$\omega_\nu(x) = |x|^{\frac{\nu}{2}} K_\nu(|x|), \quad x \in \mathbb{R}^d, \tag{97}$$

where \widehat{q} , $\widehat{\omega}_\nu$ are the Fourier transforms of q , ω_ν . The Fourier transform $\widehat{\omega}_\nu$ can be computed explicitly:

$$\widehat{\omega}_\nu(p) = \frac{c_{12}}{(1 + |p|^2)^{\frac{d}{2} + \nu}}, \quad c_{12} = \frac{\Gamma(\frac{d}{2} + \nu) 2^{\nu-1}}{\pi^{\frac{d}{2}}}. \tag{98}$$

Indeed, formula (98) follows from the Fourier inversion theorem and the following computations:

$$\begin{aligned}
\int_{\mathbb{R}^d} \frac{e^{-ipx} dp}{(1+|p|^2)^{\frac{d}{2}+\nu}} &= \int_{\mathbb{R}} \int_{\mathbb{R}^{d-1}} \frac{e^{-i|x|t} dt d\xi}{(1+t^2+|\xi|^2)^{\frac{d}{2}+\nu}} \\
&= |\mathbb{S}^{d-2}| \int_{\mathbb{R}} \int_{\mathbb{R}} \frac{e^{-i|x|t} r^{d-2} dt dr}{(1+t^2+r^2)^{\frac{d}{2}+\nu}} \\
&\stackrel{r=\sqrt{1+t^2}\tau}{=} |\mathbb{S}^{d-2}| \int_{\mathbb{R}} \frac{e^{-i|x|t} dt}{(1+t^2)^{\frac{1}{2}+\nu}} \int_0^{+\infty} \frac{\tau^{d-2} d\tau}{(1+\tau^2)^{\frac{d}{2}+\nu}} \\
&= c_{12}^{-1} |x|^\nu K_\nu(|x|), \quad x \in \mathbb{R}^d.
\end{aligned}$$

Here, it was used that

$$\begin{aligned}
\int_0^{+\infty} \frac{\tau^{d-2} d\tau}{(1+\tau^2)^{\frac{d}{2}+\nu}} &= \frac{1}{2} B\left(\frac{d-1}{2}, \nu + \frac{1}{2}\right) = \frac{1}{2} \frac{\Gamma(\frac{d-1}{2})\Gamma(\nu + \frac{1}{2})}{\Gamma(\frac{d}{2} + \nu)}, \\
|\mathbb{S}^{d-2}| &= \frac{2\pi^{\frac{d-1}{2}}}{\Gamma(\frac{d-1}{2})},
\end{aligned}$$

where B and Γ denote the beta and gamma functions.

Using (96), (98), we obtain the estimates

$$\begin{aligned}
\widehat{w}(p) &\geq \int_{|\xi| \leq 1} \frac{c_{12} |\widehat{q}(\xi)|^2}{(1+|p-\xi|)^{d+2\nu}} d\xi \geq \frac{c_1(q, \nu)}{(1+|p|)^{d+2\nu}}, \quad p \in \mathbb{R}^d, \\
c_1(q, \nu) &= \frac{c_{12}}{2^{d+2\nu}} \int_{|\xi| \leq 1} |\widehat{q}(\xi)|^2 d\xi.
\end{aligned} \tag{99}$$

Properties (23) follow from (20), (22), (96) and (99).

Lemma 1 is proved.

References

- [1] A. D. Agaltsov, T. Hohage, and R. G. Novikov, “An iterative approach to monochromatic phaseless inverse scattering,” in preparation.
- [2] T. Aktosun and P. E. Sacks, “Inverse problems on the line without phase information,” *Inverse Problems*, vol. 14, pp. 211–224, 1998.
- [3] N. V. Alexeenko, V. A. Burov, and O. D. Rumyantseva, “Solution of the three-dimensional acoustical inverse scattering problem. The modified Novikov algorithm,” *Acoustical Physics*, vol. 54, no. 3, pp. 407–419, 2008.
- [4] J. A. Barceló, C. Castro, and J. M. Reyes, “Numerical approximation of the potential in the two-dimensional inverse scattering problem,” *Inverse Problems*, vol. 32, no. 1, 2016, 015006 (19pp).

- [5] F. A. Berezin and M. A. Shubin, *The Schrödinger equation*, ser. Mathematics and its applications. Dordrecht: Kluwer Academic, 1991, vol. 66.
- [6] V. A. Burov, N. V. Alekseenko, and O. D. Rumyantseva, “Multifrequency generalization of the Novikov algorithm for the two-dimensional inverse scattering problem,” *Acoustical Physics*, vol. 56, no. 6, pp. 843–856, 2009.
- [7] K. Chadan and P. C. Sabatier, *Inverse problems in quantum scattering theory*, 2nd ed. Berlin: Springer, 1989.
- [8] G. Eskin, *Lectures on linear partial differential equations*, ser. Graduate studies in mathematics. American mathematical society, 2011, vol. 123.
- [9] L. D. Faddeev, “Uniqueness of the solution of the inverse scattering problem,” *Vest. Leningrad Univ.*, vol. 7, pp. 126–130, 1956, (in Russian).
- [10] —, “Inverse problem of quantum scattering theory. II.” *Journal of Soviet Mathematics*, vol. 5, no. 3, pp. 334–396, 1976.
- [11] L. D. Faddeev and S. P. Merkuriev, *Quantum scattering theory for several particle systems*, ser. Mathematical physics and applied mathematics. Dordrecht: Kluwer Academic, 1993, vol. 11.
- [12] P. G. Grinevich, “The scattering transform for the two-dimensional Schrödinger operator with a potential that decreases at infinity at fixed nonzero energy,” *Russian Math. Surveys*, vol. 55, no. 6, pp. 1015–1083, 2000.
- [13] P. Hähner and T. Hohage, “New stability estimates for the inverse acoustic inhomogeneous medium problem and applications,” *SIAM J. Math. Anal.*, vol. 33, no. 3, pp. 670–685, 2001.
- [14] M. I. Isayev, “Exponential instability in the inverse scattering problem on the energy interval,” *Funct. Anal. Appl.*, vol. 47, no. 3, pp. 187–194, 2013.
- [15] M. I. Isayev and R. G. Novikov, “New global stability estimates for monochromatic inverse acoustic scattering,” *SIAM J. Math. Analysis*, vol. 43, no. 5, pp. 1495–1504, 2013.
- [16] M. V. Klibanov, “Phaseless inverse scattering problems in three dimensions,” *SIAM J. Appl. Math.*, vol. 74, pp. 392–410, 2014.
- [17] M. V. Klibanov and V. G. Romanov, “Reconstruction procedures for two inverse scattering problems without the phase information,” *SIAM J. Appl. Math.*, vol. 76, no. 1, pp. 178–196, 2016.
- [18] —, “Two reconstruction procedures for a 3D phaseless inverse scattering problem for the generalized Helmholtz equation,” *Inverse Problems*, vol. 32, no. 1, 2016, 015005 (16pp).

- [19] R. G. Novikov, “Multidimensional inverse spectral problem for the equation $-\Delta\psi + (v(x) - Eu(x))\psi = 0$,” *Funct. Anal. Appl.*, vol. 22, pp. 263–272, 1988.
- [20] —, “Rapidly converging approximation in inverse quantum scattering in dimension 2,” *Physics Letters A*, vol. 238, pp. 73–78, 1998.
- [21] —, “The $\bar{\partial}$ -approach to approximate inverse scattering at fixed energy in three dimensions,” *Int. Math. Res. Pap.*, no. 6, pp. 287–349, 2005.
- [22] —, “Approximate Lipschitz stability for non-overdetermined inverse scattering at fixed energy,” *J. Inverse Ill-Posed Probl.*, vol. 26, no. 6, pp. 813–823, 2013.
- [23] —, “Inverse scattering without phase information,” in *Séminaire Laurent Schwartz — EDP et applications*. Ecole Polytechnique, 2014-2015, exp. no. 16 (13pp).
- [24] —, “Formulas for phase recovering from phaseless scattering data at fixed frequency,” *Bull. Sci. Math.*, vol. 139, no. 8, pp. 923–936, 2015.
- [25] —, “An iterative approach to non-overdetermined inverse scattering at fixed energy,” *Sbornik: Mathematics*, vol. 206, no. 1, pp. 120–134, 2015.
- [26] —, “Explicit formulas and global uniqueness for phaseless inverse scattering in multidimensions,” *The Journal of Geometric Analysis*, vol. 26, no. 1, pp. 346–359, 2016.