



HAL
open science

THE FIRST POSITIVE EIGENVALUE OF THE SUB-LAPLACIAN ON CR SPHERES

Amine Aribi, Ahmad El Soufi

► **To cite this version:**

Amine Aribi, Ahmad El Soufi. THE FIRST POSITIVE EIGENVALUE OF THE SUB-LAPLACIAN ON CR SPHERES. 2016. hal-01303936

HAL Id: hal-01303936

<https://hal.science/hal-01303936v1>

Preprint submitted on 18 Apr 2016

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

THE FIRST POSITIVE EIGENVALUE OF THE SUB-LAPLACIAN ON CR SPHERES

AMINE ARIBI AND AHMAD EL SOUFI

ABSTRACT. We prove that the first positive eigenvalue, normalized by the volume, of the sub-Laplacian associated with a strictly pseudoconvex pseudo-Hermitian structure θ on the CR sphere $\mathbb{S}^{2n+1} \subset \mathbb{C}^{n+1}$, achieves its maximum when θ is the standard contact form.

1. INTRODUCTION AND STATEMENT OF THE MAIN RESULT

According to a classical result of Hersch [15], given any Riemannian metric g on the 2-dimensional sphere \mathbb{S}^2 , the first positive eigenvalue $\lambda_1(g)$ of the Laplace-Beltrami operator Δ_g satisfies the estimate

$$\lambda_1(g)A(g) \leq \lambda_1(g_0)A(g_0) \tag{1}$$

where g_0 is the standard metric of \mathbb{S}^2 and $A(g)$ is the area of \mathbb{S}^2 with respect to g . Moreover, the equality holds in (1) if and only if g is isometric to g_0 . This result was extended to higher dimensional spheres by Ilias and the second author as follows (see [12, proposition 3.1]) : If a Riemannian metric g on the n -dimensional sphere \mathbb{S}^n is conformal to the standard metric g_0 , then

$$\lambda_1(g)V(g)^{\frac{2}{n}} \leq \lambda_1(g_0)V(g_0)^{\frac{2}{n}} \tag{2}$$

where $V(g)$ denotes the Riemannian volume of the sphere with respect to g . Again, the equality holds in (2) if and only if g is isometric to g_0 .

The aim of the present paper is to establish a version of the estimate (2) for the first positive eigenvalue of the sub-Laplacian on the CR sphere $\mathbb{S}^{2n+1} \subset \mathbb{C}^{n+1}$. Indeed, let

$$\theta_0 = \frac{i}{2} \sum_{j=1}^{n+1} (\zeta_j d\bar{\zeta}_j - \bar{\zeta}_j d\zeta_j)$$

be the standard contact form on \mathbb{S}^{2n+1} whose kernel coincides with the Levi distribution $H(\mathbb{S}^{2n+1}) = T\mathbb{S}^{2n+1} \cap J T\mathbb{S}^{2n+1}$, where J is the complex structure of \mathbb{C}^{n+1} . The set $\mathcal{P}_+(\mathbb{S}^{2n+1}) = \{f\theta_0 ; f \in C^\infty(\mathbb{S}^{2n+1}) \text{ and } f > 0\}$ contains all pseudo-Hermitian structures on \mathbb{S}^{2n+1} whose Levi form is positive definite. Given a pseudo-Hermitian structure $\theta \in \mathcal{P}_+(\mathbb{S}^{2n+1})$, we denote by $\lambda_1(\theta)$ the first positive eigenvalue of the corresponding sub-Laplacian Δ_θ , and by $V(\theta)$ the volume of \mathbb{S}^{2n+1} with respect to the volume form $\psi_\theta = \frac{1}{n!2^n} \theta \wedge (d\theta)^n$ (see the next section for precise definitions). The main result of this paper is

2010 *Mathematics Subject Classification.* 32V20, 35H20, 58J50.

Key words and phrases. CR Sphere, Sub-Laplacian, eigenvalue.

Theorem 1.1. *For every pseudo-Hermitian structure $\theta \in \mathcal{P}_+(\mathbb{S}^{2n+1})$ we have*

$$\lambda_1(\theta)V(\theta)^{\frac{1}{n+1}} \leq \lambda_1(\theta_0)V(\theta_0)^{\frac{1}{n+1}}. \quad (3)$$

The equality holds in (3) if and only if there exists a CR-automorphism γ of \mathbb{S}^{2n+1} such that $\theta = c\gamma^\theta_0$ for some constant $c > 0$, or if and only if there exist $p \in \mathbb{S}^{2n+1}$ and $t \geq 0$ such that*

$$\theta = \frac{c}{|\cosh t + \sinh t (\zeta, p)|^2} \theta_0,$$

where (\cdot, \cdot) denotes the standard Hermitian product of \mathbb{C}^{n+1} .

This result can be seen as a contribution to the program aiming to recovering the main results of spectral geometry, established for the eigenvalues of the Laplace-Beltrami operator on a compact Riemannian manifold, in the realm of CR and pseudo-Hermitian geometry. This program has motivated a lot of research in recent years and we can find significant contributions in [1, 2, 3, 4, 6, 5, 7, 8, 9, 10, 13, 14, 17, 16, 18, 19, 20, 21, 22].

2. PROOF OF THEOREM 1.1

Let M be a connected differentiable manifold of dimension $2n+1 \geq 3$. A CR structure on M is a couple $(H(M), J)$ where $H(M)$ is a $2n$ -dimensional subbundle of the tangent bundle TM , the so-called Levi distribution, endowed with a pseudo-complex operator J satisfying the following integrability condition : $\forall X, Y \in \Gamma(H(M))$,

$$[X, Y] - [JX, JY] \in \Gamma(H(M))$$

and

$$J([X, Y] - [JX, JY]) = [JX, Y] + [X, JY].$$

Real hypersurfaces of \mathbb{C}^{n+1} are the most natural examples of CR manifolds. Indeed, if $M \subset \mathbb{C}^{n+1}$ is such a hypersurface, then $H(M) := TM \cap J(TM)$ endowed with the restriction of the standard complex structure J of \mathbb{C}^{n+1} , is a CR structure on M .

If $(M, H(M), J)$ is an orientable CR manifold, then there exists a nontrivial 1-form $\theta \in \Gamma(T^*M)$ such that $\text{Ker}\theta = H(M)$. Such a 1-form, called pseudo-Hermitian structure, is of course not unique. The set of pseudo-Hermitian structures consists in all the forms $f\theta$, where f is a smooth nowhere zero function on M . To each pseudo-Hermitian structure θ we associate its Levi form L_θ defined on $H(M)$ by

$$L_\theta(X, Y) = -d\theta(JX, Y) = \theta([JX, Y]).$$

The integrability of J implies that L_θ is symmetric and J -invariant. The CR manifold M is called strictly pseudoconvex if L_θ is definite. Of course, this condition does not depend on the choice of θ (since $L_{f\theta} = fL_\theta$). In the sequel, we denote by $\mathcal{P}_+(M)$ the set of all pseudo-Hermitian structures with positive definite Levi form. Every $\theta \in \mathcal{P}_+(M)$ is in fact a contact form which induces on M the following volume form

$$\psi_\theta = \frac{1}{2^n n!} \theta \wedge (d\theta)^n.$$

The associated divergence div_θ is defined, for every smooth vector field Z on M , by

$$\mathcal{L}_Z \psi_\theta = \text{div}_\theta(Z) \psi_\theta.$$

The sub-Laplacian Δ_θ is then defined for all $u \in C^\infty(M)$, by

$$\Delta_\theta u = \operatorname{div}_\theta(\nabla^H u)$$

where $\nabla^H u \in \Gamma(H(M))$ is the horizontal vectorfield such that, $\forall X \in H(M)$, $du(X) = L_\theta(\nabla^H u, X)$. The following integration by parts formula holds for any $u, v \in C_0^\infty(M)$:

$$\int_M (\Delta_\theta u) v \psi_\theta = - \int_M L_\theta(\nabla^H u, \nabla^H v) \psi_\theta.$$

Given $\theta \in \mathcal{P}_+(M)$, there is a unique vectorfield ξ , often called Reeb vectorfield, that satisfies $\theta(\xi) = 1$ and $\xi \lrcorner d\theta = 0$. The Levi form L_θ extends to a Riemannian metric on M (the Webster metric) given by

$$g_\theta(X, Y) = L_\theta(X^H, Y^H) + \theta(X)\theta(Y)$$

with $X^H = X - \theta(X)\xi$. The corresponding Laplace-Beltrami operator Δ_{g_θ} is related to Δ_θ by the following formula (see [13])

$$\Delta_\theta = \Delta_{g_\theta} - \xi^2. \quad (4)$$

The sub-Laplacian Δ_θ is a sub-elliptic operator of order $1/2$. When M is compact, it admits a self-adjoint extension to an unbounded operator of $L^2(M, \psi_\theta)$ whose resolvent is compact (see for instance [2, Lemma 2.2]). Hence, the spectrum of $-\Delta_\theta$ is discrete and consists of a sequence of nonnegative eigenvalues of finite multiplicity $\{\lambda_k(\theta)\}_{k \geq 0}$ with $\lambda_0(\theta) = 0$. The min-max variational principle gives

$$\lambda_1(\theta) = \inf_{\int_M u \psi_\theta = 0} \frac{\int_M |\nabla^H u|_\theta^2 \psi_\theta}{\int_M u^2 \psi_\theta} \quad (5)$$

where $|\nabla^H u|_\theta^2 = L_\theta(\nabla^H u, \nabla^H u)$.

2.1. The CR Sphere. Let \mathbb{S}^{2n+1} be the unit Sphere in \mathbb{C}^{n+1}

$$\mathbb{S}^{2n+1} = \left\{ \zeta = (\zeta_1, \dots, \zeta_{n+1}) \in \mathbb{C}^{n+1}; \sum_{j=1}^{n+1} |\zeta_j|^2 = 1 \right\}$$

endowed with its standard CR-structure. The restriction to \mathbb{S}^{2n+1} of the contact form

$$\theta_0 = -\frac{i}{2} \sum_{j=1}^{n+1} (\bar{\zeta}_j d\zeta_j - \zeta_j d\bar{\zeta}_j)$$

is a pseudohermitian structure whose Reeb field $\xi = i \sum_{j=1}^{n+1} \left(\zeta_j \frac{\partial}{\partial \bar{\zeta}_j} - \bar{\zeta}_j \frac{\partial}{\partial \zeta_j} \right)$ generates the natural action of \mathbb{S}^1 on \mathbb{S}^{2n+1} . Since $d\theta_0$ is the standard Kähler form of \mathbb{C}^{n+1} , the induced Levi form on $H(\mathbb{S}^{2n+1})$ coincides with the standard metric of the sphere.

If $V^{p,q}$ is the space of harmonic polynomials of bi-degree (p, q) in \mathbb{C}^{n+1} , then, ξ acts on $V^{p,q}$ as the multiplication by $i(p - q)$ and it can be deduced, using (4), that $V^{p,q}$ is an eigenspace of Δ_{θ_0} on \mathbb{S}^{2n+1} with eigenvalue $2n(p + q) + 4pq$ (see [10, Theorem 4.1] and [22, Proposition 4.4] for details). Therefore,

$$\lambda_1(\theta_0) = 2n. \quad (6)$$

2.2. One-parameter groups of CR-automorphisms of the sphere. A differentiable map $\varphi : M \rightarrow \widetilde{M}$ between two CR manifolds is a CR map if for any $x \in M$,

$$d_x\varphi(H_x(M)) \subset H_{\varphi(x)}(\widetilde{M}) \quad \text{and} \quad d_x\varphi \circ J_x^M = J_{\varphi(x)}^{\widetilde{M}} \circ d_x\varphi. \quad (7)$$

A CR-automorphism of a CR manifold M is a diffeomorphism of M which is a CR map.

Let $e_{n+1} = (0, \dots, 0, 1) \in \mathbb{S}^{2n+1}$. The punctured sphere $\mathbb{S}^{2n+1} \setminus \{e_{n+1}\}$ can be identified with the boundary of the so-called Siegel domain $\Omega_{n+1} = \{(z, w) \in \mathbb{C}^n \times \mathbb{C} : \text{Im } w > |z|^2\} \subset \mathbb{C}^{n+1}$ through the CR diffeomorphism $\Phi : \mathbb{S}^{2n+1} \setminus \{e_{n+1}\} \rightarrow \partial\Omega_{n+1}$ given by

$$\Phi(\zeta) = \frac{1}{1 - \zeta_{n+1}} (\zeta_1, \dots, \zeta_n, i(1 + \zeta_{n+1}))$$

with

$$\Phi^{-1}(z, w) = \frac{1}{w + i} (2iz_1, \dots, 2iz_n, w - i)$$

For every $t \in \mathbb{R}$, the ‘‘dilation’’

$$H_t : \begin{array}{ccc} \partial\Omega_{n+1} & \rightarrow & \partial\Omega_{n+1} \\ (z, w) & \mapsto & (e^t z, e^{2t} w) \end{array}$$

is a CR-automorphism of $\partial\Omega_{n+1}$. We define $\gamma_t : \mathbb{S}^{2n+1} \rightarrow \mathbb{S}^{2n+1}$ by $\gamma_t(e_{n+1}) = e_{n+1}$ and, $\forall \zeta \in \mathbb{S}^{2n+1} \setminus \{e_{n+1}\}$,

$$\gamma_t(\zeta) = \Phi^{-1} \circ H_t \circ \Phi(\zeta) = \frac{1}{\cosh t + \sinh t \zeta_{n+1}} (\zeta_1, \dots, \zeta_n, \sinh t + \cosh t \zeta_{n+1})$$

or

$$\gamma_t(\zeta) = \frac{1}{\cosh t + \sinh t \zeta_{n+1}} (\zeta + (\sinh t + (\cosh t - 1) \zeta_{n+1}) e_{n+1}).$$

Lemma 2.1. *For every t , the map γ_t is a CR-automorphism of \mathbb{S}^{2n+1} which satisfies*

$$(\gamma_t)^*\theta_0 = \frac{1}{|\cosh t + \sinh t \zeta_{n+1}|^2} \theta_0.$$

Proof. Let $f_j(\zeta) = \frac{\zeta_j}{\cosh t + \sinh t \zeta_{n+1}}$, $j \leq n$, and $f_{n+1}(\zeta) = \frac{\sinh t + \cosh t \zeta_{n+1}}{\cosh t + \sinh t \zeta_{n+1}}$. Then

$$df_j = \frac{d\zeta_j}{\cosh t + \sinh t \zeta_{n+1}} - \frac{\sinh t \zeta_j d\zeta_{n+1}}{(\cosh t + \sinh t \zeta_{n+1})^2} \quad \text{and} \quad df_{n+1} = \frac{d\zeta_{n+1}}{(\cosh t + \sinh t \zeta_{n+1})^2}.$$

Therefore

$$f_j d\bar{f}_j = \frac{\zeta_j d\bar{\zeta}_j}{|\cosh t + \sinh t \zeta_{n+1}|^2} - \frac{|\zeta_j|^2 \sinh t d\bar{\zeta}_{n+1}}{|\cosh t + \sinh t \zeta_{n+1}|^2 (\cosh t + \sinh t \bar{\zeta}_{n+1})},$$

$$\bar{f}_j df_j = \frac{\bar{\zeta}_j d\zeta_j}{|\cosh t + \sinh t \zeta_{n+1}|^2} - \frac{|\zeta_j|^2 \sinh t d\zeta_{n+1}}{|\cosh t + \sinh t \zeta_{n+1}|^2 (\cosh t + \sinh t \zeta_{n+1})}$$

which gives with $\sum_{j=1}^n |\zeta_j|^2 = 1 - |\zeta_{n+1}|^2$,

$$\sum_{j=1}^n (f_j d\bar{f}_j - \bar{f}_j df_j) = \frac{1}{|\cosh t + \sinh t \zeta_{n+1}|^2} \sum_{j=1}^n (\zeta_j d\bar{\zeta}_j - \bar{\zeta}_j d\zeta_j) +$$

$$\frac{(1 - |\zeta_{n+1}|^2) \sinh t}{|\cosh t + \sinh t \zeta_{n+1}|^2} \left(\frac{d\zeta_{n+1}}{\cosh t + \sinh t \zeta_{n+1}} - \frac{d\bar{\zeta}_{n+1}}{\cosh t + \sinh t \bar{\zeta}_{n+1}} \right).$$

On the other hand,

$$f_{n+1} d\bar{f}_{n+1} - \bar{f}_{n+1} df_{n+1} = \frac{1}{|\cosh t + \sinh t \zeta_{n+1}|^2} \left(\frac{\sinh t + \cosh t \zeta_{n+1}}{\cosh t + \sinh t \bar{\zeta}_{n+1}} d\bar{\zeta}_{n+1} - \frac{(\sinh t + \cosh t \bar{\zeta}_{n+1})}{\cosh t + \sinh t \zeta_{n+1}} d\zeta_{n+1} \right)$$

Now (with $|\zeta_{n+1}|^2 = \zeta_{n+1}\bar{\zeta}_{n+1}$),

$$\frac{(1 - |\zeta_{n+1}|^2) \sinh t - (\sinh t + \cosh t \bar{\zeta}_{n+1})}{\cosh t + \sinh t \zeta_{n+1}} = -\bar{\zeta}_{n+1}$$

and

$$\frac{-(1 - |\zeta_{n+1}|^2) \sinh t + (\sinh t + \cosh t \zeta_{n+1})}{\cosh t + \sinh t \bar{\zeta}_{n+1}} = \zeta_{n+1}$$

Thus,

$$\sum_{j=1}^{n+1} (f_j d\bar{f}_j - \bar{f}_j df_j) = \frac{1}{|\cosh t + \sinh t \zeta_{n+1}|^2} \sum_{j=1}^{n+1} (\zeta_j d\bar{\zeta}_j - \bar{\zeta}_j d\zeta_j)$$

that is,

$$(\gamma_t)^* \theta_0 = \frac{i}{2} \sum_{j=1}^{n+1} (f_j d\bar{f}_j - \bar{f}_j df_j) = \frac{1}{|\cosh t + \sinh t \zeta_{n+1}|^2} \theta_0.$$

□

Let $p \in \mathbb{S}^{2n+1}$ be any point of the sphere and let $\alpha_p \in U(n+1)$ be such that $\alpha_p(p) = e_{n+1}$. The family $\gamma_t^p = \alpha_p^{-1} \circ \gamma_t \circ \alpha_p$ is a 1-parameter group of CR-automorphisms of the sphere \mathbb{S}^{2n+1} with

$$\gamma_t^p(\zeta) = \frac{1}{\cosh t + \sinh t (\zeta, p)} \{ \zeta + (\sinh t + (\cosh t - 1) (\zeta, p)) p \} \quad (8)$$

and

$$(\gamma_t^p)^* \theta_0 = \frac{1}{|\cosh t + \sinh t (\zeta, p)|^2} \theta_0. \quad (9)$$

2.3. Preparatory lemmas.

Lemma 2.2. *Let M be a strictly pseudoconvex CR manifold of dimension $2n+1$ and let $\theta, \hat{\theta} \in \mathcal{P}_+(M)$ be two pseudo-Hermitian structures with $\hat{\theta} = f\theta$, $f \in C^\infty(M)$. Then*

$$\psi_{\hat{\theta}} = f^{n+1} \psi_\theta \quad (10)$$

Proof. From $d\hat{\theta} = f d\theta + df \wedge \theta$ we deduce, by induction,

$$(d\hat{\theta})^n = f^n (d\theta)^n + \alpha_n \wedge \theta$$

where α_n is a differential form of degree $2n-1$. Thus,

$$\hat{\theta} \wedge (d\hat{\theta})^n = f\theta \wedge (f^n (d\theta)^n + \alpha_n \wedge \theta) = f^{n+1} \theta \wedge (d\theta)^n.$$

□

Lemma 2.3. *Let M be a strictly pseudoconvex CR manifold of dimension $2m + 1$ and let $\phi : M \rightarrow (\mathbb{S}^{2n+1}, \theta_0)$ be a CR map. Then, for every $\theta \in \mathcal{P}_+(M)$,*

$$\phi^* \theta_0 = \frac{1}{2m} \left(\sum_{i=1}^{2n+2} |\nabla^H \phi_i|_{\theta}^2 \right) \theta \quad (11)$$

where $\phi_1, \dots, \phi_{2n+2}$ are the Euclidean components of ϕ .

Proof. Since ϕ is a CR map, the 1-form $\phi^* \theta_0$ vanishes on $H(M)$ which implies that there exists $f \in C^\infty(M)$ such that

$$\phi^* \theta_0 = f\theta. \quad (12)$$

Differentiating, we get

$$\phi^* d\theta_0 = df \wedge \theta + f d\theta.$$

Hence, for every $X, Y \in H_x(M)$, one has $\phi^* d\theta_0(X, Y) = f d\theta(X, Y)$ and, using (7),

$$\begin{aligned} L_{\theta_0}(d\phi(X), d\phi(X)) &= d\theta_0(d\phi(X), J^{\mathbb{S}^{2n+1}} d\phi(X)) = d\theta_0(d\phi(X), d\phi(J^M X)) \\ &= \phi^* d\theta_0(X, J^M X) = f d\theta(X, J^M X) = f L_{\theta}(X, X). \end{aligned}$$

On the other hand, since L_{θ_0} coincides with the standard inner product on $H_{\phi(x)}(\mathbb{S}^{2n+1})$,

$$L_{\theta_0}(d\phi(X), d\phi(X)) = \sum_{i=1}^{2n+2} (d\phi_i(X))^2 = \sum_{i=1}^{2n+2} L_{\theta}(\nabla^H \phi_i, X)^2.$$

Thus,

$$f L_{\theta}(X, X) = \sum_{i=1}^{2n+2} L_{\theta}(\nabla^H \phi_i, X)^2.$$

Taking an L_{θ} -orthonormal basis $\{e_1, \dots, e_{2m}\}$ of $H_x(M)$, we get

$$2mf = \sum_{j=1}^{2m} \sum_{i=1}^{2n+2} L_{\theta}(\nabla^H \phi_i, e_j)^2 = \sum_{i=1}^{2n+2} L_{\theta}(\nabla^H \phi_i, \nabla^H \phi_i) = \sum_{i=1}^{2n+2} |\nabla^H \phi_i|_{\theta}^2$$

which implies (11), thanks to (12). \square

If $\phi : M \rightarrow \mathbb{R}^N$ is a map and μ is a measure on M , we denote by $\int_M \phi d\mu$ the vector $(\int_M \phi_1 d\mu, \dots, \int_M \phi_N d\mu) \in \mathbb{R}^N$, where $\phi = (\phi_1, \dots, \phi_N)$.

Lemma 2.4. *Let M be a compact manifold and let μ be a measure on M such that no open set has measure zero. If $\phi : M \rightarrow \mathbb{S}^{2n+1}$ is a non constant continuous map, then there exists a pair $(p, t) \in \mathbb{S}^{2n+1} \times [0, +\infty)$ such that*

$$\int_M \gamma_t^p \circ \phi d\mu = 0.$$

Proof. The proof uses standard arguments (see [11, 15]). We consider the map

$$\begin{aligned} F &: \mathbb{S}^{2n+1} \times (0, +\infty) \rightarrow \mathbb{B}^{2n+2} \subset \mathbb{R}^{2n+2} \\ (p, t) &\mapsto \frac{1}{V} \int_M \gamma_t^p \circ \phi d\mu \end{aligned}$$

where $V = \int_M d\mu$ and \mathbb{B}^{2n+2} is the unit Euclidean ball. Observe that (see (8)), $\forall p \in \mathbb{S}^{2n+1}$, γ_0^p is the identity map while, $\forall \zeta \neq -p$, $\gamma_t^p(\zeta)$ tends to p as $t \rightarrow +\infty$. Consequently, $F(\cdot, 0) = \frac{1}{V} \int_M \varphi d\mu$ is a constant map and $F(\cdot, t)$ tends to the identity of \mathbb{S}^{2n+1} as

$t \rightarrow +\infty$. Such a map F is necessarily onto which implies that the origin of \mathbb{R}^{2n+2} belongs to its image. \square

2.4. Proof Theorem 1.1. Let $\theta \in \mathcal{P}_+(\mathbb{S}^{2n+1})$ be a strictly pseudoconvex pseudo-Hermitian structure. We apply Lemma 2.4 to the identity map of \mathbb{S}^{2n+1} and the measure induced by ψ_θ to obtain the existence of a pair $(p, t) \in \mathbb{S}^{2n+1} \times [0, +\infty)$ such that

$$\int_{\mathbb{S}^{2n+1}} \gamma_t^p \psi_\theta = 0.$$

For simplicity, we write γ for γ_t^p . The Euclidean components $\gamma_1, \dots, \gamma_{2n+2}$ of γ satisfy $\int_{\mathbb{S}^{2n+1}} \gamma_j \psi_\theta = 0$. Hence, applying the min-max principle (5), we get for every $j \leq 2n+2$,

$$\lambda_1(\theta) \int_{\mathbb{S}^{2n+1}} \gamma_j^2 \psi_\theta \leq \int_{\mathbb{S}^{2n+1}} |\nabla^H \gamma_j|_\theta^2 \psi_\theta.$$

Summing up we obtain, with $\sum_{j \leq 2n+2} \gamma_j^2 = 1$,

$$\lambda_1(\theta) V(\theta) \leq \sum_{j \leq 2n+2} \int_{\mathbb{S}^{2n+1}} |\nabla^H \gamma_j|_\theta^2 \psi_\theta \quad (13)$$

$$\leq 2n \left(\int_{\mathbb{S}^{2n+1}} \left(\frac{1}{2n} |\nabla^H \gamma_j|_\theta^2 \right)^{n+1} \psi_\theta \right)^{\frac{1}{n+1}} V(\theta)^{1-\frac{1}{n+1}}. \quad (14)$$

Using Lemma 2.3, we see that, since γ is a CR map from $(\mathbb{S}^{2n+1}, \theta)$ to $(\mathbb{S}^{2n+1}, \theta_0)$,

$$\gamma^* \theta_0 = \left(\frac{1}{2n} \sum_{j \leq 2n+2} |\nabla^H \gamma_j|_\theta^2 \right) \theta \quad (15)$$

which gives, thanks to Lemma 2.2

$$\psi_{\gamma^* \theta_0} = \left(\frac{1}{2n} \sum_{j \leq 2n+2} |\nabla^H \gamma_j|_\theta^2 \right)^{n+1} \psi_\theta.$$

Thus

$$\lambda_1(\theta) V(\theta) \leq 2n V(\gamma^* \theta_0)^{\frac{1}{n+1}} V(\theta)^{1-\frac{1}{n+1}}.$$

Since $V(\gamma^* \theta_0) = V(\theta_0)$, we finally get

$$\lambda_1(\theta) V(\theta)^{\frac{1}{n+1}} \leq 2n V(\theta_0)^{\frac{1}{n+1}}$$

which proves the inequality of the theorem thanks to (6).

Now, if the equality holds in (3), this means that we have equality in the Cauchy-Schwarz inequality used in (14). Thus, $\sum_{j \leq 2n+2} |\nabla^H \gamma_j|_\theta^2$ is constant and, thanks to (15), θ is proportional to $\gamma^* \theta_0$ which takes the form given by (9).

Conversely, it is clear that when γ is a CR-automorphism of the sphere then $\lambda_1(\gamma^* \theta_0) = \lambda_1(\theta_0) = 2n$ and $V(\gamma^* \theta_0) = V(\theta_0)$. Hence, the equality holds in (3).

REFERENCES

- [1] Amine Aribi, Sorin Dragomir, and Ahmad El Soufi. On the continuity of the eigenvalues of a sublaplacian. Canad. Math. Bull., 57(1):12–24, 2014.
- [2] Amine Aribi, Sorin Dragomir, and Ahmad El Soufi. Eigenvalues of the sub-Laplacian and deformations of contact structures on a compact CR manifold. Differential Geom. Appl., 39:113–128, 2015.
- [3] Amine Aribi, Sorin Dragomir, and Ahmad El Soufi. A lower bound on the spectrum of the sublaplacian. J. Geom. Anal., 25(3):1492–1519, 2015.
- [4] Amine Aribi and Ahmad El Soufi. Inequalities and bounds for the eigenvalues of the sub-Laplacian on a strictly pseudoconvex CR manifold. Calc. Var. Partial Differential Equations, 47(3-4):437–463, 2013.
- [5] Elisabetta Barletta. The Lichnerowicz theorem on CR manifolds. Tsukuba J. Math., 31(1):77–97, 2007.
- [6] Elisabetta Barletta and Sorin Dragomir. On the spectrum of a strictly pseudoconvex CR manifold. Abh. Math. Sem. Univ. Hamburg, 67:33–46, 1997.
- [7] Elisabetta Barletta and Sorin Dragomir. Sublaplacians on CR manifolds. Bull. Math. Soc. Sci. Math. Roumanie (N.S.), 52(100)(1):3–32, 2009.
- [8] Der-Chen Chang and Song-Ying Li. A zeta function associated to the sub-Laplacian on the unit sphere in \mathbb{C}^n . J. Anal. Math., 86:25–48, 2002.
- [9] Shu-Cheng Chang and Hung-Lin Chiu. On the CR analogue of Obata’s theorem in a pseudohermitian 3-manifold. Math. Ann., 345(1):33–51, 2009.
- [10] Michael G. Cowling, Oldrich Klima, and Adam Sikora. Spectral multipliers for the Kohn sublaplacian on the sphere in \mathbb{C}^n . Trans. Amer. Math. Soc., 363(2):611–631, 2011.
- [11] Ahmad El Soufi and Said Ilias. Le volume conforme et ses applications d’après Li et Yau. In Séminaire de Théorie Spectrale et Géométrie, Année 1983–1984, pages VII.1–VII.15. Univ. Grenoble I, Saint-Martin-d’Hères, 1984.
- [12] Ahmad El Soufi and Said Ilias. Immersions minimales, première valeur propre du laplacien et volume conforme. Math. Ann., 275(2):257–267, 1986.
- [13] Allan Greenleaf. The first eigenvalue of a sub-Laplacian on a pseudo-Hermitian manifold. Comm. Partial Differential Equations, 10(2):191–217, 1985.
- [14] Asma Hassannezahed and Gerasim Kokarev. Sub-laplacian eigenvalue bounds on sub-riemannian manifolds. Ann. Sc. Norm. Super. Pisa (to appear).
- [15] Joseph Hersch. Quatre propriétés isopérimétriques de membranes sphériques homogènes. C. R. Acad. Sci. Paris Sér. A-B, 270:A1645–A1648, 1970.
- [16] Stefan Ivanov and Dimiter Vassilev. An Obata type result for the first eigenvalue of the sub-Laplacian on a CR manifold with a divergence-free torsion. J. Geom., 103(3):475–504, 2012.
- [17] Stefan Ivanov and Dimiter Vassilev. An Obata-type theorem on a three-dimensional CR manifold. Glasg. Math. J., 56(2):283–294, 2014.
- [18] Gerasim Kokarev. Sub-Laplacian eigenvalue bounds on CR manifolds. Comm. Partial Differential Equations, 38(11):1971–1984, 2013.
- [19] Song-Ying Li and Hing-Sun Luk. The sharp lower bound for the first positive eigenvalue of a sub-Laplacian on a pseudo-Hermitian manifold. Proc. Amer. Math. Soc., 132(3):789–798 (electronic), 2004.
- [20] Pengcheng Niu and Huiqing Zhang. Payne-Polya-Weinberger type inequalities for eigenvalues of nonelliptic operators. Pacific J. Math., 208(2):325–345, 2003.
- [21] Raphaël S. Ponge. Heisenberg calculus and spectral theory of hypoelliptic operators on Heisenberg manifolds. Mem. Amer. Math. Soc., 194(906):viii+ 134, 2008.
- [22] Nancy K. Stanton. Spectral invariants of CR manifolds. Michigan Math. J., 36(2):267–288, 1989.

UNIVERSITÉ DE TOURS, LABORATOIRE DE MATHÉMATIQUES ET PHYSIQUE THÉORIQUE, UMR-CNRS 7350, PARC DE GRANDMONT, 37200 TOURS, FRANCE.

E-mail address: Amine.Aribi@lmpt.univ-tours.fr, ahmad.elsoufi@lmpt.univ-tours.fr