



HAL
open science

An analog of Chang inversion formula for weighted Radon transforms in multidimensions

Fedor Goncharov, Roman Novikov

► **To cite this version:**

Fedor Goncharov, Roman Novikov. An analog of Chang inversion formula for weighted Radon transforms in multidimensions. 2016. hal-01304312v1

HAL Id: hal-01304312

<https://hal.science/hal-01304312v1>

Preprint submitted on 19 Apr 2016 (v1), last revised 17 May 2016 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

An analog of Chang inversion formula for weighted Radon transforms in multidimensions^{*}

Fedor O. Goncharov^{1,2}, Roman G. Novikov³

¹ Moscow Institute of Physics and Technology,

9 Institutskiy per., Dolgoprudny, Moscow Region, 141700, Russian Federation

² Institute for Information Transmission Problems (RAS),

Bolshoy Karetny per. 19, buld.1, Moscow 127051, Russian Federation

³ CNRS (UMR 7641), Centre de Mathématiques Appliquées, Ecole Polytechnique,
91128 Palaiseau, France

fedor.goncharov.01@gmail.com, novikov@cmap.polytechnique.fr

Abstract. We study weighted Radon transforms in multidimensions. Our results consist in introducing an analog of Chang approximate inversion formula for such transforms and in describing all weights for which this formula is exact.

Keywords: weighted Radon transforms, inversion formulas

1 Introduction

We consider the weighted Radon transforms R_W defined by the formula

$$R_W f(s, \theta) \stackrel{\text{def}}{=} \int_{x\theta=s} W(x, \theta) f(x) dx, \quad (1)$$
$$(s, \theta) \in \mathbb{R} \times \mathbb{S}^{n-1}, x \in \mathbb{R}^n, n \geq 2,$$

where $W = W(x, \theta)$ is the weight, $f = f(x)$ is a test function; see e.g. [1]. Such transforms arise in many domains of pure and applied mathematics; see e.g. [4], [5], [6], [8], [10], [12]. In the present work we assume that

$$W \text{ is complex - valued,}$$
$$W \in C(\mathbb{R}^n \times \mathbb{S}^{n-1}) \cap L^\infty(\mathbb{R}^n \times \mathbb{S}^{n-1}), \quad (2)$$
$$w_0(x) \stackrel{\text{def}}{=} \frac{1}{|\mathbb{S}^{n-1}|} \int_{\mathbb{S}^{n-1}} W(x, \theta) d\theta \neq 0, x \in \mathbb{R}^n,$$

where $d\theta$ is the element of standard measure on \mathbb{S}^{n-1} , $|\mathbb{S}^{n-1}|$ is the standard measure of \mathbb{S}^{n-1} .

^{*} The main part of the work was fulfilled during the stage of the first author in the Centre de Mathématiques Appliquées of Ecole Polytechnique in March-May 2016

If $W \equiv 1$, then $R = R_W$ is the classical Radon transform in \mathbb{R}^n ; see for example [6], [9], [11], [15]. Explicit inversion formulas for R were given for the first time in [15].

In dimension $n = 2$, the transforms R_W are also known as weighted ray transforms on the plane; see e.g. [10], [12]. For several important cases of W satisfying (2) for $d = 2$, explicit (and exact) inversion formulas for R_W were obtained in [2], [7], [13], [14], [16].

On the other hand, it seems that no explicit inversion formulas for R_W were given yet in the literature under assumptions (2) for $n \geq 3$, if $W \neq w_0$.

The main results of the present work consist in introducing an analog of Chang approximate (but explicit) inversion formula for R_W under assumptions (2), for $n \geq 3$, and in describing all W for which this formula is exact. These results are presented in the next section.

2 Main Results

We consider the following approximate inversion formulas for R_W under assumptions (2) in dimension $n \geq 2$:

$$f_{appr}(x) \stackrel{def}{=} \frac{(-1)^{(n-2)/2}}{2(2\pi)^{n-1}w_0(x)} \int_{\mathbb{S}^{n-1}} \mathbb{H} [R_W f]^{(n-1)}(x\theta, \theta) d\theta, \quad (3)$$

$x \in \mathbb{R}^n$, n is even,

$$f_{appr}(x) \stackrel{def}{=} \frac{(-1)^{(n-1)/2}}{2(2\pi)^{n-1}w_0(x)} \int_{\mathbb{S}^{n-1}} [R_W f]^{(n-1)}(x\theta, \theta) d\theta, \quad (4)$$

$x \in \mathbb{R}^n$, n is odd,

$$[R_W f]^{(n-1)}(s, \theta) = \frac{d^{n-1}}{ds^{n-1}} R_W f(s, \theta), \quad s \in \mathbb{R}, \theta \in \mathbb{S}^{n-1}, \quad (5)$$

$$\mathbb{H}\phi(s) \stackrel{def}{=} \frac{1}{\pi} p.v. \int_{\mathbb{R}} \frac{\phi(t)}{s-t} dt, \quad s \in \mathbb{R}. \quad (6)$$

For $W \equiv 1$ formulas (3), (4) are exact, i.e. $f_{appr} = f$, and are known as the classical Radon inversion formulas, going back to [15].

As a corollary of the classical Radon inversion formulas and definition (1), formulas (3), (4) for $W \equiv w_0$ are also exact.

Formula (3) for $n = 2$ is known as Chang approximate inversion formula for weighted Radon transforms on the plane. This explicit but approximate inversion formula was suggested for the first time in [3] for the case when

$$W(x, \theta) = \exp(-Da(x, \theta^\perp)), \quad (7)$$

$$Da(x, \theta^\perp) = \int_0^{+\infty} a(x + t\theta^\perp) dt, \quad (8)$$

where a is a non-negative sufficiently regular function on \mathbb{R}^2 with compact support, and $\theta = (\theta_1, \theta_2) \in \mathbb{S}^{n-1}$, $\theta^\perp = (\theta_2, -\theta_1)$. We recall that R_W for W given by (7), (8) is known as attenuated Radon transform on the plane and arises, in particular, in the single photon emission tomography (SPECT). In this case an explicit and simultaneously exact inversion formula for R_W was obtained for the first time in [13].

We emphasize that formulas (3), (4) are approximate, in general. In addition, the following result holds:

Theorem 1. *Let W satisfy (2). Let f_{appr} be defined by (3), (4) in terms of $R_W f$ and w_0 , $n \geq 2$. Then $f_{appr} = f$ (in the sense of distributions) on \mathbb{R}^n for all $f \in C_0(\mathbb{R}^n)$ if and only if*

$$W(x, \theta) - w_0(x) \equiv w_0(x) - W(x, -\theta), \quad x \in \mathbb{R}^n, \theta \in \mathbb{S}^{n-1}. \quad (9)$$

Here $C_0(\mathbb{R}^n)$ denotes the space of all continuous compactly supported functions on \mathbb{R}^n .

The result of Theorem 1 for $n = 2$ was obtained for the first time in [14]. Theorem 1 in the general case is proved in Section 3.

If W satisfy (2), $f \in C_0(\mathbb{R}^n)$, but the the symmetry condition (9) does not hold, i.e.

$$w_0(x) \neq \frac{1}{2} (W(x, \theta) + W(x, -\theta)), \quad \text{for some } x \in \mathbb{R}^n, \theta \in \mathbb{S}^{n-1},$$

then (3), (4) can be considered as approximate formulas for finding f from $R_W f$.

3 Proof of Theorem 1

For W satisfying (2) we also consider its symmetrization defined by

$$W_s(x, \theta) \stackrel{def}{=} \frac{1}{2} (W(x, \theta) + W(x, -\theta)), \quad x \in \mathbb{R}^n, \theta \in \mathbb{S}^{n-1}. \quad (10)$$

Using definitions (1), (10) we obtain

$$R_{W_s} f(s, \theta) = \frac{1}{2} (R_W f(s, \theta) + R_W f(-s, -\theta)). \quad (11)$$

In addition, if W satisfies (9) then

$$W_s(x, \theta) = w_0(x), \quad x \in \mathbb{R}^n, \theta \in \mathbb{S}^{n-1}. \quad (12)$$

3.1 Proof of sufficiency

The sufficiency of symmetry (9) follows from formulas (3), (4) for the exact case with $W \equiv w_0$, the identities

$$f_{appr}(x) = \frac{(-1)^{(n-2)/2}}{2(2\pi)^{n-1}w_0(x)} \int_{\mathbb{S}^{n-1}} \mathbb{H}[R_{W_s}f]^{(n-1)}(x\theta, \theta)d\theta, \quad (13)$$

for even n ,

$$f_{appr}(x) = \frac{(-1)^{(n-1)/2}}{2(2\pi)^{n-1}w_0(x)} \int_{\mathbb{S}^{n-1}} [R_{W_s}f]^{(n-1)}(x\theta, \theta)d\theta, \quad (14)$$

for odd n ,

and from the identities (11), (12).

In turn, (13) follows from the identities

$$\begin{aligned} & \int_{\mathbb{S}^{n-1}} \mathbb{H}[R_W f]^{(n-1)}(x\theta, \theta)d\theta \\ &= \frac{1}{2} \int_{\mathbb{S}^{n-1}} \left(\mathbb{H}[R_W f]^{(n-1)}(x\theta, \theta) + \mathbb{H}[R_W f]^{(n-1)}(-x\theta, -\theta) \right) d\theta \\ &= \int_{\mathbb{S}^{n-1}} \mathbb{H}[R_{W_s} f]^{(n-1)}(x\theta, \theta)d\theta. \end{aligned} \quad (15)$$

In addition, the second of the identities of (15) follows from the identities:

$$\begin{aligned} \mathbb{H}[R_{W_s}f]^{(n-1)}(s, \theta) &= \frac{1}{2\pi} p.v. \int_{\mathbb{R}} \frac{1}{s-t} \times \\ & \times \frac{d^{n-1}}{dt^{n-1}} \left[R_W f(t, \theta) + R_W f(-t, -\theta) \right] dt \\ &= \frac{1}{2} \mathbb{H}[R_W f]^{(n-1)}(s, \theta) + \frac{(-1)^{n-1}}{2\pi} p.v. \int_{\mathbb{R}} \frac{[R_W f]^{(n-1)}(-t, -\theta)}{s-t} dt; \quad (16) \\ \frac{(-1)^{n-1}}{\pi} p.v. \int_{\mathbb{R}} \frac{[R_W f]^{(n-1)}(-t, -\theta)}{s-t} dt &= -\frac{(-1)^{n-1}}{\pi} p.v. \int_{\mathbb{R}} \frac{[R_W f]^{(n-1)}(t, -\theta)}{-s-t} dt \\ &= (-1)^n \mathbb{H}[R_W f]^{(n-1)}(-s, -\theta) = \mathbb{H}[R_W f]^{(n-1)}(-s, -\theta). \quad (17) \end{aligned}$$

This concludes the proof of sufficiency for n even.

Finally, (14) follows from the identities

$$\begin{aligned} & \int_{\mathbb{S}^{n-1}} [R_W f]^{(n-1)}(x\theta, \theta) d\theta \\ &= \frac{1}{2} \int_{\mathbb{S}^{n-1}} \left([R_W f]^{(n-1)}(x\theta, \theta) + [R_W f]^{(n-1)}(-x\theta, -\theta) \right) d\theta, \end{aligned} \quad (18)$$

$$\begin{aligned} [R_{W_s} f]^{(n-1)}(t, \theta) &= \frac{1}{2} \frac{d^{n-1}}{dt^{n-1}} \left[[R_W f](t, \theta) + [R_W f](-t, -\theta) \right] \\ &= \frac{1}{2} \left[[R_W f]^{(n-1)}(t, \theta) + (-1)^{n-1} [R_W f]^{(n-1)}(-t, -\theta) \right] \\ &= \frac{1}{2} \left[[R_W f]^{(n-1)}(t, \theta) + [R_W f]^{(n-1)}(-t, -\theta) \right]. \end{aligned} \quad (19)$$

This concludes the proof of sufficiency for odd n .

3.2 Proof of necessity

Using that $f_{appr} = f$ for all $f \in C_0(\mathbb{R}^n)$ and using formulas (3), (4) for the exact case $W \equiv w_0$, we obtain

$$\int_{\mathbb{S}^{n-1}} \left(\mathbb{H} [R_W f]^{(n-1)}(x\theta, \theta) - \mathbb{H} [R_{w_0} f]^{(n-1)}(x\theta, \theta) \right) d\theta = 0 \quad (20)$$

on \mathbb{R}^n for even n ,

$$\int_{\mathbb{S}^{n-1}} [R_W f - R_{w_0} f]^{(n-1)}(x\theta, \theta) d\theta = 0 \quad (21)$$

on \mathbb{R}^n for odd n ,

for all $f \in C_0(\mathbb{R}^n)$.

Identities (15), (18), (19), (20), (21) imply the identities

$$\int_{\mathbb{S}^{n-1}} \left(\mathbb{H} [R_{W_s} f]^{(n-1)}(x\theta, \theta) - \mathbb{H} [R_{w_0} f]^{(n-1)}(x\theta, \theta) \right) d\theta = 0 \quad (22)$$

on \mathbb{R}^n for even n ,

$$\int_{\mathbb{S}^{n-1}} [R_{W_s} f - R_{w_0} f]^{(n-1)}(x\theta, \theta) d\theta = 0 \quad (23)$$

on \mathbb{R}^n for odd n ,

for all $f \in C_0(\mathbb{R}^n)$.

The necessity of symmetry (9) follows from the identities (22), (23) and the following lemmas:

Lemma 1. Let (22), (23) be valid for fixed $f \in C_0(\mathbb{R}^n)$ and W satisfying (2), $n \geq 2$. Then

$$R_{W_s} f = R_{w_0} f. \quad (24)$$

Lemma 2. Let (24) be valid for all $f \in C_0(\mathbb{R}^n)$ and fixed W satisfying (2), $n \geq 2$. Then

$$W_s = w_0. \quad (25)$$

Lemmas 1 and 2 are proved in Sections 4 and 5.

4 Proof of Lemma 1

We will use the following formulas

$$\begin{aligned} & \int_{\mathbb{R}^n} e^{i\xi x} \int_{\mathbb{S}^{n-1}} g(x\theta, \theta) d\theta dx \\ &= \frac{\sqrt{2\pi}}{|\xi|^{n-1}} \left(\hat{g} \left(|\xi|, \frac{\xi}{|\xi|} \right) + \hat{g} \left(-|\xi|, -\frac{\xi}{|\xi|} \right) \right), \end{aligned} \quad (26)$$

$$\begin{aligned} & \int_{\mathbb{R}^n} e^{i\xi x} \int_{\mathbb{S}^{n-1}} g^{(n-1)}(x\theta, \theta) d\theta dx = \int_{\mathbb{R}^n} e^{i\xi x} \int_{\mathbb{S}^{n-1}} (\theta \nabla_x)^{n-1} g(x\theta, \theta) d\theta dx \\ &= (-i)^{n-1} \sqrt{2\pi} \left(\hat{g} \left(|\xi|, \frac{\xi}{|\xi|} \right) + (-1)^{n-1} \hat{g} \left(-|\xi|, -\frac{\xi}{|\xi|} \right) \right), \end{aligned} \quad (27)$$

$$\hat{g}(\tau, \theta) = \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} e^{i\tau s} g(s, \theta) ds, \quad \tau \in \mathbb{R}, \theta \in \mathbb{S}^{n-1}, \quad (28)$$

where $g \in C(\mathbb{S}^{n-1}, L^2(\mathbb{R}))$, $\xi \in \mathbb{R}^n$. The validity of formulas (26), (27) (in the sense of distributions) follows from Theorem 1.4 of [12].

4.1 The case of odd n

Using identity (11) we get

$$g(s, \theta) = g(-s, -\theta), \quad \text{for all } s \in \mathbb{R}, \theta \in \mathbb{S}^{n-1}, \quad (29)$$

where

$$g(s, \theta) = [R_{W_s} f(s, \theta) - R_{w_0} f(s, \theta)]. \quad (30)$$

From (29), we obtain the same symmetry for the Fourier transform $\hat{g}(\cdot, \theta)$ of $g(\cdot, \theta)$:

$$\begin{aligned} \hat{g}(t, \theta) &= \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} g(s, \theta) e^{its} ds \\ &= \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} g(-s, -\theta) e^{i(-s)(-t)} ds \\ &= \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} g(s, -\theta) e^{-its} ds = \hat{g}(-t, -\theta), \quad t \in \mathbb{R}, \theta \in \mathbb{S}^{n-1}. \end{aligned} \quad (31)$$

For odd n , from identities (23), (27) it follows that

$$\hat{g}\left(|p|, \frac{p}{|p|}\right) + \hat{g}\left(-|p|, -\frac{p}{|p|}\right) = 0 \text{ in } L_{loc}^2(\mathbb{R}^n). \quad (32)$$

Using (31), (32) we obtain

$$\begin{cases} \hat{g}\left(|p|, \frac{p}{|p|}\right) = 0, \\ \hat{g}\left(-|p|, -\frac{p}{|p|}\right) = 0 \end{cases} \Leftrightarrow \hat{g} = 0 \Leftrightarrow g = 0. \quad (33)$$

Formula (24) for odd n follows from (30), (33).

4.2 The case of even n

We consider

$$g(s, \theta) = \mathbb{H}[R_{W_s}f - R_{w_0}f](s, \theta), \quad s \in \mathbb{R}, \theta \in \mathbb{S}^{n-1}, \quad (34)$$

arising in (22). Using the identity

$$\begin{aligned} \mathbb{H}[R_{W_s}f - R_{w_0}f](-s, -\theta) &= \frac{1}{\pi} p.v. \int_{\mathbb{R}} \frac{R_{W_s}f(t, -\theta) - R_{w_0}f(t, -\theta)}{-s - t} dt \\ &= \frac{1}{\pi} p.v. \int_{\mathbb{R}} \frac{R_{W_s}f(-t, -\theta) - R_{w_0}f(-t, -\theta)}{-s + t} dt \\ &= -\frac{1}{\pi} p.v. \int_{\mathbb{R}} \frac{R_{W_s}f(t, \theta) - R_{w_0}f(t, \theta)}{s - t} dt = -\mathbb{H}[R_{W_s}f - R_{w_0}f](s, \theta), \end{aligned} \quad (35)$$

we obtain

$$g(s, \theta) = -g(-s, -\theta), \quad \text{for all } s \in \mathbb{R}, \theta \in \mathbb{S}^{n-1}. \quad (36)$$

From (36), similarly with (31), we obtain the same symmetry for the Fourier transform $\hat{g}(\cdot, \theta)$ of $g(\cdot, \theta)$:

$$\hat{g}(t, \theta) = -\hat{g}(-t, -\theta), \quad t \in \mathbb{R}, \theta \in \mathbb{S}^{n-1}. \quad (37)$$

For n even, from the property of the Hilbert transform

$$\mathbb{H}[\phi^{(k)}] = (\mathbb{H}[\phi])^{(k)}, \quad \phi \in C_0^k(\mathbb{R}),$$

where this identity holds in the sense of distributions if $\phi \in C_0(\mathbb{R})$, and identities (22), (27) it follows that

$$\hat{g}\left(|p|, \frac{p}{|p|}\right) - \hat{g}\left(-|p|, -\frac{p}{|p|}\right) = 0 \text{ in } L_{loc}^2(\mathbb{R}^n). \quad (38)$$

Using (37), (38) we again obtain (33) but already for even n . Due to (33), (34) we have

$$\mathbb{H}[R_{W_s}f - R_{w_0}f] = 0. \quad (39)$$

Formula (24) for even n follows from (39), invertibility of the Hilbert transform on L^p , $p > 1$ and the fact that $R_W f \in C_0(\mathbb{R} \times \mathbb{S}^{n-1})$.

Lemma 1 is proved.

5 Proof of Lemma 2

Suppose that

$$W_s(y, \theta) - w_0(y) = z \neq 0 \quad (40)$$

for some $y \in \mathbb{R}^n$, $\theta \in \mathbb{S}^{n-1}$, $z \in \mathbb{C}$. Since W satisfies (2), then for any $\varepsilon > 0$ there exists $\delta(\varepsilon) > 0$ such that

$$\forall y' : |y' - y| < \delta \rightarrow |W_s(y', \theta) - w_0(y') - z| < \varepsilon, \quad (41)$$

for fixed y, θ .

Let $f \in C_0(\mathbb{R}^n)$, $f \geq 0$ and satisfies the conditions

$$f(y') \equiv 1, y' \in B_{\delta/2}(y), \quad (42)$$

$$\text{supp } f \subset B_{\delta}(y), \quad (43)$$

where $B_{\delta}(y)$ is the open ball with radius δ , centered at y , $\delta = \delta(\varepsilon)$, $0 < \varepsilon < |z|$. It suffices to show that

$$|R_{W_s}f(y\theta, \theta) - R_{w_0}f(y\theta, \theta)| > 0, \quad (44)$$

which contradicts the condition of the lemma.

The identity (44) follows from the formulas

$$\begin{aligned} |R_{W_s}f(y\theta, \theta) - R_{w_0}f(y\theta, \theta)| &= \left| \int_{x\theta=y\theta} f(x)(W_s(x, \theta) - w_0(x))dx \right| \\ &= \left| \int_{x\theta=y\theta} f(x)(W_s(x, \theta) - w_0(x) - z)dx + z \int_{x\theta=y\theta} f(x)dx \right| \\ &\geq |z| \int_{x\theta=y\theta} f(x)dx - \int_{x\theta=y\theta} f(x)|W_s(x, \theta) - w_0(x) - z|dx \\ &\geq (|z| - \varepsilon) \int_{x\theta=y\theta} f(x)dx > 0, \text{ for } 0 < \varepsilon < |z|. \end{aligned} \quad (45)$$

Lemma 2 is proved.

References

1. J. Boman and E. T. Quinto. *Support theorems for real-analytic Radon transforms*. Duke Mathematical J., 55(4):943–948, 1987.
2. J. Boman and J. Strömberg. Novikov’s inversion formula for the attenuated Radon transform — a new approach. *The Journal of Geometric Analysis*, 14(2):185–198, 2004.
3. L. Chang. A method for attenuation correction in radionuclide computed tomography. *Nuclear Science, IEEE Transactions on*, 25(1):638–643, 1978.
4. S. R. Deans. *The Radon transform and some of its applications*. Courier Corporation, 2007.
5. T. Durrani and D. Bisset. The Radon transform and its properties. *Geophysics*, 49(8):1180–1187, 1984.
6. I. M. Gel’fand, M. I. Graev, and N. Ya. Vilenkin. *Integral geometry and representation theory*, volume 5. Academic press, 2014.
7. S. Gindikin. A remark on the weighted Radon transform on the plane. *Inverse Problems and Imaging*, 4:649–653, 2010.
8. P. Grangeat. Mathematical framework of cone beam 3D reconstruction via the first derivative of the Radon transform. In *Mathematical methods in tomography*, pages 66–97. Springer, 1991.
9. S. Helgason. *The Radon Transform on R^n* . Springer, 2011.
10. L. A. Kunyansky. Generalized and attenuated Radon transforms: restorative approach to the numerical inversion. *Inverse Problems*, 8(5):809, 1992.
11. D. Ludwig. The Radon transform on Euclidean space. *Communications on Pure and Applied Mathematics*, 19(1):49–81, 1966.
12. F. Natterer. *The mathematics of computerized tomography*, volume 32. Siam, 1986.
13. R. G. Novikov. An inversion formula for the attenuated X-ray transformation. *Arkiv för matematik*, 40(1):145–167, 2002.
14. R. G. Novikov. Weighted Radon transforms for which Chang’s approximate inversion formula is exact. *Russian Mathematical Surveys*, 66(2):442–443, 2011.
15. J. Radon. Über die Bestimmung von Funktionen durch ihre Integralwerte längs gewisser Mannigfaltigkeiten [on the determination of functions by their integral values along certain manifolds], 75 years of Radon transform (Vienna, 1992), conf. *Proc. Lecture Notes Math. Phys., IV, Int. Press, Cambridge, MA*, pages 324–339, 1994.
16. O. Tretiak and C. Metz. The exponential Radon transform. *SIAM Journal on Applied Mathematics*, 39(2):341–354, 1980.