



HAL
open science

Shifted convolution of cusp-forms with θ -series

Guangshi Lu, Jie Wu, Wenguang Zhai

► **To cite this version:**

Guangshi Lu, Jie Wu, Wenguang Zhai. Shifted convolution of cusp-forms with θ -series. Ramanujan Journal (The), 2016, 40 (1), pp.115-133. 10.1007/s11139-015-9678-8 . hal-01313346

HAL Id: hal-01313346

<https://hal.science/hal-01313346>

Submitted on 9 May 2016

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

SHIFTED CONVOLUTION OF CUSP-FORMS WITH θ -SERIES

GUANGSHI LÜ, JIE WU & WENGUANG ZHAI

ABSTRACT. In this paper we apply simple approaches to improve a recent result due to Luo, concerning a shifted convolution sum involving the Fourier coefficients of cusp forms with those of theta series. In order to explore what should be the best possible bound, a mean square result of this shifted convolution sum is also established.

1. INTRODUCTION

Suppose $\lambda_1(n)$ and $\lambda_2(n)$ are two arithmetic functions, $b \geq 0$ is an integer. It is a classical and important problem in analytic number theory to study the shifted convolution sum

$$\sum_{n \leq x} \lambda_1(n) \lambda_2(n+b).$$

There are a large number of papers in this direction. For example, when $\lambda_1(n) = \lambda_2(n) = \tau(n)$, the Dirichlet divisor function, this problem is the so-called additive divisor problem [1, 6, 8, 9, 12, 18, 19]. For other interesting cases, see [14, 15, 7].

Recently Luo [16] investigated a mixed shifted convolution sum

$$(1.1) \quad \sum_{n \geq 1} \lambda_f(n+b) r_\ell(n) \phi(n),$$

where $\lambda_f(n)$ is the n th normalized Hecke eigenvalue of a holomorphic cusp form of weight k and level N , $b \geq 0$ is a fixed integer,

$$r_\ell(n) := |\{(n_1, \dots, n_\ell) \in \mathbb{Z}^\ell : n_1^2 + \dots + n_\ell^2 = n\}|$$

and $\phi(t)$ is a smooth function with support in $[x/2, 5x/2]$, satisfying $\phi^{(j)}(t) \ll_j (x/J)^{-j}$ for all real numbers $t \in \mathbb{R}$, all integers $j \geq 0$ and $1 \leq J \leq x^\beta$ with $\beta < 1$. In order to estimate the sum (1.1), he first established a Voronoi formula for $r_\ell(n)$ and then combined this formula with the upper bound for the Salié sum to derive

Date: December 3, 2013.

2000 Mathematics Subject Classification. 11F30, 11F11, 11F27, 11F37.

Key words and phrases. shifted convolution, cusp form, theta series.

Guangshi Lü is supported in part by the National Natural Science Foundation of China (11031004, 11171182), NCET (NCET-10-0548) and Shandong Province Natural Science Foundation for distinguished Young Scholars (JQ201102). Wenguang Zhai is supported by the National Key Basic Research Program of China (Grant No. 2013CB834201), the Natural Science Foundation of China (Grant No. 11171344) and the Natural Science Foundation of Beijing (Grant No. 1112010).

the following result [16, Theorem] : For any $\varepsilon > 0$, the inequality

$$(1.2) \quad \sum_{n \geq 1} \lambda_f(n+b)r_\ell(n)\phi(n) \ll \begin{cases} x^{\ell/2-(\ell-1)/4+\varepsilon} J^{g_\ell} & \text{if } 4 \nmid \ell \text{ or } b \neq 0 \\ x^{\ell/2-(\ell-1)/4+\varepsilon}(x^{1/4+\varepsilon} + J^{g_\ell}) & \text{if } 4 \mid \ell \text{ and } b = 0 \end{cases}$$

holds for all $x \geq 2$, where g_ℓ is the smallest integer such that $g_\ell \geq (\ell+1)/2$ [†] and the implied constant depends on f , b , ℓ and ε . With a suitable choice of $\phi(t)$ and J , the inequality (1.2) implies immediately the following result [16, Corollary] : For $\ell \geq 2$, $k \geq \ell/2 + 3$ and $\varepsilon > 0$, the inequality

$$(1.3) \quad \mathcal{S}_{f,b,\ell}(x) := \sum_{n \leq x} \lambda_f(n+b)r_\ell(n) \ll_{f,b,\ell,\varepsilon} x^{\ell/2-\vartheta_\ell+\varepsilon}$$

holds for all $x \geq 2$, where $\vartheta_\ell := (\ell-1)/(4g_\ell+4)$. In particular

$$\vartheta_2 = \frac{1}{12}, \quad \vartheta_3 = \frac{1}{6}, \quad \vartheta_4 = \frac{3}{16}, \quad \vartheta_5 = \frac{1}{4}, \quad \vartheta_6 = \frac{1}{4}, \quad \vartheta_\ell \leq \frac{\ell-1}{2\ell+6} < \frac{1}{2} \quad (\ell \geq 7).$$

The first aim of this paper is to propose a better bound for $\mathcal{S}_{f,b,\ell}(x)$.

Theorem 1. *Let f be a cusp form of weight k and level N and let $\ell \geq 2$ be an integer. For any $\varepsilon > 0$, we have*

$$(1.4) \quad \mathcal{S}_{f,b,\ell}(x) := \sum_{n \leq x} \lambda_f(n+b)r_\ell(n) \ll_{f,\ell,\varepsilon} x^{\ell/2-\vartheta_\ell+\varepsilon}$$

uniformly for $x \geq 2$ and $0 \leq b \leq x$, where

$$\vartheta_3 = \frac{1}{4}, \quad \vartheta_4 = \frac{1}{2}, \quad \vartheta_5 = \frac{1}{2}, \quad \vartheta_\ell = \frac{2}{3} \quad (\ell \geq 6).$$

In addition, if we assume $N = 1$, then (1.4) holds for $\ell = 2$ with $\vartheta_2 = \frac{1}{6}$.

Remark 1. Theorem 1 improves Luo's (1.3) in three directions; it enlarges the exponent ϑ_ℓ , relaxes the restricted condition $k \geq \ell/2 + 3$ and removes the dependence of b . Our method is completely different from that of [16]. Our new idea is to explore the regularity of $r_\ell(n)$ by the circle method in analytic number theory and Siegel's mass formula. Thanking to these classic tools of analytic number theory, we can show that the influence of $r_\ell(n)$ to the bound [22, (1.10) and Theorem 2]

$$(1.5) \quad \sum_{n \leq x} \lambda_f(n) \ll_f x^{1/3}(\log x)^{2/(\sqrt{\pi}\Gamma(5/2))-1}$$

is rather little. More precisely by using the circle method we shall give a very simple proof of $\vartheta_\ell = \frac{1}{2}$ for $\ell \geq 3$. Further when $\ell \geq 6$, with the help of Siegel's mass formula we shall prove a better exponent $\theta_\ell = \frac{2}{3}$, which means that our bound for $\mathcal{S}_{f,b,\ell}(x)$ is of the same quality as (1.5). Finally the case of $\ell = 2$ requires a more delicate consideration (see the end of Section 3).

[†]It seems that Luo's proof only could show that g_ℓ is the smallest integer such that $g_\ell > (\ell+1)/2$. This means that $\vartheta_3 = \frac{1}{8}$ and $\vartheta_5 = \frac{1}{5}$ instead of $\vartheta_3 = \frac{1}{6}$ and $\vartheta_5 = \frac{1}{4}$.

Remark 2. In fact our method allows us to consider a more general case. Let $\ell \geq 2$, $\mathbf{y} := (y_1, \dots, y_\ell) \in \mathbb{Z}^\ell$ and $\mathbf{A} = (a_{ij})$ be an integral matrix such that $a_{ii} \equiv 0 \pmod{2}$ for $1 \leq i \leq \ell$. The positive definite quadratic form $Q(\mathbf{y})$ is defined by $Q(\mathbf{y}) = \frac{1}{2}\mathbf{y}^t \mathbf{A} \mathbf{y}$. For each $n \geq 1$, define

$$r(n, Q) := |\{\mathbf{y} \in \mathbb{Z}^\ell : Q(\mathbf{y}) = n\}|.$$

Thus $r_\ell(n) = r(n, Q_0)$ with $Q_0 = y_1^2 + \dots + y_\ell^2$. Similar to $\mathcal{S}_{f,b,\ell}(x)$, we define

$$(1.6) \quad \mathcal{S}_{f,b,Q}(x) := \sum_{n \leq x} \lambda_f(n+b)r(n, Q).$$

Define \mathcal{F} to be a class of cusp forms, which consists of holomorphic cusp forms with respect to any finite volume discrete subgroup (such that ∞ is a singular cusp of width 1), any positive real weight and any multiplier systems, as well as Maass cusp forms of any weight and any level [11, 2, 4]. Since the estimate (2.1) below holds for all $f \in \mathcal{F}$ ([10, Theorem 5.3] and [11, Theorem 8.1]), our proof of Theorem 1 implies that

$$\mathcal{S}_{f,b,Q}(x) \ll_{f,Q,\varepsilon} x^{\ell/2-1/2+\varepsilon}$$

holds uniformly for $1 \leq b \leq x$, provided $\ell \geq 5$.

A natural question is what should be the best bound for $\mathcal{S}_{f,b,\ell}(x)$. It is well-known that [3, Theorem 2])

$$\sum_{n \leq x} \lambda_f(n) = \Omega_\pm \left(x^{1/4} \exp \left\{ \frac{D(\log \log x)^{1/4}}{(\log \log \log x)^{3/4}} \right\} \right)$$

where $D > 0$ a constant. A slight weaker Ω -result

$$\sum_{n \leq x} \lambda_f(n) = \Omega(x^{1/4})$$

is supported by the asymptotic formula (see [21])

$$\int_1^X \left| \sum_{n \leq x} \lambda_f(n) \right|^2 dx = C_f X^{3/2} + O(X(\log X)^2),$$

for $X \geq 2$, where C_f is a positive constant. On the other hand, we have $r_\ell(n) \asymp n^{\ell/2-1}$ ($\ell \geq 3$). From the above facts it seems reasonable to propose the following conjecture.

Conjecture. *Let f be a cusp form of weight k and level N . Let $b \geq 0$ and $\ell \geq 3$ be two integers. For any $\varepsilon > 0$, we have*

$$(1.7) \quad \mathcal{S}_{f,b,\ell}(x) \ll_{f,b,\ell,\varepsilon} x^{\ell/2-3/4+\varepsilon}$$

for $x \rightarrow \infty$.

It seems rather difficult to establish (1.7). However we can prove that the bound (1.7) is true on average for $\ell \geq 5$ (see (1.8) and (1.9) below). For this, we shall study the square mean of $\mathcal{S}_{f,b,\ell}(x)$. At this time, our method also allows us to deal with the more general case $\mathcal{S}_{f,b,Q}(x)$.

Our second aim is to prove the following asymptotic formula.

Theorem 2. *Let f be a cusp form of weight k and level N . Let $Q(\mathbf{y})$ be a positive definite quadratic form defined as above. If $\ell \geq 6$, then we have*

$$(1.8) \quad \int_1^X |\mathcal{S}_{f,b,Q}(x)|^2 dx = C_{f,b,Q} \int_1^X x^{\ell-2}(x+b)^{1/2} dx + O_{f,Q}(X^{\ell-7/12}(\log X)^{1/2}) \\ = \frac{C_{f,b,Q}}{\ell-1/2} X^{\ell-1/2} + O_{f,Q}(bX^{\ell-3/2} + X^{\ell-7/12}(\log X)^{1/2})$$

uniformly for $X \geq 2$ and $0 \leq b \leq X$, where the constant $C_{f,b,Q}$ is defined by (4.12) below, and the implied constants depend on f and Q .

Remark 3. In Section 3, we shall prove also that the inequality

$$(1.9) \quad \int_1^X |\mathcal{S}_{f,b,5}(x)|^2 dx \ll_{f,\varepsilon} X^{\ell-1/2+\varepsilon}$$

holds uniformly for $X \geq 1$ and $0 \leq b \leq X$.

2. PRELIMINARY LEMMAS

In order to prove our theorems, we need the following lemmas.

Lemma 2.1. *Let f be a cusp form of weight k and level N .*

(a) *The estimate*

$$(2.1) \quad \sum_{n \leq x} \lambda_f(n) e(\alpha n) \ll_f x^{1/2} \log x$$

holds uniformly for $\alpha \in \mathbb{R}$, where $e(u) := e^{2\pi i u}$.

(b) *For any $\varepsilon > 0$, we have*

$$(2.2) \quad \sum_{\substack{n \leq x \\ n \equiv a \pmod{q}}} \lambda_f(n) \ll_{f,\varepsilon} x^{1/2-1/6+\varepsilon}$$

uniformly for $(q, aN) = 1$ and $q \leq x^{2/3}$.

Proof. See for example, [10, Theorem 5.3] and [20]. □

Lemma 2.2. *Let f be a cusp form of weight k and level N , $(h, q) = 1$ and*

$$(2.3) \quad A(x, h/q) := \sum'_{n \leq x} \lambda_f(n) e_q(hn),$$

where $e_q(m) := e(m/q)$ and $\sum'_{n \leq x}$ means that if x is an integer, the term $n = x$ should be halved. Then for any $\varepsilon > 0$ we have

$$A(x, h/q) = \frac{q^{1/2} x^{1/4}}{\sqrt{2\pi}} \sum_{n \leq M} \frac{\lambda_f(n)}{n^{3/4}} e_q(-\bar{h}n) \cos\left(\frac{4\pi\sqrt{nx}}{q} - \frac{\pi}{4}\right) + O_{f,\varepsilon}\left(\frac{qx^{1/2+\varepsilon}}{M^{1/2}}\right)$$

uniformly for $1 \leq q \leq x$ and $1 \leq M \ll x$, where \bar{h} satisfies $h\bar{h} \equiv 1 \pmod{q}$. In particular for any $\varepsilon > 0$ we have

$$(2.4) \quad A(x, h/q) \ll_{f, \varepsilon} q^{2/3} x^{1/3+\varepsilon}$$

uniformly for $x \geq 1$ and $q \geq 1$. Here the implied constants depend on f and ε only.

Proof. The first assertion is [13, Theorem 1.1]. The bound (2.4) is trivial if $q \geq x$. When $1 \leq q \leq x$, it follows from the first assertion with $M = (q^2 x)^{1/3}$. See also, [17, (2.7)]. \square

Lemma 2.3. *Let $A(x, h/q)$ be defined as in (2.3).*

(i) *We have*

$$(2.5) \quad A_1(x, h/q) := \int_1^x A(u, h/q) \, du \ll_f q^{3/2} x^{3/4}$$

uniformly for $x \geq 1$ and $1 \leq q \leq x^{1/2}$.

(ii) *For any $\varepsilon > 0$ we have*

$$\int_1^X |A(x, h/q)|^2 \, dx = \frac{1}{(4k+2)\pi^2} \sum_{n=1}^{\infty} \frac{|\lambda_f(n)|^2}{n^{3/2}} q X^{3/2} + O_{f, \varepsilon}(q^{3/2} X^{5/4+\varepsilon} + q^2 X^{1+\varepsilon})$$

uniformly for $X \geq 1$ and $q \geq 1$.

Proof. They are (1.6.12) and (1.5.23) of [13], respectively. \square

Lemma 2.4. *Let $\ell \geq 2$, $\mathbf{y} := (y_1, \dots, y_\ell) \in \mathbb{Z}^\ell$ and $\mathbf{A} = (a_{ij})$ be an integral matrix such that $a_{ii} \equiv 0 \pmod{2}$ for $1 \leq i \leq \ell$. The positive definite quadratic form $Q(\mathbf{y})$ is defined by $Q(\mathbf{y}) = \frac{1}{2} \mathbf{y}^t \mathbf{A} \mathbf{y}$. For each $n \geq 1$, define*

$$r(n, Q) := |\{\mathbf{y} \in \mathbb{Z}^\ell : Q(\mathbf{y}) = n\}|.$$

Then for $\ell \geq 4$ we have

$$(2.6) \quad r(n, Q) = \sigma_Q n^{\ell/2-1} \sum_{q=1}^{\infty} \sum_{h=1}^q{}^* S\left(\frac{hQ}{q}\right) \frac{e_q(-hn)}{q^\ell} + O(n^{\ell/4-\delta_\ell+\varepsilon}),$$

where

$$S(Q) := \sum_{0 \leq y_1, \dots, y_\ell \leq q-1} e(Q(\mathbf{y})), \quad \sigma_Q := \frac{(2\pi)^{\ell/2}}{\Gamma(\ell/2) \sqrt{|\mathbf{A}|}}, \quad \delta_\ell := \begin{cases} \frac{1}{4} & \text{if } \ell \text{ is odd,} \\ \frac{1}{2} & \text{if } \ell \text{ is even,} \end{cases}$$

and \sum^* means the sum is over $1 \leq h \leq q$ with $(h, q) = 1$. Furthermore we have

$$(2.7) \quad S(hQ/q) \ll q^{\ell/2} \quad ((h, q) = 1),$$

Proof. See for example, Theorem 11.2 of [10]. \square

3. PROOF OF THEOREM 1

3.1. Case of $\ell \geq 3$.

Let

$$F(\alpha) := \sum_{n \leq x} \lambda_f(n+b) e(-\alpha n) \quad \text{and} \quad S(\alpha) := \sum_{|m| \leq x^{1/2}} e(\alpha m^2).$$

Then

$$\begin{aligned} \int_0^1 F(\alpha) S(\alpha)^\ell d\alpha &= \sum_{n \leq x} \lambda_f(n+b) \sum_{|m_1| \leq x^{1/2}} \cdots \sum_{|m_\ell| \leq x^{1/2}} \int_0^1 e((m_1^2 + \cdots + m_\ell^2 - n)\alpha) d\alpha \\ &= \sum_{n \leq x} \lambda_f(n+b) \sum_{|m_1| \leq x^{1/2}} \cdots \sum_{\substack{|m_\ell| \leq x^{1/2} \\ m_1^2 + \cdots + m_\ell^2 = n}} 1 \\ &= \mathcal{S}_{f,b,\ell}(x). \end{aligned}$$

Firstly Lemma 2.1 allows us to deduce

$$F(\alpha) = e(\alpha b) \sum_{b < n \leq x+b} \lambda_f(n) e(-\alpha n) \ll x^{1/2} \log x$$

uniformly for $x \geq 2$ and $0 \leq b \leq x$. On the other hand, it is easy to see that

$$\begin{aligned} \int_0^1 |S(\alpha)|^2 d\alpha &= \sum_{\substack{|m| \leq x^{1/2} \\ m^2 = n^2}} \sum_{|n| \leq x^{1/2}} 1 \ll x^{1/2}, \\ \int_0^1 |S(\alpha)|^{2d} d\alpha &= \sum_{\substack{|m_1| \leq x^{1/2}, \dots, |m_d| \leq x^{1/2} \\ m_1^2 + \cdots + m_d^2 = n_1^2 + \cdots + n_d^2}} \sum_{\substack{|n_1| \leq x^{1/2}, \dots, |n_d| \leq x^{1/2}}} 1 \leq \sum_{n \leq dx} r_d(n)^2 \ll x^{d-1} \end{aligned}$$

for $d \geq 2$. From the above estimates and the Cauchy inequality we get

$$\mathcal{S}_{f,b,3}(x) \ll x^{1/2} (\log x) \left(\int_0^1 |S(\alpha)|^2 d\alpha \int_0^1 |S(\alpha)|^4 d\alpha \right)^{1/2} \ll x^{3/2-1/4} \log x$$

and for $\ell \geq 4$

$$\begin{aligned} \mathcal{S}_{f,b,\ell}(x) &\ll x^{1/2} (\log x) \left(\int_0^1 |S(\alpha)|^4 d\alpha \int_0^1 |S(\alpha)|^{2(\ell-2)} d\alpha \right)^{1/2} \\ &\ll x^{\ell/2-1/2} \log x, \end{aligned}$$

namely we can take $\vartheta_3 = \frac{1}{4}$ and $\vartheta_\ell = \frac{1}{2}$ for $\ell \geq 4$.

3.2. Case of $\ell \geq 6$.

When $\ell \geq 6$, we can apply Lemmas 2.2 and 2.4 to improve the exponent $\vartheta_\ell = \frac{1}{2}$. From Lemma 2.4 with $Q_0(\mathbf{y}) = y_1^2 + \cdots + y_\ell^2$ we get

$$(3.1) \quad r_\ell(n) = \sigma_{Q_0} n^{\ell/2-1} \sum_{q=1}^{\infty} \sum_{h=1}^q S\left(\frac{hQ_0}{q}\right) \frac{e_q(-hn)}{q^\ell} + O_\varepsilon(n^{\ell/4-\delta_\ell+\varepsilon}).$$

In view of (2.4) in Lemma 2.2 and the bound (2.7) we easily deduce, via a simple integration by parts, that

$$\begin{aligned} \mathcal{S}_{f,b,\ell}(x) &\ll \sum_{q \geq 1} \frac{1}{q^{\ell/2}} \sum_{\substack{1 \leq h \leq q \\ (h,q)=1}} \left| \sum_{1+b \leq n \leq x+b} n^{\ell/2-1} \lambda_f(n) e_q(-hn) \right| + x^{\ell/4-\delta_\ell+1+\varepsilon} \\ &\ll_{f,\varepsilon} x^{\ell/2-2/3+\varepsilon} \sum_{q \geq 1} \frac{1}{q^{\ell/2-5/3}} + x^{\ell/4-\delta_\ell+1+\varepsilon} \\ &\ll_{f,\varepsilon} x^{\ell/2-2/3+\varepsilon} \quad (\text{recall } \ell \geq 6). \end{aligned}$$

3.3. Case of $\ell = 2$.

We need the following Lemmas.

Lemma 3.1. *Let $m \geq 2$ be a positive integer. There is an arithmetic function $h_m(n)$ such that*

$$(3.2) \quad h_m(n) = 0 \quad \text{if } \exists p \text{ such that } p \mid n \text{ and } p \nmid m,$$

$$(3.3) \quad |h_m(n)| \leq \tau(m)\tau_4(n) \quad \text{if } n \mid m^\infty,$$

$$(3.4) \quad \lambda_f(mn) = \sum_{d \mid n} h_m(d) \lambda_f(n/d),$$

where $\tau_k(n)$ denotes the number of solutions of $n = n_1 \cdots n_k$ with positive numbers n_1, \dots, n_k , and $\tau(n) := \tau_2(n)$.

Proof. As usual, we denote by $v_p(n)$ the p -adic valuation of n . By using the formula

$$\lambda_f(mn) = \prod_{p \mid d} \lambda_f(p^{v_p(n)+v_p(m)}) \prod_{p \nmid d} \lambda_f(p^{v_p(n)}),$$

we can write, for $\Re s > 1$,

$$\begin{aligned} \sum_{n=1}^{\infty} \frac{\lambda_f(mn)}{n^s} &= \prod_{p \mid m} \sum_{\nu=0}^{\infty} \frac{\lambda_f(p^{\nu+v_p(m)})}{p^{\nu s}} \prod_{p \nmid m} \sum_{\nu=0}^{\infty} \frac{\lambda_f(p^\nu)}{p^{\nu s}} \\ &= L(s, f) \prod_{p \mid m} \left(\sum_{\nu=0}^{\infty} \frac{\lambda_f(p^{\nu+v_p(m)})}{p^{\nu s}} \right) \left(\sum_{\nu=0}^{\infty} \frac{\lambda_f(p^\nu)}{p^{\nu s}} \right)^{-1}. \end{aligned}$$

This implies that there is an arithmetic function $h_m(n)$ such that (3.2) and (3.4) hold.

Next we prove (3.3). If $p \mid m$, it is easy to see that

$$h_m(p^\nu) = \begin{cases} \lambda_f(p^{v_p(m)}) & \text{if } \nu = 0, \\ \lambda_f(p^{v_p(m)+1}) - \lambda_f(p^{v_p(m)}) \lambda_f(p) & \text{if } \nu = 1, \\ \lambda_f(p^{v_p(m)+\nu}) - \lambda_f(p^{v_p(m)+\nu-1}) \lambda_f(p) + \lambda_f(p^{v_p(m)+\nu-2}) & \text{if } \nu \geq 2. \end{cases}$$

In view of Degline's bound $|\lambda_f(n)| \leq \tau(n)$, for any $n \mid m^\infty$ we have

$$\begin{aligned} |h_m(n)| &= \prod_{p^\nu \parallel n} |h_m(p^\nu)| \leq \prod_{p^\nu \parallel n} 4(v_p(m) + \nu) \\ &\leq \prod_{p^\nu \parallel n} (v_p(m) + 1) \binom{\nu + 3}{3} \leq \tau(m)\tau_4(n). \end{aligned}$$

This completes the proof. \square

Lemma 3.2. *Let f be a cusp form of weight k and level 1. Then the estimate*

$$(3.5) \quad S_f(x; a, q) := \sum_{\substack{n \leq x \\ n \equiv a \pmod{q}}} \lambda_f(n) \ll_f x^{1/3+\varepsilon}$$

holds uniformly for $x \geq 1$ and $q \geq a \geq 1$.

Proof. Note that (2.2) and (3.5) is trivial for $q > x^{2/3}$. Now suppose $q \leq x^{2/3}$. Writting $m := (a, q)$, $a = ma_1$ and $q = mq_1$ such that $(a_1, q_1) = 1$ and using (3.4) of Lemma 3.1, we have

$$\begin{aligned} S_f(x; a, q) &= \sum_{\substack{n \leq x/m \\ n \equiv a_1 \pmod{q_1}}} \lambda_f(mn) \\ &= \sum_{\substack{n_1 n_2 \leq x/m \\ n_1 n_2 \equiv a_1 \pmod{q_1}}} h_m(n_1) \lambda_f(n_2) \\ &= \sum_{n_1 \leq x/m} h_m(n_1) \sum_{\substack{n_2 \leq x/mn_1 \\ n_2 \equiv a_1 \bar{n}_1 \pmod{q_1}}} \lambda_f(n_2), \end{aligned}$$

where $n_1 \bar{n}_1 \equiv 1 \pmod{q_1}$. Noting that $(a_1 \bar{n}_1, q_1) = 1$, we can apply (2.2) with $N = 1$ and (3.2)-(3.3) of Lemma 3.1 to write

$$\begin{aligned} (3.6) \quad S_f(x; a, q) &\ll \tau(m) \left(\frac{x}{m}\right)^{1/3+\varepsilon} \sum_{n_1 \mid m^\infty} \frac{\tau_4(n_1)}{n_1^{1/3+\varepsilon}} \\ &\ll \tau(m) \left(\frac{x}{m}\right)^{1/3+\varepsilon} \prod_{p \mid m} \left(1 - \frac{1}{p^{1/3}}\right)^{-4} \\ &\ll x^{1/3+\varepsilon}. \end{aligned}$$

This completes the proof. \square

Now we prove Theorem 1 for $\ell = 2$. By the classical expression

$$r_2(n) = 4 \sum_{d \mid n} \chi(d)$$

($\chi(n)$ is the non trivial Dirichlet character modulo 4), we can write

$$(3.7) \quad \mathcal{S}_{f,b,2}(x) = 4S_1 + 4S_2 - 4S_3,$$

where

$$\begin{aligned} S_1 &:= \sum_{d \leq \sqrt{x}} \sum_{dm \leq x} \chi(d) \lambda_f(dm + b), \\ S_2 &:= \sum_{m \leq \sqrt{x}} \sum_{dm \leq x} \chi(d) \lambda_f(dm + b), \\ S_3 &:= \sum_{d \leq \sqrt{x}} \sum_{m \leq \sqrt{x}} \chi(d) \lambda_f(dm + b). \end{aligned}$$

By Lemma 3.2, we have

$$\begin{aligned} (3.8) \quad S_1 &= \sum_{d \leq \sqrt{x}} \chi(d) \sum_{\substack{n \leq x+b \\ n \equiv b \pmod{d}}} \lambda_f(n) \\ &\ll \sum_{d \leq \sqrt{x}} (x+b)^{1/3+\varepsilon} \ll x^{5/6+\varepsilon}. \end{aligned}$$

Note that $\chi(d) = 1$ if $d \equiv 1 \pmod{4}$, $\chi(d) = -1$ if $d \equiv 3 \pmod{4}$ and $\chi(d) = 0$ if $2 \mid d$, we have by Lemma 3.2 that

$$\begin{aligned} (3.9) \quad S_2 &= \sum_{m \leq \sqrt{x}} \left(\sum_{(4d+1)m \leq x} \lambda_f((4d+1)m + b) - \sum_{(4d+3)m \leq x} \lambda_f((4d+3)m + b) \right) \\ &= \sum_{m \leq \sqrt{x}} \left(\sum_{\substack{n \leq x+b \\ n \equiv m+b \pmod{4m}}} \lambda_f(n) - \sum_{\substack{n \leq x+b \\ n \equiv 3m+b \pmod{4m}}} \lambda_f(n) \right) \\ &\ll \sum_{m \leq \sqrt{x}} (x+b)^{1/3+\varepsilon} \ll x^{5/6+\varepsilon}. \end{aligned}$$

For S_3 we have by Lemma 3.2 again that

$$\begin{aligned} (3.10) \quad S_3 &= \sum_{d \leq \sqrt{x}} \chi(d) \sum_{\substack{n \leq d\sqrt{x}+b \\ n \equiv b \pmod{d}}} \lambda_f(n) \\ &\ll \sum_{d \leq \sqrt{x}} (d\sqrt{x} + b)^{1/3+\varepsilon} \ll x^{5/6+\varepsilon}. \end{aligned}$$

Now Theorem 1 for the case $\ell = 2$ follows from (3.7)-(3.10).

4. PROOF OF THEOREM 2 AND (1.9)

Suppose $Q(y)$ is the quadratic form defined in Lemma 2.4 for $\ell \geq 6$, and $Q(\mathbf{y}) = y_1^2 + \cdots + y_5^2$ for $\ell = 5$. By Deligne's inequality $|\lambda_f(n)| \leq \tau(n)$ and the bound $r(n, Q) \ll n^{\ell/2-1}$, we have, with the notation $\mathcal{L} := \log(2X)$,

$$(4.1) \quad \int_1^{\sqrt{X}} |\mathcal{S}_{f,b,Q}(x)|^2 dx \ll \int_1^{\sqrt{X}} x^\ell (\log x)^2 dx \ll X^{(\ell+1)/2} \mathcal{L}^2 \ll X^{\ell-2}.$$

It suffices for us to evaluate $\int_{\sqrt{X}}^X |\mathcal{S}_{f,b,Q}(x)|^2 dx$. For any $x \in [X^{1/2}, X] \setminus \mathbb{Z}$, Lemma 2.4 and [5, p. 256, (2.25), p. 270] allow us to write

$$(4.2) \quad \mathcal{S}_{f,b,Q}(x) = \sigma_Q \sum_{q=1}^{\infty} \sum_{h=1}^q {}^* S\left(\frac{hQ}{q}\right) \frac{e_q(bh)}{q^\ell} A_{\ell,b}(x, -h/q) + O(\delta_Q x^{\ell/4 - \delta_\ell + 1 + \varepsilon}),$$

where

$$A_{\ell,b}(x, -h/q) := \sum_{1+b \leq n \leq x+b} (n-b)^{\ell/2-1} \lambda_f(n) e_q(-hn),$$

$$\delta_Q := \begin{cases} 1 & \text{if } \ell \geq 6, \\ 0 & \text{if } \ell = 5 \text{ and } Q(\mathbf{y}) = y_1^2 + \cdots + y_5^2. \end{cases}$$

By partial summation, we can deduce, with the notation (2.3),

$$(4.3) \quad A_{\ell,b}(x, -h/q) = x^{\ell/2-1} A(x+b, -h/q) - (\ell/2-1)I$$

with

$$(4.4) \quad I := \int_b^{x+b} (u-b)^{\ell/2-2} A(u, -h/q) du.$$

It is easy to see that Lemma 2.1 implies that

$$A_{\ell,b}(x, -h/q) \ll X^{\ell/2-1/2} \mathcal{L}$$

uniformly for $1 \leq x \leq X$, $0 \leq b \leq X$ and $(h, n) = 1$. By this and (2.7), we have

$$\sum_{q > X^{1/2}} \frac{1}{q^\ell} \sum_{h=1}^q {}^* \left| S\left(\frac{hQ}{q}\right) A_{\ell,b}(x, -h/q) \right| \ll X^{\ell/4+1/2} \mathcal{L}.$$

Thus we can write (4.2) as, for $x \in [X^{1/2}, X] \setminus \mathbb{Z}$,

$$(4.5) \quad \begin{aligned} \mathcal{S}_{f,b,Q}(x) &= \sigma_Q \sum_{q \leq X^{1/2}} \sum_{h=1}^q {}^* S\left(\frac{hQ}{q}\right) \frac{e_q(bh)}{q^\ell} A_{\ell,b}(x, -h/q) \\ &\quad + O(X^{\ell/4+1/2} \mathcal{L} + \delta_Q X^{\ell/4 - \delta_\ell + 1 + \varepsilon}). \end{aligned}$$

Suppose now $1 \leq q \leq X^{1/2}$ and we estimate the integral I defined as in (4.4). If $q^2 \leq b$, by partial integration and Lemma 2.3 we have, with the notation (2.5),

$$(4.6) \quad \begin{aligned} I &= x^{\ell/2-2} A_1(x+b, -h/q) - \frac{\ell-4}{2} \int_b^{x+b} (u-b)^{\ell/2-3} A_1(u, -h/q) du \\ &\ll q^{3/2} X^{\ell/2-5/4}. \end{aligned}$$

If $q^2 > b$, we can write

$$(4.7) \quad \begin{aligned} I &= \int_b^{q^2} A(u, -h/q) (u-b)^{\ell/2-2} du + \int_{q^2}^{x+b} A(u, -h/q) (u-b)^{\ell/2-2} du \\ &\ll q^{\ell-1} \log(2q) + q^{3/2} X^{\ell/2-5/4} \ll q^{3/2} X^{\ell/2-5/4} \mathcal{L}, \end{aligned}$$

where we estimated the first integral by Lemma 2.1 and the second as in (4.6). In view of (4.6), (4.7) and (2.7), the contribution of I to the last sum in (4.5) is

$$\begin{aligned} &\ll X^{\ell/2-5/4} \mathcal{L} \sum_{q \leq X^{1/2}} q^{-(\ell-5)/2} \\ &\ll R_\ell(X) := \begin{cases} X^{\ell/2-1+\varepsilon} & \text{if } \ell \geq 6, \\ X^{\ell/2-3/4+\varepsilon} & \text{if } \ell = 5. \end{cases} \end{aligned}$$

Since $\ell/4 - \delta_\ell + 1 \leq \ell/2 - 1$ ($\ell \geq 6$), thus we can write (4.5) as, for $x \in [X^{1/2}, X] \setminus \mathbb{Z}$,

$$(4.8) \quad \mathcal{S}_{f,b,Q}(x) = \sigma_Q x^{\ell/2-1} \sum_{q \leq X^{1/2}} \sum_{h=1}^q S\left(\frac{hQ}{q}\right) \frac{e_q(bh)}{q^\ell} A(x+b, -h/q) + O(R_\ell(X)).$$

Now we split the sum over $q \leq X^{1/2}$ into two parts according to $q \leq X^{1/6}$ or $X^{1/6} < q \leq X^{1/2}$. In view of $X^{1/2} \leq x$, we can apply Lemma 2.2 to the sum $\sum_{q \leq X^{1/6}}$ by taking $M = x + b$. It is easy to see that the contribution of the error term to $\mathcal{S}_{f,b,Q}(x)$ is $O(X^{\ell/2-1+\varepsilon})$ if $\ell \geq 6$ and $O(X^{19/12})$ if $\ell = 5$. Thus the formula (4.8) becomes

$$(4.9) \quad \mathcal{S}_{f,b,Q}(x) = S_1(x) + S_2(x) + O(R_\ell(X)),$$

where

$$\begin{aligned} S_1(x) &:= \frac{\sigma_Q}{\sqrt{2\pi}} x^{\ell/2-1} (x+b)^{1/4} \sum_{q \leq X^{1/6}} \sum_{h=1}^q \sum_{n \leq x+b} \psi(h, n, q) \cos(g_{n,q}(x)), \\ S_2(x) &:= \sigma_Q x^{\ell/2-1} \sum_{X^{1/6} < q \leq X^{1/2}} \sum_{h=1}^q S\left(\frac{hQ}{q}\right) \frac{e_q(bh)}{q^\ell} A(x+b, -h/q), \end{aligned}$$

and

$$\psi(h, n, q) := \frac{\lambda_f(n) e_q(bh - \bar{h}n)}{n^{3/4} q^{\ell-1/2}} S\left(\frac{hQ}{q}\right), \quad g_{n,q}(x) := \frac{4\pi \sqrt{n(x+b)}}{q} - \frac{\pi}{4}.$$

We first evaluate the integral $\int_{\sqrt{X}}^X |S_1(x)|^2 dx$. By the elementary formula

$$\cos u \cos v = \frac{\cos(u-v) + \cos(u+v)}{2},$$

we can write

$$(4.10) \quad |S_1(x)|^2 = S_{11}(x) + S_{12}(x) + S_{13}(x),$$

where $h_j \overline{h_j} \equiv 1 \pmod{q_j}$ for $j = 1, 2$, and

$$\begin{aligned}
S_{11}(x) &:= \left(\frac{\sigma_Q}{2\pi}\right)^2 x^{\ell-2} (x+b)^{1/2} \sum_{q_1 \leq X^{1/6}} \sum_{h_1=1}^{q_1*} \sum_{n_1 \leq x+b} \sum_{q_2 \leq X^{1/6}} \sum_{h_2=1}^{q_2*} \sum_{n_2 \leq x+b} \\
&\quad \overline{\psi(h_1, n_1, q_1) \psi(h_2, n_2, q_2)}, \\
S_{12}(x) &:= \left(\frac{\sigma_Q}{2\pi}\right)^2 x^{\ell-2} (x+b)^{1/2} \sum_{q_1 \leq X^{1/6}} \sum_{h_1=1}^{q_1*} \sum_{n_1 \leq x+b} \sum_{q_2 \leq X^{1/6}} \sum_{h_2=1}^{q_2*} \sum_{n_2 \leq x+b} \\
&\quad \overline{\psi(h_1, n_1, q_1) \psi(h_2, n_2, q_2)} \cos(g_{n_1, q_1}(x) - g_{n_2, q_2}(x)), \\
S_{13}(x) &:= \left(\frac{\sigma_Q}{2\pi}\right)^2 x^{\ell-2} (x+b)^{1/2} \sum_{q_1 \leq X^{1/6}} \sum_{h_1=1}^{q_1*} \sum_{n_1 \leq x+b} \sum_{q_2 \leq X^{1/6}} \sum_{h_2=1}^{q_2*} \sum_{n_2 \leq x+b} \\
&\quad \overline{\psi(h_1, n_1, q_1) \psi(h_2, n_2, q_2)} \cos(g_{n_1, q_1}(x) + g_{n_2, q_2}(x)).
\end{aligned}$$

The relation $\sqrt{n_1}/q_1 = \sqrt{n_2}/q_2$ implies that n_1 and n_2 have the same square-free factor. We write $n_j = d_j^2 m$ with $\mu(m) \neq 0$ for $j = 1, 2$. Recalling (2.7), Deligne's well-known bound $|\lambda_f(n)| \leq \tau(n)$ and the trivial inequality $\tau(uv) \leq \tau(u)\tau(v)$, we have

$$\begin{aligned}
(4.11) \quad \overline{\psi(h_1, n_1, q_1) \psi(h_2, n_2, q_2)} &\ll \frac{\tau(d_1^2 m) \tau(d_2^2 m)}{(d_1 d_2 m)^{3/2} (q_1 q_2)^{\ell/2 - 1/2}} \\
&\ll \frac{\tau(d_1)^2 \tau(d_2)^2 \tau(m)^2}{(d_1 d_2 m)^{3/2} (q_1 q_2)^{\ell/2 - 1/2}}.
\end{aligned}$$

The above estimate (4.11) implies that the infinite series

$$(4.12) \quad C_{f,b,Q} := \left(\frac{\sigma_Q}{2\pi}\right)^2 \sum_{q_1 \geq 1} \sum_{1 \leq h_1 \leq q_1} \sum_{n_1 \geq 1} \sum_{q_2 \geq 1} \sum_{1 \leq h_2 \leq q_2} \sum_{n_2 \geq 1} \overline{\psi(h_1, n_1, q_1) \psi(h_2, n_2, q_2)}$$

$\sqrt{n_1}/q_1 = \sqrt{n_2}/q_2$

converges absolutely provided $\ell \geq 4$ and

$$(4.13) \quad C_{f,b,Q} \ll_{f,Q} 1 \quad (\forall b).$$

If a quadruple (q_1, q_2, n_1, n_2) satisfies $\sqrt{n_1}/q_1 = \sqrt{n_2}/q_2$ but is not included in $S_{11}(x)$, then we must have that $q_1 > X^{1/6}$, or $q_2 > X^{1/6}$, or $n_1 > x + b \geq X^{1/2}$, or $n_2 > x + b \geq X^{1/2}$. Thus $n_1 q_2^2 = n_2 q_1^2 > X^{1/3}$. In view of the following elementary estimates

$$\begin{aligned}
(4.14) \quad t_\ell(m) &:= \sum_{m=nq^2} \tau(n) q^{-\ell/2+3} \leq \tilde{t}_\ell(m) := \begin{cases} \tau_3(m) & \text{if } \ell \geq 6, \\ m^{1/4} \tau_3(m) & \text{if } \ell = 5, \end{cases} \\
&\sum_{n \leq u} \tau_3(n)^2 \ll u (\log u)^8 \quad (u \geq 2),
\end{aligned}$$

we can derive, for all $x \in [X^{1/2}, X] \setminus \mathbb{Z}$,

$$\begin{aligned}
 (4.15) \quad \left| \frac{S_{11}(x)}{x^{\ell-2}(x+b)^{1/2}} - C_{f,b,Q} \right| &\ll \sum_{\substack{n_1, n_2, q_1, q_2 \\ n_2 q_1^2 = n_1 q_2^2 > X^{1/3}}} \frac{\tau(n_1)\tau(n_2)}{(n_1 q_2^2 n_2 q_1^2)^{3/4} (q_1 q_2)^{\ell/2-3}} \\
 &\ll \sum_{m > X^{1/3}} \frac{t_\ell(m)^2}{m^{3/2}} \leq \sum_{m > X^{1/3}} \frac{\tilde{t}_\ell(m)^2}{m^{3/2}} \\
 &\ll \begin{cases} X^{-1/6} \mathcal{L}^8 & \text{if } \ell \geq 6, \\ X^{-1/12} \mathcal{L}^8 & \text{if } \ell = 5. \end{cases}
 \end{aligned}$$

On the other hand, we have, in view of (4.13),

$$C_{f,b,Q} \int_{\sqrt{X}}^X x^{\ell-2}(x+b)^{1/2} dx = C_{f,b,Q} \int_1^X x^{\ell-2}(x+b)^{1/2} dx + O(X^{\ell/2})$$

uniformly for $X \geq 1$ and $0 \leq b \leq X$. And hence we obtain

$$(4.16) \quad \int_{\sqrt{X}}^X S_{11}(x) dx = C_{f,b,Q} \int_1^X x^{\ell-2}(x+b)^{1/2} dx + \begin{cases} O(X^{\ell-2/3} \mathcal{L}^8) & \text{if } \ell \geq 6, \\ O(X^{\ell-7/12} \mathcal{L}^8) & \text{if } \ell = 5. \end{cases}$$

By the first derivative test, (4.11) and (2.7), we have (note $\ell \geq 5$)

$$\begin{aligned}
 (4.17) \quad \int_{X^{1/2}}^X |S_{13}(x)| dx &\ll X^{\ell-1} \sum_{\substack{n_1, n_2 \leq x+b \\ q_1, q_2 \leq X^{1/6}}} \frac{\tau(n_1)\tau(n_2)}{(n_1 n_2)^{3/4} (q_1 q_2)^{\ell/2-3/2}} \frac{1}{\sqrt{n_1}/q_1 + \sqrt{n_2}/q_2} \\
 &\ll X^{\ell-1} \sum_{\substack{n_1, n_2 \leq x+b \\ q_1, q_2 \leq X^{1/6}}} \frac{\tau(n_1)\tau(n_2)}{n_1 n_2 (q_1 q_2)^{\ell/2-2}} \\
 &\ll X^{\ell-5/6} \mathcal{L}^6,
 \end{aligned}$$

where we used the inequality $u + v \geq 2\sqrt{uv}$ for $u, v > 0$.

Similarly by the first derivative test again

$$(4.18) \quad \int_{X^{1/2}}^X |S_{12}(x)| dx \ll X^{\ell-1} (\Sigma_{11} + \Sigma_{12}),$$

where

$$\begin{aligned}
 \Sigma_{11} &:= \sum_{\substack{n_1, n_2 \leq x+b, q_1, q_2 \leq X^{1/6} \\ |\sqrt{n_1}/q_1 - \sqrt{n_2}/q_2| \geq \frac{1}{10} (\sqrt{n_1 n_2}/q_1 q_2)^{1/2}}} \frac{\tau(n_1)\tau(n_2)}{(n_1 n_2)^{3/4} (q_1 q_2)^{\ell/2-3/2}} \frac{1}{|\sqrt{n_1}/q_1 - \sqrt{n_2}/q_2|}, \\
 \Sigma_{12} &:= \sum_{\substack{n_1, n_2 \leq x+b, q_1, q_2 \leq X^{1/6} \\ 0 < |\sqrt{n_1}/q_1 - \sqrt{n_2}/q_2| < \frac{1}{10} (\sqrt{n_1 n_2}/q_1 q_2)^{1/2}}} \frac{\tau(n_1)\tau(n_2)}{(n_1 n_2)^{3/4} (q_1 q_2)^{\ell/2-3/2}} \frac{1}{|\sqrt{n_1}/q_1 - \sqrt{n_2}/q_2|}.
 \end{aligned}$$

As in the argument of (4.17) we get easily that (recall $\ell \geq 5$)

$$(4.19) \quad \Sigma_{11} \ll \sum_{n_1, n_2 \leq x+b, q_1, q_2 \leq X^{1/6}} \frac{\tau(n_1)\tau(n_2)}{n_1 n_2 (q_1 q_2)^{\ell/2-2}} \ll X^{1/6} \mathcal{L}^6.$$

If $|\sqrt{n_1}/q_1 - \sqrt{n_2}/q_2| < \frac{1}{10}(\sqrt{n_1 n_2}/q_1 q_2)^{1/2}$ holds, then it follows that

$$\sqrt{n_1} q_2 \asymp \sqrt{n_2} q_1 \quad \text{and} \quad \sqrt{n_1} q_2 + \sqrt{n_2} q_1 \asymp (n_1 n_2)^{1/4} (q_1 q_2)^{1/2}.$$

So in view of these facts, by a similar argument as in (4.15) we have, for $\ell \geq 6$,

$$\begin{aligned} \Sigma_{12} &\ll \sum_{\substack{q_1, q_2 \leq X^{1/6}; n_1, n_2 \leq x+b \\ n_1 q_2^2 - n_2 q_1^2 \neq 0}} \frac{d(n_1) d(n_2)}{(n_1 n_2 q_1^2 q_2^2)^{1/2} (q_1 q_2)^{\ell/2-4} |n_1 q_2^2 - n_2 q_1^2|} \\ &\ll X^{1/3} \sum_{\substack{m_1, m_2 \leq X^{1/6}(X+b) \\ m_1 \neq m_2}} \frac{d_3(m_1) d_3(m_2)}{(m_1 m_2)^{1/2} |m_1 - m_2|} \\ &\ll X^{1/3} \sum_{1 < h \leq X^{1/6}(X+b)} \frac{1}{h} \sum_{m \leq X^{1/6}(X+b)} \frac{d_3(m) d_3(m+h)}{(m(m+h))^{1/2}}. \end{aligned}$$

By applying the Cauchy inequality to the sum over m and (4.14), we can deduce

$$(4.20) \quad \Sigma_{12} \ll X^{1/3} \mathcal{L}^9 \quad (\ell \geq 6).$$

When $\ell = 5$, the similar estimate with $X^{1/2}$ in place of $X^{1/3}$ holds. Inserting (4.19) and (4.20) into (4.18), we find that

$$(4.21) \quad \int_{\sqrt{X}}^X |S_{12}(x)| dx \ll R_\ell^*(X) := \begin{cases} X^{\ell-2/3} \mathcal{L}^9 & \text{if } \ell \geq 6, \\ X^{\ell-1/2} \mathcal{L}^9 & \text{if } \ell = 5. \end{cases}$$

Combining (4.16), (4.21) and (4.17) with (4.10) yields

$$(4.22) \quad \int_{\sqrt{X}}^X |S_1(x)|^2 dx = C_{f,b,Q} \int_1^X x^{\ell-2} (x+b)^{1/2} dx + O(R_\ell^*(X)).$$

Finally we estimate $\int_{\sqrt{X}}^X |S_2(x)|^2 dx$. By (2.7) and Cauchy's inequality, it follows

$$\begin{aligned} |S_2(x)|^2 &\ll x^{\ell-2} \left(\sum_{X^{1/6} < q \leq X^{1/2}} q^{-\ell/2+1} |A(x+b, -h/q)| \right)^2 \\ &\ll X^{\ell-2} \mathcal{L} \sum_{X^{1/6} < q \leq X^{1/2}} q^{-\ell+3} |A(x+b, -h/q)|^2. \end{aligned}$$

From this and Lemma 2.3(ii), we deduce (recall $\ell \geq 5$ and $0 \leq b \leq X$)

$$(4.23) \quad \begin{aligned} \int_{\sqrt{X}}^X |S_2(x)|^2 dx &\ll X^{\ell-2} \mathcal{L} \sum_{X^{1/6} < q \leq X^{1/2}} q^{-\ell+3} (qX^{3/2} + q^{3/2} X^{5/4+\varepsilon} + q^2 X^{1+\varepsilon}) \\ &\ll \begin{cases} X^{\ell-2/3} \mathcal{L} & \text{if } \ell \geq 6, \\ X^{\ell-1/2+\varepsilon} & \text{if } \ell = 5. \end{cases} \end{aligned}$$

On the other hand, (4.22) implies immediately

$$\int_{\sqrt{X}}^X |S_1(x)|^2 dx \ll X^{\ell-1/2}.$$

In view of this and (4.23), a simple application of Cauchy's inequality yields

$$(4.24) \quad \int_{\sqrt{X}}^X |S_1(x)S_2(x)| dx \ll \begin{cases} X^{\ell-7/12}\mathcal{L}^{1/2} & \text{if } \ell \geq 6, \\ X^{\ell-1/2+\varepsilon} & \text{if } \ell = 5. \end{cases}$$

Now the first assertion of Theorem 2 and (1.9) follow from (4.1), (4.9), (4.22), (4.23) and (4.24). The second is an immediate consequence by noticing that

$$(x+b)^{1/2} = x^{1/2} + O(bx^{-1/2})$$

valid uniformly for $x \geq 1$ and $b \geq 1$.

$$\begin{aligned} \sum_{p_1+p_2+p_3=N} \tau(p_1) &= \sum_{p_1+p_2+p_3=N} \tau(N-p_2-p_3) \\ &\ll N^{15/2} \exp\{-C\sqrt{\log N}\} \end{aligned}$$

REFERENCES

- [1] J.-M. Deshouillers & H. Iwaniec, *An additive divisor problem*, J. London Math. Soc. (2) **26** (1982), no. 1, 1–14.
- [2] W. Duke & H. Iwaniec, *Bilinear forms in the Fourier coefficients of half-integral weight cusp forms and sums over primes*, Math. Ann. **286** (1990), 783–802.
- [3] J. L. Hafner & A. Ivić, *On sums of Fourier coefficients of cusp forms*, Enseign. Math. (2) **35** (1989), no. 3-4, 375–382.
- [4] G. Harcos & P. Michel, *The subconvexity problem for Rankin-Selberg L-functions and equidistribution of Heegner points. II*, Invent. Math. **163** (2006), 581–655.
- [5] G. H. Hardy, *Representation of a number as the sum of squares*, Trans. Amer. Math. Soc. **17** (1920), 255–284.
- [6] D. R. Heath-Brown, *The twelfth power moment of the Riemann zeta-function*, Quart. J. Math. (Oxford) **29** (1978), 443–462.
- [7] R. Holowinsky, *A sieve method for shifted convolution sums*, Duke Math. J. **146** (2009), no. 3, 401–448.
- [8] A. E. Ingham, *Some asymptotic formulae in the theory of numbers*, J. London Math. Soc. **2** (1927), 202–208.
- [9] A. Ivić & Y. Motohashi, *On some estimates involving the binary additive divisor problem*, Quart. J. Math. Oxford **46** (1995), 471–483.
- [10] H. Iwaniec, *Topics in classical automorphic forms*, Graduate Studies in Mathematics, vol. 17, American Mathematical Society, Providence, Rhode Island, 1997.
- [11] H. Iwaniec, *Spectral methods of automorphic forms*, Amer. Math. Soc., Second edition, 2002.
- [12] M. Jutila, *The additive divisor problem and exponential sums*, Advances in number theory, Oxford Univ. Press, New York, 1993, 113–135.
- [13] M. Jutila, *A Method in the Theory of Exponential Sums*, Tata Inst. of Fundamental Research, Bombay Lectures No. 80, Springer-Verlag, Berlin-Heidelberg-New York, 1987.
- [14] M. Jutila, *The additive divisor problem and its analogs for Fourier coefficients of cusp forms. I*, Math. Z. **223** (1996), no. 3, 435–461.

- [15] M. Jutila, *The additive divisor problem and its analogs for Fourier coefficients of cusp forms. II*, Math. Z. **225** (1997), 625–637.
- [16] Wenzhi Luo, *Shifted convolution of cusp-forms with θ -series*, Abh. Math. Semin. Univ. Hambg. **81** (2011), 45–53.
- [17] Guangshi Lü, *The average value of Fourier coefficients of cusp forms in arithmetic progressions*, J. Number Theory **129** (2009), 477–487.
- [18] T. Meurman, *On the binary additive divisor problem*, Number theory (Turku, 1999), 223–246, de Gruyter, Berlin, 2001.
- [19] Y. Motohashi, *The binary additive divisor problem*, Ann. Sci. Ec. Norm. Supér. **27** (1994), 529–572.
- [20] Robert A. Smith, *Fourier coefficients of modular forms over arithmetic progressions. I, II*, With remarks by M. R. Murty, C. R. Math. Rep. Acad. Sci. Canada **15** (1993), no. 2-3, 85–90, 91–98.
- [21] A. Walfisz, *Über Gitterpunkte in mehrdimensional Ellipsoiden, VIII*, Trudy Tbil. Math. Inst. **5** (1938), 1–65.
- [22] J. Wu, *Power sums of Hecke eigenvalues and applications*, Acta Arith. **137** (2009), 333–344.

GUANGSHI LÜ, SCHOOL OF MATHEMATICS, SHANDONG UNIVERSITY, JINAN, SHANDONG 250100, CHINA

E-mail address: `gslv@sdu.edu.cn`

JIE WU, SCHOOL OF MATHEMATICS, SHANDONG UNIVERSITY, JINAN, SHANDONG 250100, CHINA. CNRS, INSTITUT ÉLIE CARTAN DE LORRAINE, UMR 7502, 54506 VANDŒUVRE-LÈS-NANCY, FRANCE

Current address: Université de Lorraine, Institut Élie Cartan de Lorraine, UMR 7502, 54506 Vandœuvre-lès-Nancy, France

E-mail address: `jie.wu@univ-lorraine.fr`

WENGUANG ZHAI, DEPARTMENT OF MATHEMATICS, CHINA UNIVERSITY OF MINING AND TECHNOLOGY, BEIJING 100083, CHINA

E-mail address: `zhaiwg@hotmail.com`