



**HAL**  
open science

# HOROSPHERE TOPOLOGY

Filippo Bracci, Hervé Gaussier

► **To cite this version:**

| Filippo Bracci, Hervé Gaussier. HOROSPHERE TOPOLOGY. 2016. hal-01315332v3

**HAL Id: hal-01315332**

**<https://hal.science/hal-01315332v3>**

Preprint submitted on 18 Sep 2016 (v3), last revised 5 Apr 2018 (v6)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# HOROSPHERE TOPOLOGY

FILIPPO BRACCI\* AND HERVÉ GAUSSIER

ABSTRACT. We introduce a prime end-type theory on complete Kobayashi hyperbolic manifolds using horosphere sequences. This allows to introduce a new notion of boundary—new even in the unit disc in the complex space—the horosphere boundary, and a topology on the manifold together with its horosphere boundary, the horosphere topology. We prove that a bounded strongly pseudoconvex domain endowed with the horosphere topology is homeomorphic to its Euclidean closure, while for the polydisc such a horosphere topology is not even Hausdorff and is different from the Gromov topology. We use this theory to study boundary behavior of univalent maps from bounded strongly pseudoconvex domains. Among other things, we prove that every univalent map of the unit ball whose image is bounded and convex, extends as a homeomorphism up to the closure. Such a result, relying in an essential way on our theory and on the Gromov hyperbolicity theory, is completely new, dealing with non smooth domains.

## CONTENTS

1. Introduction	1
2. Carathéodory prime ends theory vs. Horosphere topology in the unit disc	4
3. Admissible sequences, Busemann sequences and horosphere boundary	8
4. Horosphere topology, impression and principal part	11
5. Strongly pseudoconvex domains	13
6. Convex domains	20
6.1. Convex domains biholomorphic to strongly pseudoconvex domains	22
6.2. Convex domains biholomorphic to strongly convex domains	25
7. Horosphere boundary versus Gromov boundary	30
8. Boundary behavior	35
References	37

## 1. INTRODUCTION

Carathéodory prime ends theory is one of the most powerful tools for studying boundary behavior of univalent functions in the unit disc  $\mathbb{D} \subset \mathbb{C}$ . Given a simply connected domain  $\Omega \subset \mathbb{C}$ , one can define an abstract boundary  $\partial_C \Omega$ , the Carathéodory boundary, whose points, called prime ends, are given as the set of equivalent classes of null chains (see, e.g., [14, 16, 34] or Section 2). Then one can give a natural topology, the Carathéodory topology, to the space  $\hat{\Omega}_C := \Omega \cup \partial_C \Omega$ . It turns out that  $\hat{\mathbb{D}}_C$  is homeomorphic to the Euclidean closure  $\overline{\mathbb{D}}$  and, if  $f : \mathbb{D} \rightarrow \Omega$  is a biholomorphism, then  $f$  extends to a homeomorphism from  $\hat{\mathbb{D}}_C$  to  $\hat{\Omega}_C$ . The link between the Carathéodory boundary and the boundary behavior of univalent functions

---

2010 *Mathematics Subject Classification.* 32F27, 32F45, 32H40, 32Q55, 32T15, 53C23, 30E25, 30F25.

*Key words and phrases.* Invariant distances, Horospheres, Prime ends theory, Gromov hyperbolicity, Boundary extension, Pseudoconvex domains.

\*Supported by the ERC grant “HEVO - Holomorphic Evolution Equations” n. 277691.

is provided by impressions and principal parts of prime ends. In particular, if the impression of each prime end of  $\Omega$  is just one point, the map  $f$  extends continuously.

Carathéodory prime ends theory is defined by using Euclidean objects (the null chains, which are sequences of Jordan arcs ending at the frontier of  $\Omega$ ), and the ultimate reason why it works is because univalent mappings in  $\mathbb{D}$  are (quasi-)conformal. Indeed, Carathéodory's theory can be generalized to domains in  $\mathbb{R}^n$  for quasi-conformal mappings (see [37] and [25]). Other generalizations of the prime ends theory in metric spaces have been studied in [5]. The problem with those generalizations to higher dimension, is that they cannot be applied to univalent maps without adding some extra hypotheses (in general, a univalent map in higher dimension is not quasi-conformal).

Different types of boundaries were introduced in other contexts. For instance, the Gromov boundary was introduced by M. Gromov [24] in hyperbolic groups with the purpose of a better understanding of the growth of some groups at infinity. The construction is valid in hyperbolic metric spaces, namely metric spaces in which geodesic triangles are thin. In most situations, the space is assumed geodesic, meaning that any two points may be joined by a geodesic, and proper, meaning that closed balls are compact. We recall that a geodesic ray, in a geodesic metric space  $(X, d)$ , is an isometry from  $[0, +\infty[$  to  $X$  such that the length of the segment  $\gamma([0, t])$  is equal to  $d(\gamma(0), \gamma(t))$  for every  $t \geq 0$ . Two geodesic rays  $\gamma_1$  and  $\gamma_2$  are equivalent if there is some  $c > 0$  such that  $d(\gamma_1(t), \gamma_2(t)) \leq c$  for every  $t \geq 0$ .

The Gromov boundary of  $X$  is the quotient of the set of geodesic rays whose origin is some fixed base point  $x \in X$  by that equivalence relation. Moreover, it does not depend on the chosen base point. See Section 7 for the precise definitions.

In this paper we introduce a completely new prime ends theory defined via horospheres related to sequences. Horospheres have been used pretty much in geometric function theory in one and several variables, especially for studying iteration theory, Julia's Lemma, Denjoy-Wolff theorems (see, e.g., [1, 3, 4, 9, 10, 35, 19] and references therein), and they are a particular instance of a general notion of horospheres in locally complete metric spaces, see [15, 6]. In complex geometry, horospheres defined by using complex geodesics are sometimes called Busemann horospheres. In strongly convex domains with smooth boundary, thanks to Lempert's theory [27], horospheres turn out to be level sets of a pluricomplex Poisson kernel [11, 12] and in the unit ball  $\mathbb{B}^n$  they are just ellipsoids internally tangent to the boundary of the ball at one point.

Our point of view is different than in the previous works. To be precise, let  $M$  be a Kobayashi complete hyperbolic manifold, let  $K_M$  denote its Kobayashi distance, and let  $x \in M$ . Given a compactly divergent sequence  $\{u_n\}$  in  $M$ , and  $R > 0$ , we define the *horosphere* relative to  $\{u_n\}$  with radius  $R$  by

$$E_x(\{u_n\}, R) := \{w \in M : \limsup_{n \rightarrow \infty} [K_M(w, u_n) - K_M(x, u_n)] < \frac{1}{2} \log R\}.$$

The sequence is *admissible* if  $E_x(\{u_n\}, R) \neq \emptyset$  for all  $R > 0$ . We introduce an equivalence relation on the set of all admissible sequences by declaring equivalent two admissible sequences if every horosphere relative to one sequence is contained in a horosphere of the other and vice versa (see Section 3 for precise statements and definitions). The *horosphere boundary*  $\partial_H M$  is the set of equivalence classes of admissible sequences. Let  $\hat{M} := M \cup \partial_H M$ . Then, using horospheres, we define a topology on  $\hat{M}$  which induces on  $M$  its natural topology. We call *horosphere topology* such a topology (see Section 4). Since this topology is defined via the Kobayashi distance  $K_M$ , it turns out that if  $F : M \rightarrow N$  is a biholomorphism then  $F$  extends naturally to a homeomorphism  $\hat{F} : \hat{M} \rightarrow \hat{N}$ .

The first main result of the paper, is the following generalization of Carathéodory's theorem (see Theorem 5.10):

**Theorem 1.1.** *Let  $D \subset \mathbb{C}^N$  be a bounded strongly pseudoconvex domain with  $C^3$  boundary. Then  $\hat{D}$  endowed with the horosphere topology is homeomorphic to  $\overline{D}$  (closure in  $\mathbb{C}^N$ ) endowed with the Euclidean topology.*

The somewhat unnatural technical assumption on  $C^3$  regularity of the boundary of  $D$  is needed in order to apply some theory of complex geodesics in strongly convex domains.

As a matter of fact, the horosphere boundary of the polydisc is not Hausdorff; hence, we have another proof of the well known fact that strongly pseudoconvex domains cannot be biholomorphic to polydiscs. More generally, using this result, we can prove that there is no holomorphic isometric embeddings of a polydisc into any strongly pseudoconvex domain.

As in Carathéodory's prime ends theory, for domains in  $\mathbb{C}^N$  (or more generally in  $\mathbb{C}\mathbb{P}^N$ ), we can define *horosphere impressions* and *horosphere principal parts*. In particular, for a bounded strongly pseudoconvex domain  $D \subset \mathbb{C}^N$  with  $C^3$  boundary, the horosphere impressions always reduce to one point at the boundary. Therefore, if  $F : D \rightarrow \Omega$  is a biholomorphism, the limit of  $F$  at a point  $p \in \partial D$  is given by the impression of  $\hat{F}(\underline{x}_p)$ , where  $\underline{x}_p$  is the point of  $\partial_H D$  corresponding to  $p$  under the isomorphism of Theorem 1.1. In particular,  $F$  extends continuously on  $\overline{D}$  if and only if the impressions of each point of  $\partial_H \Omega$  is just one point in  $\partial \Omega$ —and this also gives another proof of homeomorphic extension of biholomorphisms among strongly pseudoconvex domains. On the other hand, non-tangential limits (in fact, a larger notion of limits which we call E-limits) can be controlled using the principal part of horospheres, similarly to what happens in Carathéodory's theory for the principal parts of prime ends (see Section 8).

We apply the horosphere theory to study biholomorphisms  $F : D \rightarrow \Omega$  from a strongly pseudoconvex domain  $D$  to a convex domain  $\Omega$ . If the domain  $\Omega$  is strongly convex with smooth boundary, then Fefferman's theorem [17] implies that the map  $F$  extends as a diffeomorphism from  $\overline{D}$  to  $\overline{\Omega}$ . In case  $\Omega$  has no boundary regularity, some other conditions on the behavior of the Kobayashi distance can be useful to obtain continuous extension (see [39] where A. Zimmer proves the continuous extension of Kobayashi isometric embeddings of a bounded convex domain with  $C^{1,\alpha}$  boundary into a strictly  $\mathbb{C}$ -convex domain with  $C^{1,\alpha}$  boundary, [8] and references therein). However, if nothing is assumed on  $\Omega$ , very little is known, even when  $D = \mathbb{B}^n$ , the unit ball. In fact, a conjecture of Muir and Suffridge [31, 32] states that if  $F : \mathbb{B}^n \rightarrow \mathbb{C}^N$  is univalent and its image is bounded and convex, then  $F$  extends continuously to  $\partial \mathbb{B}^n$ .

Using in an essential way the theory developed in Section 6 and the theory of Gromov hyperbolicity, not only we give an affirmative answer to the Muir and Suffridge conjecture, but also we prove homeomorphic extension without any additional assumption (see Corollary 8.3):

**Theorem 1.2.** *Let  $D \subset \mathbb{C}^N$  be a bounded strongly convex domain with  $C^3$  boundary, for instance,  $D = \mathbb{B}^n$ . Let  $F : D \rightarrow \Omega$  be a biholomorphism. If  $\Omega$  is a bounded convex domain, then  $F$  extends as a homeomorphism from  $\overline{D}$  to  $\overline{\Omega}$ .*

The same result holds for univalent mappings from bounded strongly pseudoconvex domains whose image is bounded and *strictly  $\mathbb{C}$ -linearly convex*, see Section 8.

We point out that such results are completely new since they deal with non necessarily smooth domains. Moreover, conceptually, it seems that classical theories considered before in extension problems do not provide sufficient tools in order to prove the homeomorphic extension in our situation; our approach relies in an essential way on the horosphere topology and on the hyperbolicity theory; it gives an example of a deep interaction between metric properties and topological properties of a complex manifold.

As a spin off result of our work, we prove a Denjoy-Wolff Theorem for bounded convex domains  $D \subset \mathbb{C}^N$  biholomorphic to strongly convex domains and for strictly  $\mathbb{C}$ -linearly convex domains biholomorphic to bounded strongly pseudoconvex domains. Namely, we prove that if  $f$  is a holomorphic self-map of  $D$  without fixed points then the sequence of its iterates converges to exactly one boundary point (see Proposition 8.7).

We also compare the horosphere boundary we introduced with the Gromov boundary. For strongly pseudoconvex domains, by [7, Theorem 1.4], the Gromov boundary is homeomorphic to the Euclidean boundary, thus, by Theorem 1.1, homeomorphic to the horosphere boundary. Therefore, we examine in details the interesting case of the bidisc. It is standard that the bidisc is not Gromov hyperbolic. We prove that the topology of the Gromov boundary of the bidisc is not trivial while the horosphere topology of the bidisc is trivial, thus the two boundaries are not homeomorphic.

The plan of the paper is the following. In Section 2, we consider the case of simply connected domains in  $\mathbb{C}$ , both to explain the ideas underlying the horosphere theory in some simple case like the unit disc and to compare horosphere topology with the Carathéodory topology. In Section 3 we introduce the notion of admissible sequences and define the horosphere boundary. In Section 4, we introduce the horosphere topology for a general complete hyperbolic complex manifold, and the notions of impressions and principal parts for domains in  $\mathbb{C}^N$ . In Section 5, we turn our attention to the case of strongly pseudoconvex domains, we relate horosphere sequences with Abate's big and small horospheres, and we prove Theorem 1.1. Next, in Section 6 we concentrate on convex domains. After proving some preliminary results on horospheres for hyperbolic convex domains, we consider convex domains biholomorphic to strongly pseudoconvex domains and then bounded convex domains biholomorphic to strongly convex domains. In Section 7, we face the natural question of comparing horosphere boundaries and Gromov's different notions of boundaries. Finally, in Section 8, we prove our extensions results using the theory we developed in Section 6 and Gromov's theory of hyperbolic metric spaces.

*Acknowledgements.* The authors wish to thank Andrew Zimmer for fruitful conversations.

## 2. CARATHÉODORY PRIME ENDS THEORY VS. HOROSPHERE TOPOLOGY IN THE UNIT DISC

In this section we introduce the horosphere topology for simply connected domains in  $\mathbb{C}$  and compare this theory with the classical theory of Carathéodory. The section does not contain any material which will be used later on, does not contain any proof of stated facts, and can be harmlessly skipped. However, the horosphere theory we introduce is new also in dimension one. Moreover, looking first at an easy case like the unit disc might simplify comprehension of the several complex variables case. Therefore, we decided to add this section.

We start by briefly recalling Carathéodory's prime ends theory (we refer to [16, 14, 34] for details). Let  $D \subset \mathbb{C}$  be a simply connected domain. A *cross-cut* is a Jordan arc or a Jordan curve in  $\overline{D}$  such that its interior belongs to  $D$  and whose two end points belong to  $\partial D$  (if the domain  $D$  is unbounded, we consider its closure in the Riemann sphere  $\mathbb{C}\mathbb{P}^1$ ). Every cross-cut  $C$  of  $D$  divides  $D$  into two connected components by Jordan's theorem, and we say that  $C$  separates two sets  $A, B \subset D$  if  $A$  belongs to one connected component of  $D \setminus C$  and  $B$  to the other. A *null chain*  $(C_n)$  is a sequence of cross-cuts such that  $C_j \cap C_k = \emptyset$  for  $j \neq k$ , each  $C_n$  separates  $C_{n+1} \cap D$  from  $C_0 \cap D$  in  $D$  for all  $n \geq 1$  and  $\text{diam}(C_n) \rightarrow 0$ , where  $\text{diam}(C_n)$  denotes the diameter in the Euclidean metric or in the spherical metric if the domain is unbounded. For  $n \geq 1$  we denote by  $V_n$  the interior part of  $C_n$ , that is, the connected component of  $D \setminus C_n$  which does not contain  $C_0 \cap D$ .

We say that two null chains  $(C_n)$  and  $(C'_n)$  are equivalent if there exists  $n_0 \in \mathbb{N}$  such that for every  $n > n_0$ ,  $n \in \mathbb{N}$  there exists  $m \in \mathbb{N}$  such that  $V_m \subset V'_n$  and  $V'_m \subset V_n$  (here  $V_n$  is the interior part of  $C_n$  and  $V'_n$  is the interior part of  $C'_n$ ). An equivalence class of null chains is called a *prime end*. The set of all prime ends is denoted by  $\partial_C D$  and it is called the *Carathéodory boundary* of  $D$ . Let  $\hat{D}_C := D \cup \partial_C D$ .

We give a topology on  $\hat{D}_C$  as follows. A sequence  $\{z_n\} \subset D$  converges to a prime end  $\underline{x}^C \in \partial_C D$  if there exists a null chain  $(C_n)$  representing  $\underline{x}^C$  such that for every  $N \in \mathbb{N}$  the sequence  $\{z_n\}$  is eventually contained in  $V_N$ . A sequence  $\{\underline{x}_m^C\} \subset \partial_C D$  converges to  $\underline{x}^C \in \partial_C D$  if there exist null chains  $(C_n^m)$  representing  $\underline{x}_m^C$  and a null chain  $(C_n)$  representing  $\underline{x}^C$  such that for every  $N \in \mathbb{N}$  there exists  $m_0 \in \mathbb{N}$

such that for each  $m \geq m_0$  the sequence  $(C_n^m)$  is eventually contained in  $V_N$ . On  $D$  we keep the Euclidean topology. The topology generated by the previous definitions is called the *Carathéodory topology*. Two main results of Carathéodory theory are the following:

- $\hat{\mathbb{D}}_C$  endowed with the Carathéodory topology is homeomorphic to the closed unit disc  $\overline{\mathbb{D}}$  endowed with the Euclidean topology.
- If  $D_1, D_2$  are two simply connected domains,  $f : D_1 \rightarrow D_2$  a biholomorphism, then  $f$  extends to a homeomorphism  $\hat{f}_C : \hat{D}_{1C} \rightarrow \hat{D}_{2C}$ .

Given a prime end  $\underline{x}^C \in \partial_C D$ , the *prime end impression* is defined by

$$I_D^C(\underline{x}^C) := \bigcap_{n \geq 0} \overline{V_n},$$

where  $\{V_n\}$  is the interior part of any null chain  $(C_n)$  representing  $\underline{x}$  and the closure has to be understood in  $\mathbb{C}\mathbb{P}^1$  in case of unbounded domains. Note that,  $p \in I_D^C(\underline{x}^C)$  if and only if there exists a sequence  $\{z_n\} \subset D$  such that  $z_n \rightarrow p$  in the Euclidean topology and  $z_n \rightarrow \underline{x}^C$  in the Carathéodory topology.

Let  $f : \mathbb{D} \rightarrow D \subset \mathbb{C}$  be a biholomorphism,  $\zeta \in \partial\mathbb{D}$ . We denote by  $\Gamma(f; \zeta)$  the cluster set of  $f$  at  $\zeta$ , namely,  $p \in \Gamma(f; \zeta)$  if there exists a sequence  $\{z_n\} \subset \mathbb{D}$  converging to  $\zeta$  such that  $f(z_n) \rightarrow p$ . Now, the point  $\zeta$  corresponds to a point  $\underline{x}_\zeta^C \in \partial_C \mathbb{D}$ . A null chain representing  $\underline{x}_\zeta^C$  is given by  $(C_n)$  with  $C_n := \{z \in \overline{\mathbb{D}} : |z - \zeta| = r_n\}$ , where  $\{r_n\}$  is any strictly decreasing sequence of positive numbers converging to 0. Therefore, if  $\{z_n\} \subset \mathbb{D}$  converges to  $\zeta$  in the Euclidean topology, then it also converges to  $\underline{x}_\zeta^C$  in the Carathéodory topology. Now, one can choose the sequence  $\{r_n\}$  in such a way that  $(f(C_n))$  is a null chain in  $D$  which represents  $\hat{f}_C(\underline{x}_\zeta^C)$ . The sequence  $\{f(z_n)\}$  belongs eventually to the interior part of each  $\hat{C}_n$ , hence, it accumulates to points in the prime ends impression of  $\hat{f}_C(\underline{x}_\zeta^C)$ . Namely,

$$\Gamma(f; \zeta) = I_D^C(\hat{f}_C(\underline{x}_\zeta^C)).$$

In particular, a biholomorphism  $f : \mathbb{D} \rightarrow D$  extends continuously to  $\partial\mathbb{D}$  if and only if the impression of each prime end is one point.

The *principal part*  $\Pi_D^C(\underline{x}^C)$  of a prime end  $\underline{x}^C \in \partial_C D$ , is defined as follows. A point  $p \in \partial D$  belongs to  $\Pi_D^C(\underline{x}^C)$  if for every open neighborhood  $U$  of  $p$  there exists a null chain  $(C_n)$  representing  $\underline{x}^C$  such that  $C_n \subset U$  for all  $n \in \mathbb{N}$ . Given a biholomorphism  $f : \mathbb{D} \rightarrow D$  and  $\zeta \in \partial\mathbb{D}$ , let denote by  $\Gamma_{NT}(f; \zeta)$  the cluster set of  $f$  along non-tangential sequences. Namely,  $p \in \partial\mathbb{D}$  belongs to  $\Gamma_{NT}(f; \zeta)$  if there exists a sequence  $\{z_n\} \subset \mathbb{D}$ , converging non-tangentially at  $\zeta$  such that  $f(z_n) \rightarrow p$ . Then it holds

$$\Gamma_{NT}(f; \zeta) = \Pi_D^C(\hat{f}_C(\underline{x}_\zeta^C)).$$

Null chains are Euclidean objects, and the fact that a biholomorphism maps null chains almost into null chains, allowing to extend the map as a homeomorphism on the Carathéodory boundary, relies strongly on quasi-conformality of biholomorphisms. On the other hand, once this is done, and the homeomorphism between  $\hat{\mathbb{D}}_C$  and  $\overline{\mathbb{D}}$  is proved, the relation between impressions and unrestricted limits comes almost for free.

Here we take a dual point of view. We define a boundary and a topology via the intrinsic hyperbolic distance, in such a way that the homeomorphic extension of biholomorphisms to the newly defined boundary comes for free, but on the other hand, the price we have to pay is that horospheres impressions are less immediate to understand than prime ends impressions in Carathéodory theory.

In order to give some lights on the construction we present in the next sections, we describe here the horosphere topology for the unit disc and simply connected domains in  $\mathbb{C}$ . A *horosphere* of vertex  $\zeta \in \partial\mathbb{D}$

and radius  $R > 0$  in  $\mathbb{D}$  is given by

$$E(\zeta, R) := \{z \in \mathbb{D} : \frac{|\zeta - z|^2}{1 - |z|^2} < R\}.$$

It is a disc of radius  $R/(R + 1)$  contained in  $\mathbb{D}$  and tangent to  $\partial\mathbb{D}$  at  $\zeta$ . One can easily show that

$$(2.1) \quad E(\zeta, R) = \{z \in \mathbb{D} : \lim_{w \rightarrow \zeta} [K_{\mathbb{D}}(z, w) - K_D(0, w)] < \frac{1}{2} \log R\},$$

where  $K_{\mathbb{D}}$  denotes the Poincaré distance in  $\mathbb{D}$ . Equation (2.1) is however not yet suitable for being considered in other simply connected domains, essentially because it is related to the point  $\zeta$  in the Euclidean boundary of  $\mathbb{D}$ , which, from an intrinsic point of view of Poincaré distance, does not exist. Therefore, instead of considering limits to a given boundary point, we consider sequences  $\{z_n\} \subset \mathbb{D}$  such that  $\liminf_{n \rightarrow \infty} K_{\mathbb{D}}(z_n, 0) = \infty$  (namely, we consider sequences which, from an Euclidean point of view, go to the boundary). For  $R > 0$  we define

$$(2.2) \quad E^{\mathbb{D}}(\{u_n\}, R) := \{z \in \mathbb{D} : \limsup_{n \rightarrow \infty} [K_{\mathbb{D}}(z, u_n) - K_{\mathbb{D}}(0, u_n)] < \frac{1}{2} \log R\}.$$

By (2.1), if the cluster set of  $\{u_n\}$  is more than one point, there exists  $R > 0$  such that  $E(\{u_n\}, R) = \emptyset$ , while, if  $\{u_n\}$  converges to  $\zeta \in \partial\mathbb{D}$ , then

$$(2.3) \quad E^{\mathbb{D}}(\{u_n\}, R) = E(\zeta, R)$$

for all  $R > 0$ . Therefore, morally, we replace  $\zeta \in \partial\mathbb{D}$  with sequences  $\{u_n\}$  such that  $E^{\mathbb{D}}(\{u_n\}, R) = E(\zeta, R)$  for all  $R > 0$ . To be more formal, we say that a compactly divergent sequence  $\{u_n\} \subset \mathbb{D}$  is *admissible* provided  $E^{\mathbb{D}}(\{u_n\}, R) \neq \emptyset$  for all  $R > 0$ .

If  $D \subset \mathbb{C}$  is a simply connected domain, we can define admissible sequences in  $D$  using the same token: a sequence  $\{u_n\} \subset D$  is admissible if  $\liminf_{n \rightarrow \infty} K_D(x, u_n) = \infty$  and

$$E_x^D(\{u_n\}, R) := \{z \in \mathbb{D} : \limsup_{n \rightarrow \infty} [K_D(z, w) - K_D(0, w)] < \frac{1}{2} \log R\}$$

is not empty for all  $R > 0$ . Here  $K_D$  is the Poincaré distance on  $D$  and  $x \in D$  is a fixed point, whose choice does not play any substantial role.

Then we define an equivalence relation on admissible sequences by declaring  $\{u_n\}$  equivalent to  $\{v_n\}$  if for every  $R > 0$  there exist  $R', R'' > 0$  such that

$$E_x^D(\{u_n\}, R') \subset E_x^D(\{v_n\}, R) \subset E_x^D(\{u_n\}, R'').$$

In case of the unit disc, two admissible sequences  $\{u_n\}$  and  $\{v_n\}$  are then equivalent if and only if they converge to the same boundary point.

We denote by  $\partial_H D$  the set of all equivalence classes of admissible sequences in  $D$  and we call it the *horosphere boundary* of  $D$ . Let  $\hat{D} := D \cup \partial_H D$ . We want to give a topology on  $\hat{D}$  in such a way that on  $D$  it coincides with the Euclidean topology and  $\hat{D}$  is homeomorphic to  $\overline{\mathbb{D}}$ .

We start with the last requirement. As we said, if  $\underline{x} \in \partial_H \mathbb{D}$ , every admissible sequence  $\{u_n\}$  representing  $\underline{x}$  converges to the same point  $\zeta_{\underline{x}} \in \partial\mathbb{D}$ . Thus, it is natural to associate  $\underline{x}$  to the point  $\zeta_{\underline{x}}$ . Moreover, by (2.3), we can also think of  $\underline{x}$  as the family of horospheres  $E(\zeta_{\underline{x}}, R)$ ,  $R > 0$ . Now, it is easy to see that a sequence of points  $\{\zeta_j\} \in \partial\mathbb{D}$  converges to  $\zeta \in \partial\mathbb{D}$  if and only if for every  $R > 0$  there exists  $n_R \in \mathbb{N}$  such that for all  $n \geq n_R$  it holds  $E(\zeta_j, R) \cap E(\zeta, R) \neq \emptyset$ . This is exactly the definition we can exploit in the general case: let  $D \subset \mathbb{C}$  be a simply connected domain. A sequence  $\{\underline{x}_j\} \subset \partial_H D$  converges to  $\underline{x} \in \partial_H D$  if there exist admissible sequences  $\{u_n^j\}$  representing  $\underline{x}_j$  for every  $j$ , and an admissible sequence  $\{u_n\}$  representing  $\underline{x}$  such that for every  $R > 0$  there exists  $m_R \in \mathbb{N}$  such that  $E_x^D(\{u_n^j\}, R) \cap E_x^D(\{u_n\}, R) \neq \emptyset$  for all  $j \geq m_R$ .

Now, we consider convergence from inside  $\mathbb{D}$  to the horosphere boundary. It is easy to see that a sequence  $\{z_n\} \subset \mathbb{D}$  converges to  $\zeta \in \partial\mathbb{D}$  if there exists a sequence of points  $\{\zeta_n\} \subset \partial\mathbb{D}$  (not necessarily all different each other) such that for every  $R > 0$  there exists  $m_R > 0$  such that  $E(\zeta, R) \cap E(\zeta_n, R) \neq \emptyset$  and  $z_j \in E(\zeta_n, R)$  for all  $j \geq m_R$ . Thus, we can transform this observation into the general definition: a sequence  $\{z_j\} \subset D$  converges to  $\underline{x} \in \partial_H D$  if there exist admissible sequences  $\{u_n^j\} \subset D$  and an admissible sequence  $\{u_n\} \subset D$  representing  $\underline{x}$  such that for every  $R > 0$  there exists  $m_R \in \mathbb{N}$  such that  $E_x^D(\{u_n^j\}, R) \cap E_x^D(\{u_n\}, R) \neq \emptyset$  and  $z_j \in E_x^D(\{u_n^j\}, R)$  for all  $j \geq m_R$ .

With the previous definitions, we gave a meaning to the notion of convergence in  $\hat{D}$ . In this way we can define the *horosphere topology* on  $\hat{D}$  by declaring a subset  $C \subset \hat{D}$  to be closed if, whenever  $\{z_n\} \subset C$  converges in the sense described above to  $z \in \hat{D}$ , then  $z \in C$ .

By the above discussion, it is clear that  $\hat{\mathbb{D}}$  is homeomorphic to  $\overline{\mathbb{D}}$ . Moreover, let  $f : \mathbb{D} \rightarrow D \subset \mathbb{C}$  be a biholomorphism. Since the map  $f$  is an isometry between  $K_{\mathbb{D}}$  and  $K_D$  and all the constructions are made in terms of hyperbolic distance, it follows at once that  $f$  extends to a homeomorphism  $\hat{f} : \hat{\mathbb{D}} \rightarrow \hat{D}$ .

Being completely intrinsic, the definition of horosphere boundary and horosphere topology is suitable to be generalized to any complete hyperbolic complex manifold. However, for the same reason, it is harder to relate it to boundary limits, but it can be done.

Let  $D \subset \mathbb{C}$  be a simply connected domain. Given  $\underline{x} \in \partial_H D$ , we say that  $p \in I_D^H(\underline{x})$  if there exists a sequence  $\{z_n\} \subset D$  such that  $z_n \rightarrow p$  in the Euclidean topology and  $z_n \rightarrow \underline{x}$  in the horosphere topology. We call  $I_D^H(\underline{x})$  the *horosphere impression* of  $\underline{x}$ .

If  $f : \mathbb{D} \rightarrow D$  is a biholomorphism, and  $\zeta \in \partial\mathbb{D}$ , let  $\underline{x}_\zeta \in \partial_H \mathbb{D}$  be the point given by the homeomorphism between  $\overline{\mathbb{D}}$  and  $\hat{\mathbb{D}}$ . If  $\{z_n\}$  is a sequence in  $\mathbb{D}$  converging to  $\underline{x}_\zeta$  in the horosphere topology of  $\hat{\mathbb{D}}$ , then the sequence  $\{f(z_n)\}$  converges to  $\hat{f}(\underline{x}_\zeta)$  in the horosphere topology of  $\hat{D}$ . Hence,

$$\Gamma(f; \zeta) = I_D^H(\hat{f}(\underline{x}_\zeta)).$$

This implies in particular that  $I_D^H(\hat{f}(\underline{x}_\zeta)) = I_D^C(\hat{f}_C(\underline{x}_\zeta^C))$ .

We can also define a *horosphere principal part*. Let  $\underline{x} \in \partial_H D$ . We say that a sequence  $E$ -converges to  $\underline{x}$  if there exists an admissible sequence  $\{u_n\} \subset D$  representing  $\underline{x}$  such that  $\{z_n\}$  is eventually contained in  $E_x^D(\{u_n\}, R)$  for every  $R > 0$ . Clearly the definition does not depend on the admissible sequence  $\{u_n\}$  chosen. Also, it is evident that if  $\{z_n\}$  is  $E$ -converging to  $\underline{x}$  then it is also converging to  $\underline{x}$  in the horosphere topology. For instance, all sequences in  $\mathbb{D}$  which converge non-tangentially to one point  $\zeta \in \partial\mathbb{D}$ , are also  $E$ -converging to  $\underline{x}_\zeta$ . However, there exist  $E$ -convergent sequences in  $\mathbb{D}$  which do not converge non-tangentially to a boundary point. Then we let the *horosphere principal part*  $\Pi_D^H(\underline{x})$  of a point  $\underline{x} \in \partial_H D$  be the set of points  $p \in \partial D$  such that there exists a sequence  $\{z_n\}$  subset  $D$  converging to  $p$  and  $E$ -converging to  $\underline{x}$ . It is clear that  $\Pi_D^H(\underline{x}) = \bigcap_{R>0} \overline{E_x^D(\{u_n\}, R)}$  where  $\{u_n\}$  is any admissible sequence representing  $\underline{x}$ .

Now, if  $f : \mathbb{D} \rightarrow D$  is a biholomorphism, and  $\zeta \in \partial\mathbb{D}$ , let  $\Gamma_E(f; \zeta)$  be the set of points  $p \in \partial D$  such that there exists a sequence  $\{z_n\}$   $E$ -convergent to  $\underline{x}_\zeta$  such that  $f(z_n) \rightarrow p$ . Since all the notions are defined via intrinsic distance, it follows at once that

$$\Gamma_E(f; \zeta) = \Pi_D^H(\hat{f}(\underline{x}_\zeta)) = \bigcap_{R>0} \overline{E_x^D(\{f(u_n)\}, R)}.$$

Note that  $\Gamma_{NT}(f; \zeta) \subseteq \Gamma_E(f; \zeta)$ , hence,

$$\Pi_D^C(\hat{f}_C(\underline{x}_\zeta^C)) \subseteq \Pi_D^H(\hat{f}(\underline{x}_\zeta)).$$

In [20], Gaier and Pommerenke constructed examples of univalent mappings  $f : \mathbb{D} \rightarrow \mathbb{C}$  for which, in our notation,  $\Pi_D^C(\hat{f}_C(\underline{x}_\zeta^C)) \neq \Pi_D^H(\hat{f}(\underline{x}_\zeta))$ . On the other hand, Twomey [36] proved that if  $f : \mathbb{D} \rightarrow \mathbb{C}$  is starlike then  $\Gamma_E(f; \zeta)$  consists of one point for every  $\zeta \in \partial\mathbb{D}$ . Therefore, in this case,  $\Pi_D^C(\hat{f}_C(\underline{x}_\zeta^C)) = \Pi_D^H(\hat{f}(\underline{x}_\zeta))$ .



### 3. ADMISSIBLE SEQUENCES, BUSEMANN SEQUENCES AND HOROSPHERE BOUNDARY

Let  $M$  be a (connected) complex manifold of complex dimension  $N$ . We denote by  $K_M$  the Kobayashi distance on  $M$ . We assume that  $M$  is (Kobayashi) complete hyperbolic, namely that the metric space  $(M, K_M)$  is complete.

Let  $\{u_n\}$  be a sequence of points in  $M$ . For  $R > 0$  and  $x \in M$ , we denote by  $E_x(\{u_n\}, R)$  the (sequence) horosphere defined by

$$E_x(\{u_n\}, R) := \{w \in M \mid \limsup_{n \rightarrow \infty} (K_M(w, u_n) - K_M(x, u_n)) < \frac{1}{2} \log R\}.$$

**Lemma 3.1.** *Let  $x, y \in M$ . Let  $\{u_n\}$  be a sequence in  $M$ . Then there exist  $\alpha, \beta > 0$  such that for all  $R > 0$*

$$E_y(\{u_n\}, \alpha R) \subset E_x(\{u_n\}, R) \subset E_y(\{u_n\}, \beta R).$$

*Proof.* Let  $w \in E_x(\{u_n\}, R)$ . Let  $\frac{1}{2} \log \beta := \limsup_{n \rightarrow \infty} [K_M(x, u_n) - K_M(y, u_n)]$ . Note that  $\beta \in (0, \infty)$  since

$$-\infty < -K_M(x, y) \leq \frac{1}{2} \log \beta \leq K_M(x, y) < +\infty.$$

Then

$$\begin{aligned} \limsup_{n \rightarrow \infty} [K_M(w, u_n) - K_M(y, u_n)] &\leq \limsup_{n \rightarrow \infty} [K_M(w, u_n) - K_M(x, u_n)] \\ &\quad + \limsup_{n \rightarrow \infty} [K_M(x, u_n) - K_M(y, u_n)] < \frac{1}{2} \log R + \frac{1}{2} \log \beta, \end{aligned}$$

hence  $w \in E_y(\{u_n\}, \beta R)$ .

Similarly, if  $\frac{1}{2} \log \alpha := -\limsup_{n \rightarrow \infty} [K_M(y, u_n) - K_M(x, u_n)]$ , we obtain  $E_y(\{u_n\}, \alpha R) \subset E_x(\{u_n\}, R)$ .  $\square$

**Definition 3.2.** Let  $M$  be a complete hyperbolic manifold. Let  $x \in M$ . A sequence  $\{u_n\}$  is *admissible* if

- (i)  $\liminf_{n \rightarrow \infty} K_M(x, u_n) = \infty$ ,
- (ii)  $\forall R > 0, E_x(\{u_n\}, R) \neq \emptyset$ .

We denote by  $\Lambda_M$  the set of admissible sequences.

By Lemma 3.1, the definition of admissible sequence does not depend on the base point  $x$  chosen to define horospheres.

The construction of admissible sequences is valid for any geodesic, proper, complete metric space. The existence of admissible sequences in that general situation is given by the following proposition, communicated to the authors by Andrew Zimmer. In case  $M$  is a complete hyperbolic manifold for the Kobayashi metric, the existence of a geodesic joining any two points comes from the Hopf-Rinow Theorem in locally compact, complete length spaces. We state Proposition 3.3 for complete Kobayashi hyperbolic manifolds, to keep the context of the paper.

**Proposition 3.3.** *Let  $M$  be a complete hyperbolic manifold and let  $x \in M$ . Then every sequence  $\{u_n\}$  of points in  $M$ , such that  $\lim_{n \rightarrow \infty} K_M(u_n, x) = +\infty$ , admits an admissible subsequence. In particular,  $\Lambda_M \neq \emptyset$ .*

*Proof.* Consider any sequence  $\{u_n\}$  converging to infinity for the Kobayashi distance. The function  $b_n : M \ni z \mapsto K_M(u_n, z) - K_M(u_n, 0)$  is 1-Lipschitz for every  $n$ . According to the Ascoli-Arzelà Theorem, the sequence  $\{b_n\}$  admits a subsequence  $\{b_{\varphi(n)}\}$  that converges, uniformly on compact subsets of  $M$ , to

some function  $b$ . For every  $n \geq 0$ , let  $\gamma_{\varphi(n)} : [0, T_n] \rightarrow M$  be a real geodesic joining 0 to  $u_{\varphi(n)}$ . Then by the definition of  $b_n$  we get:

$$\begin{aligned} -\forall 0 \leq t \leq T_{\varphi(n)}, b_{\varphi(n)}(\gamma_{\varphi(n)}(t)) &= K_D(\gamma_{\varphi(n)}(T_{\varphi(n)}), \gamma_{\varphi(n)}(t)) - K_D(\gamma_{\varphi(n)}(T_{\varphi(n)}), \gamma_{\varphi(n)}(0)) \\ &= -K_D(\gamma_{\varphi(n)}(t), \gamma_{\varphi(n)}(0)) \\ &= -t. \end{aligned}$$

Since  $(M, K_M)$  is a proper metric space, it follows from the Ascoli-Arzelà Theorem that the sequence  $\{\gamma_{\varphi(n)}\}$  admits a subsequence  $\{\gamma_{\sigma(n)}\}$  that converges, locally uniformly, to some  $\gamma$ . In particular we get  $b(\gamma(t)) = -t$  for every  $t \geq 0$ .

Let us fix  $r > 0$  and consider a real number  $t_r$  satisfying  $t_r > -\frac{1}{2} \log(r)$ . Then for sufficiently large  $n$  we have:

$$b_{\sigma(n)}(\gamma(t_r)) < \frac{1}{2} \log(r),$$

meaning that  $\gamma(t_r) \in E_x(\{u_{\sigma(n)}\}, r)$ . □

We collect here some properties of horospheres:

**Proposition 3.4.** *Let  $M$  be a complete hyperbolic manifold and let  $x \in M$ . Let  $\{u_n\}$  be an admissible sequence. Then*

- (1)  $E_x(\{u_n\}, R)$  is open for every  $R > 0$ .
- (2) If  $0 < R < R'$  then  $E_x(\{u_n\}, R) \subset E_x(\{u_n\}, R')$ .
- (3)  $\bigcap_{R>0} E_x(\{u_n\}, R) = \emptyset$ .

*Proof.* (1) If  $w \in E_x(\{u_n\}, R)$ , then there exists  $0 < R' < R$  such that

$$\limsup_{n \rightarrow \infty} (K_M(w, u_n) - K_M(x, u_n)) = \frac{1}{2} \log R'.$$

Hence, if  $z \in B_K(w, \delta)$ , the Kobayashi ball of center  $w$  and radius  $\delta = \frac{1}{2} \log \frac{R}{R'}$ , it follows

$$\begin{aligned} K_M(z, u_n) - K_M(x, u_n) &= K_M(z, u_n) - K_M(w, u_n) + K_M(w, u_n) - K_M(x, u_n) \\ &\leq K_M(z, w) + K_M(w, u_n) - K_M(x, u_n), \end{aligned}$$

and then

$$\limsup_{n \rightarrow \infty} [K_M(z, u_n) - K_M(x, u_n)] < \frac{1}{2} \log \frac{R}{R'} + \frac{1}{2} \log R' = \frac{1}{2} \log R.$$

Therefore,  $B_K(w, \delta) \subset E_x(\{u_n\}, R)$ , and the horosphere is open.

(2) is obvious.

(3) Since for every  $w \in M$ ,

$$-K_M(x, w) \leq K_M(w, u_n) - K_M(x, u_n),$$

it follows that if  $w \in \bigcap_{R>0} E_x(\{u_n\}, R)$  then  $K_M(x, w) = \infty$ , a contradiction. □

A less obvious property is the following property which states somewhat that horospheres go “uniformly at the boundary”:

**Proposition 3.5.** *Let  $M$  be a complete hyperbolic manifold and let  $x \in M$ . Then for every compact set  $K \subset M$  there exists  $R_0 > 0$  such that*

$$E_x(\{u_n\}, R) \cap K = \emptyset$$

for all  $R \leq R_0$  and all admissible sequence  $\{u_n\} \subset M$ .

*Proof.* Assume by contradiction this is not the case. Then for every  $N \in \mathbb{N}$  there exist  $z_N \in K$  and an admissible sequence  $\{u_n^N\} \subset M$  such that  $z_N \in E_x(\{u_n^N\}, 1/N)$ . Up to subsequences, we can assume that  $\{z_N\}$  converges to  $z_0 \in K$  and that for all  $N \in \mathbb{N}$  it holds  $K_D(z_N, z_0) < 1$ . Then, using the triangle inequality, for all  $n \in \mathbb{N}$  we have

$$\begin{aligned} -K_D(z_0, x) &\leq \limsup_{n \rightarrow \infty} [K_D(z_0, u_n^N) - K_D(x, u_n^N)] \\ &\leq \limsup_{n \rightarrow \infty} [K_D(z_N, z_0) + K_D(z_N, u_n^N) - K_D(x, u_n^N)] \\ &\leq 1 + \limsup_{n \rightarrow \infty} [K_D(z_N, u_n^N) - K_D(x, u_n^N)] < 1 + \frac{1}{2} \log \frac{1}{N}, \end{aligned}$$

a contradiction.  $\square$

Another natural class of sequences is the following:

**Definition 3.6.** Let  $M$  be a complete hyperbolic manifold,  $x \in M$ . A sequence  $\{u_n\} \subset M$  is called a *Busemann sequence* if

- (i)  $\liminf_{n \rightarrow \infty} K_M(x, u_n) = \infty$ ,
- (ii) for all  $z \in M$  the limit  $\lim_{n \rightarrow \infty} [K_M(z, u_n) - K_M(x, u_n)]$  exists.

We denote by  $\mathcal{B}_M$  the set of Busemann sequences.

It is not difficult to show that the definition of Busemann sequence does not depend on the base point  $x \in M$  chosen.

Busemann sequences have been introduced (under the name *horosphere sequences*) in balanced bounded convex domains in [26], and later studied in bounded convex domains in [9]. The proof of the following result is an adaptation of the ideas contained in those papers.

**Proposition 3.7.** *Let  $M$  be a complete hyperbolic manifold,  $x \in M$ .*

- (1) *If  $\{u_n\} \subset M$  satisfies  $\lim_{n \rightarrow \infty} K_M(x, u_n) = \infty$ , then there exists a subsequence  $\{u_{n_k}\}$  of  $\{u_n\}$  which is a Busemann sequence. Therefore,  $\mathcal{B}_M \neq \emptyset$ .*
- (2) *If  $\{u_n\} \subset M$  is an admissible sequence, then there exists a subsequence  $\{u_{n_k}\}$  of  $\{u_n\}$  which is a Busemann admissible sequence and  $E_x(\{u_n\}, R) \subseteq E_x(\{u_{n_k}\}, R)$  for all  $R > 0$ .*
- (3) *If for every two points  $z, w \in M$ ,  $z \neq w$  there exists a complex geodesic (i.e., an analytic disc which is an isometry between  $K_{\mathbb{D}}$  and  $K_D$ ) which contains  $z, w$  then every Busemann sequence is an admissible sequence.*

*Proof.* (1) Let  $\{z_m\}_{m \in \mathbb{N}}$  be a dense set in  $M$ . Using a standard diagonal argument one can find a subsequence  $\{u_{n_j}\}$  such that  $f_j(z) := K_D(z, u_{n_j}) - K_D(z, u_{n_j})$  has the property that  $\lim_{j \rightarrow \infty} f_j(z_m)$  exists for all  $m \in \mathbb{N}$ . If  $z \in M$  and  $\{z_{m_k}\}$  is a subsequence converging to  $z$ , by the triangle inequality it follows

$$|f_j(z) - f_j(z_{m_k})| \leq K_D(z, z_{m_k}).$$

From this it is easy to see that  $\lim_{j \rightarrow \infty} f_j(z)$  exists, hence  $\{u_{n_j}\} \in \mathcal{B}_M$ .

(2) If  $\{u_n\} \in \Lambda_M$ , then by (1) we can extract a Busemann subsequence  $\{u_{n_k}\}$ . Let  $R > 0$  and let  $z \in E_x(\{u_n\}, R)$ . Then

$$\limsup_{k \rightarrow \infty} [K_D(z, u_{n_k}) - K_D(z, u_{n_k})] \leq \limsup_{n \rightarrow \infty} [K_D(z, u_n) - K_D(z, u_n)] < \frac{1}{2} \log R,$$

which proves that  $z \in E_x(\{u_{n_k}\}, R)$ .

(3) Assume that every two points of  $M$  belongs to a complex geodesic and let  $\{u_n\} \in \mathcal{B}_M$ . Fix  $R > 0$  and let  $0 < r < \min\{1, R\}$ . Let  $B_k(x, -\frac{1}{2} \log r)$  be the Kobayashi ball of center  $x$  and radius  $-\frac{1}{2} \log r > 0$ . Since  $K_D(x, \{u_n\}) \rightarrow \infty$ , we can assume that  $u_n \notin B_k(x, -\frac{1}{2} \log r)$  for all  $n \in \mathbb{N}$ . By hypothesis  $x$  and

$u_n$  are contained in a complex geodesic. Hence, there exists  $q_n$  in the same complex geodesic such that  $k_D(x, q_n) = -\frac{1}{2} \log r$  and

$$K_D(x, u_n) = K_D(x, q_n) + K_D(q_n, u_n) = -\frac{1}{2} \log r + K_D(q_n, u_n).$$

Note that  $q_n \in \partial B_k(x, -\frac{1}{2} \log r)$  and the latter is compact. Hence we can extract a converging subsequence  $\{q_{n_k}\}$  converging to some  $q \in \partial B_k(x, -\frac{1}{2} \log r)$ . Now, taking into account that  $\{u_n\}$  is a Busemann sequence, we have

$$\begin{aligned} \limsup_{n \rightarrow \infty} [K_D(q, u_n) - K_D(x, u_n)] &= \lim_{k \rightarrow \infty} [K_D(q, u_{n_k}) - K_D(x, u_{n_k})] \\ &\leq \frac{1}{2} \log r + \lim_{k \rightarrow \infty} K_D(q, q_{n_k}) = \frac{1}{2} \log r, \end{aligned}$$

that is  $q \in E_x(\{u_n\}, R)$ . Hence  $\{u_n\}$  is admissible.  $\square$

**Definition 3.8.** Let  $M$  be a complete hyperbolic manifold and let  $x \in M$ .

- (i) Two sequences  $\{u_n\}, \{v_n\}$  in  $\Lambda_M$  are *equivalent* and we write  $\{u_n\} \sim \{v_n\}$  if for every  $R > 0$  there exist  $R' > 0$  and  $R'' > 0$  such that

$$E_x(\{u_n\}, R') \subseteq E_x(\{v_n\}, R), \quad E_x(\{v_n\}, R'') \subseteq E_x(\{u_n\}, R).$$

- (ii) We denote by  $\partial_H M$  the *horosphere boundary* of  $M$  defined by  $\partial_H M := \Lambda_M / \sim$ .

It is easy to see that  $\sim$  is an equivalence relation on  $\Lambda_M$ . Moreover, by Lemma 3.1, the definition of equivalence admissible sequences is independent of the base point  $x$ .

**Remark 3.9.** By Proposition 3.3, the horosphere boundary of a complete hyperbolic manifold is always not empty.

#### 4. HOROSPHERE TOPOLOGY, IMPRESSION AND PRINCIPAL PART

Let  $M$  be a complete hyperbolic complex manifold,  $x \in M$ , and let  $\partial_H M$  be its horosphere boundary. Let  $\hat{M} := M \cup \partial_H M$ . We define a topology on  $\hat{M}$ , which coincides with the topology of  $M$  on  $M$ , as follows:

**Definition 4.1.** A sequence  $\{y_m\} \subset \partial_H M$  converges to  $\underline{y} \in \partial_H M$  if there exist admissible sequences  $\{u_n^m\}_{n \in \mathbb{N}}$  with  $[\{u_n^m\}] = y_m$  and an admissible sequence  $\{u_n\}$  with  $[\{u_n\}] = \underline{y}$  with the property that for every  $R > 0$  there exists  $m_R \in \mathbb{N}$  such that

$$(4.1) \quad E_x(\{u_n^m\}, R) \cap E_x(\{u_n\}, R) \neq \emptyset \quad \forall m \geq m_R.$$

By Lemma 3.1, the previous definition of convergence is independent of the base point  $x$ .

**Definition 4.2.** A sequence  $\{z_m\} \subset M$  converges to  $\underline{y} \in \partial_H M$  if there exist  $\{u_n\}, \{v_n^j\}_{j \in \mathbb{N}} \subset \Lambda_M$  with  $[\{u_n\}] = \underline{y}$  and with the property that for all  $R > 0$  there exists  $m_R \in \mathbb{N}$  such that  $z_m \in E_x(\{v_n^m\}, R)$ , and  $E_x(\{v_n^m\}, R) \cap E_x(\{u_n\}, R) \neq \emptyset$  for all  $m \geq m_R$ .

It is clear from Lemma 3.1 that this definition does not depend on the base point  $x$ .

**Remark 4.3.** Note that, if the  $\{v_n^j\}$  are admissible sequences as in Definition 4.2, and  $\underline{y}_j := [\{v_n^j\}]$ , then  $\{\underline{y}_j\}$  converges to  $\underline{y}$  in the sense of Definition 4.1.

A sequence  $\{z_n\} \subset M \subset \hat{M}$  converges to  $z \in M$  if  $\lim_{n \rightarrow \infty} K_M(z_n, z) = 0$ . Now we can define a topology on  $\hat{M}$  as follows:

**Definition 4.4.** A subset  $C \subset \hat{M}$  is closed if for every sequence  $\{z_n\} \subset C$  converging to  $z \in \hat{M}$ , it follows that  $z \in C$ .

We call such a topology the *horosphere topology* of  $M$  and we denote it by  $\mathcal{T}_H(\hat{M})$ .

**Remark 4.5.** By definition, a sequence  $\{z_n\} \subset \partial_H M$  can converge only to points which belong to  $\partial_H M$  in the horosphere topology.

**Remark 4.6.** A relatively compactly sequence  $\{z_n\} \subset M$  can not converge to a point  $\underline{y} \in \partial_H M$  by Proposition 3.5. Therefore the horosphere topology restricted on  $M$  coincides with the topology of  $M$ .

If  $M, N$  are two complete hyperbolic complex manifolds and  $F : M \rightarrow N$  is a biholomorphism, then  $F$  induces a bijective map  $\hat{F} : \hat{M} \rightarrow \hat{N}$  as follows. If  $z \in M$  then  $\hat{F}(z) = F(z)$ . If  $\underline{y} \in \partial_H M$ , let  $\{u_n\}$  be an admissible sequence in  $M$  which represents  $\underline{y}$ . Since  $F$  is an isometry between  $\bar{K}_M$  and  $K_N$ , it follows easily that  $\{F(u_n)\}$  is an admissible sequence in  $N$  and therefore it represents a point  $\hat{F}(\underline{y}) \in \partial_H N$ . Clearly, since  $F$  is an isometry between the Kobayashi distance of  $M$  and that of  $N$ , such a point does not depend on the admissible sequence chosen to represent it.

Moreover, by the same token, it follows that  $\hat{F}^{-1} = \widehat{F^{-1}}$  and  $\hat{F}$  and  $\widehat{F^{-1}}$  map closed sets onto closed sets in the horosphere topology. Therefore we have

**Theorem 4.7.** Let  $M, N$  be complete hyperbolic complex manifolds. Let  $F : M \rightarrow N$  be a biholomorphism. Then  $F$  extends to a homeomorphism  $\hat{F} : \hat{M} \rightarrow \hat{N}$ .

In case of domains in  $\mathbb{C}^N$  one can relate convergence to the horosphere boundary in the horosphere topology with (Euclidean) convergence to the Euclidean boundary. This can be done in two natural ways using horosphere topology, and gives rise to the notion of horosphere impression and horosphere principal part. These notions, as it will be clear later on, are strictly related to extension of univalent maps from bounded strongly pseudoconvex domains.

As a matter of notation, if  $D \subset \mathbb{C}^N$  is a domain, we denote by  $\overline{D}^{\mathbb{C}P^N}$  its closure in  $\mathbb{C}P^N$ .

**Definition 4.8.** Let  $D$  be a complete hyperbolic domain in  $\mathbb{C}^n$ . Let  $\underline{x} \in \partial_H D$ . We say that a point  $p \in \overline{D}^{\mathbb{C}P^N}$  belongs to the *horosphere impression* of  $\underline{x}$ , denoted by  $I_D^H(\underline{x})$ , if there exists a sequence  $\{w_n\} \subset D$  such that  $\{w_n\}$  converges to  $\underline{x}$  in the horosphere topology and  $\{w_n\}$  converges to  $p$  in  $\mathbb{C}P^N$ .

**Remark 4.9.** By Proposition 3.5, for every  $\underline{x} \in \partial_H D$  it holds  $I_D^H(\underline{x}) \cap D = \emptyset$ .

In order to define the horosphere principal part, we premise a general definition of convergence in the horosphere topology, which, somehow, replace the notion of non-tangential limits:

**Definition 4.10.** Let  $M$  be a complete hyperbolic manifold,  $x \in M$ . Let  $\underline{x} \in \partial_H M$ . We say that a sequence  $\{z_n\} \subset M$  is *E-converging* to  $\underline{x}$ , and we write

$$E - \lim_{n \rightarrow \infty} z_n = \underline{x},$$

if for one—and hence any—admissible sequence  $\{u_n\}$  representing  $\underline{x}$ , the sequence  $\{z_n\}$  is eventually contained in  $E_x(\{u_n\}, R)$  for all  $R > 0$ .

Note that, according to the definition of horosphere topology, if  $\{z_n\}$  is E-converging to  $\underline{x}$  then, in particular, it is converging to  $\underline{x}$  in the horosphere topology.

We are now ready to define the *horosphere principal part*:

**Definition 4.11.** Let  $D$  be a complete hyperbolic domain in  $\mathbb{C}^n$ . Let  $\underline{x} \in \partial_H D$ . We say that a point  $p \in \overline{D}^{\mathbb{C}P^N}$  belongs to the *horosphere principal part* of  $\underline{x}$ , denoted by  $\Pi_D^H(\underline{x})$ , if there exists a sequence  $\{w_n\} \subset D$  such that  $\{w_n\}$  E-converges to  $\underline{x}$  and  $\{w_n\}$  converges to  $p$  in  $\mathbb{C}P^N$ .

Clearly,  $\Pi_D^H(\underline{x}) \subset I_D^H(\underline{x})$  for all  $\underline{x} \in \partial_H D$ . By the very definition we have

**Lemma 4.12.** *Let  $D$  be a complete hyperbolic domain in  $\mathbb{C}^n$ ,  $x \in D$ . Let  $\underline{x} \in \partial_H D$ . Then*

$$I_D^H(\underline{x}) = \bigcap_{R>0} \overline{E_x(\{u_n\}, R)}^{\mathbb{C}P^N},$$

where  $\{u_n\} \subset D$  is any admissible sequence representing  $\underline{x}$ .

## 5. STRONGLY PSEUDOCONVEX DOMAINS

Abate (see [1, 3]) defined the small and big horospheres as follows:

**Definition 5.1.** Let  $D \subset \mathbb{C}^N$  be a domain and let  $p \in \partial D$ . Let  $R > 0$  and  $x \in D$ . The *small horosphere*  $E_x(p, R)$  of vertex  $p$  and radius  $R$  is defined by

$$E_x(p, R) := \{w \in D : \limsup_{z \rightarrow p} [K_D(w, z) - K_D(x, z)] < \frac{1}{2} \log R\}.$$

The *big horosphere*  $F_x(p, R)$  of vertex  $p$  and radius  $R$  is defined by

$$F_x(p, R) := \{w \in D : \liminf_{z \rightarrow p} [K_D(w, z) - K_D(x, z)] < \frac{1}{2} \log R\}.$$

The following result follows from [1, Lemma 1.1, Thm. 1.7] and [3, Thm. 2.6.47]

**Theorem 5.2.** *Let  $D \subset \mathbb{C}^N$  be a bounded strongly pseudoconvex domain with  $C^2$  boundary. Let  $p \in \partial D$  and  $x \in D$ . Then*

- (1) for every  $R > 0$ ,  $E_x(p, R) \subset F_x(p, R)$ ,
- (2) for every  $R' > R > 0$ ,  $E_x(p, R) \subset E_x(p, R')$ ,  $F_x(p, R) \subset F_x(p, R')$ ,
- (3)  $\bigcap_{R>0} F_x(p, R) = \emptyset$ ,
- (4) for every  $R > 0$ ,  $\overline{F_x(p, R)} \cap \partial D = \{p\}$ .

Moreover, if  $D$  is a strongly convex domain with  $C^3$  boundary, then for every  $R > 0$  it holds  $E_x(p, R) = F_x(p, R)$ .

In what follows, we need the following lemma.

**Lemma 5.3.** *Let  $D \subset \mathbb{C}^N$  be a bounded strongly pseudoconvex domain with  $C^2$  boundary. Let  $V \subset \partial D$  be a closed set and  $x \in D$ . Then for every open neighborhood  $U$  of  $V$  there exists  $R_0 > 0$  such that*

$$\bigcup_{p \in V} F_x(p, R) \subset U \cap D \quad \text{for all } 0 < R \leq R_0.$$

In particular, if  $V, V' \subset \partial D \setminus \{p\}$  are two closed sets such that  $V \cap V' = \emptyset$  then there exists  $R_0 > 0$  such that, for all  $0 < R < R_0$  it holds  $\bigcup_{p \in V} F_x(p, R) \cap \bigcup_{p \in V'} F_x(p, R) = \emptyset$ .

*Proof.* Let  $R > 0$ . First, note that

$$F_x(V, R) := \{z \in D : \liminf_{w \rightarrow V} [K_D(z, w) - K_D(x, w)] < \frac{1}{2} \log R\} = \bigcup_{p \in V} F_x(p, \frac{1}{2} \log R).$$

Indeed, if  $z \in F_x(V, R)$ , then there exists a sequence  $\{w_m\}$  such that

$$\lim_{m \rightarrow \infty} [K_D(z, w_m) - K_D(x, w_m)] = \liminf_{w \rightarrow V} [K_D(z, w) - K_D(x, w)].$$

Up to subsequences, we can assume that  $\{w_m\}$  converges to  $p \in V$ . Therefore,

$$\liminf_{w \rightarrow p} [K_D(z, w) - K_D(x, w)] \leq \lim_{m \rightarrow \infty} [K_D(z, w_m) - K_D(x, w_m)] < \frac{1}{2} \log R,$$

hence,  $z \in F_x(p, R)$ . Conversely, if  $z \in F_x(p, R)$  then

$$\liminf_{w \rightarrow V} [K_D(z, w) - K_D(x, w)] \leq \liminf_{w \rightarrow p} [K_D(z, w) - K_D(x, w)] < \frac{1}{2} \log R,$$

proving that  $z \in F_x(V, R)$ .

Now, we argue by contradiction. Let assume there exists an open neighborhood  $U$  of  $V$  such that, for all  $n \in \mathbb{N}$  there exists  $z_n \in F_x(V, \frac{1}{n}) \cap (D \setminus U)$ . Up to subsequence, we can assume that  $z_n \rightarrow z_0 \in \overline{D \setminus U}$ . Arguing as in the proof of Proposition 3.5, it is not hard to see that  $z_0 \notin D$ . Therefore,  $z_0 \in \partial D$ . Fix  $R > 0$ . Since  $F_x(V, R) \subset F_x(V, R')$  for all  $0 < R < R'$ , it follows that  $z_n \in F_x(V, R)$  for all  $n > 1/R$ . In particular, by Theorem 5.2.(4),

$$z_0 \in \overline{F_x(V, R)} \cap \partial D = \bigcup_{p \in V} \overline{F_x(p, R)} \cap \partial D = V.$$

But then  $z_0 \in V$ , again, a contradiction.  $\square$

As a matter of notation, if  $D \subset \mathbb{C}^N$  is a domain and  $z \in D$  let  $\delta(z)$  denote the distance of  $z$  from  $\partial D$ .

We prove the following optimal localization principle for the Kobayashi distance on strongly pseudoconvex domains. An associated result was given by Z. Balogh and M. Bonk in [7]. Our more precise statement relies on their approach. After a first draft of our paper was completed, N. Nikolov informed us that one can obtain a similar estimate using the techniques he developed in [33].

**Lemma 5.4.** *Let  $D \subset \mathbb{C}^N$  be a bounded strongly pseudoconvex domain with  $C^2$  boundary. Let  $p \in \partial D$ . Let  $U$  be an open neighborhood of  $p$ . Then there exists an open neighborhood  $W \subset U$  of  $p$  and a constant  $T \geq 1$  such that for every  $z, w \in W \cap D$  it holds*

$$(5.1) \quad K_{U \cap D}(w, z) - K_D(w, z) \leq \frac{1}{2} \log T.$$

*Proof.* First of all, notice that, given any open set  $U$  of  $p$ , there exists an open neighborhood  $\tilde{U} \subset U$  of  $p$  such that  $\tilde{U} \cap D$  is a strongly pseudoconvex domain with  $C^2$  boundary. If (5.1) holds for  $\tilde{U}$ , then for every  $z, w \in W \cap D$

$$K_{U \cap D}(w, z) - K_D(w, z) \leq K_{\tilde{U} \cap D}(w, z) - K_D(w, z) \leq \frac{1}{2} \log T,$$

and hence (5.1) holds for  $U$  as well.

Therefore, without loss of generality, we can assume that  $U \cap D$  is a strongly pseudoconvex domain with  $C^2$  boundary.

For  $x, y \in \partial D$ , let  $d_H^{\partial D}(x, y)$  denote the Carnot-Carathéodory distance between  $x$  and  $y$ . The distance  $d_H^{\partial D}(x, y)$  is defined as follows (see, e.g., [7]). A piecewise  $C^1$ -smooth curve  $\alpha : [0, 1] \rightarrow \partial D$  is *horizontal* provided  $\alpha'(t) \in T_{\alpha(t)}^{\mathbb{C}} \partial D$  for almost every  $t$ . The set of horizontal curves is denoted by  $\mathcal{H}(\partial D)$ . For every  $x, y \in \partial D$  there exist horizontal curves joining  $x$  and  $y$ . Let

$$\rho_D(z) := \begin{cases} -\delta(z) & \text{for } z \in D \\ \delta(z) & \text{for } z \in \mathbb{C}^N \setminus D \end{cases}$$

Then  $\rho_D$  is  $C^2$  on an open neighborhood of  $\partial D$ . Let  $L_{\rho_D}$  be the Levi form of  $\rho_D$ . The Levi-length of a horizontal curve  $\alpha$  is defined as

$$\ell_L^D(\alpha) := \int_0^1 (L_{\rho_D}(\alpha(t); \alpha'(t)))^{1/2} dt.$$

Then

$$d_H^{\partial D}(x, y) := \inf \{ \ell_L^D(\alpha) : \alpha \in \mathcal{H}(\partial D), \alpha(0) = x, \alpha(1) = y \}.$$

Let  $W'$  be an open neighborhood of  $p \in \partial D$  such that for every  $z \in W' \cap D$  there exists a unique point  $\pi_D(z) \in \partial D$  such that

$$\delta(z) = |z - \pi_D(z)|.$$

For  $z, w \in W' \cap D$  define

$$g_D(z, w) := 2 \log \left[ \frac{d_H^{\partial D}(\pi_D(z), \pi_D(w)) + \max\{\sqrt{\delta(z)}, \sqrt{\delta(w)}\}}{\sqrt{\delta(z)\delta(w)}} \right].$$

By [7, Corollary 1.3] there exists  $C_D \geq 0$  such that for all  $z, w \in W' \cap D$  it holds

$$(5.2) \quad g_D(z, w) - C_D \leq K_D(z, w) \leq g_D(z, w) + C_D.$$

Similarly, there exists  $C_{U \cap D} \geq 0$  such that for all  $z, w \in W' \cap D$  it holds

$$(5.3) \quad g_{U \cap D}(z, w) - C_{U \cap D} \leq K_{U \cap D}(z, w) \leq g_{U \cap D}(z, w) + C_{U \cap D}.$$

By (5.2) and (5.3), for all  $z, w \in W' \cap D$  we have

$$(5.4) \quad K_{U \cap D}(w, z) - K_D(w, z) \leq g_{U \cap D}(z, w) - g_D(z, w) + C_D + C_{U \cap D}.$$

Now, up to shrinking  $W'$  is necessary, we can assume that for all  $z \in W' \cap D$  it holds

$$\delta^{U \cap D}(z) = \delta(z),$$

where  $\delta^{U \cap D}(z)$  denotes the distance from  $z$  to the boundary of  $U \cap D$ . With such an assumption,  $\pi(z) := \pi_D(z) = \pi_{U \cap D}(z)$ . We claim that there exists a open neighborhood  $W \subseteq W'$  such that for all  $z, w \in W \cap D$  it holds

$$(5.5) \quad d_H^{\partial D}(\pi(z), \pi(w)) = d_H^{\partial(U \cap D)}(\pi(z), \pi(w)).$$

Once this is proved, it follows from (5.4) that for all  $z, w \in W \cap D$

$$K_{U \cap D}(w, z) - K_D(w, z) \leq \frac{1}{2} \log T := C_D + C_{U \cap D},$$

and the lemma is proved.

In order to prove (5.5), we argue as follows. By the ‘‘box-ball estimate’’ (see [7, eq. (3.1)]) there exist  $A_1, A_2 > 0$  such that

$$A_1|x - y| \leq d_H^{\partial D}(x, y) \leq A_2|x - y|^{1/2},$$

for all  $x, y \in \partial D$ . In particular this implies that there exists an open neighborhood  $W_1 \subset W'$  of  $p$  such that for every  $x, y \in W_1 \cap \partial D$  and for every horizontal curve  $\alpha$  joining  $x$  and  $y$  such that  $\alpha([0, 1]) \cap (\partial D \setminus W') \neq \emptyset$  it holds

$$(5.6) \quad \ell_L^D(\alpha) \geq d_H^{\partial D}(x, y) + 1.$$

Similarly, that there exists an open neighborhood  $W_2 \subset W'$  of  $p$  such that for every  $x, y \in W_2 \cap \partial D$  and for every horizontal curve  $\alpha$  joining  $x$  and  $y$  such that  $\alpha([0, 1]) \cap (\partial(U \cap D) \setminus W') \neq \emptyset$  it holds

$$(5.7) \quad \ell_L^{U \cap D}(\alpha) \geq d_H^{\partial(U \cap D)}(x, y) + 1.$$

Let  $W \subset W_1 \cap W_2$  be an open neighborhood of  $p$  such that  $\pi(W \cap D) \subset\subset (W_1 \cap W_2) \cap \partial D$ . Let  $z, w \in W \cap D$ . Let  $0 < \epsilon < 1$ . Let  $\alpha$  be a horizontal curve joining  $\pi(z)$  and  $\pi(w)$  such that

$$\ell_L^D(\alpha) \leq d_H^{\partial D}(\pi(z), \pi(w)) + \epsilon.$$

By (5.6),  $\alpha([0, 1]) \subset W'$ . Since  $\rho_D(z) = \rho_{U \cap D}(z)$  for all  $z \in W' \cap D$ , it follows that  $\ell_L^{U \cap D}(\alpha) = \ell_L^D(\alpha)$ . Hence

$$d_H^{\partial(U \cap D)}(\pi(z), \pi(w)) \leq \ell_L^{U \cap D}(\alpha) = \ell_L^D(\alpha) \leq d_H^{\partial D}(\pi(z), \pi(w)) + \epsilon,$$



and by the arbitrariness of  $\epsilon$ , it follows that  $d_H^{\partial(U \cap D)}(\pi(z), \pi(w)) \leq d_H^{\partial D}(\pi(z), \pi(w))$ . A similar argument gives the opposite inequality, and the proof is completed.  $\square$

In what follows, we need also the following boundary estimates for the Kobayashi distance, see, e.g., [18], [3, Thm. 2.3.56], [3, Thm. 2.3.54] and [7, p. 530-531]. Note that, in those references, the estimates are proved for a given point in  $\partial D$ . However, the compactness of  $\partial D$  allows easily to get uniform estimates.

**Lemma 5.5.** *Let  $D \subset \mathbb{C}^N$  be a bounded domain with  $C^2$  boundary. Then there exist  $C > 0$  and  $\epsilon_0 > 0$  such that for every  $p \in \partial D$  and for every  $z, w \in D \cap B(p, \epsilon_0)$  it holds:*

$$(5.8) \quad K_D(w, z) \leq \frac{1}{2} \log \left( 1 + \frac{|w - z|}{\delta(w)} \right) + \frac{1}{2} \log \left( 1 + \frac{|w - z|}{\delta(z)} \right) + C,$$

where  $|w - z|$  denotes the Euclidean distance between  $w$  and  $z$  and  $\delta(z)$  the distance from  $z$  to  $\partial D$ .

Moreover, if  $D$  is strongly pseudoconvex, there exist  $\epsilon_0 > 0$  and  $C' > 0$  such that for all  $p \in \partial D$ ,  $x \notin D \cap B(p, \epsilon_0)$  and  $z \in D \cap B(p, \epsilon_0)$  it holds

$$(5.9) \quad K_D(x, z) \geq C' - \frac{1}{2} \log(\delta(z)).$$

**Definition 5.6.** Let  $D \subset \mathbb{C}^N$  be a domain. A *cône region*  $\mathcal{C}(p, \alpha, \epsilon)$  in  $D$  of vertex  $p \in \partial D$ , aperture  $\alpha > 1$  and size  $\epsilon > 0$  is

$$\mathcal{C}(p, \alpha, \epsilon) := \{z \in D : |z - p| < \min\{\alpha\delta(z), \epsilon\}\}.$$

**Proposition 5.7.** *Let  $D \subset \mathbb{C}^N$  be a bounded strongly pseudoconvex domain with  $C^2$  boundary,  $x \in D$ . Let  $\alpha > 1$  and  $R > 0$ . Then there exists  $\epsilon > 0$  such that for every  $p \in \partial D$  and for every sequence  $\{u_n\}$  converging to  $p$  it holds*

$$(5.10) \quad \mathcal{C}(p, \alpha, \epsilon) \subset E_x(\{u_n\}, R).$$

In particular, every sequence in  $D$  converging to a boundary point is admissible.

*Proof.* Let  $\{u_n\} \subset D$  be a sequence converging to  $p \in \partial D$ .

Let  $\epsilon_0 > 0$  be given by (5.8) and (5.9) and such that  $x \notin D \cap B(p, \epsilon_0)$ . Then for  $w \in D \cap B(p, \epsilon_0)$  and  $n$  large enough, we have:

$$(5.11) \quad K_D(w, u_n) - K_D(x, u_n) \leq \frac{1}{2} \log \left( 1 + \frac{|u_n - w|}{\delta(w)} \right) + \frac{1}{2} \log(\delta(u_n) + |u_n - w|) + C - C'.$$

Therefore,

$$(5.12) \quad \limsup_{n \rightarrow \infty} [K_D(w, u_n) - K_D(x, u_n)] \leq C - C' + \frac{1}{2} \log \left( 1 + \frac{|p - w|}{\delta(w)} \right) + \frac{1}{2} \log |p - w|.$$

Now, fix  $\alpha > 1$ . Given  $\epsilon \in (0, \epsilon_0]$ , for every  $w \in \mathcal{C}(p, \alpha, \epsilon)$  it holds

$$\limsup_{n \rightarrow \infty} [K_D(w, u_n) - K_D(x, u_n)] \leq C - C' + \frac{1}{2} \log(1 + \alpha) + \frac{1}{2} \log \epsilon.$$

Hence, given  $R > 0$ , there exists  $\epsilon \in (0, \epsilon_0]$  (depending on  $R$  and  $\alpha$  but not on  $p$  and  $\{u_n\}$ ), such that (5.10) holds.

In particular,  $E_x(\{u_n\}, R) \neq \emptyset$  for all  $R > 0$ . Since  $D$  is complete hyperbolic, it follows that  $\liminf_{n \rightarrow \infty} K_M(x, u_n) = \infty$ . Hence  $\{u_n\}$  is admissible.  $\square$

For strongly pseudoconvex domains we can say more about equivalent admissible sequence:

**Proposition 5.8.** *Let  $D \subset \mathbb{C}^N$  be a bounded strongly pseudoconvex domain with  $C^3$  boundary. If  $\{u_n\}, \{v_n\} \subset D$  are two sequences converging to  $p$  then they are admissible and  $\{u_n\} \sim \{v_n\}$ .*

*Proof.* The two sequences are admissible by Proposition 5.7.

Fix  $R > 0$ . We want to show that there exist  $R', R'' > 0$  such that  $E_x(\{u_n\}, R') \subset E_x(\{v_n\}, R)$  and  $E_x(\{v_n\}, R'') \subset E_x(\{u_n\}, R)$ . We are going to prove the existence of  $R'$ , a similar argument holds for  $R''$ .

We claim that for every open neighborhood  $W$  of  $p$ , there exists  $r_0 > 0$  such that

$$(5.13) \quad \overline{E_x(\{u_n\}, r)} \subset (W \cap D) \cup \{p\},$$

for all  $0 < r < r_0$  (and similarly for  $\{v_n\}$ ). Indeed, for every  $r > 0$ ,  $E_x(\{u_n\}, r) \subset F_x(p, r)$ , where  $F_x(p, r)$  denotes Abate's big horosphere. Hence, the claim follows from Lemma 5.3.

Now choose  $U$  to be an open neighborhood of  $p$  such that  $U \cap D$  is biholomorphic to a strongly convex domain with  $C^3$  boundary. Let  $W \subset U$  be the open neighborhood of  $p$  given by Lemma 5.4.

Since the equivalence relation among admissible sequences is independent of the base point, we can assume with no loss of generality that  $x \in W \cap D$ . Moreover, since both  $\{u_n\}$  and  $\{v_n\}$  are eventually contained in  $W$ , we can also assume that  $\{u_n\}, \{v_n\} \subset W \cap D$ .

For  $r > 0$ , let  $E_x^{U \cap D}(p, r)$  denote the small horosphere of vertex  $p$  and radius  $r$  in  $U \cap D$ . Since clearly  $E_x^{U \cap D}(p, r) \subset E_x^{U \cap D}(\{u_n\}, r) \subset F_x^{U \cap D}(p, r)$  and by Theorem 5.2,  $E_x^{U \cap D}(p, r) = F_x^{U \cap D}(p, r)$ , it follows that for every  $r > 0$

$$(5.14) \quad E_x^{U \cap D}(p, r) = E_x^{U \cap D}(\{u_n\}, r) = E_x^{U \cap D}(\{v_n\}, r).$$

Now we claim that for every  $r > 0$  such that  $E_x(\{v_n\}, r) \subset W$  there exists  $r' > 0$  such that

$$(5.15) \quad E_x^{U \cap D}(p, r') \subset E_x(\{v_n\}, r).$$

Conversely, for every  $r > 0$  such that  $E_x^{U \cap D}(p, r) \subset W$  there exists  $r' > 0$  such that

$$(5.16) \quad E_x(\{u_n\}, r') \subset E_x^{U \cap D}(p, r).$$

Assuming the claims, the proof ends as follows. By (5.13), there exists  $r > 0$  such that  $E_x(\{v_n\}, r) \subset E_x(\{v_n\}, R) \cap W$ . By (5.15) there exists  $r' > 0$  such that  $E_x^{U \cap D}(p, r') \subset E_x(\{v_n\}, r)$ , and by (5.16) there exists  $R' > 0$  such that  $E_x(\{u_n\}, R') \subset E_x^{U \cap D}(p, r')$ . Hence,  $E_x(\{u_n\}, R') \subset E_x(\{v_n\}, R)$  as needed.

In order to prove (5.15), let  $r > 0$  be such that  $E_x(\{v_n\}, r) \subset W$ . Let  $T \geq 1$  be given by Lemma 5.4. Let  $r_T \in (0, r]$  be such that  $E_x^{U \cap D}(p, T^{-1}r_T) \subset W$ . By (5.14),  $E_x^{U \cap D}(p, T^{-1}r_T) = E_x^{U \cap D}(\{v_n\}, T^{-1}r_T)$ . Hence, for every  $z \in E_x^{U \cap D}(p, T^{-1}r_T)$  it holds

$$\begin{aligned} & \limsup_{n \rightarrow \infty} [K_D(z, v_n) - K_D(x, v_n)] \leq \limsup_{n \rightarrow \infty} [K_{U \cap D}(z, v_n) - K_D(x, v_n)] \\ & \leq \limsup_{n \rightarrow \infty} [K_{U \cap D}(z, v_n) - K_{U \cap D}(x, v_n)] + \limsup_{n \rightarrow \infty} [K_{U \cap D}(x, v_n) - K_D(x, v_n)] \\ & \stackrel{(5.1)}{\leq} \frac{1}{2} \log(T^{-1}r_T) + \frac{1}{2} \log T = \frac{1}{2} \log r_T, \end{aligned}$$

which shows that  $z \in E_x(\{v_n\}, r_T) \subseteq E_x(\{v_n\}, r)$ . Namely,  $E_x^{U \cap D}(p, T^{-1}r_T) \subset E_x(\{v_n\}, r)$  and (5.15) is proved with  $r' = T^{-1}r_T$ .

In order to prove (5.16), let  $r > 0$  be such that  $E_x^{U \cap D}(p, r) \subset W$ . Let  $T \geq 1$  be given by Lemma 5.4. Let  $r_T \in (0, r]$  be such that  $E_x(\{u_n\}, T^{-1}r_T) \subset W$ . Then for every  $z \in E_x(\{u_n\}, T^{-1}r_T)$  it holds

$$\begin{aligned} & \limsup_{n \rightarrow \infty} [K_{U \cap D}(z, u_n) - K_{D \cap U}(x, u_n)] \leq \limsup_{n \rightarrow \infty} [K_{U \cap D}(z, u_n) - K_D(x, u_n)] \\ & \leq \limsup_{n \rightarrow \infty} [K_{U \cap D}(z, u_n) - K_D(z, u_n)] + \limsup_{n \rightarrow \infty} [K_D(z, u_n) - K_D(x, u_n)] \\ & \stackrel{(5.1)}{\leq} \frac{1}{2} \log T + \frac{1}{2} \log(T^{-1}r_T) = \frac{1}{2} \log r_T, \end{aligned}$$

which, by (5.14), shows that  $z \in E_x^{U \cap D}(p, r_T) \subseteq E_x^{U \cap D}(p, r)$ . Namely,  $E_x(\{u_n\}, T^{-1}r_T) \subset E_x^{U \cap D}(p, r)$ , and (5.16) is proved with  $r' = T^{-1}r_T$ .  $\square$

A consequence of the previous proposition is given by the following:

**Proposition 5.9.** *Let  $D \subset \mathbb{C}^N$  be a bounded strongly pseudoconvex domain with  $C^3$  boundary. Let  $\{u_n\} \subset D$  be an admissible sequence. Then there exists  $p \in \partial D$  such that  $\lim_{n \rightarrow \infty} u_n = p$ . Moreover, every admissible sequence  $\{v_n\}$  which is equivalent to  $\{u_n\}$  converges to  $p$ . Finally, for  $p \in \partial D$ , let  $\mathcal{A}_p$  denote the set of all sequences in  $D$  which converges to  $p$ . Then, for all  $R > 0$*

- (1)  $F_x(p, R) = \cup_{\{u_n\} \in \mathcal{A}_p} E_x(\{u_n\}, R)$ ;
- (2)  $E_x(p, R) = \cap_{\{u_n\} \in \mathcal{A}_p} E_x(\{u_n\}, R) \neq \emptyset$ .

*Proof.* Since  $D$  is complete hyperbolic, Property (1) in Definition 3.2 implies that all accumulation points of  $\{u_n\}$  are contained in  $\partial D$ . Let  $p \in \partial D$  be one of such points. Let  $\{u_{n_k}\}$  be a subsequence of  $\{u_n\}$  such that  $u_{n_k} \rightarrow p$ . Let  $z \in E_x(\{u_n\}, R)$  for some  $R > 0$ . Then

$$\begin{aligned} \liminf_{w \rightarrow p} [K_D(z, w) - K_D(x, w)] &\leq \limsup_{k \rightarrow \infty} [K_D(z, u_{n_k}) - K_D(x, u_{n_k})] \\ &\leq \limsup_{n \rightarrow \infty} [K_D(z, u_n) - K_D(x, u_n)] \leq \frac{1}{2} \log R. \end{aligned}$$

Hence  $z \in F_x(p, R)$ . This implies that  $E_x(\{u_n\}, R) \subset F_x(p, R)$ . In particular, if  $q \in \partial D$  were another accumulation point of  $\{u_n\}$  different from  $p$ , it would hold

$$E_x(\{u_n\}, R) \subset F_x(p, R) \cap F_x(q, R) \quad \forall R > 0,$$

contradicting Lemma 5.3. Hence  $\lim_{n \rightarrow \infty} u_n = p$ .

Now, let  $\{v_n\}$  be another admissible sequence which is equivalent to  $\{u_n\}$ . From what we just proved, there exists  $q \in \partial D$  such that  $\lim_{n \rightarrow \infty} v_n = q$ . If  $q \neq p$ , let  $R_0 > 0$  be given by Lemma 5.3 such that  $F_x(p, R) \cap F_x(q, R) = \emptyset$  for all  $0 < R < R_0$ . Fix  $0 < R < R_0$ . By the previous argument,  $E_x(\{v_n\}, R) \subset F_x(q, R)$ . Since  $\{u_n\}$  is equivalent to  $\{v_n\}$  this implies that there exists  $R' > 0$  such that  $E_x(\{u_n\}, R') \subset E_x(\{v_n\}, R) \subset F_x(q, R)$ . Now, if  $R' \geq R$ , it follows that  $E_x(\{u_n\}, R) \subset F_x(q, R)$ , giving a contradiction since we proved that  $E_x(\{u_n\}, R) \subset F_x(p, R)$ . If  $R' < R$ , again we obtain a contradiction since  $E_x(\{u_n\}, R') \subset E_x(\{u_n\}, R) \subset F_x(p, R)$ .

In order to prove (1), we already saw that  $\cup_{\{u_n\} \in \mathcal{A}_p} E_x(\{u_n\}, R) \subset F_x(p, R)$ . Let  $z \in F_x(p, R)$ . By the very definition, there exists a sequence  $\{u_n\} \subset D$  converging to  $p$  such that  $\lim_{n \rightarrow \infty} [K_D(z, u_n) - K_D(x, u_n)] = \frac{1}{2} \log R'$  for some  $R' < R$ . By Proposition 5.8,  $\{u_n\}$  is admissible and  $z \in E_x(\{u_n\}, R)$ , hence  $F_x(p, R) \subset \cup_{\{u_n\} \in \mathcal{A}_p} E_x(\{u_n\}, R)$ .

(2) the proof is similar and we omit it. The fact that  $E_x(p, R) \neq \emptyset$  follows then immediately from (5.10).  $\square$

The main result of this section is the following:

**Theorem 5.10.** *Let  $D \subset \mathbb{C}^N$  be a bounded strongly pseudoconvex domain with  $C^3$  boundary. Then  $\hat{D}$  endowed with the horosphere topology is homeomorphic to  $\overline{D}$  (closure of  $D$  in  $\mathbb{C}^N$ ) endowed with the Euclidean topology.*

*Proof.* We define a map  $\Theta : \hat{D} \rightarrow \overline{D}$  as follows. If  $z \in D \subset \hat{D}$  then we just set  $\Theta(z) = z$ . If  $\underline{y} \in \partial_H D$ , by Proposition 5.9 all admissible sequences representing  $\underline{y}$  converge to a same point  $p \in \partial D$ . Then we set  $\Theta(\underline{y}) := p$ . The map  $\Theta$  is bijective by Proposition 5.8.

In order to prove that  $\Theta$  is a homeomorphism, we show that given  $C \subset \overline{D}$  then  $C$  is closed in the Euclidean topology if and only if  $\Theta^{-1}(C) \subset \hat{D}$  is closed in the horosphere topology.

Assume that  $C$  is closed in the Euclidean topology. In order to show that  $\Theta^{-1}(C)$  is closed in the horosphere topology, we need to show that if  $\{\xi_n\}$  is a sequence in  $\Theta^{-1}(C)$  converging to  $\xi \in \hat{D}$ , then  $\xi \in \Theta^{-1}(C)$ . There are three cases to examine, from which the general case follows immediately:

Case 1. If  $\{\Theta^{-1}(z_n)\}$  is a sequence in  $\Theta^{-1}(C) \cap D$  converging to  $\Theta^{-1}(z) \in D$ , then by the very definition of  $\Theta$  and since the topology induced on  $D \subset \hat{D}$  by the horosphere topology is the Euclidean topology, it follows easily that  $\Theta^{-1}(z) \in \Theta^{-1}(C)$ .

Case 2. Let now  $\{\underline{y}_m\}$  be a sequence in  $\Theta^{-1}(C) \cap \partial_H D$ , converging in the horosphere topology to  $\underline{y} \in \partial_H D$  (by Remark 4.5 a sequence on the horosphere boundary can only converge to a point on the horosphere boundary). Let  $p_m := \Theta(\underline{y}_m) \in C$ ,  $m \in \mathbb{N}$  and let  $p := \Theta(\underline{y})$ . By the definition of convergence in the horosphere topology, there exist horosphere sequences  $\{u_n^m\}_{n \in \mathbb{N}}$  with  $[u_n^m] = \underline{y}_m$  and  $\{u_n\}$  with  $[u_n] = \underline{y}$  such that for every  $R > 0$  there exists  $m_R \in \mathbb{N}$  such that  $E_x(\{u_n^m\}, R) \cap E_x(\{u_n\}, R) \neq \emptyset$  for  $m > m_R$ . By the definition of  $\Theta$ , it holds  $\lim_{n \rightarrow \infty} u_n^m = p_m$  and  $\lim_{n \rightarrow \infty} u_n = p$ . Thus, by Proposition 5.9.(1), it follows that

$$F_x(p, R) \cap F_x(p_m, R) \neq \emptyset$$

for all  $m > m_R$ . If  $\{p_m\}$  contained a subsequence not converging to  $p$ , say  $\{p_{m_k}\}$ , taking  $V := \{p\}$  and  $V' := \overline{\{p_{m_k}\}}$ , we would get a contradiction with Lemma 5.3. Therefore,  $\lim_{m \rightarrow \infty} p_m = p$ . Since  $C$  is closed, this implies that  $p \in C$  as well, hence  $\Theta^{-1}(p) = \underline{y}$  belongs to  $\Theta^{-1}(C)$ .

Case 3. Finally, assume  $\{z_n = \Theta^{-1}(z_n)\} \subset \Theta^{-1}(C) \cap D$  converges to  $\underline{y} \in \partial_H D$ . Let  $p = \Theta(\underline{y}) \in \partial D$ . By Definition 4.2, and by Proposition 5.9.(1), it follows that there exists a sequence  $\{p_n\} \subset \partial D$  such that for all  $R > 0$  there exists  $n_R \in \mathbb{N}$  such that  $z_n \in F_x(p_n, R)$  and  $F_x(p_n, R) \cap F_x(p, R) \neq \emptyset$  for  $n \geq n_R$ . As before, this implies that  $\lim_{n \rightarrow \infty} p_n = p$ . Thus, given an open set  $U$  containing  $p$ , there exists  $J \in \mathbb{N}$  such that  $\{p_n\}_{n \geq J} \subset U$ . Let  $V = \{p_n\}_{n \geq J} \cup \{p\}$ . The set  $V$  is closed, and by Lemma 5.3, there exists  $R_0 > 0$  such that  $\bigcup_{n \geq J} F_x(p_n, R) \subset U \cap D$  for all  $0 < R < R_0$ . Therefore,  $\{z_n\}$  is eventually contained in  $U$ . By the arbitrariness of  $U$ , it follows that  $\lim_{n \rightarrow \infty} z_n = p$ . Since  $C$  is closed, this implies that  $p \in C$  and hence  $\underline{y} = \Theta^{-1}(p) \in \Theta^{-1}(C)$ .

Hence, if  $C$  is closed in the Euclidean topology then  $\Theta^{-1}(C)$  is closed in the horosphere topology.

Assume that  $\Theta^{-1}(C)$  is closed in the horosphere topology. We have to show that if  $\{\xi_n\} \subset C$  is a sequence converging to  $\xi \in \overline{D}$  then  $\xi \in C$ . As before, we distinguish three cases.

Case 1. If  $\{\xi_n\}, \xi \subset D$ , then the statement is clearly true.

Case 2. Let  $\{p_m\}$  be a sequence in  $C \cap \partial D$  converging to  $p \in \partial D$ . Let  $\underline{y}_m := \Theta^{-1}(p_m) \in \Theta^{-1}(C) \cap \partial_H D$  and let  $\underline{y} := \Theta^{-1}(p) \in \partial_H D$ .

We want to show that  $\{\underline{y}_m\}$  converges to  $\underline{y}$  in the horosphere topology. Once we proved this, since  $\Theta^{-1}(C)$  is closed, it follows that  $\underline{y} \in \Theta^{-1}(C)$  and hence,  $p = \Theta(\underline{y}) \in C$ .

To show this, for every  $m \in \mathbb{N}$ , let  $\{v_n^m\}_{n \in \mathbb{N}}$  be a sequence converging to  $p_m$ , and let  $\{v_n\}$  be a sequence converging to  $p$ . By Proposition 5.9,  $[v_n^m] = \underline{y}_m$  and  $[v_n] = \underline{y}$ . Fix  $R > 0$ . By (5.10), there exists  $\varepsilon > 0$  such that the cone region  $\mathcal{C}(q, 2, \varepsilon) \subset E_x(\{w_n\}, R)$  for all  $q \in \partial D$  and all sequences  $\{w_n\}$  converging to  $q$ . Since  $\partial D$  is  $C^1$ , there exists an open neighborhood  $V$  of  $p$  such that for all  $q \in V \cap \partial D$  it holds  $\mathcal{C}(p, 2, \varepsilon) \cap \mathcal{C}(q, 2, \varepsilon) \neq \emptyset$ . Since  $\{p_m\} \subset V$  eventually, it follows that  $E_x(\{v_n^m\}, R) \cap E_x(\{v_n\}, R) \neq \emptyset$  for all  $m$  sufficiently large. By definition, this means that  $\{\underline{y}_m\}$  converges to  $\underline{y}$  in the horosphere topology, as needed.

Case 3. Let  $\{z_m\}$  be a sequence in  $C \cap D$  converging to  $p \in \partial D$ . Let  $\underline{y} := \Theta^{-1}(p)$ . As before, to conclude that  $p \in C$ , it is enough to show that  $\{z_m = \Theta^{-1}(z_m)\}$  converges to  $\underline{y}$  in the horosphere topology. To show this, fix  $\alpha > 1$  and for  $N \in \mathbb{N}$ , let  $\varepsilon_N > 0$  be given by Proposition 5.7 such that for every  $q \in \partial D$  and for every sequence  $\{w_n\}$  converging to  $q$  it holds  $\mathcal{C}(q, \alpha, \varepsilon_N) \subset E_x(\{w_n\}, \frac{1}{N})$ .

Since  $\{z_m\}$  converges to  $p$  and  $\partial D$  is  $C^2$ , we can assume with no loss of generality that for all  $m \in \mathbb{N}$  there exists a unique  $p_m \in \partial D$  such that  $|z_m - p_m| = \delta(z_m)$ . Note that  $z_m \in \mathcal{C}(p_m, \alpha, r)$  for all  $r > \delta(z_m)$ .

For each  $m \in \mathbb{N}$  let  $\{v_n^m\} \subset D$  be a sequence converging to  $p_m$  and let  $\{u_n\} \subset D$  be a sequence converging to  $p$ . By Proposition 5.9,  $[u_n] = \underline{y}$ .

Fix  $N \in \mathbb{N}$ . Since  $\delta(z_m) \rightarrow 0$ , there exists  $m_N^1 \in \mathbb{N}$  such that  $z_m \in \mathcal{C}(p_m, \alpha, \varepsilon_N) \subset E_x(\{v_n^m\}, \frac{1}{N})$  for all  $m \geq m_N^1$ . Moreover, since  $\{p_m\}$  converges to  $p$ , there exists  $m_N^2 \in \mathbb{N}$  such that  $\mathcal{C}(p_m, \alpha, \varepsilon_N) \cap \mathcal{C}(p, \alpha, \varepsilon_N) \neq \emptyset$  for all  $m \geq m_N^2$ , which implies  $E_x(\{v_n^m\}, \frac{1}{N}) \cap E_x(\{u_n\}, \frac{1}{N}) \neq \emptyset$  for all  $m \geq m_N^2$ .

Now let  $R > 0$  be given. Let  $N \in \mathbb{N}$ ,  $N > 1/R$  and let  $m_R := \max\{m_N^1, m_N^2\}$ . Then for all  $m \geq m_R$  it holds  $z_m \in E_x(\{v_n^m\}, \frac{1}{N}) \subset E_x(\{v_n^m\}, R)$  and

$$\emptyset \neq E_x(\{v_n^m\}, \frac{1}{N}) \cap E_x(\{u_n\}, \frac{1}{N}) \subset E_x(\{v_n^m\}, R) \cap E_x(\{u_n\}, R),$$

which means that  $\{z_m\}$  converges to  $\underline{y}$  in the horosphere topology, and we are done.

Hence, if  $\Theta^{-1}(C)$  is closed in the horosphere topology then  $C$  is closed in the Euclidean topology.  $\square$

We end this section by proving the following result which will be useful to study boundary behavior:

**Proposition 5.11.** *Let  $D \subset \mathbb{C}^N$  be a bounded strongly pseudoconvex domain with  $C^3$  boundary. Let  $\underline{x} \in \partial_H D$ . Then there exists  $p_{\underline{x}} \in \partial D$  such that  $I_D^H(\underline{x}) = \{p_{\underline{x}}\}$ . Moreover, if  $\underline{y} \in \partial_H D$  then  $p_{\underline{x}} = p_{\underline{y}}$  if and only if  $\underline{x} = \underline{y}$ .*

*Proof.* By Theorem 5.10, there is a homeomorphism  $\Theta : \hat{D} \rightarrow \overline{D}$  (where  $\overline{D}$  is the closure of  $D$  in  $\mathbb{C}^N$ ). Let  $p_{\underline{x}} := \Theta(\underline{x})$ . Recall also that  $\Theta(z) = z$  for  $z \in D$ .

We have to show that, if  $\{w_n\} \subset D$  is a sequence converging to  $\underline{x}$  in the horosphere topology, then  $\{w_n\}$  converges to  $p_{\underline{x}}$  in the Euclidean topology. This follows at once by Theorem 5.10, because  $\{w_n\}$  converges to  $\underline{x}$  in the horosphere topology if and only if  $\{\Theta(w_n) = w_n\}$  converges to  $p_{\underline{x}}$  in the Euclidean topology. This also shows the last statement of the proposition.  $\square$

## 6. CONVEX DOMAINS

In this section we consider convex domains in  $\mathbb{C}^n$ . Here we mean convex in the real geometrical sense, that is,  $D \subset \mathbb{C}^N$  is convex if for every two points  $p, q \in D$  the real segment  $[p, q]$  joining  $p$  and  $q$  is contained in  $D$ .

By [13], a (possibly unbounded) convex domain in  $\mathbb{C}^N$  is complete hyperbolic if and only if it is biholomorphic to a bounded domain of  $\mathbb{C}^N$ , in particular, a convex domain is hyperbolic if and only if it is complete hyperbolic.

We start with the following result:

**Proposition 6.1.** *Let  $D \subset \mathbb{C}^N$  be a hyperbolic convex domain. Let  $x \in D$ . Let  $\{u_n\}$  be an admissible sequence in  $D$ . Then for every  $R > 0$  the horosphere  $E_x(\{u_n\}, R)$  is convex.*

*Proof.* Let  $B(0, N) := \{z \in \mathbb{C}^N : |z| < N\}$ ,  $N \in \mathbb{N}$ . Let  $D_N := D \cap B(0, N)$ . Then  $D_N$  is a bounded convex domain and its Kobayashi distance  $k_{D_N}$  is a convex function (see [3, Prop. 2.3.46]). Passing to the limit, it turns out that  $k_D$  is a convex function as well.

Now, let  $\{u_n\} \subset D$  be an admissible sequence. Let  $R > 0$  and let  $z, w \in E_x(\{u_n\}, R)$ . Then, for  $s \in (0, 1)$  we have

$$K_D(sz + (1-s)w, u_n) - K_D(x, u_n) \leq \max\{K_D(z, u_n), K_D(w, u_n)\} - K_D(x, u_n).$$

Taking the limsup as  $n \rightarrow \infty$ , this implies that  $sz + (1-s)w \in E_x(\{u_n\}, R)$ , which is thus convex.  $\square$

**Definition 6.2.** Let  $D \subset \mathbb{C}^N$  be a bounded convex domain,  $x \in D$ . Let  $\{u_n\}$  be an admissible sequence. For  $R > 0$  let

$$\Pi(\{u_n\}, R) := \overline{E_x(\{u_n\}, R)} \cap \partial D.$$

Note that, if  $\{u_n\}$  is an admissible sequence and denoting  $[\{u_n\}] \in \partial_H D$ , by Lemma 4.12 it holds

$$\Pi_D^H([\{u_n\}]) = \bigcap_{R>0} \overline{E_x(\{u_n\}, R)} = \bigcap_{R>0} \Pi(\{u_n\}, R).$$

**Remark 6.3.** Note that  $\Pi(\{u_n\}, R)$  and  $\Pi_D^H([\{u_n\}])$  are nonempty sets and  $\Pi_D^H([\{u_n\}])$  is convex. Indeed, this follows at once from Proposition 6.1, Proposition 3.4.(2) and since  $\overline{E_x(\{u_n\}, R)} \subset \overline{E_x(\{u_n\}, R')}$  for all  $0 < R < R'$ . Moreover, while  $\Pi(\{u_n\}, R)$  depends in general on the base point  $x$ , the horosphere principal part,  $\Pi_D^H([\{u_n\}])$ , does not.

**Lemma 6.4.** Let  $D \subset \mathbb{C}^N$  be a bounded convex domain. Let  $\{u_n\}$  be an admissible sequence. Let  $\mathcal{M}$  denote the set of all convergent subsequences extracted from  $\{u_n\}$ . Then

$$(6.1) \quad \Pi_D^H([\{u_n\}]) = \bigcap_{\{v_n\} \in \mathcal{M}} \Pi_D^H([\{v_n\}]).$$

Moreover, if  $\{u_n\}$  is convergent to  $p \in \partial D$ , then  $p \in \Pi_D^H([\{u_n\}])$ .

*Proof.* It is easy to check that for all  $R > 0$ ,

$$(6.2) \quad E_x(\{u_n\}, R) = \bigcap_{\{v_n\} \in \mathcal{M}} E_x(\{v_n\}, R).$$

In order to prove (6.1), by (6.2), it is clear that  $\Pi_D^H([\{u_n\}]) \subseteq \bigcap_{\{v_n\} \in \mathcal{M}} \Pi_D^H([\{v_n\}])$ . Conversely, let  $q \in \bigcap_{\{v_n\} \in \mathcal{M}} \Pi_D^H([\{v_n\}])$ , let  $R > 0$  and let  $z \in E_x(\{u_n\}, R)$ . Then  $z \in E_x(\{v_n\}, R)$  for all  $\{v_n\} \in \mathcal{M}$  by (6.2). Since  $E_x(\{v_n\}, R)$  is convex by Proposition 6.1, the segment  $tz + (1-t)q$ ,  $0 < t \leq 1$  belongs to  $E_x(\{v_n\}, R)$  for all  $\{v_n\} \in \mathcal{M}$ , thus it belongs to  $E_x(\{u_n\}, R)$ , proving that  $q \in \Pi_D^H([\{u_n\}])$ .

In case  $\{u_n\}$  converges to  $p \in \partial D$ , arguing similar to [4, Lemma 2.3], one can prove that  $p \in \Pi_D^H([\{u_n\}])$ . We give a sketch of the proof for the reader convenience. Let  $t \in (0, 1)$  and define  $f_t^n(w) := tw + (1-t)u_n$ . Note that  $f_t^n : D \rightarrow D$  is holomorphic, hence  $f_t^n$  decreases the Kobayashi distance of  $D$ . Moreover,  $f_t^n(u_n) = u_n$  and  $\lim_{n \rightarrow \infty} f_t^n(z) = tz + (1-t)p \in D$ . Fix  $R > 0$ . Let  $z \in E_x(\{u_n\}, R)$ . Then

$$\begin{aligned} & \limsup_{n \rightarrow \infty} [K_D(tz + (1-t)p, u_n) - K_D(x, u_n)] \\ & \leq \limsup_{n \rightarrow \infty} [K_D(f_t^n(z), f_t^n(u_n)) - K_D(x, u_n)] + \limsup_{n \rightarrow \infty} [K_D(tz + (1-t)p, f_t^n(z))] < \frac{1}{2} \log R. \end{aligned}$$

Therefore  $tz + (1-t)p \in E_x(\{u_n\}, R)$ , hence  $p \in \Pi(\{u_n\}, R)$  for all  $R > 0$  and thus  $p \in \Pi_D^H([\{u_n\}])$ .  $\square$

In general, if  $\{u_n\} \subset D$  is an admissible sequence which does not converge and  $p \in \partial D$  is in the cluster set of  $\{u_n\}$ , it does not hold  $p \in \Pi_D^H([\{u_n\}])$ , as the following example shows:

**Example 6.5.** Let  $D = \mathbb{D}^2$  be the bidisc. Consider the sequences  $\{v_n\}$ ,  $\{w_n\}$  given by  $v_n = (0, 1 - 1/n)$  and  $w_n = (1 - 1/n, 0)$ , for every  $n \in \mathbb{N}, n \geq 1$ . Since  $K_{\mathbb{D}^2}((z_1, z_2), (w_1, w_2)) = \max\{K_{\mathbb{D}}(z_1, w_1), K_{\mathbb{D}}(z_2, w_2)\}$ , it follows easily that for all  $R > 0$

$$E_{(0,0)}(\{v_n\}, R) = \mathbb{D} \times E_0^{\mathbb{D}}(1, R),$$

where  $E_0^{\mathbb{D}}(1, R) = \{\zeta \in \mathbb{D} : |1 - \zeta|^2 / (1 - |\zeta|^2) < R\}$ . Hence  $\Pi_D^H([\{v_n\}]) = \overline{\mathbb{D}} \times \{1\}$ . On the other hand,

$$E_{(0,0)}(\{w_n\}, R) = E_0^{\mathbb{D}}(1, R) \times \mathbb{D},$$

and  $\Pi_D^H([\{w_n\}]) = \{1\} \times \overline{\mathbb{D}}$ .

Let  $\{u_n\}$  be the sequence defined by  $u_{2n-1} = v_n, u_{2n} = w_n, n \geq 1$ . Hence  $\{u_n\}$  is admissible, since, for  $R > 0$

$$E_{(0,0)}(\{u_n\}, R) = E_{(0,0)}(\{v_n\}, R) \cap E_{(0,0)}(\{w_n\}, R) = E_0^{\mathbb{D}}(1, R) \times E_0^{\mathbb{D}}(1, R).$$

Then  $\Pi_D^H(\{\{u_n\}\}) = \{(1, 1)\}$ , and all points in the cluster set of  $\{u_n\}$ , namely  $(1, 0)$  and  $(0, 1)$ , do not belong to  $\Pi_D^H(\{\{u_n\}\})$ .

The previous example shows also that  $\Pi(\{u_n\}, R)$  is not convex in general.

**6.1. Convex domains biholomorphic to strongly pseudoconvex domains.** We examine now the case of convex domains (with no regularity assumption on the boundary) biholomorphic to strongly pseudoconvex domains. We start with the following general result:

**Lemma 6.6.** *Let  $M$  be a complex manifold biholomorphic to a bounded strongly pseudoconvex domain with  $C^3$  boundary,  $x \in M$ . Let  $\{u_n\}$  be an admissible sequence in  $M$ . Let  $\{z_n\} \subset M$  be a non relatively compact sequence which is eventually contained in  $E_x(\{u_n\}, R)$  for some  $R > 0$ . Then  $\{z_n\}$  is admissible and it is equivalent to  $\{u_n\}$ .*

*Proof.* Let  $F : D \rightarrow M$  be a biholomorphism between a bounded strongly pseudoconvex domain with  $C^3$  boundary and  $M$ .

The sequence  $\{F^{-1}(u_n)\}$  is admissible in  $D$ . By Proposition 5.9, there exists  $p \in \partial D$  such that  $\{F^{-1}(u_n)\}$  converges to  $p$ . Since  $\{z_n\}$  is eventually contained in  $E_x^M(\{u_n\}, R)$ , then  $\{F^{-1}(z_n)\}$  is eventually contained in  $E_{F^{-1}(x)}^D(\{F^{-1}(u_n)\}, R)$ . Moreover,  $\{F^{-1}(z_n)\}$  is not relatively compact in  $D$ , as  $\{z_n\}$  is not in  $M$ . Therefore every limit of  $\{F^{-1}(z_n)\}$  has to be contained in

$$(6.3) \quad \overline{E_{F^{-1}(x)}^D(\{F^{-1}(u_n)\}, R)} \cap \partial D \subset \overline{E_{F^{-1}(x)}^D(p, R)} \cap \partial D = \{p\}$$

by Theorem 5.2.(4).

Therefore  $\{F^{-1}(z_n)\}$  is converging to  $p$ . By Proposition 5.8, the sequence  $\{F^{-1}(z_n)\}$  is admissible in  $D$  and equivalent to  $\{F^{-1}(u_n)\}$ . Hence  $\{z_n\}$  is admissible in  $M$  and equivalent to  $\{u_n\}$ .  $\square$

As a corollary we have the following result

**Proposition 6.7.** *Let  $D \subset \mathbb{C}^N$  be a bounded convex domain. Assume  $D$  is biholomorphic to a bounded strongly pseudoconvex domain with  $C^3$  boundary. Let  $\underline{x} \in \partial_H D$ . Let  $\{u_n\}$  be any admissible sequence in  $D$  representing  $\underline{x}$ . Then for every  $R > 0$  it holds*

$$(6.4) \quad \Pi(\{u_n\}, R) = \Pi_D^H(\underline{x}).$$

*Proof.* By definition,  $\Pi_D^H(\underline{x}) \subseteq \Pi(\{u_n\}, R)$ . Let now  $p \in \Pi(\{u_n\}, R)$ . Fix  $x \in D$ . Then there exists a sequence  $\{p_n\} \subset E_x(\{u_n\}, R)$  such that  $\lim_{n \rightarrow \infty} p_n = p$ . By Lemma 6.6,  $\{p_n\}$  is an admissible sequence in  $D$  and it is equivalent to  $\{u_n\}$ . By Lemma 6.4 it follows then that  $p \in \Pi_D^H(\underline{x})$ .  $\square$

The previous result allows to generalize Lemma 5.3 to convex domains:

**Lemma 6.8.** *Let  $D \subset \mathbb{C}^N$  be a bounded convex domain. Assume  $D$  is biholomorphic to a bounded strongly pseudoconvex domain with  $C^3$  boundary. Let  $\{\underline{x}_j\} \subset \partial_H D$  be a sequence and let  $V = \bigcup_j \Pi_D^H(\underline{x}_j)$ . Then for every open neighborhood  $U$  of  $V$  there exists  $R_0 > 0$  such that for every  $0 < R < R_0$  and for every admissible sequence  $\{u_n^j\}$  representing  $\underline{x}_j$  for some  $j$ , it holds*

$$E_x(\{u_n^j\}, R) \subset U.$$

*Proof.* We argue by contradiction. Let  $\mathcal{A}_j$  denote the set of all admissible sequences in  $D$  which represent  $\underline{x}_j$ . If the result is not true, for every  $n \in \mathbb{N}$  there exists  $z_n \in D$  such that

$$z_n \in \bigcup_j \bigcup_{\{u_m\} \in \mathcal{A}_j} E(\{u_m\}, \frac{1}{n}) \cap (D \setminus U).$$

We can assume that  $z_n \rightarrow z_0$  for some  $z_0 \in \overline{D}$ . By Proposition 3.5,  $z_0 \in \partial D$ . In particular, given  $R > 1/n$ , it holds  $z_n \in \bigcup_j \bigcup_{\{u_m\} \in \mathcal{A}_j} E(\{u_m\}, R)$  for every  $n \in \mathbb{N}$ . Hence,

$$z_0 \in \overline{\bigcup_j \bigcup_{\{u_m\} \in \mathcal{A}_j} E(\{u_m\}, R) \cap \partial D} = V,$$

by Proposition 6.7. Therefore,  $z_0 \in V \cap (\partial D \setminus U) = \emptyset$ , a contradiction.  $\square$

As in the strongly pseudoconvex case, the previous lemma allows to relate Euclidean topology with horosphere topology:

**Corollary 6.9.** *Let  $D \subset \mathbb{C}^N$  be a bounded convex domain. Assume  $D$  is biholomorphic to a bounded strongly pseudoconvex domain with  $C^3$  boundary. Also, assume that for every  $\underline{x} \in \partial_H D$  there exists  $p_{\underline{x}} \in \partial D$  such that  $\Pi_D^H(\underline{x}) = \{p_{\underline{x}}\}$ . If  $\{\underline{x}_j\} \subset \partial_H D$  is a sequence converging to  $\underline{x} \in \partial_H D$  in the horosphere topology, then  $\lim_{j \rightarrow \infty} p_{\underline{x}_j} = p_{\underline{x}}$  in the Euclidean topology.*

*Proof.* Assume the conclusion of the corollary is not true and, eventually up to extracting subsequences, assume that  $p_{\underline{x}_j} \rightarrow q$  for some  $q \in \partial D \setminus \{p_{\underline{x}}\}$ . Let  $V = \overline{\bigcup_j \{p_{\underline{x}_j}\}}$ . Let  $U_0, U_1$  be two open sets in  $\mathbb{C}^N$  such that  $V \subset U_0$ ,  $\{p_{\underline{x}}\} \subset U_1$  and  $U_0 \cap U_1 = \emptyset$ . By Lemma 6.8, there exists  $R_0 > 0$  such that for all  $0 < R < R_0$  and all admissible sequences  $\{u_n^j\}$  representing  $\underline{x}_j$  it holds  $E_x(\{u_n^j\}, R) \subset U_0$  and moreover, for all admissible sequences  $\{u_n\}$  representing  $\underline{x}$  it holds  $E_x(\{u_n\}, R) \subset U_1$ . But then, (4.1) can never be satisfied for  $0 < R < R_0$ , and  $\underline{x}_j$  cannot converge to  $\underline{x}$  in the horosphere topology. A contradiction.  $\square$

**Theorem 6.10.** *Let  $D \subset \mathbb{C}^N$  be a bounded convex domain. Assume  $D$  is biholomorphic to a bounded strongly pseudoconvex domain with  $C^3$  boundary. Also, assume that for every  $\underline{x} \in \partial_H D$  there exists  $p_{\underline{x}} \in \partial D$  such that  $\Pi_D^H(\underline{x}) = \{p_{\underline{x}}\}$ . Then for every  $\underline{x} \in \partial_H D$  it holds*

$$I_D^H(\underline{x}) = \Pi_D^H(\underline{x}) = \{p_{\underline{x}}\}.$$

*Proof.* We argue by contradiction. Let  $\{w_n\} \subset D$  be a sequence converging to  $\underline{x}$  in the horosphere topology. Assume that  $\{w_n\}$  converges to  $q \in \partial D$  in the Euclidean topology, with  $q \neq p_{\underline{x}}$ . By definition of convergence in the horosphere topology, there exist admissible sequences  $\{u_n^j\}$  and  $\{u_n\}$  in  $D$  such that  $\{u_n\}$  represents  $\underline{x}$ , and for every  $R > 0$  there exists  $m_R$  such that for every  $m \geq m_R$  it holds  $w_m \in E_x(\{u_n^m\}, R)$  and  $E_x(\{u_n^m\}, R) \cap E_x(\{u_n\}, R) \neq \emptyset$ . Let  $\underline{x}_m := [\{u_n^m\}] \in \partial_H D$ . By Remark 4.3, the sequence  $\{\underline{x}_j\}$  converges in the horosphere topology to  $\underline{x}$ . Therefore, by Corollary 6.9, the sequence  $\{p_{\underline{x}_j}\}$  converges to  $p_{\underline{x}}$  in the Euclidean topology.

Let  $U_0$  and  $U_1$  be two open sets in  $\mathbb{C}^N$  such that  $q \in U_0$ ,  $p_{\underline{x}} \in U_1$  and  $U_0 \cap U_1 = \emptyset$ . Without loss of generality, we can assume that  $\{w_n\} \subset U_0$  and  $\{p_{\underline{x}_j}\} \subset U_1$ . By Lemma 6.8, there exists  $R_0 > 0$  such that  $E_x(\{u_n^m\}, R) \subset U_1$  for all  $0 < R < R_0$  and all  $m \in \mathbb{N}$ .

Therefore, given  $0 < R < R_0$ , for  $m > m_R$ , we have  $w_m \in U_0 \cap E_x(\{u_n^m\}, R) \subset U_0 \cap U_1 = \emptyset$ , a contradiction.  $\square$

As shown by the previous result, it is important to see which bounded convex domains biholomorphic to bounded strongly pseudoconvex domains have the property that  $\Pi_D^H(\underline{x})$  is a point for every  $\underline{x} \in \partial_H D$ . We conjecture that this is always the case, but presently we are able to prove it for bounded strictly  $\mathbb{C}$ -linearly



convex domains and in case of convex domains biholomorphic to strongly convex domains. In order to state the result, we need a definition:

**Definition 6.11.** Let  $D \subset \mathbb{C}^N$  be a bounded convex domain,  $p \in \partial D$ . A *complex supporting functional* at  $p$  is a  $\mathbb{C}$ -linear map  $\sigma : \mathbb{C}^N \rightarrow \mathbb{C}$  such that  $\operatorname{Re} \sigma(z) < \operatorname{Re} \sigma(p)$  for all  $z \in D$ . A *complex supporting hyperplane* for  $D$  at  $p$  is an affine complex hyperplane  $L$  of the form  $L = p + \ker \sigma = \{z \in \mathbb{C}^N : \sigma(z) = \sigma(p)\}$  where  $\sigma$  is a complex supporting functional at  $p$ . Let  $\mathcal{L}_p$  denote the set of all complex supporting hyperplanes at  $p$ . We set

$$\operatorname{Ch}(p) = \bigcap_{L \in \mathcal{L}_p} L \cap \overline{D}.$$

Clearly,  $\operatorname{Ch}(p)$  is a closed convex set containing  $p$ .

**Definition 6.12.** A bounded convex domain  $D \subset \mathbb{C}^N$  is *strictly  $\mathbb{C}$ -linearly convex* if for every  $p \in \partial D$  it holds  $\operatorname{Ch}(p) = \{p\}$ .

**Proposition 6.13.** Let  $D \subset \mathbb{C}^N$  be a bounded strictly  $\mathbb{C}$ -linearly convex domain. Assume  $D$  is biholomorphic to a bounded strongly pseudoconvex domain with  $C^3$  boundary. Then for every  $\underline{x} \in \partial_H D$  there exists a unique  $p \in \partial D$  such that

$$(6.5) \quad I_D^H(\underline{x}) = \Pi_D^H(\underline{x}) = \{p\}.$$

*Proof.* If we prove that  $\Pi_D^H(\underline{x}) = \{p\}$ , then the result follows from Theorem 6.10.

Let  $\underline{x} \in \partial_H D$ . Suppose by contradiction that  $p, q \in \Pi_D^H(\underline{x})$ . By hypothesis, there exist  $\Omega \subset \mathbb{C}^N$  a bounded strongly pseudoconvex domain with  $C^3$  boundary and a biholomorphism  $F : \Omega \rightarrow D$ .

Let  $\{u_n\}$  be an admissible sequence representing  $\underline{x}$ . For every  $R > 0$ ,  $\Pi(\{u_n\}, R) = \Pi_D^H(\underline{x})$  by (6.4). Therefore, since  $\Pi_D^H(\underline{x})$  is convex, the real segment  $[p, q]$  joining  $p$  and  $q$  is contained in  $\Pi(\{u_n\}, R)$ .

Let  $v = p - q$  and let  $L := \mathbb{C}(p - q) + q$ . There are two possibilities: either  $L \cap D \neq \emptyset$  or  $L \cap D = \emptyset$ .

In case  $L \cap D \neq \emptyset$ , since  $\cup_{R>0} E_x^D(\{u_n\}, R) = D$ , there exists  $R > 0$  such that  $E_x^D(\{u_n\}, R) \cap L \neq \emptyset$ . By convexity of  $E_x^D(\{u_n\}, R)$  (see Proposition 6.1),  $\Delta := L \cap E_x^D(\{u_n\}, R)$  is a convex domain in  $L$  whose boundary contains the segment  $[p, q]$ . Hence by the uniformization theorem, there exists a biholomorphism  $\varphi : \mathbb{D} \rightarrow \Delta$  and, by the Schwarz reflection principle, there exists an arc  $A \subset \partial \mathbb{D}$  such that  $\varphi$  extends analytically on  $A$  and  $\varphi(A) = [p, q]$ . Consider the map  $F^{-1}|_{\Delta} : \Delta \rightarrow \Omega$ . Since it is not constant, there exists a linear projection  $\pi : \mathbb{C}^N \rightarrow \mathbb{C}$  such that  $\pi \circ F_{\Delta}^{-1} : \Delta \rightarrow \mathbb{C}$  is not constant. Therefore, the map  $g := \pi \circ F^{-1} \circ \varphi : \mathbb{D} \rightarrow \mathbb{C}$  is a holomorphic bounded map. We claim that there exists  $a \in \mathbb{C}$  such that

$$(6.6) \quad \lim_{r \rightarrow 1^-} g(r\zeta) = a \quad \forall \zeta \in A,$$

which, by Fatou's lemma, implies that  $g$  is constant, reaching a contradiction.

In order to prove (6.6), we just note that if  $\{w_n\} \subset \Delta$  is a sequence converging to a point  $\zeta \in [p, q]$ , since  $\{w_n\} \subset E_x^D(\{u_n\}, R)$ , it follows that  $\{F^{-1}(w_n)\}$  is a non-relatively compact sequence in  $\Omega$  contained in the horosphere  $E_{F^{-1}(x)}^{\Omega}(\{F^{-1}(u_n)\}, R)$  and thus there exists a point  $u \in \partial \Omega$  (which depends only on  $\{F^{-1}(u_n)\}$ ) such that  $\{F^{-1}(w_n)\}$  converges to  $u$  (see (6.3)). From this, (6.6) follows at once.

Next, assume that  $L \cap D = \emptyset$ . Then  $[p, q] \subseteq L \cap \overline{D} = L \cap \partial D$ . Let  $\xi \in (p, q)$ . Let  $H \in \mathcal{L}_{\xi}$  be a complex supporting hyperplane for  $D$  at  $\xi$ . If  $\sigma : \mathbb{C}^N \rightarrow \mathbb{C}$  is a complex supporting functional such that  $H = \{z \in \mathbb{C}^N : \sigma(z - \xi) = 0\}$  then  $\sigma(p - q) = 0$ , since  $t(p - q) + \xi \in \partial D$  for  $t \in \mathbb{R}$ ,  $|t| \ll 1$ . This proves that  $L \subset H$ . But then,  $[p, q] \in \overline{D} \cap H$ , and by the arbitrariness of  $H$ , it follows that  $[p, q] \subset \operatorname{Ch}(\xi)$ , contradicting  $D$  being strictly  $\mathbb{C}$ -linearly convex.

Therefore,  $\Pi_D^H(\underline{x})$  consists of one point.  $\square$

The hypothesis of  $D$  being strictly  $\mathbb{C}$ -linearly convex in the previous proposition is not necessary, as the following example shows:

**Example 6.14.** Let  $D := \{(z_1, z_2) \in \mathbb{C}^2 : \operatorname{Re} z_1 > (\operatorname{Re} z_2)^2\}$ . It is easy to see that  $D$  is convex, and it is biholomorphic to the Siegel domain  $\mathbb{H}^2 := \{(w_1, w_2) \in \mathbb{C}^2 : \operatorname{Re} w_1 > |w_2|^2\}$  via the map  $(w_1, w_2) \mapsto (w_1 + 2w_2^2, w_2)$ . The Siegel domain  $\mathbb{H}$  is nothing but the unbounded realization of the ball  $\mathbb{B}^2$  via the generalized Cayley transform  $(z_1, z_2) \mapsto (\frac{1+z_1}{1-z_1}, \frac{z_2}{1-z_1})$ , hence, there exists a biholomorphism  $F : \mathbb{B}^2 \rightarrow D$  which extends as a homeomorphism on  $\partial\mathbb{B}^2 \setminus \{(1, 0)\}$ . In particular,  $F^{-1}$  is continuous in a neighborhood of the point  $(0, 0)$ , and  $F^{-1}(0, 0) = (-1, 0)$ . The sequence  $\{w_n\} := \{(\frac{1}{n}, 0)\}$  converges to  $(0, 0)$ , and thus  $\{F^{-1}(w_n)\}$  converges to  $(-1, 0)$  and it is admissible. It turns out that  $\{w_n\}$  is admissible and  $E_x^D(\{w_n\}, R) = \{(-1, 0)\}$  for all  $R > 0$  (here, as usual,  $x \in D$  is any chosen point). Therefore  $\Pi_H^D(\{u_n\}) = \{(-1, 0)\}$ . On the other hand, there is only one complex supporting hyperplane for  $D$  at  $(0, 0)$ , that is,  $H = \{(z_1, z_2) \in \mathbb{C}^2 : z_1 = 0\}$ . Therefore

$$\operatorname{Ch}(0, 0) = \partial D \cap H = \{(0, ti) : t \in \mathbb{R}\}.$$

**6.2. Convex domains biholomorphic to strongly convex domains.** The aim of this subsection is to prove Proposition 6.13 for convex domains biholomorphic to strongly convex domains without any assumption on  $\mathbb{C}$ -strictly convexity.

We first recall some notions of the Gromov hyperbolicity theory.

**Definition 6.15.** Let  $(D, d)$  be a metric space.

- A curve  $\gamma : [a, b] \rightarrow D$  is a *geodesic* if  $\gamma$  is an isometry for the usual distance function on  $[a, b] \subset \mathbb{R}$ , i.e.,  $d(\gamma(t_1), \gamma(t_2)) = |t_1 - t_2|$  for all  $t_1, t_2 \in [a, b]$ . We call  $\gamma([a, b])$  a *geodesic segment*.
- The metric space  $(D, d)$  is said to be a *geodesic metric space* if any two points in  $D$  are connected by a geodesic.
- A *geodesic triangle* in  $D$  is a union of images of three geodesics  $\gamma_i : [a_i, b_i] \rightarrow D$ ,  $i = 1, 2, 3$ , such that  $\gamma_i(b_i) = \gamma_{i+1}(a_{i+1})$  where the indices are taken modulo 3. The image of each  $\gamma_i$  is called a *side* of the geodesic triangle.
- A geodesic metric space  $(D, d)$  is *Gromov hyperbolic* or  $\delta$ -hyperbolic if there exists  $\delta > 0$  such that for any geodesic triangle in  $D$  the image of every side is contained in the  $\delta$ -neighborhood of union of the other two sides.
- Let  $A \geq 1$  and  $B > 0$ . We say that  $\gamma : [a, b] \rightarrow D$  is a  $(A, B)$  *quasi-geodesic* if for every  $t, t' \in ]a, b[$  we have:

$$\frac{1}{A}|t - t'| - B \leq d(\gamma(t), \gamma(t')) \leq A|t - t'| + B.$$

Also, we need to prove some preliminaries lemmas:

**Lemma 6.16.** Let  $D \subset \mathbb{C}^N$  be a convex domain biholomorphic to a bounded strongly pseudoconvex domain. Let  $\{p_j\}$  and  $\{q_j\}$  be two sequences in  $D$  which converge to two different boundary points. Then

$$(6.7) \quad \lim_{j \rightarrow \infty} K_D(p_j, q_j) = \infty.$$

*Proof.* Assume (6.7) is not true. Then, up to subsequences, we can assume that there exists  $C > 0$  such that for all  $j$

$$(6.8) \quad K_D(p_j, q_j) \leq C.$$

Since  $D$  is a complete hyperbolic convex domain, for every  $j$  there exists a complex geodesic  $\varphi_j : \mathbb{D} \rightarrow D$  such that  $\varphi_j(0) = p_j$  and  $\varphi_j(t_j) = q_j$  for some  $t_j \in (0, 1)$ , see [13, Lemma 3.3]. By (6.8) it follows

$$K_{\mathbb{D}}(0, t_j) = K_D(\varphi_j(p_j), \varphi_j(q_j)) = K_D(p_j, q_j) \leq C,$$

hence, there exists  $c \in (0, 1)$  such that  $t_j < c$  for all  $j$ , and we can assume without loss of generality that  $t_j \rightarrow t_0$  for some  $t_0 \in [0, 1)$ . Being  $D$  taut, up to subsequences, we can also assume that  $\{\varphi_j\}$  converges

uniformly on compacta to a holomorphic map  $\varphi : \mathbb{D} \rightarrow \overline{D}$ . Moreover,

$$\varphi(t_0) = \lim_{j \rightarrow \infty} \varphi_j(t_j) = \lim_{j \rightarrow \infty} q_j = q,$$

and similarly,  $\varphi(0) = p$ . This implies that  $t_0 \neq 0$  and  $\varphi$  is not constant. In particular,  $\partial D$  contains (non-constant) analytic discs. But  $D$  is Gromov hyperbolic (with respect to the Kobayashi distance) since it is biholomorphic to a bounded strongly pseudoconvex domain (see [7]) and by [38, Thm. 3.1],  $\partial D$  cannot contain (non-constant) analytic discs, a contradiction.  $\square$

**Lemma 6.17.** *Let  $D \subset \mathbb{C}^N$  be a bounded convex domain and let  $p \in \partial D$ . Let  $V \subset \mathbb{C}^N$  be an open set such that  $\text{Ch}(p) \subset V$  and  $D \setminus V \neq \emptyset$ . Then for every  $M > 0$  there exists an open set  $V' \subset \mathbb{C}^N$  such that  $\text{Ch}(p) \subset V' \subset V$  and such that for every  $z \in V' \cap D$  it holds  $K_D(z, D \setminus V) \geq M$ .*

*In particular,  $\lim_{n \rightarrow \infty} K_D(z_n, D \setminus V) = +\infty$  for every sequence  $\{z_n\} \subset D$  which accumulates on  $\text{Ch}(p)$ .*

*Proof.* Let  $H := \cap_{L \in \mathcal{L}_p} L$ . Then  $H$  is a complex affine space of dimension  $k \in \{0, \dots, n-1\}$ . In particular there exist  $n-k$  complex affine hyperplanes  $L_1, \dots, L_{n-k} \in \mathcal{L}_p$  in general position such that  $H = \cap_{j=1}^{n-k} L_j$ . For every  $j$  there exists a vector  $\underline{n}_j \in \mathbb{C}^N$  such that  $|\underline{n}_j| = 1$ ,  $L_j = \{z \in \mathbb{C}^N : \langle z - p, \underline{n}_j \rangle = 0\}$  and  $D \subset \{z \in \mathbb{C}^N : \text{Re} \langle z - p, \underline{n}_j \rangle < 0\}$ . We set

$$\psi(z) := \prod_{j=1}^{n-k} [1 - \langle z - p, \underline{n}_j \rangle]^{-1}.$$

It is easy to check that  $\psi : D \rightarrow \mathbb{D}$  is holomorphic and continuous up to  $\partial D$  and  $|\psi(z)| = 1$  for  $z \in \overline{D}$  if and only if  $\psi(z) = 1$  if and only if  $z \in \text{Ch}(p)$ .

Denote by  $\mathbb{D}(1, r)$  the Euclidean disc of center 1 and radius  $r > 0$  and by  $K_{\mathbb{D}}$  the Poincaré distance in  $\mathbb{D}$ . One can find  $1 > \epsilon > \epsilon' > 0$  such that  $\psi^{-1}(\mathbb{D}(1, \epsilon)) \cap D \subset V \cap D$  and for all  $\zeta \in \mathbb{D}(1, \epsilon') \cap \mathbb{D}$  it holds

$$(6.9) \quad K_{\mathbb{D}}(\zeta, \mathbb{D} \setminus \mathbb{D}(1, \epsilon)) \geq M.$$

Set  $V' = \psi^{-1}(\mathbb{D}(1, \epsilon')) \cap V$ . By construction,  $\text{Ch}(p) \subset V' \subset V$ .

Let  $w \in D \setminus V$  and let  $z \in V' \cap D$ . Note that  $\psi(w) \in \mathbb{D} \setminus \mathbb{D}(1, \epsilon)$  and  $\psi(z) \in \mathbb{D}(1, \epsilon') \cap \mathbb{D}$ . Hence, by (6.9) it holds  $K_D(z, w) \geq K_{\mathbb{D}}(\psi(w), \psi(z)) \geq M$ , and the proof is completed.  $\square$

**Lemma 6.18.** *Let  $D$  be a hyperbolic convex domain in  $\mathbb{C}^n$ . Let  $x \in D$  and let  $p \in \partial D$ . Then there exist  $A, B > 0$  such that for every sequence  $\{z_j\}$  of points in  $D$  converging to  $p$ , the line segment  $[x, z_j]$  is a  $(A, B)$  quasi-geodesic.*

*Proof.* The proof of Lemma 6.18 is a modification of the proof of [21, Proposition 4.2]. For convenience of the reader we present it, adapted to our situation.

For every  $j \geq 1$ , we parametrize  $[x, z_j]$  with respect to Kobayashi arc length, meaning that we consider a piecewise  $\mathcal{C}^1$  map  $\gamma_j : [0, t_j] \rightarrow D$  such that  $\gamma_j([0, t_j]) = [x, z_j]$  and the Kobayashi length  $l_D^K(\gamma_j([0, t]))$  of the line segment  $\gamma_j([0, t])$  is equal to  $t$ , for every  $j \geq 1$  and for every  $0 \leq t \leq t_j$ . We notice that  $\lim_{j \rightarrow \infty} t_j = +\infty$ .

By modifying  $D$  by a complex affine linear map if necessary, we may assume that  $p = 0$  and that  $D$  is contained in the real half space  $\mathbb{H}_n := \{(z', z_n) \in \mathbb{C}^n : \text{Re}(z_n) < 0\}$ . Moreover, by changing the real hyperplane if necessary, we may assume that there is  $t^1 > 0$ , sufficiently small, such that the interval  $\{t\nu, 0 < t \leq t^1\}$  is contained in  $D$  where  $\nu = (0' - 1) \in \mathbb{C}^{n-1} \times \mathbb{C}$ .

For a point  $y \in D$ , we denote by  $\delta(y, \partial D)$  the Euclidean distance from  $y$  to  $\partial D$  and by  $\delta(y, \partial D \cap \mathbb{C}\nu)$  the Euclidean distance from  $y$  to  $\partial D$  along the complex line  $\mathbb{C}\nu$ .

We choose  $t_0 > 0$  such that for every  $j \geq 1$  and for every  $q_j \in \gamma_j([t_0, t_j])$  there is some point  $q'_j \in D \cap \{t\nu, 0 < t < t^1\}$  with  $Re((q'_j)_n) = Re((q_j)_n)$  where  $(q'_j)_n$  and  $(q_j)_n$  denote the  $n$ -th components of  $q'_j$  and  $q_j$ .

It is enough to prove the following

*Claim.*  $\gamma_j|_{[t_0, t_j]}$  is a  $(A_1, B_1)$  quasi-geodesic for some suitable constants  $A_1 > 1, B_1 > 0$  independent of  $j$ .

Indeed suppose for the moment that the Claim is proved.

- Let  $0 < t < t_0 < t'$  and let  $j \geq 1$ . Since  $|t' - t| = l_D^K(\gamma_j([t, t']))$  we have:

$$\begin{aligned} K_D(\gamma_j(t), \gamma_j(t')) &\leq l_D^K(\gamma_j([t, t'])) \\ &\leq |t' - t| \\ &\leq t_0 + |t_0 - t'| \\ &\leq t_0 + A_1 K_D(\gamma_j(t_0), \gamma_j(t')) + A_1 B_1 \\ &\leq t_0 + A_1 K_D(\gamma_j(t), \gamma_j(t')) + A_1 B_1. \end{aligned}$$

For the fourth inequality we use the fact that  $\gamma_j|_{[t_0, t_j]}$  is a  $(A_1, B_1)$  quasi-geodesic.

- Let  $t < t' < t_0$ . Since  $D$  is complete hyperbolic (see [13]) then the Kobayashi balls are compact. Moreover the points  $\gamma_j(t)$  and  $\gamma_j(t')$  satisfy  $K_D(x, \gamma_j(t)) \leq l_D^K(\gamma_j([0, t])) = t \leq t_0$ , and hence they are in the closed Kobayashi ball centered at  $x$  with radius  $t_0$ . Since the topologies defined by the Kobayashi distance and the Euclidean distance are equivalent on the closed Kobayashi ball centered at  $x$  with radius  $t_0$ , there exist  $A_2 > 1$  and  $B_2 > 0$  independent of  $j \geq 1$  such that for every  $j \geq 1$ :

$$K_D(\gamma_j(t), \gamma_j(t')) \leq l_D^K(\gamma_j([t, t'])) = |t' - t| \leq A_2 K_D(\gamma_j(t), \gamma_j(t')) + B_2.$$

- Let  $t_0 < t < t'$ . We use directly the Claim to obtain:

$$K_D(\gamma_j(t), \gamma_j(t')) \leq l_D^K(\gamma_j([t, t'])) \leq A_1 K_D(\gamma_j(t), \gamma_j(t')) + B_1.$$

If we set  $A := \max(A_1, A_2)$  and  $B := \max(t_0 + A_1 B_1, B_1)$  then it follows from the three cases above that  $\gamma_j$  is a  $(A, B)$  quasi-geodesic with  $A, B > 0$ , independent of  $j \geq 1$ .

Hence we need to prove that there exist  $A_1, B_1 > 0$  such that the line segment  $\gamma_j([t_0, t_j])$  is a  $(A_1, B_1)$  quasi-geodesic for every  $j$ , where  $A_1$  and  $B_1$  depend only on  $D$ , on  $x$  and on  $p$ .

For every  $j \geq 1$ , let  $\lambda_j : [0, s_j] \rightarrow D$  denote the line segment joining  $q_j$  to  $q'_j$  and parametrized by Kobayashi arc length. We can prove exactly as in the proof of [21, Proposition 4.2] that there is  $C_1 > 0$ , not depending on  $j$ , such that:

$$l_D^K(\lambda_j([0, s_j])) \leq C_1 \frac{|q_j - q'_j|}{\delta_D(q'_j)}.$$

By the choice of  $\nu$  there exists  $C_2 > 0$ , independent of  $j$ , such that for every  $j \geq 1$ :  $|q_j - q'_j| \leq C_2 \delta_D(q'_j)$ .

The two previous inequalities imply

$$l_D^K(\lambda_j([0, s_j])) \leq C_1 C_2,$$

which is equivalent to Inequality (4.1) in [21].

In particular, to prove the result, it is sufficient to prove that there exist  $A_2, B_2 > 0$  such that the line segment  $[x, q'_j]$  is a  $(A_2, B_2)$  quasi-geodesic, for every  $j \geq 1$ .

Since  $K_D(x, t^1\nu)$  is finite, the line segment  $[x, t^1\nu]$  is a quasi-geodesic. It remains to prove that there exist  $A_2, B_2 > 0$  such that the line segment  $[t^1\nu, q'_j]$  is a  $(A_2, B_2)$  quasi-geodesic, for every  $j \geq 1$ .

Let  $\gamma : \mathbb{R}^+ \rightarrow D$ , parametrizing the interval  $]0, t^1\nu]$  by the Kobayashi length, with  $\gamma(0) = t^1\nu$ .

Since  $D$  is convex, there exists  $C > 1$  such that for every  $s \in \mathbb{R}^+$ :

$$\delta(\gamma(s), \partial D \cap \mathbb{C}\nu) \geq \delta(\gamma(s), \partial D) \geq \frac{1}{C} \delta(\gamma(s), p).$$

Hence, we have for every  $0 < t < t'$ :

$$\begin{aligned} K_D(\gamma(t), \gamma(t')) &\leq l_D^K(\gamma(t), \gamma(t')) \leq \int_t^{t'} \frac{|\gamma'(s)|}{\delta(\gamma(s), \partial D \cap \mathbb{C}\nu)} ds \\ &\leq C \int_t^{t'} \frac{|\gamma'(s)|}{\delta(\gamma(s), p)} ds \\ &\leq C \int_t^{t'} \frac{|\gamma'(s)|}{\delta(\gamma(s), \partial \mathbb{H}_n \cap \mathbb{C}\nu)} ds \\ &\leq CK_{H_{\mathbb{H}_n}}(\gamma(t), \gamma(t')) \\ &\leq CK_D(\gamma(t), \gamma(t')). \end{aligned}$$

The third inequality uses the fact that  $\delta(\gamma(s), \partial \mathbb{H}_n \cap \mathbb{C}\nu) \leq \delta(\gamma(s), p)$  since  $p \in \partial \mathbb{H}_n \cap \mathbb{C}\nu$ . The fourth inequality uses the fact that  $l_{\mathbb{H}_n}^K(\gamma(t), \gamma(t')) = K_{\mathbb{H}_n}(\gamma(t), \gamma(t'))$  since  $\gamma(\mathbb{R}^+)$  is contained in the real inward normal of  $\mathbb{H}_n$  at 0. The fifth inequality uses the fact that  $D$  is contained in  $\mathbb{H}_n$ .

This proves that the line segment  $[t\nu, t'\nu]$  is a  $(C, c)$  quasi-geodesic, for every  $c > 0$  and for every  $j \geq 1$ : the Claim, and hence Lemma 6.18, is proved.  $\square$

We will need the following statement (see also Lemma 3.3. in [22]).

**Lemma 6.19.** *Let  $D \subset \mathbb{C}^N$  be a convex domain biholomorphic to a bounded strongly convex domain with  $C^3$  boundary. Then for every couple of points  $z_0, w_0 \in D$  there exists a unique real geodesic for the Kobayashi distance which joins  $z_0$  and  $w_0$ .*

*Proof.* Since  $D$  is biholomorphic to a bounded strongly convex domain, Lempert's theory [27, 28, 29] (see also [3]) implies that for every  $z_0, w_0 \in D$ ,  $z_0 \neq w_0$ , there exists a unique complex geodesic whose image contains  $z_0, w_0$ . In other words, there exists  $\varphi : \mathbb{D} \rightarrow D$  holomorphic such that  $K_{\mathbb{D}}(\zeta, \eta) = K_D(\varphi(\zeta), \varphi(\eta))$  for all  $\zeta, \eta \in \mathbb{D}$  and  $z_0, w_0 \in \varphi(\mathbb{D})$  and, moreover, if  $\tilde{\varphi} : \mathbb{D} \rightarrow D$  is any holomorphic map such that there exist  $\zeta_0, \zeta_1 \in \mathbb{D}$  with  $\tilde{\varphi}(\zeta_0) = z_0$ ,  $\tilde{\varphi}(\zeta_1) = w_0$  and  $K_D(z_0, w_0) = K_{\mathbb{D}}(\zeta_0, \zeta_1)$ , then there exists an automorphism  $\theta : \mathbb{D} \rightarrow \mathbb{D}$  such that  $\tilde{\varphi} = \varphi \circ \theta$ . If  $\varphi : \mathbb{D} \rightarrow D$  is a complex geodesic, there exists a holomorphic retraction  $\rho : D \rightarrow D$ , called the Lempert projection, such that  $\rho \circ \varphi = \varphi$ ,  $\rho(D) = \varphi(\mathbb{D})$ . In what follows we will use the following fact: if  $z_0 \in \varphi(\mathbb{D})$  and  $w \in D \setminus \varphi(\mathbb{D})$  then

$$(6.10) \quad K_D(z_0, \rho(w)) = K_D(\rho(z_0), \rho(w)) < K_D(z_0, w).$$

In order to prove inequality (6.10) we can assume that  $D$  is a bounded strongly convex domain with  $C^3$  boundary. Let  $\varphi : \mathbb{D} \rightarrow D$  be a complex geodesic and let  $\rho : D \rightarrow \varphi(\mathbb{D})$  be the associated Lempert projection. Fix  $R > 0$  and  $z_0 \in \varphi(\mathbb{D})$ . Since  $D$  is assumed to be strongly convex, the Kobayashi ball  $B_K(z_0, R)$  of center  $z_0$  and radius  $R > 0$  is strongly convex as well, and the Lempert projection  $\rho$  has affine fibers. Taking into account that  $\rho$  contracts the Kobayashi distance  $K_D$ , it follows that for every  $\zeta \in \mathbb{D}$

with  $K_D(\varphi(\zeta), z_0) = R$  it holds  $\rho^{-1}(\varphi(\zeta)) \cap \overline{B_K(z_0, R)} = \{\varphi(\zeta)\}$ . From this, (6.10) follows at once by considering  $\zeta \in \mathbb{D}$  with  $\varphi(\zeta) = \rho(w)$ .

Now, let  $z_0, w_0 \in D$ . Hence, there exists a unique complex geodesic  $\varphi : \mathbb{D} \rightarrow D$  such that  $z_0, w_0 \in \varphi(\mathbb{D})$ . Since  $\varphi$  is an isometry between  $K_{\mathbb{D}}$  and  $K_D$ , it follows that there exists a real geodesic for  $K_D$ , call it  $\gamma : [0, a] \rightarrow \varphi(\mathbb{D})$ ,  $\gamma(0) = z_0, \gamma(a) = w_0$  for  $a = k_D(z_0, w_0)$ , which is contained in  $\varphi(\mathbb{D})$ . Moreover,  $\gamma$  is the only real geodesic joining  $z_0$  and  $w_0$  contained in  $\varphi(\mathbb{D})$ .

Let assume that  $\tilde{\gamma} : [0, a] \rightarrow D$ , is another real geodesic for  $K_D$  such that  $\tilde{\gamma}(0) = z_0$  and  $\tilde{\gamma}(a) = w_0$  and  $\tilde{\gamma}([0, a]) \neq \gamma([0, a])$ . Then the image of  $\tilde{\gamma}$  is not contained in  $\varphi(\mathbb{D})$ . Hence, there exists  $t \in (0, a)$  such that  $z_1 := \tilde{\gamma}(t) \notin \varphi(\mathbb{D})$ . Then, since  $\tilde{\gamma}$  is a real geodesic for the Kobayashi distance,

$$K_D(z_0, w_0) = K_D(z_0, z_1) + K_D(z_1, w_0).$$

Now, let  $\rho : D \rightarrow \varphi(\mathbb{D})$  be the Lempert projection associated with  $\varphi$ . Since  $\rho$  contracts the Kobayashi distance, by (6.10) it follows:

$$\begin{aligned} K_D(z_0, w_0) &= K_D(z_0, z_1) + K_D(z_1, w_0) > k_D(\rho(z_0), \rho(z_1)) + k_D(\rho(z_1), \rho(w_0)) \\ &\geq K_D(\rho(z_0), \rho(w_0)) = K_D(z_0, w_0), \end{aligned}$$

a contradiction, and the result is proved.  $\square$

**Theorem 6.20.** *Let  $D \subset \mathbb{C}^N$  be a bounded convex domain biholomorphic to a bounded strongly convex domain with  $C^3$  boundary. Then for every  $\underline{x} \in \partial_H D$  there exists a unique  $p \in \partial D$  such that*

$$(6.11) \quad I_D^H(\underline{x}) = \Pi_D^H(\underline{x}) = \{p\}.$$

*Proof.* If we prove that  $\Pi_D^H(\underline{x}) = \{p\}$ , then the result follows from Theorem 6.10.

Assume by contradiction that there exists  $q \in \Pi_D^H(\underline{x})$  with  $p \neq q$ . Fix  $x \in D$ .

Since  $\Pi_D^H(\underline{x})$  is convex, the real segment  $[p, q] \subset \Pi_D^H(\underline{x})$ . Let  $\{u_n\}$  be an admissible sequence which represents  $\underline{x}$ . By (6.4), the segment  $[p, q] \subset \Pi(\{u_n\}, R)$  for all  $R > 0$ . In particular, if we take  $R > 1$  so that  $x \in E_x(\{u_n\}, R)$ , the real segments  $[x, p]$  and  $[x, q]$  are contained in  $E_x(\{u_n\}, R)$  (by convexity, see Proposition 6.1). Set  $p_j := (1 - \frac{1}{j})p + \frac{1}{j}x$  and  $q_j := (1 - \frac{1}{j})q + \frac{1}{j}x$ ,  $j \geq 1$ . Then both sequences  $\{p_j\}$  and  $\{q_j\}$  are contained in  $E_x(\{u_n\}, R)$ .

By Lemma 6.18, there exist  $A, B > 0$  such that for all  $j \geq 1$ , the segments  $[x, p_j]$  and  $[x, q_j]$  are  $(A, B)$ -quasi-geodesics.

By Lemma 6.6, the sequences  $\{p_j\}, \{q_j\}$  are admissible and equivalent to  $\{u_n\}$ .

By hypothesis, there exist  $\Omega \subset \mathbb{C}^N$  a bounded strongly pseudoconvex domain with  $C^3$  boundary and a biholomorphism  $F : \Omega \rightarrow D$ . Then the sequences  $\{F^{-1}(p_j)\}$  and  $\{F^{-1}(q_j)\}$  belong to the same horosphere  $E_{F^{-1}(x)}(\{F^{-1}(u_n)\}, R)$  and are equivalent. Also, both sequences are equivalent to  $\{F^{-1}(u_n)\}$ . Hence, it follows from Proposition 5.9 that the sequences  $\{F^{-1}(p_j)\}, \{F^{-1}(q_j)\}$  and  $\{F^{-1}(u_j)\}$  converge to the same boundary point  $p \in \partial\Omega$ . Now according to Theorem 5.2 we have:  $E_{F^{-1}(x)}(p, R) = E_{F^{-1}(x)}(p, R)$ . This gives by Proposition 5.9:

$$E_{F^{-1}(x)}(\{F^{-1}(u_n)\}, R) = E_{F^{-1}(x)}(\{F^{-1}(q_j)\}, R) = E_{F^{-1}(x)}(\{F^{-1}(p_j)\}, R)$$

or equivalently:

$$E_x(\{u_n\}, R) = E_x(\{q_j\}, R) = E_x(\{p_j\}, R).$$

Fix  $R' > R$ . Let  $j \in \mathbb{N}$ . Then there exists  $n_j \in \mathbb{N}$  such that for all  $n \geq n_j$  it holds  $K_D(p_j, q_n) - K_D(x, q_n) \leq \frac{1}{2} \log R'$ . In particular, we can find a sequence  $\{n_j\} \subset \mathbb{N}$  converging to  $\infty$  such that for all  $j \in \mathbb{N}$  it holds

$$K_D(p_j, q_{n_j}) - K_D(x, q_{n_j}) \leq \frac{1}{2} \log R'.$$

Up to relabelling, we still denote  $\{q_{n_j}\}$  by  $\{q_j\}$ . With a similar argument, exchanging the role of  $\{p_j\}$  and  $\{q_j\}$  (in fact, of  $\{q_{n_j}\}$ ), we find a subsequence of  $\{p_j\}$  (which we still denote by  $\{p_j\}$ ) such that for all  $j \in \mathbb{N}$  it holds  $K_D(p_j, q_j) - K_D(x, p_j) \leq \frac{1}{2} \log R'$ . In the end, we prove that we can choose  $\{p_j\}, \{q_j\}$  in such a way that there exists a constant  $C > 0$  such that for all  $j \in \mathbb{N}$  it holds

$$(6.12) \quad \begin{aligned} K_D(p_j, q_j) &\leq k_D(p_j, x) + C, \\ K_D(p_j, q_j) &\leq k_D(q_j, x) + C. \end{aligned}$$

By Lemma 6.19, for every  $j \in \mathbb{N}$  there exists a unique real geodesic  $\gamma_j$  for the Kobayashi distance which joins  $p_j$  with  $q_j$ .

We claim that  $\Gamma_j := [x, p_j] \cup \gamma_j$  is a quasi-geodesic joining  $x$  with  $q_j$ . Indeed, denoting by  $\ell(\Gamma_j)$  its hyperbolic length, by (6.12), we have

$$\ell(\Gamma_j) \leq AK_D(x, p_j) + B + K_D(p_j, q_j) \leq AK_D(x, p_j) + B + K_D(x, q_j) + C.$$

Moreover, by (6.12) and the triangle inequality,

$$K_D(x, p_j) \leq K_D(x, q_j) + K_D(q_j, p_j) \leq 2K_D(x, q_j) + C.$$

Hence,

$$\ell(\Gamma_j) \leq (2A + 1)K_D(x, q_j) + C(A + 1) + B,$$

proving that  $\Gamma_j$  is a  $(2A + 1, C(A + 1) + B)$ -quasi-geodesic for all  $j \in \mathbb{N}$ .

Since  $(D, K_D)$  is Gromov hyperbolic by [7, Theorem 1.4], by Gromov's result (see [23], 6. Théorème page 82), for every  $A, B > 0$  fixed, there exists  $M > 0$  such that for every  $(A, B)$ -quasi-geodesic  $\xi$  there exists a (real) geodesic  $\hat{\xi}$  for  $K_D$  such that  $\xi$  belongs to the hyperbolic neighborhood

$$\mathcal{N}_M(\hat{\xi}) := \{z \in D : \exists w \in \hat{\xi} : K_D(w, z) < M\}.$$

In particular, in our setting, given the unique real geodesic  $\eta_j$  joining  $x$  and  $q_j$ , the two quasi-geodesics  $[x, q_j]$  and  $\Gamma_j$  belong to some hyperbolic  $\frac{M}{2}$ -neighborhood of  $\eta_j$ , for some  $M > 0$  and for all  $j \in \mathbb{N}$ . Hence, by the triangle inequality, for all  $j \in \mathbb{N}$ , it holds  $\Gamma_j \subset \mathcal{N}_M([x, q_j])$ . This implies that  $K_D(p_j, [x, q_j]) < M$  for every  $j \in \mathbb{N}$ . Thus, there exists a sequence  $\{w_j\}$  such that  $w_j \in [x, q_j]$  and for all  $j \in \mathbb{N}$  it holds

$$(6.13) \quad K_D(p_j, w_j) < 2M.$$

The sequence  $\{w_j\}$  has to be relatively compact in  $D$  by Lemma 6.16. But this contradicts Lemma 6.17.  $\square$

## 7. HOROSPHERE BOUNDARY VERSUS GROMOV BOUNDARY

Different types of topological boundaries may be defined in the general context of metric spaces. For instance the construction of the Gromov boundary, using geodesic rays or sequences, is based on the following

**Definition 7.1.** Let  $(X, d)$  be a metric space and let  $w \in X$  be a base point.

- (i) Given  $x, y \in X$ , the Gromov product of  $x$  and  $y$  with respect to  $w$  is defined by  $(x, y)_w := \frac{1}{2}(d(x, w) + d(y, w) - d(x, y))$ .
- (ii) A geodesic ray  $\gamma : [0, +\infty) \rightarrow X$ , with  $\gamma(0) = w$ , is an isometry from  $([0, +\infty), |\cdot|)$  to  $(X, d)$  where  $|\cdot|$  denotes the absolute value on  $\mathbb{R}$ ; namely if  $l(\gamma([s, t]))$  denotes the length of the curve  $\gamma([s, t])$ , we have:

$$\forall s, t \geq 0, l(\gamma([s, t])) = d(\gamma(s), \gamma(t)) = |t - s|.$$

- (iii) Two geodesic rays  $\gamma$  and  $\tilde{\gamma}$  are equivalent (we write  $\gamma \sim_r \tilde{\gamma}$ ) if they are at finite Hausdorff distance from each other, i.e., if there exists  $\exists C > 0$  such that for all  $t \geq 0$ , it holds  $d(\gamma(t), \tilde{\gamma}(t)) \leq C$ . The relation  $\sim_r$  is an equivalence relation on the set of geodesic rays.

- (iv) A sequence  $\{x^\nu\}_\nu$  of points in  $X$  tends to infinity if  $\lim_{\nu, \mu \rightarrow \infty} (x^\nu, x^\mu)_w = +\infty$ .
- (v) Two sequences  $\{x^\nu\}_\nu$  and  $\{y^\nu\}_\nu$  of points in  $X$  are equivalent (we write  $\{x^\nu\} \sim_s \{y^\nu\}$ ) if  $\lim_{\nu \rightarrow \infty} (x^\nu, y^\nu)_w = +\infty$ .

There are different ways to define the Gromov boundaries, using geodesic rays, or using sequences. Both are equivalent when  $X$  is a geodesic, proper, hyperbolic (in the sense of Gromov) metric space.

**Definition 7.2.** Let  $(X, d)$  be a geodesic, proper, metric space.

- (i) The Gromov boundary  $\partial_G^r X$  (with respect to rays) is defined by

$$\partial_G^r X := E^r / \sim_r$$

where  $E^r$  denotes the set of geodesic rays.

- (ii) Let  $p \in \partial_G^r X$  and let  $r > 0$ . We consider

$$V(p, r) := \{q \in \partial_G^r X : \exists \gamma \in p, \exists \tilde{\gamma} \in q, \liminf_{s, t \rightarrow \infty} (\gamma(t), \tilde{\gamma}(s))_w \geq r\}.$$

Then  $(V(p, r))_{p \in \partial_G^r X, r > 0}$  is a basis of open neighborhoods for the Gromov topology  $\mathcal{T}_G^r(\partial_G^r X)$  on  $\partial_G^r X$ .

**Remark 7.3.** In the previous definitions it is not necessary that the domain be (Gromov) hyperbolic. However, if  $(X, d)$  is not (Gromov) hyperbolic then the Gromov boundary  $\partial_G^r(X)$  may not be Hausdorff.

One may also define the Gromov boundary using equivalent sequences:

**Definition 7.4.** Let  $(X, d)$  be a proper, hyperbolic (in the sense of Gromov) metric space. The Gromov boundary  $\partial_G^s X$  (with respect to sequences) is defined by

$$\partial_G^s X := E^s / \sim_s$$

where  $E^s$  denotes the set of sequences converging to infinity.

**Remark 7.5.** • In case  $(X, d)$  is proper, geodesic, (Gromov) hyperbolic, there exists a bijection between  $\partial_G^r X$  and  $\partial_G^s X$ .

- If  $(X, d)$  is not (Gromov) hyperbolic then  $\sim_s$  may not be an equivalence relation. This is the case for the bidisc endowed with its Kobayashi distance  $K_{\mathbb{D}^2}$ :  $\{(1 - 1/n, 0)\} \sim_s \{(0, 1 - 1/n)\}$ ,  $\{(0, 1 - 1/n)\} \sim_s \{(-1 + 1/n, 0)\}$  but  $\{(1 - 1/n, 0)\} \not\sim_s \{(-1 + 1/n, 0)\}$ . In particular we can not define  $\partial_G^s \mathbb{D}^2$ .

It is natural to study the relations between the Gromov boundary  $\partial_G^r M$  and the horosphere boundary  $\partial_H M$  for a complete (Kobayashi) hyperbolic manifold  $(M, K_M)$ . Indeed, since  $(M, K_M)$  is a length space, it follows from the Hopf-Rinow Theorem that any two points can be joined by a geodesic segment.

Let  $\mathbb{D}^2 := \{(z_1, z_2) \in \mathbb{C}^n : |z_1| < 1, |z_2| < 1\}$  be the bidisc in  $\mathbb{C}^n$ . The remaining part of this section is dedicated to prove the following

**Proposition 7.6.** The horosphere boundary and the Gromov boundary of the bidisc, endowed with their respective topologies, are not homeomorphic.

We first start with a description of the horospheres of sequences converging to some boundary point in  $\partial \mathbb{D}^2$ . In order to avoid burdening notations, in this section, we simply write  $E(\{u_n\}, R)$  to denote the horospheres of an admissible sequence  $\{u_n\}$  in  $\mathbb{D}^2$  with respect to the base point  $(0, 0)$ .

**Lemma 7.7.** Every sequence  $\{u_n = (u_n^1, u_n^2)\} \subset \mathbb{D}^2$  converging to a point  $(p_1, p_2) \in \partial \mathbb{D}^2$  is admissible. Moreover,



- (i) if  $p = (e^{it}, p_2) \in \partial\mathbb{D}^2$  with  $t \in \mathbb{R}$  and  $|p_2| < 1$  (respectively  $p = (p_1, e^{it}) \in \partial\mathbb{D}^2$  with  $|p_1| < 1$ ,  $t \in \mathbb{R}$ ) for every  $R > 0$  it holds

$$E(\{u_n\}, R) = E_{\mathbb{D}}(e^{it}, R) \times \mathbb{D}$$

(respect.,  $E(\{u_n\}, R) = \mathbb{D} \times E_{\mathbb{D}}(e^{it}, R)$ ).

- (ii) If  $p = (e^{it_1}, e^{it_2}) \in \partial\mathbb{D}^2$ ,  $t_1, t_2 \in \mathbb{R}$ , let  $T_1 := \limsup_n \left( \frac{1-|u_n^1|^2}{1-|u_n^2|^2} \right)$  and  $T_2 := \limsup_n \left( \frac{1-|u_n^2|^2}{1-|u_n^1|^2} \right)$ .

Then, for every  $R > 0$ :

- (a)  $(T_1 > 1, T_2 > 1) \Rightarrow E(\{u_n\}, R) = E_{\mathbb{D}}(e^{it_1}, R) \times E_{\mathbb{D}}(e^{it_2}, R)$ ,
- (b)  $(T_1 > 1, T_2 \leq 1) \Rightarrow E(\{u_n\}, R) = E_{\mathbb{D}}(e^{it_1}, R/T_2) \times E_{\mathbb{D}}(e^{it_2}, R)$ ,
- (c)  $(T_1 \leq 1, T_2 > 1) \Rightarrow E(\{u_n\}, R) = E_{\mathbb{D}}(e^{it_1}, R) \times E_{\mathbb{D}}(e^{it_2}, R/T_1)$

with the convention  $E_{\mathbb{D}}(e^{it_1}, +\infty) = \mathbb{D}$ .

*Proof.* (i). Let  $p = (e^{it}, p_2) \in \partial\mathbb{D}^2$  with  $|p_2| < 1$ . Then for  $R > 0$ , for  $w = (w_1, w_2) \in \mathbb{D}$  and for sufficiently large  $n$ :

$$K_{\mathbb{D}^2}(u_n, w) - K_{\mathbb{D}^2}(u_n, 0) = K_{\mathbb{D}}(u_n^1, w_1) - K_{\mathbb{D}}(u_n^1, 0).$$

Hence  $E_{\mathbb{D}^2}(\{u_n\}, R) = E_{\mathbb{D}}(\{u_n^1\}, R) \times \mathbb{D}$ .

(ii). By definition

$$K_{\mathbb{D}^2}(u_n, w) - K_{\mathbb{D}^2}(u_n, 0) = \max(K_{\mathbb{D}}(u_n^1, w_1), K_{\mathbb{D}}(u_n^2, w_2)) - \max(K_{\mathbb{D}}(u_n^1, 0), K_{\mathbb{D}}(u_n^2, 0)).$$

Following [3, pp.264-266], we obtain

$$(7.1) \quad \begin{aligned} & \limsup_{n \rightarrow \infty} (K_{\mathbb{D}^2}(u_n, w) - K_{\mathbb{D}^2}(u_n, 0)) \\ &= \frac{1}{2} \log \left( \max \left( \frac{|w_1 - e^{it_1}|^2}{1 - |w_1|^2} \min(1, T_2), \frac{|w_2 - e^{it_2}|^2}{1 - |w_2|^2} \min(1, T_1) \right) \right). \end{aligned}$$

Hence

$$w \in E(\{u_n\}, R) \Leftrightarrow \begin{cases} \frac{|w_1 - e^{it_1}|^2}{1 - |w_1|^2} \min(1, T_2) < R \\ \frac{|w_2 - e^{it_2}|^2}{1 - |w_2|^2} \min(1, T_1) < R \end{cases}$$

The conclusion follows directly from that equivalence.  $\square$

Now we show that, in fact, one can only consider admissible sequences:

**Proposition 7.8.** *Let  $\{u_n\}$  be an admissible sequence in  $\mathbb{D}^2$ . Then  $\{u_n\}$  is equivalent to one and only one of the following sequences:*

- (1)  $\{w_n^{(1)}(p_1, p_2) := (p_1(1 - \frac{1}{n}), p_2(1 - \frac{1}{n}))\}_{n \in \mathbb{N}}$  for some  $p_1, p_2 \in \partial\mathbb{D}$ ,
- (2)  $\{w_n^{(2)}(p) := (p(1 - \frac{1}{n}), 0)\}_{n \in \mathbb{N}}$  for some  $p \in \partial\mathbb{D}$ ,
- (3)  $\{w_n^{(3)}(p) := (0, p(1 - \frac{1}{n}))\}_{n \in \mathbb{N}}$  for some  $p \in \partial\mathbb{D}$ .

*In particular, every admissible sequence in  $\mathbb{D}^2$  is equivalent to an admissible sequence in  $\mathbb{D}^2$  which converges to a point in  $\partial\mathbb{D}^2$ .*

*Proof.* First of all, it is clear by Lemma 7.7 that the horospheres of  $\{w_n^{(1)}(p_1, p_2)\}$  are  $E_{\mathbb{D}}(p_1, R) \times E_{\mathbb{D}}(p_2, R)$ , the horospheres of  $\{w_n^{(2)}(p)\}$  are  $E(p, R) \times \mathbb{D}$  and of  $\{w_n^{(3)}(p)\}$  are  $\mathbb{D} \times E(p, R)$ , for all  $R > 0$ . Hence the sequences  $\{w_n^{(1)}(p_1, p_2)\}$ ,  $\{w_n^{(2)}(p)\}$  and  $\{w_n^{(3)}(p)\}$  are not equivalent.

Then, as in (6.2), we have

$$(7.2) \quad E(\{u_n\}, R) = \bigcap_{\{v_n\} \in \mathcal{M}} E(\{v_n\}, R),$$

where  $\mathcal{M}$  denotes the set of all subsequences of  $\{u_n\}$  converging to a boundary point. In particular, every sequence  $\{v_n\} \subset \mathcal{M}$  is admissible (by (7.2) or by Lemma 7.7). Note that (7.2) implies that if  $\{v_n\}, \{\tilde{v}_n\} \in \mathcal{M}$  then for every  $R > 0$  it holds

$$(7.3) \quad E(\{v_n\}, R) \cap E(\{\tilde{v}_n\}, R) \neq \emptyset.$$

The horospheres of converging sequences are described in Lemma 7.7. There are different cases to be considered:

*Case 1.* Assume that there exist  $p_1, p_2 \in \partial\mathbb{D}$ ,  $\{v_n\}, \{\tilde{v}_n\} \in \mathcal{M}$ , (possibly  $\{v_n\} = \{\tilde{v}_n\}$ ) and  $T \in (0, 1]$  such that

$$E(\{v_n\}, R) \cap E(\{\tilde{v}_n\}, R) = E_{\mathbb{D}}(p_1, R) \times E_{\mathbb{D}}(p_2, R/T)$$

for every  $R > 0$ .

We claim that  $\{u_n\}$  is equivalent to the sequence  $\{w_n^{(1)}(p_1, p_2)\}$ . Indeed, taking into account (7.3) and the list in Lemma 7.7, we see that if  $\{z_n\} \in \mathcal{M}$ , then either  $E(\{z_n\}, R) = E_{\mathbb{D}}(p_1, R) \times E_{\mathbb{D}}(p_2, R/Q)$  or  $E(\{z_n\}, R) = E_{\mathbb{D}}(p_1, R/Q) \times E_{\mathbb{D}}(p_2, R)$  for some  $Q \in [0, 1]$  (with the usual convention that, if  $Q = 0$  then  $E_{\mathbb{D}}(p_1, R/Q) = \mathbb{D}$ ). In any case,  $E(\{w_n^{(1)}(p_1, p_2)\}, R) \subset E(\{z_n\}, R)$ . Therefore from (7.2),

$$E_{\mathbb{D}}(p_1, R) \times E_{\mathbb{D}}(p_2, R) \subset E(\{u_n\}, R) \subset E_{\mathbb{D}}(p_1, R) \times E_{\mathbb{D}}(p_2, R/T).$$

In particular, it is easy to see that for every  $R > 0$  there exists  $R' > 0$  such that  $E(\{w_n^{(1)}(p_1, p_2)\}, R) \subset E(\{u_n\}, R) \subset E(\{w_n^{(1)}(p_1, p_2)\}, R')$ . Hence  $\{u_n\}$  is equivalent to  $\{w_n^{(1)}(p_1, p_2)\}$ .

*Case 2.* Assume that there exist  $p_1, p_2 \in \partial\mathbb{D}$ ,  $\{v_n\}, \{\tilde{v}_n\} \in \mathcal{M}$ , (possibly  $\{v_n\} = \{\tilde{v}_n\}$ ) and  $T \in (0, 1]$  such that

$$E(\{v_n\}, R) \cap E(\{\tilde{v}_n\}, R) = E_{\mathbb{D}}(p_1, R/T) \times E_{\mathbb{D}}(p_2, R)$$

for every  $R > 0$ .

In this case the argument goes exactly as in Case 1, and  $\{u_n\}$  is equivalent to  $\{w_n^{(1)}(p_1, p_2)\}$ .

*Case 3.* There are no sequences  $\{v_n\}, \{\tilde{v}_n\}$  in  $\mathcal{M}$  as in Case 1. Hence, by Lemma 7.7, for all  $\{v_n\} \in \mathcal{M}$  there exists  $p_{\{v_n\}} \in \partial\mathbb{D}$  such that either  $E(\{v_n\}, R) = E_{\mathbb{D}}(p_{\{v_n\}}, R) \times \mathbb{D}$ , or  $E(\{v_n\}, R) = \mathbb{D} \times E_{\mathbb{D}}(p_{\{v_n\}}, R)$  for all  $R > 0$ . Moreover, since we are excluding Case 1 and Case 2, if for some  $\{v_n\} \in \mathcal{M}$  it holds  $E(\{v_n\}, R) = E_{\mathbb{D}}(p_{\{v_n\}}, R) \times \mathbb{D}$ , then for all  $\{\tilde{v}_n\} \in \mathcal{M}$  it holds  $E(\{\tilde{v}_n\}, R) = E_{\mathbb{D}}(p_{\{\tilde{v}_n\}}, R) \times \mathbb{D}$ . By (7.3), there exists  $p \in \partial\mathbb{D}$  such that  $p_{\{v_n\}} = p$  for all  $\{v_n\} \in \mathcal{M}$ , hence  $E(\{u_n\}, R) = E(p, R) \times \mathbb{D}$ , and  $\{u_n\}$  is equivalent to  $\{u_n\}$  is equivalent to  $\{w_n^{(2)}(p)\}$ .

Similarly, if  $E(\{v_n\}, R) = \mathbb{D} \times E_{\mathbb{D}}(p_{\{v_n\}}, R)$ , it follows that  $\{u_n\}$  is equivalent to  $\{w_n^{(3)}(p)\}$ .  $\square$

Now we are ready to prove the main result of this section:

**Proposition 7.9.** *The horosphere topology of  $\partial_H\mathbb{D}^2$  induced from  $\widehat{\mathbb{D}^2}$  is trivial.*

*Proof.* We are going to show that the only non empty closed subset of  $\partial_H\mathbb{D}^2$  is  $\partial_H\mathbb{D}^2$ . In order to do that, we prove that the closure in  $\partial_H\mathbb{D}^2$  of the class of any admissible sequence is  $\partial_H\mathbb{D}^2$ .

Since by Proposition 7.8 every admissible sequence is equivalent to  $\{w_n^{(1)}(p_1, p_2)\}$ ,  $\{w_n^{(2)}(p)\}$  or  $\{w_n^{(3)}(p)\}$ , we only need to consider those sequences.

In fact, the following observations yield immediately the result:

- A) for every  $p_1, p_2 \in \partial\mathbb{D}$ , the closure of the point  $[\{w_n^{(1)}(p_1, p_2)\}] \in \partial_H\mathbb{D}^2$  contains the points  $[w_n^{(2)}(p_1)]$  and  $[w_n^{(3)}(p_2)]$ .

- B) for every  $p \in \partial\mathbb{D}$ , the closure of the point  $[\{w_n^{(2)}(p)\}] \in \partial_H\mathbb{D}^2$  contains the points  $[\{w_n^{(1)}(p, q_2)\}]$  for all  $q_2 \in \partial\mathbb{D}$ .
- C) for every  $p \in \partial\mathbb{D}$ , the closure of the point  $[\{w_n^{(3)}(p)\}] \in \partial_H\mathbb{D}^2$  contains the points  $[\{w_n^{(1)}(q_1, p)\}]$  for all  $q_1 \in \partial\mathbb{D}$ .

Indeed, it follows from A), B) and C) that the closure of the point  $[\{w_n^{(1)}(p_1, p_2)\}] \in \partial_H\mathbb{D}^2$  also contains the points  $[\{w_n^{(1)}(q_1, q_2)\}]$  for all  $q_1, q_2 \in \partial\mathbb{D}$ , the closure of the point  $[\{w_n^{(2)}(p_1)\}] \in \partial_H\mathbb{D}^2$  also contains the points  $[\{w_n^{(2)}(q_1)\}]$  for all  $q_1 \in \partial\mathbb{D}$ , and the closure of the point  $[\{w_n^{(3)}(p_2)\}] \in \partial_H\mathbb{D}^2$  also contains the points  $[\{w_n^{(3)}(q_2)\}]$  for all  $q_2 \in \partial\mathbb{D}$ .

We only show A), the proofs of B) and C) being similar. By Lemma 7.7, it follows that for all  $R > 0$ ,

$$E(\{w_n^{(1)}(p_1, p_2)\}, R) \cap E(\{w_n^{(2)}(p_1)\}, R) \neq \emptyset.$$

Hence, if we define the constant sequence  $\{x_m\}$  of elements in  $\partial_H\mathbb{D}^2$  by  $x_m := [\{w_n^{(1)}(p_1, p_2)\}]$  for every  $m \in \mathbb{N}$ , then according to Definition 4.1 the sequence  $\{x_m\}$  converges to  $[\{w_n^{(2)}(p_1)\}] \in \partial_H\mathbb{D}^2$  when  $m$  tends to infinity. Hence  $[\{w_n^{(2)}(p_1)\}]$  belongs to the closure of  $[\{w_n^{(1)}(p_1, p_2)\}]$ . Similarly,  $[\{w_n^{(3)}(p_2)\}]$  belongs to the closure of  $[\{w_n^{(1)}(p_1, p_2)\}]$ .  $\square$

As an application of Theorem 4.7, Theorem 5.10 and Proposition 7.9 we obtain a different proof of the well-known result due to H. Poincaré

**Proposition 7.10.** *There is no biholomorphism between the unit ball and the bidisc in  $\mathbb{C}^2$ .*

*Proof.* A biholomorphism between the unit ball and the bidisc would extend as a homeomorphism between their horosphere boundaries. This is not possible by Propositions 5.10 and 7.9.  $\square$

In fact, one can use the previous arguments to show that the restriction of the horosphere topology of the polydisc  $\mathbb{D}^n \subset \mathbb{C}^N$  to  $\partial_H\mathbb{D}^n$  for every  $n > 1$  is not Hausdorff. Hence, by Theorem 4.7 and Theorem 5.10 one can see also that there exist no biholomorphisms between any bounded strongly pseudoconvex domain and the polydisc in  $\mathbb{C}^N$ ,  $n > 1$ .

In order to compare the Gromov boundary with the horosphere boundary of  $\mathbb{D}^2$ , we will prove now the following

**Proposition 7.11.** *The topology of  $\partial_G^r\mathbb{D}^2$  is not trivial.*

*Proof.* We choose the base point  $(0, 0) \in \mathbb{D}^2$ . Let  $a : (-\infty, +\infty) \rightarrow \mathbb{D}$  be the geodesic for the hyperbolic distance of  $\mathbb{D}$  whose image is the segment  $(-1, 1)$  and satisfying  $\lim_{t \rightarrow -\infty} a(t) = -1$ ,  $\lim_{t \rightarrow +\infty} a(t) = 1$ .

Let  $(1, 0)_G$  be the point of  $\partial_G^r\mathbb{D}^2$  represented by the geodesic ray  $\alpha(t) := (a(t), 0)$ ,  $t \geq 0$ , and let  $(-1, 0)_G$  be the point of  $\partial_G^r\mathbb{D}^2$  represented by the geodesic ray  $\beta(s) := (a(-s), 0)$ ,  $s \geq 0$ .

Since  $K_{\mathbb{D}^2}((z_1, z_2), (w_1, w_2)) = \max_{j=1,2} \{K_{\mathbb{D}}(z_j, w_j)\}$ , it follows that every geodesic ray  $\gamma : [0, +\infty) \rightarrow \mathbb{D}^2$  such that  $\gamma(0) = (0, 0)$  is of the form  $\gamma(t) = (f(t), g(t))$ , where either  $f$  or  $g$  (or both) is a geodesic in  $\mathbb{D}$ . Hence, if  $\gamma(t) = (f_1(t), f_2(t))$  is a geodesic ray in  $\mathbb{D}^2$  equivalent to  $(1, 0)_G$ , it follows that  $f_1 = a$  and  $f_2([0, +\infty))$  is relatively compact in  $\mathbb{D}$ . Thus, every geodesic ray representing  $(1, 0)_G$  is of the form  $(a(t), f(t))$  where  $\sup_{t \in [0, +\infty)} |f(t)| < 1$  and  $K_{\mathbb{D}}(0, f(t)) \leq t$  for every  $t \geq 0$ . Similarly, if  $\gamma$  is a geodesic ray equivalent to  $(-1, 0)_G$  it follows that  $\gamma(s) = (a(-s), g(s))$ , with  $\sup_{t \in [0, +\infty)} |g(t)| < 1$  and  $K_{\mathbb{D}}(0, g(t)) \leq t$  for every  $t \geq 0$ .

Now, let  $\gamma^+(t) := (a(t), f(t))$  be a geodesic ray representing  $(1, 0)_G$  and  $\gamma^-(s) := (a(-s), g(s))$  be a geodesic ray representing  $(-1, 0)_G$ . The Gromov product between  $\gamma^+$  and  $\gamma^-$  with respect to  $(0, 0)$  is:

$$\begin{aligned} 2(\gamma^+(t), \gamma^-(s))_{(0,0)} &= 2((a(t), f(t)), (a(-s), g(s)))_{(0,0)} \\ &= K_{\mathbb{D}^2}((a(t), f(t)), (0, 0)) + K_{\mathbb{D}^2}((a(-s), g(s)), (0, 0)) \\ &\quad - K_{\mathbb{D}^2}((a(t), f(t)), (a(-s), g(s))) \\ &= t + s - \max\{K_{\mathbb{D}}(a(t), a(-s)), K_{\mathbb{D}}(f(t), g(s))\} \\ &= t + s - \max\{s + t, K_{\mathbb{D}}(f(t), g(s))\} \leq 0. \end{aligned}$$

Hence, the Gromov product is always 0.

Therefore, for any couple of geodesic rays  $\gamma^+$  representing  $(1, 0)_G$  and  $\gamma^-$  representing  $(-1, 0)_G$  it holds

$$\liminf_{t,s \rightarrow +\infty} (\gamma^+(t), \gamma^-(s)) = 0.$$

This implies that for every  $r > 0$  the point  $(-1, 0)_G$  does not belong to the open set  $V((1, 0)_G, r)$ . Hence, for every  $r > 0$ , the set  $\partial_G^r \mathbb{D}^2 \setminus V((1, 0)_G, r)$  is a non empty closed set which does not contain  $(1, 0)_G$ . In particular, the Gromov topology on  $\partial_G^r \mathbb{D}^2$  is not trivial.  $\square$

Proposition 7.6 follows at once from Proposition 7.9 and Proposition 7.11.

## 8. BOUNDARY BEHAVIOR

In this section we apply the results developed so far to study boundary behavior of univalent mappings defined on strongly pseudoconvex smooth domains. If  $F : D \rightarrow \mathbb{C}^N$  is a holomorphic map and  $p \in \partial D$ , we denote by  $\Gamma(F; p)$  the cluster set of  $F$  at  $p$ , that is,

$$\Gamma(F; p) := \{q \in \mathbb{C}^N : \exists \{w_n\} \subset D, F(w_n) \rightarrow q\}.$$

We start by the following result:

**Theorem 8.1.** *Let  $D \subset \mathbb{C}^N$  be a bounded strongly pseudoconvex domain with  $C^3$  boundary. Let  $F : D \rightarrow \Omega$  be a biholomorphism. Let  $p \in \partial D$ . Then there exists  $\underline{x} \in \partial_H \Omega$  such that*

$$\Gamma(F; p) = I_{\Omega}^H(\underline{x}).$$

*In particular, if there exists  $q \in \partial \Omega$  such that  $I_{\Omega}^H(\underline{x}) = \{q\}$ , then  $\lim_{z \rightarrow p} F(z) = q$ .*

*Proof.* By Theorem 4.7,  $F$  defines a homeomorphism  $\hat{F} : \hat{D} \rightarrow \hat{\Omega}$ . By Theorem 5.10, there is a homeomorphism  $\Theta : \hat{D} \rightarrow \overline{D}$  such that  $\Theta(z) = z$  for  $z \in D$ . Therefore, a sequence  $\{z_n\} \subset D$  converges to  $p$  if and only if  $\{\Theta^{-1}(z_n)\}$  converges to  $\Theta^{-1}(p) \in \partial_H D$ . Let  $\underline{x} := \hat{F}(\Theta^{-1}(p))$ . Hence, the sequence  $\{F(z_n) = \hat{F}(\Theta^{-1}(z_n))\}$  converges to  $\underline{x}$  if and only if  $\{z_n\}$  converges to  $p$ . From this the result follows.  $\square$

A first application is the following well known result (see, [17]):

**Corollary 8.2.** *Let  $D \subset \mathbb{C}^N$  be a bounded strongly pseudoconvex domain with  $C^3$  boundary. Let  $F : D \rightarrow \Omega$  be a biholomorphism. If  $\Omega$  is a bounded strongly pseudoconvex domain with  $C^3$  boundary then  $F$  extends to a homeomorphism from  $\overline{D}$  to  $\overline{\Omega}$ .*

*Proof.* It is enough to apply Theorem 8.1 and Proposition 5.11 to both  $F$  and  $F^{-1}$ .  $\square$

Another application of our theory gives a positive answer to Conjecture 3.1.(a) in [31], in fact, proving not only continuous extension, but extension as homeomorphism:

**Corollary 8.3.** *Let  $D \subset \mathbb{C}^N$  be a bounded strongly convex domain with  $C^3$  boundary. Let  $F : D \rightarrow \Omega$  be a biholomorphism. If  $\Omega$  is a bounded convex domain, then  $F$  extends as a homeomorphism from  $\overline{D}$  to  $\overline{\Omega}$ .*

*Proof.* By Theorem 6.20 and Theorem 8.1, for every  $p \in \partial D$  the limit  $F(p) := \lim_{z \rightarrow p} F(z)$  exists. In order to see that the map  $F : \overline{D} \rightarrow \overline{\Omega}$  is continuous, we have only to show that if  $\{p_j\} \subset \partial D$  is a sequence converging to  $p \in \partial D$ , then  $F(p_j) \rightarrow F(p)$ . Indeed, by Proposition 5.11, there exist  $\underline{x}_j, \underline{x} \in \partial_H D$  such that  $I_D^H(\underline{x}_j) = \{p_j\}$ ,  $I_D^H(\underline{x}) = \{p\}$  and  $\{\underline{x}_j\}$  converges to  $\underline{x}$  in the horosphere topology of  $D$ . Hence  $\{\hat{F}(\underline{x}_j)\}$  converges to  $\{\hat{F}(\underline{x})\}$  in the horosphere topology of  $\Omega$ . Moreover,  $\Pi_\Omega^H(\hat{F}(\underline{x}_j)) = I_\Omega^H(\hat{F}(\underline{x}_j)) = \{F(p_j)\}$  and similarly  $\Pi_\Omega^H(\hat{F}(\underline{x})) = I_\Omega^H(\hat{F}(\underline{x})) = \{F(p)\}$ . Hence,  $F(p_j) \rightarrow F(p)$  by Corollary 6.9.

Therefore,  $F : \overline{D} \rightarrow \overline{\Omega}$  is continuous. Since  $\overline{D}$  is compact, in order to prove that  $F$  is a homeomorphism, we only need to prove that it is injective.

We argue by contradiction. Assume  $p_0, p_1 \in \partial D$  and  $q := F(p_0) = F(p_1) \in \partial \Omega$ . Let  $\{u_n\} \subset D$  be a sequence converging to  $p_0$  and  $\{v_n\} \subset D$  be a sequence converging to  $p_1$ . Then  $\{u_n\}$  and  $\{v_n\}$  are admissible by Proposition 5.7 and not equivalent by Proposition 5.9. Let  $x \in D$  and  $R > 0$  be such that  $V := \overline{E_x^D(\{u_n\}, R) \cap E_x^D(\{v_n\}, R)} \neq \emptyset$ . Note that  $V$  is open and relatively compact in  $D$  because  $\overline{E_x^D(\{u_n\}, R) \cap \partial D} = \{p_0\}$  and  $\overline{E_x^D(\{v_n\}, R) \cap \partial D} = \{p_1\}$ .

Since  $F$  is a biholomorphism it maps horospheres onto horospheres. Moreover,  $F(E_x^D(\{u_n\}, R)) = E_{F(x)}^\Omega(\{F(u_n)\}, R)$  and  $F(E_x^D(\{v_n\}, R)) = E_{F(x)}^\Omega(\{F(v_n)\}, R)$  are convex by Proposition 6.1. Hence,

$$F(V) = F(E_x^D(\{u_n\}, R) \cap E_x^D(\{v_n\}, R)) = E_{F(x)}^\Omega(\{F(u_n)\}, R) \cap E_{F(x)}^\Omega(\{F(v_n)\}, R)$$

is open, convex and relatively compact in  $\Omega$ .

Now,  $q \in \overline{E_{F(x)}^\Omega(\{F(u_n)\}, R) \cap E_{F(x)}^\Omega(\{F(v_n)\}, R)}$ . Let  $z_0 \in F(V)$ . Since the two horospheres are both open, convex, and  $z_0$  is contained in both horospheres, the real segment  $\gamma := \{tz_0 + (1-t)q : t \in (0, 1]\}$  as well is contained in both horospheres. That is,  $\gamma \subset F(V)$ . But then  $F(V)$  is not relatively compact in  $\Omega$ , a contradiction.  $\square$

With a similar argument, using Proposition 6.13 instead of Theorem 6.20, we have

**Corollary 8.4.** *Let  $D \subset \mathbb{C}^N$  be a bounded strongly pseudoconvex domain with  $C^3$  boundary. Let  $F : D \rightarrow \Omega$  be a biholomorphism. If  $\Omega$  is a strictly  $\mathbb{C}$ -linearly bounded convex domain, then  $F$  extends as a homeomorphism from  $\overline{D}$  to  $\overline{\Omega}$ .*

Now we consider non-tangential limits. In fact, in our theory, the right notion to consider is that of *E-limits*. Let  $D \subset \mathbb{C}^N$  is a bounded strongly pseudoconvex domain with  $C^3$  boundary. If  $p \in \partial D$ , by Proposition 5.11, there exists  $\underline{x}_p \in \partial_H D$  such that  $I_D^H(\underline{x}_p) = \{p\}$ . Given a map  $f : \mathbb{D} \rightarrow \mathbb{C}^N$ , we denote by  $\Gamma_E(f; p)$  the cluster set of  $f$  at  $p$  along sequences  $E$ -converging to  $\underline{x}_p$  (see Definition 4.10), namely,

$$\Gamma_E(f; p) = \{q \in \mathbb{C}^N : \exists \{z_n\} \subset D : E - \lim_{n \rightarrow \infty} z_n = \underline{x}_p, f(z_n) \rightarrow q\}.$$

A slight modification of the proof of Theorem 8.1, taking into account that  $F$  maps sequences  $E$ -converging to  $\underline{y}$  to sequences  $E$ -converging to  $\hat{F}(\underline{y})$  and Lemma 4.12, gives the following:

**Theorem 8.5.** *Let  $D \subset \mathbb{C}^N$  be a bounded strongly pseudoconvex domain with  $C^3$  boundary. Let  $F : D \rightarrow \Omega$  be a biholomorphism,  $x \in \Omega$ . Let  $p \in \partial D$ . Then there exists  $\underline{x} \in \partial_H \Omega$  such that*

$$\Gamma_E(f; p) = II_\Omega^H(\underline{x}) = \bigcap_{R>0} \overline{E_x(\{u_n\}, R)}^{\mathbb{C}P^N},$$

where  $\{u_n\} \subset D$  is any admissible sequence representing  $\underline{x}$ .

Finally, let  $D \subset \mathbb{C}^N$  is a bounded strongly pseudoconvex domain with  $C^3$  boundary and  $p \in \partial D$ . If  $f : D \rightarrow \mathbb{C}^N$  is a map, we denote by  $\Gamma_{NT}(f; p)$  the cluster set of  $f$  along sequences converging to  $p$

non-tangentially. By Proposition 5.7,

$$\Gamma_{NT}(f; p) \subseteq \Gamma_E(f; p) \subseteq \Gamma(f; p).$$

In particular, by Theorem 8.5 we have:

**Corollary 8.6.** *Let  $D \subset \mathbb{C}^N$  be a bounded strongly pseudoconvex domain with  $C^3$  boundary. Let  $F : D \rightarrow \Omega$  be a biholomorphism and assume that for every  $\underline{x} \in \partial_H \Omega$  the horosphere principal part  $\Pi_{\Omega}^H(\underline{x})$  consists of one point. Then for every  $p \in \partial D$  the non-tangential limit  $\angle \lim_{z \rightarrow p} F(z)$  exists.*

As a spin off result of our work, we prove the following Wolff-Denjoy theorem, which gives a (partial) positive answer to a conjecture in [4] (see [4, Rmk. 3.3]). Let  $D \subset \mathbb{C}^N$  be a bounded domain. Let  $f : D \rightarrow D$  be holomorphic. A point  $q \in \overline{D}$  belongs to the target set  $T(f)$  of  $f$  if there exist a sequence  $\{k_m\} \subset \mathbb{N}$  converging to  $\infty$  and  $z \in D$  such that  $\lim_{m \rightarrow \infty} f^{k_m}(z) = q$ .

**Proposition 8.7.** *Let  $D \subset \mathbb{C}^N$  be a bounded convex domain. Assume that either  $D$  is biholomorphic to a strongly convex domain with  $C^3$  boundary or  $D$  is  $\mathbb{C}$ -strictly linearly convex and biholomorphic to a bounded strongly pseudoconvex domain with  $C^3$  boundary. Let  $f : D \rightarrow D$  be holomorphic without fixed points in  $D$ . Then there exists  $p \in \partial D$  such that  $T(f) = \{p\}$ .*

*Proof.* Since  $f$  has no fixed point in  $D$  then  $\{f^k\}$  is compactly divergent (see [1, 2]). By [4, Lemma 3.10] there exists a Busemann admissible sequence  $\{u_n\}$  such that for every  $R > 0$ ,

$$(8.1) \quad f(E_x(\{u_n\}, R)) \subset E_x(\{u_n\}, R).$$

Hence the result follows from either Theorem 6.20 or Proposition 6.13.  $\square$

**Remark 8.8.** *As we already pointed out, there exist convex domains biholomorphic to the unit ball which are not  $\mathbb{C}$ -strictly linearly convex, but for which the Denjoy-Wolff theorem holds by Proposition 8.7. We conjecture that in fact the result in Proposition 8.7 holds for every bounded convex domain  $D \subset \mathbb{C}^N$  whose boundary does not contain non-constant analytic discs.*

## REFERENCES

- [1] M. Abate, *Horospheres and iterates of holomorphic maps*. Math. Z. 198 (1988), 225-238.
- [2] M. Abate, *Iteration theory, compactly divergent sequences and commuting holomorphic maps*. Ann. Scuola Norm. Sup. Pisa 18 (1991), 167-191.
- [3] M. Abate, *Iteration theory of holomorphic maps on taut manifolds*. Research and Lecture Notes in Mathematics. Complex Analysis and Geometry, Mediterranean Press, Rende, 1989.
- [4] M. Abate, J. Raissy, *Wolff-Denjoy theorems in nonsmooth convex domains*. Ann. Mat. Pura Appl. (4) 193 (2014), 1503-1518.
- [5] T. Adamowicz, A. Björn, J. Björn, N. Shanmugalingam, *Prime ends for domains in metric spaces*. Adv. Math. 238 (2013), 459-505.
- [6] W. Balmann, M. Gromov, V. Schroeder, *Manifolds of Nonpositive Curvature*. Birkhäuser, Basel (1985).
- [7] Z. M. Balogh, M. Bonk, *Gromov hyperbolicity and the Kobayashi metric on strictly pseudoconvex domains*. Comment. Math. Helv. 75 (2000), 504-533.
- [8] G. Bharali, A. Zimmer, *Goldilocks domains, a weak notion of visibility, and applications*. ArXiv:1602.01742v1
- [9] M. Budzyńska, *The Denjoy-Wolff theorem in  $\mathbb{C}^N$* . Nonlinear Analysis 75 (2012), 22-29.
- [10] M. Budzyńska, T. Kuczumow, S. Reich, *Theorems of Denjoy-Wolff type*. Ann. Mat. Pura Appl. (4) 192 (2013), 621-648.
- [11] F. Bracci, G. Patrizio, *Monge-Ampère foliations with singularities at the boundary of strongly convex domains*. Math. Ann. 332 (2005), 499-522.
- [12] F. Bracci, G. Patrizio, S. Trapani, *The pluricomplex Poisson kernel for strongly convex domains*. Trans. Am. Math. Soc. 361 (2009), 979-1005.
- [13] F. Bracci, A. Saracco, *Hyperbolicity in unbounded convex domains*. Forum Math. 21 (2009), 815-826.
- [14] E.F. Collingwood, A.J. Lohwater, *The theory of cluster sets*. Cambridge Tracts in Mathematics and Mathematical Physics, No. 56 Cambridge Univ. Press, Cambridge, 1966.
- [15] P. Eberlein, B. O'Neill, *Visibility manifolds*. Pac. J. Math. 46 (1973), 45-109.

- [16] D. B. A. Epstein, *Prime ends*. Proc. London Math. Soc. (3) 42 (1981), 385-414.
- [17] C. Fefferman, *The Bergman kernel and biholomorphic mappings of pseudoconvex domains*. Invent. Math. 26 (1974), 1-65.
- [18] F. Forstnerič, J.-P. Rosay, *Localization of the Kobayashi metric and the boundary continuity of proper holomorphic mappings*. Math. Ann. 279 (1987), 239-252.
- [19] C. Frosini, *Busemann functions and Julia-Wolff-Carathéodory theorem on polydiscs*. Adv. Geom. 10 (2010), 435-463.
- [20] D. Gaier, Ch. Pommerenke, *On the boundary behavior of conformal maps*. Michigan Math. J. 14 (1967), 79-82.
- [21] H. Gaussier, H. Seshadri, *On the Gromov hyperbolicity of convex domains in  $\mathbb{C}^N$* . ArXiv:1312.0368.
- [22] H. Gaussier, H. Seshadri, *Totally geodesic discs in strongly convex domains*. Math. Z. 274 (2013), 185-197.
- [23] E. Ghys, P. de La Harpe, *Sur les groupes hyperboliques d'après Mikhael Gromov*. Progress in Mathematics **83**, Birkhauser.
- [24] M. Gromov, *Hyperbolic groups. Essays in group theory*. Math. Sci. Res. Inst. Publ., 8, Springer, New York (1987), 75-263.
- [25] H. Hamada, G. Kohr, *On some classes of bounded univalent mappings in several complex variables*. Manuscripta Math. 131 (2010), 487-502.
- [26] J. Kapeluszny, T. Kuczumow, S. Reich, *The Denjoy-Wolff theorem in the open unit ball of a strictly convex Banach space*. Adv. Math. 143 (1999), 111-123.
- [27] L. Lempert, *La métrique de Kobayashi et la représentation des domaines sur la boule*. Bull. Soc. Math. Fr. 109 (1981), 427-474.
- [28] L. Lempert, *Holomorphic retracts and intrinsic metrics in convex domains*. Analysis Math. 8 (1982), 257-261.
- [29] L. Lempert, *Intrinsic distances and holomorphic retracts*. Complex Analysis and Applications, Proc. Int. Conf., Varna/Bulg. 1981 (1984), 341-364.
- [30] P. R. Mercer, *Complex geodesics and iterates of holomorphic maps on convex domains in  $\mathbb{C}^N$* . Trans. Amer. Math. Soc. 338 (1993), 201-211.
- [31] J. R. Muir, Jr., T. J. Suffridge, *Unbounded convex mappings of the ball in  $\mathbb{C}^N$* . Proc. Amer. Math. Soc. 129 (2001), 3389-3393.
- [32] J. R. Muir, Jr., T. J. Suffridge, *A generalization of half-plane mappings to the ball in  $\mathbb{C}^N$* . Trans. Amer. Math. Soc. 359 (2007), 1485-1498.
- [33] N. Nikolov, *Comparison of invariant functions on strongly pseudoconvex domains* J. Math. Anal. Appl., 421, (2015), 180-185.
- [34] Ch. Pommerenke, *Boundary behaviour of conformal mappings*. Springer-Verlag (1992).
- [35] S. Reich, D., Shoikhet, *Nonlinear Semigroups, Fixed Points, and Geometry of Domains in Banach Spaces*. Imperial College Press, London (2005).
- [36] J. B. Twomey, *Tangential Limits of Starlike Univalent Functions*. Proc. Amer. Math. Soc., 97 (1986), 49-54.
- [37] J. Väisälä, *Lectures on n-Dimensional Quasiconformal Mappings*. Lecture Notes in Math. 229, Springer-Verlag (1971).
- [38] A. M. Zimmer, *Gromov hyperbolicity and the Kobayashi metric on convex domains of finite type*. To appear in Math. Ann. ArXiv:1405.2858.
- [39] A. M. Zimmer, *Characterizing polynomial domains by their automorphism group*. ArXiv:1506.07852.

FILIPPO BRACCI

DIPARTIMENTO DI MATEMATICA

UNIVERSITÀ DI ROMA "TOR VERGATA"

VIA DELLA RICERCA SCIENTIFICA, 1

00133 ROMA

ITALY

E-mail address: fbracci@mat.uniroma2.it herve.haussier@univ-grenoble-alpes.fr

HERVÉ GAUSSIER

UNIV. GRENOBLE ALPES, IF, F-38000 GRENOBLE, FRANCE

CNRS, IF, F-38000 GRENOBLE, FRANCE