



HAL
open science

Commutative and non-commutative bialgebras of quasi-posets and applications to Ehrhart polynomials

Loïc Foissy

► **To cite this version:**

Loïc Foissy. Commutative and non-commutative bialgebras of quasi-posets and applications to Ehrhart polynomials. *Advances in Pure and Applied Mathematics*, 2019, 10 (1), pp.27-63. 10.1515/apam-2016-0051 . hal-01321986v2

HAL Id: hal-01321986

<https://hal.science/hal-01321986v2>

Submitted on 14 Nov 2016

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Commutative and non-commutative bialgebras of quasi-posets and applications to Ehrhart polynomials

Loïc Foissy

*Fédération de Recherche Mathématique du Nord Pas de Calais FR 2956
Laboratoire de Mathématiques Pures et Appliquées Joseph Liouville
Université du Littoral Côte d'Opale-Centre Universitaire de la Mi-Voix
50, rue Ferdinand Buisson, CS 80699, 62228 Calais Cedex, France*

Email: foissy@lmpa.univ-littoral.fr

Abstract

To any poset or quasi-poset is attached a lattice polytope, whose Ehrhart polynomial we study from a Hopf-algebraic point of view. We use for this two interacting bialgebras on quasi-posets. The Ehrhart polynomial defines a Hopf algebra morphism taking its values in $\mathbb{Q}[X]$; we deduce from the interacting bialgebras an algebraic proof of the duality principle, a generalization and a new proof of a result on B-series due to Whright and Zhao, using a monoid of characters on quasi-posets, and a generalization of Faulhaber's formula.

We also give non-commutative versions of these results: polynomials are replaced by packed words. We obtain in particular a non-commutative duality principle.

Keywords. Ehrhart polynomials; Quasi-posets; Characters monoids; Interacting bialgebras

AMS classification. 16T30; 06A11

Contents

1	Bialgebras in cointeraction	5
1.1	Definition	5
1.2	Monoids actions	6
1.3	Polynomial morphisms	8
2	Examples from quasi-posets	13
2.1	Definition	13
2.2	First coproduct	14
2.3	Second coproduct	16
2.4	Cointeractions	19
3	Ehrhart polynomials	21
3.1	Definition	22
3.2	Recursive computation of ehr and ehr^{str}	24
3.3	Characterization of quasi-posets by packed words	26
3.4	Link with linear extensions	27

4	Characters associated to ehr and ehr^{str}	28
4.1	The character λ	28
4.2	The character α^{str}	31
4.3	The character α and the duality principle	32
4.4	A link with Bernoulli numbers	34
5	Noncommutative version	35
5.1	Reminders on packed words	35
5.2	Hopf algebra morphisms in WQSym	35
5.3	The non-commutative duality principle	38
5.4	Compatibility with the other product and coproduct	40
5.5	Restriction to posets	44

Introduction

Let P be a lattice polytope, that is to say that all its vertices are in \mathbb{Z}^n . The Ehrhart polynomial $ehr_P^{cl}(X)$ is the unique polynomial such that, for all $k \geq 1$, $ehr_P^{cl}(k)$ is the number of points in $\mathbb{Z}^n \cap kP$, where kP is the image of P by the homothety of center 0 and ratio k . For example, if S is the square $[0, 1]^n$ and T is the triangle of vertices $(0, 0)$, $(1, 0)$ and $(1, 1)$:


$$ehr_S^{cl}(X) = (X + 1)^2, \quad ehr_T^{cl}(X) = \frac{(X + 1)(X + 2)}{2}.$$

These polynomials satisfy the reciprocity principle: for all $k \geq 1$, $(-1)^{dim(P)} ehr^{cl}(-k)$ is the number of points of $\mathbb{Z}^n \cap k\dot{P}$, where \dot{P} is the interior of P . For example:

$$ehr_S^{cl}(-X) = (X - 1)^2, \quad ehr_T^{cl}(-X) = \frac{(X - 1)(X - 2)}{2}.$$

We refer to [2] for general results on Ehrhart polynomials.

It turns out that these polynomials appear in the theory of B-series (B for Butcher [4]), as explained in [3, 6]. We now consider rooted trees:



If t is a rooted tree, we orient its edges from the root to the leaves. If i, j are two vertices of t , we shall write $i \xrightarrow{t} j$ if there is an edge from i to j in t .

To any rooted tree t , whose vertices are indexed by $1 \dots n$, we associate a lattice polytope $pol(t)$ in a following way:

$$pol(t) = \left\{ (x_1, \dots, x_n) \in [0, 1]^n \mid \forall 1 \leq i, j \leq n, (i \xrightarrow{t} j) \implies (x_i \leq x_j) \right\}$$

For example, if $t = \mathbf{1}$, indexed as $\mathbf{1}_1^1$, then $pol(t) = T$.

We can consider the Ehrhart polynomial $ehr_{pol(t)}^{cl}(X)$, which we shall simply denote by $ehr_t^{cl}(X)$: for all $k \geq 1$,

$$ehr_t^{cl}(k) = \# \left\{ (x_1, \dots, x_n) \in \{0, \dots, k\}^n \mid \forall 1 \leq i, j \leq n, (i \xrightarrow{t} j) \implies (x_i \leq x_j) \right\}.$$

Note that ehr_t^{cl} does not depend on the indexation of the vertices of t . By the duality principle:

$$(-1)^n ehr_t^{cl}(-k) = \# \left\{ (x_1, \dots, x_n) \in \{1, \dots, k - 1\}^n \mid \forall 1 \leq i, j \leq n, (i \xrightarrow{t} j) \implies (x_i < x_j) \right\}.$$

A B-series is a formal series indexed by rooted trees, of the form:

$$\sum_t a_t \frac{t}{\text{aut}(t)} = a_{\bullet} + a_{\mathbf{1}} \mathbf{1} + a_{\mathbf{V}} \frac{\mathbf{V}}{2} + a_{\mathbf{1}\mathbf{1}} \mathbf{1}\mathbf{1} + \dots,$$

where $\text{aut}(t)$ is the number of automorphisms of t . The following B-series is of special importance in numerical analysis:

$$E = \sum_t \frac{1}{t!} \frac{t}{\text{aut}(t)} = \bullet + \frac{1}{2} a_{\mathbf{1}} \mathbf{1} + \frac{1}{3} \frac{\mathbf{V}}{2} + \frac{1}{6} \mathbf{1}\mathbf{1} + \dots,$$

where $t!$ is the tree factorial (see definition 33). This series is the formal solution of an ordinary differential equation, describing the flow of a vector field. The set of B-series is given a group structure by a substitution operation, which is dually represented by the contraction-extraction coproduct defined in [5]. The inverse of E is called the backward error analysis:

$$E^{-1} = \sum_t \lambda_t \frac{t}{\text{aut}(t)!}.$$

Wright and Zhao [19] proved that these coefficients λ_t are related to Ehrhart polynomials:

$$\lambda_t = (-1)^{|t|} \frac{d \text{ehr}_t^{\text{cl}}(X)}{dX} \Big|_{X=-1}.$$

We shall in this text study Ehrhart polynomial attached to quasi-posets in a combinatorial Hopf-algebraic way. A quasi-poset P is a pair (A, \leq_P) , where A is a finite set and \leq_P is a reflexive and transitive relation on A . The isoclasses of quasi-posets are represented by their Hasse graphs:

$$\mathbf{1}; \quad \bullet; \quad \dots, \mathbf{1}, \dots_2; \quad \dots, \dots_1, \dots_2, \mathbf{V}, \mathbf{\Lambda}, \mathbf{1}\mathbf{1}, \mathbf{1}^2, \mathbf{1}_2, \dots_3; \dots$$

In particular, rooted trees can be seen as quasi-posets. For any quasi-poset $P = (\{1, \dots, n\}, \leq_P)$, the polytope associated to P is:

$$\text{pol}(P) = \{(x_1, \dots, x_n) \in [0, 1]^n \mid \forall 1 \leq i, j \leq n, (i \leq_P j) \implies (x_i \leq x_j)\}.$$

We put $\text{ehr}_P(X) = \text{ehr}_{\text{pol}(P)}^{\text{cl}}(X-1)$; note the translation by -1 , which will give us objects more suitable to our purpose. In other words, for all $k \geq 1$:

$$\text{ehr}_P(k) = \#\{(x_1, \dots, x_n) \in \{1, \dots, k\}^n \mid \forall 1 \leq i, j \leq n, (i \leq_P j) \implies (x_i \leq x_j)\}.$$

We also define a polynomial $\text{ehr}_P^{\text{str}}(X)$ such that for all $k \geq 1$:

$$\text{ehr}_P^{\text{str}}(k) = \#\{(x_1, \dots, x_n) \in \{1, \dots, k\}^n \mid \forall 1 \leq i, j \leq n, (i \leq_P j \text{ and not } j \leq_P i) \implies (x_i < x_j)\}.$$

See definition 21 and proposition 22 for more details. These polynomials can be inductively computed, with the help of the minimal elements of P (proposition 25).

We shall consider two products m and \downarrow , and two coproducts Δ and δ on the space $\mathcal{H}_{\mathbf{qp}}$ generated by isoclasses of quasi-posets. The coproduct Δ , defined in [10, 11] by restriction to open and closed sets of the topologies associated to quasi-posets, makes $(\mathcal{H}_{\mathbf{qp}}, m, \Delta)$ a graded, connected Hopf algebra, and $(\mathcal{H}_{\mathbf{qp}}, \downarrow, \Delta)$ an infinitesimal bialgebra; the coproduct δ , defined in [8] by an extraction-contraction operation, makes $(\mathcal{H}_{\mathbf{qp}}, m, \delta)$ a bialgebra. Moreover, δ is also a right coaction of $(\mathcal{H}_{\mathbf{qp}}, m, \delta)$ over $(\mathcal{H}_{\mathbf{qp}}, m, \Delta)$, and $(\mathcal{H}_{\mathbf{qp}}, m, \Delta)$ becomes a Hopf algebra in the category of $(\mathcal{H}_{\mathbf{qp}}, m, \delta)$ -comodules, which we summarize telling that $(\mathcal{H}_{\mathbf{qp}}, m, \Delta)$ and $(\mathcal{H}_{\mathbf{qp}}, m, \delta)$

are two bialgebras in cointeraction (definition 1). For example, the bialgebras $(\mathbb{K}[X], m, \Delta)$ and $(\mathbb{K}[X], m, \delta)$ where m is the usual product of $\mathbb{K}[X]$ and Δ, δ are the coproducts defined by

$$\Delta(X) = X \otimes 1 + 1 \otimes X, \quad \delta(X) = X \otimes X,$$

are two cointeracting bialgebras.

Ehrhart polynomials $ehr_P(X)$ and $ehr_P^{str}(X)$ can now be seen as maps from $\mathcal{H}_{\mathbf{qp}}$ to $\mathbb{K}[X]$, and both are Hopf algebra morphisms from $(\mathcal{H}_{\mathbf{qp}}, m, \Delta)$ to $(\mathbb{K}[X], m, \Delta)$ (theorem 23); we shall prove in theorem 37 that ehr^{str} is the unique morphism from $\mathcal{H}_{\mathbf{qp}}$ to $\mathbb{K}[X]$ compatible with both bialgebra structures on $\mathcal{H}_{\mathbf{qp}}$ and $\mathbb{K}[X]$. Using the cointeraction between the two bialgebra structures on $\mathcal{H}_{\mathbf{qp}}$, we show that the monoid $M_{\mathbf{qp}}$ of characters of $(\mathcal{H}_{\mathbf{qp}}, m, \delta)$ acts on the set $E_{\mathcal{H}_{\mathbf{qp}} \rightarrow \mathbb{K}[X]}$ of Hopf algebra morphisms from $(\mathcal{H}_{\mathbf{qp}}, m, \Delta)$ to $\mathbb{K}[X]$ (lemma 5). Moreover, there exists a particular homogeneous morphism $\phi_0 \in E_{\mathcal{H}_{\mathbf{qp}} \rightarrow \mathbb{K}[X]}$ such that for all quasi-poset P :

$$\phi_0(P) = \lambda_P X^{cl(P)} = \frac{\mu_P}{cl(P)!} X^{cl(P)},$$

where μ_P is the number of heap-orderings of P and $cl(P)$ is the number of equivalence classes of the equivalence associated to the quasi-order of P (proposition 32). This formula simplifies if P is a rooted tree: in this case,

$$\phi_0(P) = \frac{1}{P!} X^{|P|}.$$

We prove that there exist characters α and α^{str} in $M_{\mathbf{qp}}$, such that for any quasi-poset P :

$$ehr_P(X) = \sum_{\sim \triangleleft P} \frac{\mu_{P/\sim}}{cl(\sim)!} \alpha_{P|\sim} X^{cl(\sim)}, \quad ehr_P^{str}(X) = \sum_{\sim \triangleleft P} \frac{\mu_{P/\sim}}{cl(\sim)!} \alpha_{P|\sim}^{str} X^{cl(\sim)},$$

where the sum is over a certain family of equivalence relations \sim on the set of vertices of V , $P|\sim$ is a restriction operation and P/\sim is a contraction operation. Applied to corollas, this gives Faulhaber's formula. We prove that α^{str} is the inverse of the character λ associated to ϕ_0 (theorem 37), which is a generalization, as well as a Hopf-algebraic proof, of Wright and Zhao's result. We also give an algebraic proof of the duality principle (theorem 38), and we define a Hopf algebra automorphism $\theta : (\mathcal{H}_{\mathbf{qp}}, m, \Delta) \rightarrow (\mathcal{H}_{\mathbf{qp}}, m, \Delta)$ with the help of the cointeraction of the two bialgebra structures on $\mathcal{H}_{\mathbf{qp}}$, satisfying $ehr^{str} \circ \theta = ehr$ (proposition 39).

We propose non-commutative versions of these results in the last section of the paper. Here, (isoclasses of) quasi-posets are replaced by quasi-posets on sets $[n] = \{1, \dots, n\}$, making a Hopf algebra $\mathcal{H}_{\mathbf{QP}}$, in cointeraction with $(\mathcal{H}_{\mathbf{qp}}, m, \delta)$, and $\mathbb{K}[X]$ is replaced by the Hopf algebra of packed words **WQSym** [15]. We define two surjective Hopf algebra morphisms EHR and EHR^{str} from $\mathcal{H}_{\mathbf{QP}}$ to **WQSym** (proposition 41), generalizing ehr and ehr^{str} . The automorphism θ is generalized as a Hopf algebra automorphism $\Theta : \mathcal{H}_{\mathbf{QP}} \rightarrow \mathcal{H}_{\mathbf{QP}}$, such that $EHR^{str} \circ \Theta = EHR$ (proposition 42), and we formulate a non-commutative duality principle

Examples of bialgebras in interaction can be found in [5] (for rooted trees) and in [14] (for various families of graphs). Another example is given by the algebra $\mathbb{K}[X]$, with its usual product m , and the two coproducts defined by:

$$\Delta(X) = X \otimes 1 + 1 \otimes X, \quad \delta(X) = X \otimes X.$$

The bialgebras $(\mathbb{K}[X], m, \Delta)$ and $(\mathbb{K}[X], m, \delta)$ are in cointeraction, via the coaction $\rho = \delta$. Identifying $\mathbb{K}[X] \otimes \mathbb{K}[X]$ and $\mathbb{K}[X, Y]$:

$$\Delta(P)(X, Y) = P(X + Y), \quad \delta(P)(X, Y) = P(XY).$$

Remark. If A and B are in cointeraction, the coaction of B on A is an algebra morphism.

Proposition 2 *Let A and B be two bialgebras in cointeraction. We assume that A is a Hopf algebra, with antipode S . Then S is a morphism of B -comodules, that is to say:*

$$\rho \circ S = (S \otimes Id) \circ \rho$$

Proof. We work in the space $End_{\mathbb{K}}(A, A \otimes B)$. As $A \otimes B$ is an algebra and A is a coalgebra, it is an algebra for the convolution product \circledast :

$$\forall f, g \in End_{\mathbb{K}}(A, A \otimes B), f \circledast g = m_{A \otimes B} \circ (f \otimes g) \circ \Delta_A.$$

Its unit is:

$$\eta : \begin{cases} A & \longrightarrow & A \otimes B \\ a & \longrightarrow & \varepsilon(a)1_A \otimes 1_B. \end{cases}$$

We consider three elements in this algebra, respectively ρ , $F_1 = (S \otimes Id) \circ \rho$ and $F_2 = \rho \circ S$. Firstly:

$$\begin{aligned} (F_1 \circledast \rho)(a) &= S((a^{(1)})_1)(a^{(2)})_1 \otimes (a^{(1)})_0(a^{(2)})_0 \\ &= S((a_1)^{(1)})(a_1)^{(2)} \otimes a_0 \\ &= \varepsilon_A(a_1)1_A \otimes a_0 \\ &= \varepsilon_A(a)1_A \otimes 1_B \\ &= \eta(a). \end{aligned}$$

Secondly:

$$\begin{aligned} (\rho \circledast F_2)(a) &= (a^{(1)})_1 S(a^{(2)})_1 \otimes (a^{(1)})_0 (S(a^{(2)}))_0 \\ &= \varepsilon_A(a)(1_A)_1 \otimes (1_A)_0 \\ &= \varepsilon_A(a)1_A \otimes 1_B \\ &= \eta(a). \end{aligned}$$

We obtain that $F_1 \circledast \rho = \rho \circledast F_2 = \eta$, so $F_1 = F_1 \circledast \eta = F_1 \circledast \rho \circledast F_2 = \eta \circledast F_2 = F_2$. \square

1.2 Monoids actions

Proposition 3 *Let A and B be two bialgebras in cointeraction, through the coaction ρ . We denote by M_A and M_B the monoids of characters of respectively A and B . Then B acts on A by monoid endomorphisms, via the map:*

$$\leftarrow : \begin{cases} M_A \times M_B & \longrightarrow & M_A \\ (\phi, \lambda) & \longrightarrow & \phi \leftarrow \lambda = (\phi \otimes \lambda) \circ \rho. \end{cases}$$

Proof. We denote by $*$ the convolution product of M_B and by \star the convolution product of M_A . As $\rho : A \rightarrow A \otimes B$ is an algebra morphism, \leftarrow is well-defined. Let $\phi \in M_A$, $\lambda_1, \lambda_2 \in M_B$.

$$\begin{aligned} (\phi \leftarrow \lambda_1) \leftarrow \lambda_2 &= (\phi \otimes \lambda_1 \otimes \lambda_2) \circ (\rho \otimes Id) \circ \rho \\ &= (\phi \otimes \lambda_1 \otimes \lambda_2) \circ (Id \otimes \Delta_B) \circ \rho \\ &= \phi \leftarrow (\lambda_1 * \lambda_2). \end{aligned}$$

So \leftarrow is an action. Let $\phi_1, \phi_2 \in M_A$, $\lambda \in M_B$. For all $a \in A$:

$$\begin{aligned} ((\phi_1 \star \phi_2) \leftarrow \lambda)(a) &= (\phi_1 \otimes \phi_2 \otimes \lambda) \circ (\Delta_A \otimes Id) \circ \rho(a) \\ &= (\phi_1 \otimes \phi_2 \otimes \lambda)((a_0)^{(1)} \otimes (a_0)^{(2)} \otimes a_1) \\ &= (\phi_1 \otimes \phi_2 \otimes \lambda)((a^{(1)})_0 \otimes (a^{(2)})_0 \otimes (a^{(1)})_1 (a^{(2)})_1) \\ &= \phi_1((a^{(1)})_0) \lambda((a^{(1)})_1) \phi_2((a^{(2)})_0) \lambda((a^{(2)})_1) \\ &= (\phi_1 \leftarrow \lambda)(a^{(1)}) (\phi_2 \leftarrow \lambda)(a^{(2)}) \\ &= ((\phi_1 \leftarrow \lambda) \star (\phi_2 \leftarrow \lambda))(a). \end{aligned}$$

So \leftarrow is an action by monoid endomorphisms. □

Example. We take $A = (\mathbb{K}[X], m, \Delta)$, $B = (\mathbb{K}[X], m, \delta)$ and $\rho = \delta$. We consider the map:

$$ev : \begin{cases} \mathbb{K} & \rightarrow & \mathbb{K}[X]^* \\ \lambda & \rightarrow & \begin{cases} \mathbb{K}[X] & \rightarrow & \mathbb{K} \\ P(X) & \rightarrow & ev_\lambda(P) = P(\lambda). \end{cases} \end{cases}$$

Then ev is an isomorphism from $(\mathbb{K}, +)$ to (M_A, \star) and from (\mathbb{K}, \cdot) to $(M_B, *)$. Moreover, for all $\lambda, \mu \in \mathbb{K}$:

$$ev_\lambda \leftarrow ev_\mu = ev_{\lambda\mu}.$$

Proposition 4 *Let A and B be two bialgebras in cointeraction, through the coaction ρ .*

1. *Let H be a bialgebra. We denote by M_B the monoid of characters of B and by $E_{A \rightarrow H}$ the set of bialgebra morphisms from A to H . Then M_B acts on $E_{A \rightarrow H}$ via the map:*

$$\leftarrow : \begin{cases} E_{A \rightarrow H} \times M_B & \rightarrow & E_{A \rightarrow H} \\ (\phi, \lambda) & \rightarrow & \phi \leftarrow \lambda = (\phi \otimes \lambda) \circ \rho \end{cases}$$

2. *Let H_1 and H_2 be two bialgebras and let $\theta : H_1 \rightarrow H_2$ be a bialgebra morphism. For all $\phi \in E_{A \leftarrow H_1}$, for all $\lambda \in M_B$, in $E_{A \leftarrow H_2}$:*

$$\theta \circ (\phi \leftarrow \lambda) = (\theta \circ \phi) \leftarrow \lambda.$$

3. *if $\lambda, \mu \in M_B$, in $E_{A \rightarrow A}$:*

$$(Id \leftarrow \lambda) \circ (Id \leftarrow \mu) = Id \leftarrow (\lambda * \mu).$$

The following map is an injective monoid morphism:

$$\begin{cases} (M_B, *) & \rightarrow & (E_{A \rightarrow A}, \circ) \\ \lambda & \rightarrow & Id \leftarrow \lambda. \end{cases}$$

Proof. 1. For all $\phi \in E_{A \leftarrow B}$, $\lambda \in M_B$, $\phi \leftarrow \lambda : A \rightarrow H \otimes \mathbb{K} = H$. As ϕ , λ and ρ are algebra morphisms, by composition $\phi \leftarrow \lambda$ is an algebra morphism. Let $a \in A$.

$$\begin{aligned}
\Delta_H(\phi \leftarrow \lambda(a)) &= \Delta_H(\phi(a_0)\lambda(a_1)) \\
&= \lambda(a_1)\Delta_H \circ \phi(a_1) \\
&= \lambda(a_1)\phi(a_0)^{(1)} \otimes \phi(a_0)^{(2)} \\
&= \lambda(a_1)\phi((a_0)^{(1)}) \otimes \phi((a_0)^{(2)}) \\
&= \lambda((a^{(1)})_1(a^{(2)})_1)\phi((a^{(1)})_0) \otimes \phi((a^{(2)})_0) \\
&= \lambda((a^{(1)})_1)\lambda((a^{(2)})_1)\phi((a^{(1)})_0) \otimes \phi((a^{(2)})_0) \\
&= \phi((a^{(1)})_0)\lambda((a^{(1)})_1) \otimes \phi((a^{(2)})_0)\lambda((a^{(2)})_1) \\
&= \phi \leftarrow \lambda(a^{(1)}) \otimes \phi \leftarrow \lambda(a^{(2)}) \\
&= ((\phi \leftarrow \lambda) \otimes (\phi \leftarrow \lambda)) \circ \Delta_A(a).
\end{aligned}$$

So $\phi \leftarrow \lambda \in E_{A \rightarrow H}$.

Let $\phi \in E_{A \rightarrow H}$. For all $a \in A$, $\phi \leftarrow \eta \circ \varepsilon(a) = \phi(a_0)\varepsilon(a_1) = \phi(a)$. Let $\lambda, \mu \in M_B$.

$$(\phi \leftarrow \lambda) \leftarrow \mu = (\phi \otimes \lambda \otimes \mu) \circ (\rho \otimes Id) \circ \rho = (\phi \otimes \lambda \otimes \mu) \circ (Id \otimes \Delta_B) \circ \rho = \phi \leftarrow (\lambda * \mu).$$

So \leftarrow is indeed an action of M_B on $E_{A \rightarrow H}$.

2. Let $a \in H$.

$$(\theta \circ \phi) \leftarrow \lambda(a) = \theta \circ \phi(a_1)\lambda(a_0) = \theta(\phi(a_1)\lambda(a_0)) = \theta(\phi \leftarrow \lambda(a)) = \theta \circ (\phi \leftarrow \lambda)(a).$$

So $(\theta \circ \phi) \leftarrow \lambda = \theta \circ (\phi \leftarrow \lambda)$.

3. Consequently, if $\lambda, \mu \in M_B$, in $E_{A \rightarrow A}$:

$$(Id \leftarrow \lambda) \circ (Id \leftarrow \lambda) = (Id \leftarrow \lambda) \leftarrow \mu = Id \leftarrow (\lambda * \mu).$$

If $Id \leftarrow \lambda = Id$, then, composing by ε' , we obtain $\varepsilon' * \lambda = \varepsilon'$, so $\lambda = \varepsilon'$. □

Example. We take $A = (\mathbb{K}[X], m, \Delta)$, $B = (\mathbb{K}[X], m, \delta)$ and $\rho = \delta$. In $E_{A \rightarrow A}$, for any $\lambda \in \mathbb{K}$:

$$Id \leftarrow ev_\lambda(X) = ev_\lambda(X)X = \lambda X,$$

so for any $P \in \mathbb{K}[X]$, $(Id \leftarrow ev_\lambda)(P) = P(\lambda X)$. In this case, the monoids $(M_B, *)$ and $(E_{A \rightarrow A}, \circ)$ are isomorphic.

1.3 Polynomial morphisms

In this section, we deal with a family (A, m, Δ, δ) such that:

1. (A, m, Δ) is a graded, connected Hopf algebra. As a graded algebra, it is isomorphic to the symmetric algebra $S(V)$, where V is a graded subspace of A .
2. (A, m, δ) is a bialgebra.
3. (A, m, Δ) and (A, m, δ) are in cointeraction, through the coaction δ .
4. $V_1 = A_1$ has a basis $(g_i)_{i \in I}$ such that:

$$\forall i \in I, \delta(g_i) = g_i \otimes g_i.$$

We shall denote by J the set of sequences $\alpha = (\alpha_i)_{i \in I}$ with a finite support. For all $\alpha \in J$, we put $g_\alpha = \prod_{i \in I} g_i^{\alpha_i}$. These are group-like elements of (A, m, δ) .

5. For all $n \geq 2$, V_n can be decomposed as:

$$V_n = \bigoplus_{i \in I, \alpha \in J} V_n(g_i, g_\alpha),$$

such that for all $x \in V_n(g_i, g_\alpha)$:

$$\delta(x) - g_i \otimes x - x \otimes g_\alpha \in S(V_1 \oplus \dots \oplus V_{n-1})^{\otimes 2}.$$

The counit of (A, m, Δ) will be denoted by ε and the counit of (A, m, δ) by ε' . We denote by M_B the monoid of characters of (A, m, δ) .

Remark. If $x \in V_n(g_i, g_\alpha)$, as $\varepsilon'(g_i) = \varepsilon'(g_\alpha) = 1$, necessarily, $\varepsilon'(x) = 0$ and:

$$\delta(x) - g_i \otimes x - x \otimes g_\alpha \in \text{Ker}(\varepsilon')^{\otimes 2}.$$

Lemma 5 *Let $\lambda \in M_B$. It has an inverse in M_B if, and only if, for all $i \in I$, $\lambda(g_i) \neq 0$.*

Proof. \implies . Let μ be the inverse of λ in M_B . For all $i \in I$, $\lambda * \mu(g_i) = \lambda(g_i)\mu(g_i) = \varepsilon'(g_i) = 1$, so $\lambda(g_i) \neq 0$.

\impliedby . We define two characters $\mu, \nu \in M_B$ by inductively defining $\mu_n = \mu|_{V_n}$ and $\nu_n = \nu|_{V_n}$. For $n = 1$, we put $\mu_1(g_i) = \nu_1(g_i) = \lambda(g_i)^{-1}$. Let us assume that g_1, \dots, g_{n-1} are already defined, with $n \geq 2$. If $x \in V_n(g_i, g_\alpha)$, we put:

$$\delta(x) - g_i \otimes x - x \otimes g_\alpha = \sum x'_k \otimes x''_k \in S(V_1 \oplus \dots \oplus V_{n-1})^{\otimes 2}.$$

Hence, for all k , $\mu(x'_k)$ and $\nu(x''_k)$ are defined. We put:

$$\begin{aligned} \mu_n(x) &= \prod_{i \in I} \frac{1}{\lambda(g_i)^{\alpha_i}} \left(\varepsilon'(x) - \mu(g_i)\lambda(x) - \sum \mu(x'_k)\lambda(x''_k) \right), \\ \nu_n(x) &= \frac{1}{\lambda(g_i)} \left(\varepsilon'(x) - \lambda(x)\nu(g_\alpha) - \sum \lambda(x'_k)\nu(x''_k) \right). \end{aligned}$$

Consequently, $\mu, \nu \in M_B$ and for all $x \in V$, $\mu * \lambda(x) = \lambda * \nu(x) = \varepsilon'(x)$, so $\mu * \lambda = \lambda * \nu = \varepsilon'$, and finally $\mu = \mu * (\lambda * \nu) = (\mu * \lambda) * \nu = \nu$, so λ is invertible in M_B . \square

Lemma 6 *Let C be a graded, connected Hopf algebra. We denote by $C_+ = \text{Ker}(\varepsilon_C)$ its augmentation ideal. Let $\lambda : C_+ \rightarrow \mathbb{K}$ be any linear map. There exists a unique coalgebra morphism $\phi : C \rightarrow \mathbb{K}[X]$ such that:*

$$\forall x \in C_+, \frac{d\phi(x)}{dX}(0) = \lambda(x).$$

Moreover:

1. ϕ is homogeneous if, and only if, for all $n \geq 2$, $\lambda(C_n) = (0)$.
2. ϕ is a Hopf algebra morphism if, and only if, for all $x, y \in C_+$, $\lambda(xy) = 0$.

Proof. Let $\pi : \mathbb{K}[X] \rightarrow \text{Vect}(X)$ be the canonical projection. For any $P \in \mathbb{K}[X]$:

$$\pi(P) = \frac{dP}{dX}(0)X.$$

Existence. We define $\phi|_{C_n}$ by induction on n . For $n = 0$, we put $\phi(1) = 1$. Let us assume that $\phi|_{C_0 \oplus \dots \oplus C_{n-1}}$ is defined such that for all $x \in C_0 \oplus \dots \oplus C_{n-1}$,

$$\Delta \circ \phi(x) = (\phi \otimes \phi) \circ \Delta(x), \quad \pi \circ \phi(x) = \lambda(x)X.$$

Let $x \in C_n$. As $\tilde{\Delta}(x) \in (C_1 \oplus \dots \oplus C_{n-1})^{\otimes 2}$, we can consider the element

$$y = (\phi \otimes \phi) \circ \tilde{\Delta}(x) \in \mathbb{K}[X]_+^{\otimes 2}.$$

We put:

$$y = \sum_{i,j \geq 1} a_{i,j} \frac{X^i}{i!} \otimes \frac{X^j}{j!}.$$

Moreover:

$$\begin{aligned} \tilde{\Delta} \otimes Id(y) &= ((\tilde{\Delta} \circ \phi) \otimes \phi) \circ \tilde{\Delta}(x) \\ &= (\phi \otimes \phi \otimes \phi) \circ (\tilde{\Delta} \otimes Id) \circ \tilde{\Delta}(x) \\ &= (\phi \otimes \phi \otimes \phi) \circ (Id \otimes \tilde{\Delta}) \circ \tilde{\Delta}(x) \\ &= (\phi \otimes (\tilde{\Delta} \circ \phi)) \circ \tilde{\Delta}(x) \\ &= (Id \otimes \tilde{\Delta})(y). \end{aligned}$$

Hence:

$$\sum_{i,j,k \geq 1} a_{i+j,k} \frac{X^i}{i!} \otimes \frac{X^j}{j!} \otimes \frac{X^k}{k!} = \sum_{i,j,k \geq 1} a_{ij+k} \frac{X^i}{i!} \otimes \frac{X^j}{j!} \otimes \frac{X^k}{k!}.$$

For all $i, j, k \geq 1$, $a_{i+j,k} = a_{i,j+k}$, so there exist scalars a_n such that for all $i, j \geq 1$, $a_{i,j} = a_{i+j}$. We obtain that:

$$y = \sum_{n \geq 2} a_n \left(\sum_{i,j \geq 1, i+j=n} \frac{X^i}{i!} \otimes \frac{X^j}{j!} \right) = \sum_{n \geq 2} a_n \tilde{\Delta} \left(\frac{X^n}{n!} \right).$$

We then put $\phi(x) = \sum_{n \geq 2} a_n \frac{X^n}{n!} + \lambda(x)X$.

We obtain in this way a coalgebra morphism such that $\pi \circ \phi(x) = \lambda(x)X$ for all $x \in C$.

Unicity. Let $\phi, \psi : C \rightarrow \mathbb{K}[X]$ be coalgebra morphisms such that $\pi \circ \phi = \pi \circ \psi$. Let us prove that $\phi(x) = \psi(x)$ for all $x \in A_n$, $n \geq 0$, by induction on n . For $n = 0$, as the unique group-like element of $\mathbb{K}[X]$ is 1, $\phi(1) = \psi(1) = 1$. Let us assume the result at all ranks $k < n$. Then, if $x \in A_n$:

$$\tilde{\Delta} \circ \phi(x) = (\phi \otimes \phi) \circ \tilde{\Delta}(x) = (\psi \otimes \psi) \circ \tilde{\Delta}(x) = \tilde{\Delta} \circ \psi(x),$$

so $\phi(x) - \psi(x) \in \text{Ker}(\tilde{\Delta}) = \text{Vect}(X)$. Hence, $0 = \pi \circ \phi(x) - \pi \circ \psi(x) = \phi(x) - \psi(x)$.

1. \implies . If ϕ is homogeneous, then, for all $n \geq 2$, $\phi(C_n) \subseteq \text{Vect}(X^n)$, so:

$$\pi \circ \phi(C_n) = \lambda(C_n)X = (0).$$

1. \impliedby . Let us go back to the construction of ϕ is the *Existence* part. If $n \geq 1$, $x \in C_1$, then $\phi(x) = \lambda(x)X$ is homogeneous of degree 1. If $n \geq 2$, then, by homogeneity, $a_{i,j} = 0$ if $i + j \neq n$, so $a_k = 0$ if $n \neq k$, and $\phi(x) = a_n \frac{X^n}{n!}$: ϕ is homogeneous.

2. \implies . Let $x, y \in C_+$. Then $\phi(x), \phi(y) \in \mathbb{K}[X]_+ = X\mathbb{K}[X]$, so $\phi(xy) = \phi(x)\phi(y) \in X^2\mathbb{K}[X]$ and $\pi \circ \phi(xy) = \lambda(xy)X = 0$.

2. \Leftarrow . Let us consider $\phi_1 = m \circ (\phi \otimes \phi)$ and $\phi_2 = \phi \circ m$. By composition, they are both coalgebra morphisms from the graded bialgebra $C \otimes C$ to $\mathbb{K}[X]$. Moreover, if $x, y \in C_+$:

$$\begin{aligned} \pi \circ \phi_1(1 \otimes y) &= \pi \circ \phi(y), & \pi \circ \phi_2(1 \otimes y) &= \pi \circ \phi(y), \\ \pi \circ \phi_1(x \otimes 1) &= \pi \circ \phi(x), & \pi \circ \phi_2(x \otimes 1) &= \pi \circ \phi(x), \\ \pi \circ \phi_1(x \otimes y) &= \pi(\phi(x)\phi(y)) = 0, & \pi \circ \phi_2(x \otimes y) &= \lambda(xy)X = 0. \end{aligned}$$

So, for all $z \in (C \otimes C)_+ = (\mathbb{K}1 \otimes C_+) \oplus (C_+ \otimes \mathbb{K}1) \oplus (C_+ \otimes C_+)$, $\pi \circ \phi_1(z) = \pi \circ \phi_2(z)$. By the *Unicity* part, $\phi_1 = \phi_2$, so ϕ is an algebra morphism. \square

Remark. If $x \in V_1$, $\phi(x) = \lambda(x)X$.

Theorem 7 *Under the hypotheses 1–5, there exists a unique homogeneous Hopf algebra morphism $\phi_0 : (A, m, \Delta) \rightarrow (\mathbb{K}[X], m, \Delta)$ such that:*

$$\forall x \in A_1, \phi_0(x) = \varepsilon'(x)X.$$

Moreover, there exists a unique character $\lambda_0 \in M_B$, invertible in M_B , such that:

$$\forall n \geq 0, \forall x \in A_n, \phi_0(x) = \lambda_0(x)X^n.$$

Proof. Existence. Let $\lambda \in A^*$ such that $\lambda(x) = \varepsilon'(x)$ if $x \in A_1$ and $\lambda(x) = 0$ if $x \in A_n$, $n \geq 2$. We denote by ϕ_0 the unique coalgebra morphism such that $\frac{d\phi_0(x)}{dX}(0) = \lambda(x)$ for all $x \in A_+$. By the first point of lemma 6, ϕ_0 is homogeneous. As $A_+^2 \subseteq A_{\geq 2}$, by the second point of lemma 6, ϕ_0 is an algebra morphism.

Unicity. If ϕ is such a morphism, by the first point of lemma 6, for all $x \in A_n$, $n \geq 2$, $\pi \circ \phi(x) = 0$; hence, for all $x \in A_+$, $\pi \circ \phi_0 = \pi \circ \phi$. By the unicity in lemma 6, $\phi = \phi_0$.

By homogeneity of ϕ_0 , for all $x \in A_n$, there exists a unique scalar $\lambda_0(x) \in \mathbb{K}$ such that:

$$\phi_0(x) = \lambda_0(x)X^n.$$

If $x \in A_m$, $y \in A_n$, $xy \in A_{m+n}$ and then:

$$\phi(xy) = \lambda_0(xy)X^{n+m} = \phi(x)\phi(y) = \lambda_0(x)\lambda_0(y)X^{n+m},$$

so $\lambda_0 \in M_B$. For all $i \in I$, $\phi(g_i) = \varepsilon'(g_i)X = X$, so $\lambda_0(g_i) = 1$. By lemma 5, λ_0 is an invertible element of M_B . \square

Theorem 8 *Under the hypotheses 1–5, the following map is a bijection:*

$$\theta : \begin{cases} M_B & \longrightarrow E_{A \rightarrow \mathbb{K}[X]} \\ \lambda & \longrightarrow \phi_0 \leftarrow \lambda. \end{cases}$$

Moreover, if $\phi = \theta(\lambda)$, with $\lambda \in M_B$, then for all $x \in V$,

$$\lambda(x) = \frac{d\phi(x)}{dX}(0).$$

Proof. Let $\lambda \in M_B$, and $\phi = \phi_0 \leftarrow \lambda$. For all $i \in I$, $\phi(g_i) = X\lambda(g_i)$, so $\frac{d\phi(g_i)}{dX}(0) = \lambda(g_i)$. If $n \geq 2$ and $x \in V_n(g_i, g_\alpha)$, we put:

$$\delta(x) = g_i \otimes x + \sum x'_k \otimes x''_k,$$

with for all k , x'_k homogeneous of degree ≥ 2 or homogeneous of degree 1, with $\varepsilon'(x'_k) = 0$. For all k , $\pi \circ \phi_0(x'_k) = 0$. We obtain:

$$\pi \circ \phi(x) = \pi \left(\lambda(x)X + \sum \phi_0(x'_k)\lambda(x''_k) \right) = \lambda(x)X.$$

By the unicity in lemma 6, ϕ is injective. If $\psi \in E_{A \rightarrow \mathbb{K}[X]}$, we define $\lambda \in M_B$ by:

$$\forall x \in V, \lambda(x) = \frac{d\psi(x)}{dX}(0).$$

We put $\phi = \phi_0 \leftarrow \lambda$. Then for all $x \in V$, $\frac{d\phi(x)}{dX}(0) = \frac{d\psi(x)}{dX}(0) = \lambda(x)$. If $x \in A_+^2$, by the second point of lemma 6, $\frac{d\phi(x)}{dX}(0) = \frac{d\psi(x)}{dX}(0) = 0$. Finally, as V generates A , $A_+ = V + A_+^2$, and for all $x \in A_+$, $\frac{d\phi(x)}{dX}(0) = \frac{d\psi(x)}{dX}(0)$. By the unicity in lemma 6, $\phi = \psi$, so θ is surjective. \square

Corollary 9 *Under the hypotheses 1–5, for any $\mu \in M_B$, there exists a unique Hopf algebra morphism $\phi : A \rightarrow \mathbb{K}[X]$, such that:*

$$\forall x \in A, \phi(x)(1) = \mu(1).$$

This morphism is: $\phi_0 \leftarrow (\lambda_0^{-1} * \mu)$.*

Proof. Let ϕ be a Hopf algebra morphism from A to $\mathbb{K}[X]$. By theorem 8, there exists $\lambda \in M_B$ such that $\phi = \phi_0 \leftarrow \lambda$. Let $x \in A$. We write $x = \sum x'_k \otimes x''_k$, with, for all k , x'_k homogeneous of degree n_k . Then:

$$\phi(x)(1) = \left(\sum \lambda_0(x'_k)X^{n_k}\lambda(x''_k) \right)_{|X=1} = \sum \lambda_0(x'_k)\lambda(x''_k) = \lambda_0 * \lambda(x).$$

So ϕ satisfies the required conditions if, and only if, $\lambda_0 * \lambda = \mu$, if, and only if, $\lambda = \lambda_0^{*-1} * \mu$, as λ_0 is invertible. So such a ϕ exists and is unique. \square

Corollary 10 *Under the hypotheses 1–5, there exists a unique morphism $\phi_1 : A \rightarrow \mathbb{K}[X]$ such that:*

1. ϕ_1 is a Hopf algebra morphism from (A, m, Δ) to $(\mathbb{K}[X], m, \Delta)$.
2. ϕ_1 is a bialgebra morphism from (A, m, δ) to $(\mathbb{K}[X], m, \delta)$.

Moreover, $\phi_1 = \phi_0 \leftarrow \lambda_0^{*-1}$ and, for all $x \in A$, $\phi_1(x)(1) = \varepsilon'(x)$.

Proof. *Unicity.* If such a ϕ_1 exists, then for all $x \in A$, $\phi_1(x)(1) = \varepsilon' \circ \phi_1(x) = \varepsilon'(x)$. By corollary 9, $\phi_1 = \phi_0 \leftarrow (\lambda_0^{*-1} * \varepsilon') = \phi_0 \leftarrow \lambda_0^{*-1}$.

Existence. Let ϕ_1 be the unique Hopf algebra morphism in $E_{A \rightarrow \mathbb{K}[X]}$ such that for all $x \in A$, $\phi_1(x)(1) = \varepsilon'(x)$. Recall that we identify $\mathbb{K}[X] \otimes \mathbb{K}[X]$ with $\mathbb{K}[X, Y]$; for all $P \in \mathbb{K}[X]$:

$$\Delta(P)(X, Y) = P(X + Y), \quad \delta(P)(X, Y) = P(XY).$$

Let us fix $x \in A$. Let us prove that for all $k, l \in \mathbb{N}^*$, $\delta \circ \phi_1(x)(k, l) = (\phi_1 \otimes \phi_1) \circ \Delta(x)(k, l)$ by induction on k . First, observe that $\delta \circ \phi_1(x)(k, l) = \phi_1(x)(kl)$. If $k = 1$, then:

$$(\phi_1 \otimes \phi_1) \circ \Delta(x)(1, l) = \phi_1((\varepsilon' \otimes Id) \circ \Delta(x))(l) = \phi_1(x)(l).$$

Let us assume the result at rank k . Then:

$$\begin{aligned}
(\phi_1 \otimes \phi_1) \circ \Delta(x)(k+1, l) &= (\Delta \otimes Id) \circ (\phi_1 \otimes \phi_1) \circ \delta(x)(k, 1, l) \\
&= (\phi_1 \otimes \phi_1 \otimes \phi_1) \circ (\Delta \otimes Id) \circ \delta(x)(k, 1, l) \\
&= (\phi_1 \otimes \phi_1 \otimes \phi_1) \circ m_{2,4}^3 \circ (\delta \otimes \delta) \circ \Delta(x)(k, 1, l) \\
&= (\phi_1 \otimes \phi_1 \otimes \phi_1 \otimes \phi_1) \circ (\delta \otimes \delta) \circ \Delta(x)(k, l, 1, l) \\
&= (\phi_1 \otimes \phi_1) \circ \Delta(x)(kl, l) \\
&= \Delta \circ \phi_1(x)(kl, l) \\
&= \phi_1(x)(kl + l) \\
&= \phi_1(x)((k+1)l).
\end{aligned}$$

So the result is true for all $k, l \geq 1$. Hence, $(\phi_1 \otimes \phi_1) \circ \delta(x) = \delta \circ \phi_1(x)$: ϕ_1 is a bialgebra morphism from (A, m, δ) to $(\mathbb{K}[X], m, \delta)$. \square

2 Examples from quasi-posets

2.1 Definition

Definition 11 1. Let A be a set finite set. A quasi-order on A is a transitive, reflexive relation \leq on A . If \leq is a quasi-order on A , we shall say that (A, \leq) is a quasi-poset. If P is a quasi-poset:

(a) Its isoclass is denoted by $[P]$.

(b) \sim_P is defined by:

$$\forall a, b \in A, a \sim_P b \text{ if } (a \leq b \text{ and } b \leq a).$$

It is an equivalence on A .

(c) $\bar{A} = A / \sim_P$ is given an order by:

$$\forall a, b \in A, \bar{a} \leq \bar{b} \text{ if } a \leq b.$$

The poset (\bar{A}, \leq) is denoted by \bar{P} .

(d) The cardinality of \bar{P} is denoted by $cl(P)$.

2. Let $n \in \mathbb{N}$.

(a) The set of quasi-posets which underlying set is $[n] = \{1, \dots, n\}$ is denoted by $\mathbf{QP}(n)$.

(b) The set of posets which underlying set is $[n]$ is denoted by $\mathbf{P}(n)$.

(c) The set of isoclasses of quasi-posets of cardinality n is denoted by $\mathbf{qp}(n)$.

(d) The set of isoclasses of quasi-posets of cardinality n is denoted by $\mathbf{p}(n)$.

We put:

$$\begin{aligned}
\mathbf{QP} &= \bigsqcup_{n \geq 0} \mathbf{QP}(n), & \mathbf{P} &= \bigsqcup_{n \geq 0} \mathbf{P}(n), & \mathbf{qp} &= \bigsqcup_{n \geq 0} \mathbf{qp}(n), & \mathbf{p} &= \bigsqcup_{n \geq 0} \mathbf{p}(n), \\
\mathcal{H}_{\mathbf{QP}} &= \text{Vect}(\mathbf{QP}), & \mathcal{H}_{\mathbf{P}} &= \text{Vect}(\mathbf{P}), & \mathcal{H}_{\mathbf{qp}} &= \text{Vect}(\mathbf{qp}) & \mathcal{H}_{\mathbf{p}} &= \text{Vect}(\mathbf{p}).
\end{aligned}$$

As posets are quasi-posets, there are canonical injections from $\mathcal{H}_{\mathbf{P}}$ into $\mathcal{H}_{\mathbf{QP}}$ and from $\mathcal{H}_{\mathbf{p}}$ into $\mathcal{H}_{\mathbf{qp}}$. Moreover, the map $P \rightarrow \bar{P}$ induces surjective maps from $\mathcal{H}_{\mathbf{QP}}$ to $\mathcal{H}_{\mathbf{P}}$ and from

$\mathcal{H}_{\mathbf{QP}}$ to $\mathcal{H}_{\mathbf{P}}$, both denoted by ξ . The map $P \rightarrow [P]$ induces maps $\sqcup : \mathcal{H}_{\mathbf{QP}} \rightarrow \mathcal{H}_{\mathbf{QP}}$ and $\sqcup : \mathcal{H}_{\mathbf{P}} \rightarrow \mathcal{H}_{\mathbf{P}}$. The following diagram commutes:

$$\begin{array}{ccccc}
\mathcal{H}_{\mathbf{QP}} & \xrightarrow{\sqcup} & \mathcal{H}_{\mathbf{QP}} & & \\
& \searrow \xi & & \nearrow \xi & \\
& & \mathcal{H}_{\mathbf{P}} & \xrightarrow{\sqcup} & \mathcal{H}_{\mathbf{P}} \\
& \nearrow Id & & \searrow Id & \\
\mathcal{H}_{\mathbf{P}} & \xrightarrow{\sqcup} & \mathcal{H}_{\mathbf{P}} & &
\end{array} \tag{1}$$

We shall represent any element P of \mathbf{QP} by the Hasse graph of \overline{P} , indicating on the vertices the elements of the corresponding equivalence class. For example, the elements of $\mathbf{QP}(n)$, $n \leq 3$, are:

$$\begin{aligned}
& 1; \quad \cdot_1; \quad \cdot_1 \cdot_2, \mathbf{1}_1^2, \mathbf{1}_2^1, \cdot_1 \cdot_2; \quad \cdot_1 \cdot_2 \cdot_3, \cdot_1 \mathbf{1}_2^3, \cdot_1 \mathbf{1}_3^2, \cdot_2 \mathbf{1}_1^3, \cdot_2 \mathbf{1}_3^1, \cdot_3 \mathbf{1}_1^2, \cdot_3 \mathbf{1}_2^1, \cdot_1 \cdot_2 \cdot_3, \cdot_2 \cdot_1 \cdot_3, \cdot_3 \cdot_1 \cdot_2, \\
& {}^2\mathbf{V}_1^3, {}^1\mathbf{V}_2^3, {}^1\mathbf{V}_3^2, {}^2\mathbf{\Lambda}_3^1, {}^1\mathbf{\Lambda}_3^2, {}^1\mathbf{\Lambda}_3^3, \mathbf{1}_1^3, \mathbf{1}_1^2, \mathbf{1}_2^3, \mathbf{1}_2^1, \mathbf{1}_3^3, \mathbf{1}_3^1, \mathbf{1}_3^2, \mathbf{1}_1^{2,3}, \mathbf{1}_2^{1,3}, \mathbf{1}_3^{1,2}, \mathbf{1}_{2,3}^1, \mathbf{1}_{1,3}^2, \mathbf{1}_{1,2}^3, \cdot_1 \cdot_2 \cdot_3.
\end{aligned}$$

We shall represent any element $P \in \mathbf{qp}$ by the Hasse graph of \overline{P} , indicating on the vertices the cardinality of the corresponding equivalence class, if this cardinality is not equal to 1. For example, the elements of $\mathbf{qp}(n)$, $n \leq 3$, are:

$$1; \quad \cdot; \quad \dots, \mathbf{1}, \cdot_2; \quad \dots, \cdot \mathbf{1}, \cdot_2; \quad \mathbf{V}, \mathbf{\Lambda}, \mathbf{1}, \mathbf{1}^2, \mathbf{1}_2, \cdot_3.$$

2.2 First coproduct

By Alexandroff's theorem [1, 18], finite quasi-posets are in bijection with finite topological spaces. Let us recall the definition of the topology attached to a quasi-poset.

Definition 12 1. Let $P = (A, \leq)$ be a quasi-poset. An open set of P is a subset O of A such that:

$$\forall i, j \in A, (i \in O \text{ and } i \leq j) \implies (j \in O).$$

The set of open sets of P (the topology associated to P) is denoted by $\text{top}(P)$.

2. Let $P = (A, \leq)$ be a quasi-poset and $B \subseteq A$. We denote by $P|_B$ the quasi-poset $(B, \leq|_B)$.
3. Let $P = (A, \leq_P)$ be a quasi-poset. We assume that A is also given a total order \leq : for example, A is a subset of \mathbb{N} . If the cardinality of A is n , there exists a unique increasing bijection f from $[n]$, with its usual order, to (A, \leq) . We denote by $\text{Std}(P)$ the quasi-poset, element of $\mathbf{QP}(n)$, defined by:

$$\forall i, j \in [n], i \leq_{\text{Std}(P)} j \iff f(i) \leq_P f(j).$$

Proposition 13 1. We define a product m on $\mathcal{H}_{\mathbf{QP}}$ in the following way: if $P \in \mathbf{QP}(k)$, $Q \in \mathbf{QP}(l)$, then $PQ = m(P, Q) \in \mathbf{QP}(k+l)$ and

$$\begin{aligned}
\forall i, j \in [k+l], i \leq_{PQ} \iff & (1 \leq i, j \leq k \text{ and } i \leq_P j) \\
& \text{or } (k+1 \leq i, j \leq k+l \text{ and } i-k \leq_Q j-k).
\end{aligned}$$

2. We define a second product \downarrow on $\mathcal{H}_{\mathbf{QP}}$ in the following way: if $P \in \mathbf{QP}(k)$, $Q \in \mathbf{QP}(l)$, then $PQ = m(P, Q) \in \mathbf{QP}(k+l)$ and

$$\begin{aligned}
\forall i, j \in [k+l], i \leq_{PQ} \iff & (1 \leq i, j \leq k \text{ and } i \leq_P j) \\
& \text{or } (k+1 \leq i, j \leq k+l \text{ and } i-k \leq_Q j-k) \\
& \text{or } (1 \leq i \leq k < j \leq k+l).
\end{aligned}$$

3. We define a coproduct Δ on $\mathcal{H}_{\mathbf{QP}}$ in the following way:

$$\forall P \in \mathbf{QP}(n), \Delta(P) = \sum_{O \in \text{top}(P)} \text{Std}(P|_{[n] \setminus O}) \otimes \text{Std}(P|_O).$$

Then $(\mathcal{H}_{\mathbf{QP}}, m, \Delta)$ is a non-commutative, non-cocommutative Hopf algebra, and $(\mathcal{H}_{\mathbf{QP}}, \downarrow, \Delta)$ is an infinitesimal bialgebra.

Proof. See [10, 11]. □

Examples. If $\{a, b\} = \{1, 2\}$ and $\{i, j, k\} = \{1, 2, 3\}$:

$$\begin{aligned} \Delta(\bullet_1) &= \bullet_1 \otimes 1 + 1 \otimes \bullet_1, \\ \Delta(\mathbf{1}_a^b) &= \mathbf{1}_a^b \otimes 1 + 1 \otimes \mathbf{1}_a^b + \bullet_a \otimes \bullet_b, \\ \Delta({}^j\mathbf{V}_i^k) &= {}^j\mathbf{V}_i^k \otimes 1 + 1 \otimes {}^j\mathbf{V}_i^k + \mathbf{1}_i^j \otimes \bullet_k + \mathbf{1}_i^k \otimes \bullet_j + \bullet_i \otimes \bullet_{j \bullet k}, \\ \Delta({}_j\mathbf{A}_k^i) &= {}_j\mathbf{A}_k^i \otimes 1 + 1 \otimes {}_j\mathbf{A}_k^i + \bullet_j \otimes \mathbf{1}_k^i + \bullet_k \otimes \mathbf{1}_j^i + \bullet_{j \bullet k} \otimes \bullet_i, \\ \Delta(\mathbf{1}_i^j) &= \mathbf{1}_i^j \otimes 1 + 1 \otimes \mathbf{1}_i^j + \bullet_i \otimes \mathbf{1}_j^i + \mathbf{1}_i^j \otimes \bullet_k. \end{aligned}$$

Remark. This Hopf algebraic structure is compatible with the morphisms of (1), that is to say:

1. $\mathcal{H}_{\mathbf{P}}$ is a Hopf subalgebra of $\mathcal{H}_{\mathbf{QP}}$.
2. observe that:
 - If (P_1, P_2) and (Q_1, Q_2) are pairs of isomorphic quasi-posets, then $P_1 Q_1$ and $P_2 Q_2$ are isomorphic.
 - If P_1 and P_2 are isomorphic quasi-posets of $\mathbf{QP}(n)$, and if $\phi : [n] \rightarrow [n]$ is an isomorphism from P_1 to P_2 , then the topology associated to P_2 is the image by ϕ of the topology associated to P_1 and for any subset I of P_1 , $\phi|_I$ is an isomorphism from $(P_1)|_I$ to $(P_2)|_{\phi(I)}$.

Consequently, the surjective map $\llbracket \cdot \rrbracket : \mathcal{H}_{\mathbf{QP}} \rightarrow \mathcal{H}_{\mathbf{qp}}$ is compatible with the product and the coproduct: $\mathcal{H}_{\mathbf{qp}}$ inherits a Hopf algebra structure. Its product is the disjoint union of quasi-posets. For any quasi-poset $P = (A, \leq_P)$:

$$\Delta(\llbracket P \rrbracket) = \sum_{O \in \text{top}(P)} \llbracket P|_{A \setminus O} \rrbracket \otimes \llbracket P|_O \rrbracket.$$

3. $\mathcal{H}_{\mathbf{p}}$ is a Hopf subalgebra of $\mathcal{H}_{\mathbf{qp}}$.
4. All the morphisms in (1) are Hopf algebra morphisms.

Definition 14 1. We shall say that a finite quasi-poset $P = (A, \leq_P)$ is connected if its associated topology is connected.

2. For any finite quasi-poset P , we denote by $cc(P)$ the number of connected components of its associated topology.

It is well-known that P is connected if, and only if, the Hasse graph of \overline{P} is connected. Any quasi-poset P can be decomposed as the disjoint union of its connected components; in an algebraic setting, $\mathcal{H}_{\mathbf{qp}}$ is generated as a polynomial algebra by the connected quasi-posets. This is not true in $\mathcal{H}_{\mathbf{QP}}$: for example, $\mathbf{1}_1^2$ is both not connected and indecomposable in $\mathcal{H}_{\mathbf{QP}}$.

2.3 Second coproduct

Definition 15 Let $P = (A, \leq_P)$ be a quasi-poset and let \sim be an equivalence on A .

1. We define a second quasi-order $\leq_{P|\sim}$ on A by the relation:

$$\forall x, y \in A, x \leq_{P|\sim} y \text{ if } (x \leq_P y \text{ and } x \sim y).$$

2. We define a third quasi-order $\leq_{P/\sim}$ on A as the transitive closure of the relation \mathcal{R} defined by:

$$\forall x, y \in A, x \mathcal{R} y \text{ if } (x \leq_P y \text{ or } x \sim y).$$

3. We shall say that \sim is P -compatible and we shall denote $\sim \triangleleft P$ if the two following conditions are satisfied:

- The restriction of P to any equivalence class of \sim is connected.
- The equivalences $\sim_{P/\sim}$ and \sim are equal. In other words:

$$\forall x, y \in A, (x \leq_{P/\sim} y \text{ and } y \leq_{P/\sim} x) \implies x \sim y;$$

note that the converse assertion trivially holds.

Remarks.

1. $P|\sim$ is the disjoint union of the restriction of \leq_P to the equivalence classes of \sim .

2. Let $x, y \in P$. Then $x \leq_{P/\sim} y$ if there exist $x_1, x'_1, \dots, x_k, x'_k \in A$ such that:

$$x \leq_P x_1 \sim x'_1 \leq_P \dots \leq_P x_k \sim x'_k \leq_P y.$$

3. If $\sim \triangleleft P$, then:

- (a) The equivalence classes of $\sim_{P/\sim}$ are the equivalence classes of \sim and are included in a connected component of P . This implies that the connected components of P/\sim are the connected components of P . Consequently:

$$cl(P/\sim) = cl(\sim), \quad cc(P/\sim) = cc(P), \quad (2)$$

where $cl(\sim)$ is the number of equivalence classes of \sim .

- (b) If $x \sim_P y$ and $x \sim y$, then $x \sim_{P|\sim} y$: the equivalence classes of $\sim_{P|\sim}$ are the equivalence classes of \sim_P ; the connected components of $P|\sim$ are the equivalence classes of \sim . Consequently:

$$cl(P|\sim) = cl(P), \quad cc(P|\sim) = cl(\sim). \quad (3)$$

Definition 16 Let $P \in \mathbf{QP}$. We shall say that P is discrete if $[\overline{P}] = \cdot^{cl(P)}$.

In other words, P is discrete if, and only if, $\sim_P = \leq_P$.

Definition 17 We define a second coproduct δ on $\mathcal{H}_{\mathbf{QP}}$ in the following way: for all $P \in \mathbf{QP}$,

$$\delta(P) = \sum_{\sim \triangleleft P} (P/\sim) \otimes (P|\sim).$$

Then $(\mathcal{H}_{\mathbf{QP}}, m, \delta)$ is a bialgebra. Its counit ε' is given by:

$$\forall P \in \mathbf{QP}, \varepsilon'(P) = \begin{cases} 1 & \text{if } P \text{ is discrete,} \\ 0 & \text{otherwise.} \end{cases}$$

Proof. Firstly, let us prove the compatibility of δ and m . Let $P = (A, \leq_P)$ and $Q = (B, \leq_Q)$ be two elements of \mathbf{QP} . Let \sim be an equivalence relation on P . We denote by \sim' and \sim'' the restriction of \sim to P and Q . Then:

- If $\sim \triangleleft PQ$, then as the equivalence classes of \sim are connected, they are included in A or in B . Consequently, if $x \in A$ and $y \in B$, x and y are not equivalent for \sim . Moreover, $\sim' \triangleleft P$ and $\sim'' \triangleleft Q$, and:

$$PQ| \sim = (P| \sim')(Q| \sim''), \quad PQ/ \sim = (P/ \sim')(Q/ \sim'').$$

- Conversely, if $\sim' \triangleleft P, \sim'' \triangleleft Q$ and for all $x \in A, y \in B$, x and y are not \sim -equivalent, then $\sim \triangleleft PQ$.

Hence:

$$\begin{aligned} \delta(PQ) &= \sum_{\sim \triangleleft PQ} (PQ/ \sim) \otimes (PQ| \sim) \\ &= \sum_{\sim' \triangleleft P, \sim'' \triangleleft Q} (P/ \sim')(Q/ \sim'') \otimes (P| \sim')(Q| \sim'') \\ &= \delta(P)\delta(Q). \end{aligned}$$

Let us now prove the coassociativity of δ . Let $P \in \mathbf{QP}$.

First step. We put:

$$A = \{(r, r') \mid r \triangleleft P, r' \triangleleft P/r\}, \quad B = \{(s, s') \mid s \triangleleft P, s' \triangleleft P|s\}.$$

We consider the maps:

$$F : \begin{cases} A & \longrightarrow B \\ (r, r') & \longrightarrow (r', r), \end{cases} \quad G : \begin{cases} B & \longrightarrow A \\ (s, s') & \longrightarrow (s', s). \end{cases}$$

F is well-defined: we put $(s, s') = (r', r)$. The equivalence classes of s are the equivalence classes of r' , so are P -connected. If $x \sim_{P|s} y$, there exist $x_1, x'_1, \dots, x_k, s'_k$ and $y_1, y'_1, \dots, y_l, y_l$ such that:

$$x \leq_P x_1 r' x'_1 \leq_P \dots \leq_P x_k r' x'_k \leq_P y, \quad y \leq_P y_1 r' y'_1 \leq_P \dots \leq_P y_l r' y'_l \leq_P x.$$

Hence:

$$x \leq_{P/r} x_1 r' x'_1 \leq_{P/r} \dots \leq_{P/r} x_k r' x'_k \leq_{P/r} y, \quad y \leq_{P/r} y_1 r' y'_1 \leq_{P/r} \dots \leq_{P/r} y_l r' y'_l \leq_{P/r} x.$$

So $x \sim_{P/r} y$. As $r' \triangleleft P/r, x \sim_P y: s \triangleleft P$.

Let us assume that $xs'y$. Then xry , so, as $r \triangleleft y$, there exists a path from x to y in the Hasse graph of P , made of vertices all r -equivalent to x and y . If x' and y' are two elements of this path, Then $x'r'y'$, so $x' \leq_{G/r} y'$ and finally $x' \leq_{(P/r)/r'} y'$. As $r' \triangleleft P/r, x'r'y'$, so xsy . So the elements of this path are all $P|s$ -equivalent: the equivalence classes of s' are $P|s$ -connected.

Let us assume that $x \sim_{(P|s)/s'} y$. There exist $x_1, x'_1, \dots, x_k, x'_k$ and $y_1, y'_1, \dots, y_l, y'_l$ such that:

$$x \leq_{P|r'} x_1 r x'_1 \leq_{P|r'} \dots \leq_{P|r'} x_k r x'_k \leq_{P|r'} y, \quad y \leq_{P|r'} y_1 r y'_1 \leq_{P|r'} \dots \leq_{P|r'} y_l r y'_l \leq_{P|r'} x.$$

Then:

$$x \leq_P x_1 r x'_1 \leq_P \dots \leq_P x_k r x'_k \leq_P y, \quad y \leq_P y_1 r y'_1 \leq_P \dots \leq_P y_l r y'_l \leq_P x,$$

So $x \leq_{P/r} y$ and $y \leq_{P/r} x$. As $r \triangleleft P, xry$, so $xs'y$: we obtain that $s' \triangleleft P|s$.

G is well-defined: let $(s, s') \in B$ and let us put $G(s, s') = (r, r')$. The equivalence classes of r are $P|s$ -connected, so are P -connected. Let us assume that $x \sim_{P/r} y$. There exist $x_1, x'_1, \dots, x_k, x'_k$ and $y_1, y'_1, \dots, y_l, y'_l$ such that:

$$x \leq_P x_1 s' x'_1 \leq_P \dots \leq_P x_k s' x'_k \leq_P y, \quad y \leq_P y_1 s' y'_1 \leq_P \dots \leq_P y_l s' y'_l \leq_P x.$$

As the equivalence classes of s' are $P|s$ -connected, all this elements are in the same connected component of $P|s$, so are s -equivalent:

$$x \leq_{P|s} x_1 s' x'_1 \leq_{P|s} \dots \leq_{P|s} x_k s' x'_k \leq_{P|s} y, \quad y \leq_{P|s} y_1 s' y'_1 \leq_{P|s} \dots \leq_{P|s} y_l s' y'_l \leq_{P|s} x.$$

Hence, $x \sim_{(P|s)/s'} y$, so as $s' \triangleleft P | s$, $x s' y$, so $x r y$: $r \triangleleft P$.

The equivalence classes of r' are the equivalence classes of s , so are P -connected and therefore P/r -connected. Let us assume that $x \sim_{(P/r)/r'} y$. Note that if $x' s' y'$, then x' and y' are in the same connected component of $P|s$, so $x' s y'$. By the definition of $\leq_{P/s'}$ as a transitive closure, using this observation, we obtain:

$$x \leq_P x_1 s x'_1 \leq_P \dots \leq_P x_k s x'_k \leq_P y, \quad y \leq_P y_1 s y'_1 \leq_P \dots \leq_P y_l s y'_l \leq_P x.$$

So $x \sim_{P/s} y$. As $s \triangleleft P$, $x s y$, so $x r' y$: $r' \triangleleft P/r$.

Clearly, F and G are inverse bijections.

Second step. Let $(r, r') \in A$ and let $F(r, r') = (s, s')$. Note that if $x r y$, then $x / \sim_{P/r} y$, so $x / \sim_{(P/r)/r'} y$, so $x r' y$ as $r' \triangleleft P/r$. Then:

$$\begin{aligned} \leq_{(P/r)/r'} &= \text{transitive closure of } ((x r' y) \text{ or } (x \leq_{P/r} y)) \\ &= \text{transitive closure of } ((x r' y) \text{ or } (x \leq_P y) \text{ or } (x \leq_r y)) \\ &= \text{transitive closure of } ((x r' y) \text{ or } (x \leq_P y)) \\ &= \text{transitive closure of } ((x s y) \text{ or } (x \leq_P y)) \\ &= \leq_{P/s}. \end{aligned}$$

So $P/s = (P/r)/r'$.

$$\begin{aligned} \leq_{(P|s)/s'} &= \text{transitive closure of } ((x s' y) \text{ or } (x \leq_{P|s} y)) \\ &= \text{transitive closure of } ((x r y) \text{ or } (x \leq_{P|r'} y)) \\ &= \text{transitive closure of } ((x r y) \text{ or } ((x \leq_P y) \text{ and } (x r' y))) \\ &= \text{transitive closure of } (((x r y) \text{ or } (x \leq_P y)) \text{ and } ((s r y) \text{ or } (x r' y))) \\ &= \text{transitive closure of } ((x \leq_{P/r} y) \text{ and } (s r' y)) \\ &= \leq_{(P/r)|r'}. \end{aligned}$$

So $(P|s)/s' = (P/r)|r'$. For all x, y :

$$\begin{aligned} x \leq_{(P|s)/s'} y &\iff (x \leq_{P|s} y) \text{ and } (x s' y) \\ &\iff (x \leq_P y) \text{ and } x s y \text{ and } (x s' y) \\ &\iff (x \leq_P y) \text{ and } x r' y \text{ and } (x r y) \\ &\iff (x \leq_P y) \text{ and } (x r y) \\ &\iff x \leq_{P|r} y. \end{aligned}$$

So $(P|s)|s' = P|r$. Finally:

$$\begin{aligned} (\delta \otimes Id) \circ \delta(P) &= \sum_{(r, r') \in A} (P/r)/r' \otimes (P/r)|r' \otimes P|r \\ &= \sum_{(s, s') \in B} P/s \otimes (P|s)/s' \otimes (P|s)|s' \\ &= (Id \otimes \delta) \circ \delta(P). \end{aligned}$$

So $\mathcal{H}_{\mathbf{QP}}$ is a bialgebra.

Let P be a quasi-poset. If P is discrete, then $\delta(P) = P \otimes P$, so $(\varepsilon' \otimes Id) \circ \delta(P) = (Id \otimes \varepsilon') \circ \Delta(P) = P$. If P is not discrete, there are three types of relations $\sim \triangleleft P$:

1. The equivalence classes of \sim are the connected components of P : in this case, $P| \sim = P$ and $P/ \sim = P_1$ is discrete.
2. $\sim = \sim_P$: in this case, $P/ \sim = P$ and $P| \sim = P_2$ is discrete.
3. \sim is not one of two preceding relations: in this case, nor P/ \sim , nor $P| \sim$ is discrete.

So:

$$\delta(P) - P_1 \otimes P - P \otimes P_2 \in Ker(\varepsilon') \otimes Ker(\varepsilon'),$$

which implies that $(\varepsilon' \otimes Id) \circ \delta(P) = (Id \otimes \varepsilon') \circ \Delta(P) = P$. □

Examples. If $\{a, b\} = \{1, 2\}$ and $\{i, j, k\} = \{1, 2, 3\}$:

$$\begin{aligned} \delta(\cdot_1) &= \cdot_1 \otimes \cdot_1, \\ \delta(\mathbf{1}_a^b) &= \mathbf{1}_a^b \otimes \cdot_{a \cdot b} + \cdot_{a, b} \otimes \mathbf{1}_a^b, \\ \delta({}^j \mathbf{V}_i^k) &= {}^j \mathbf{V}_i^k \otimes \cdot_{i \cdot j \cdot k} + \mathbf{1}_{i, j}^k \otimes \mathbf{1}_i^j \cdot k + \mathbf{1}_{i, k}^j \otimes \mathbf{1}_i^k \cdot j + \cdot_{i, j, k} \otimes {}^j \mathbf{V}_i^k, \\ \delta({}_j \mathbf{\Lambda}_k^i) &= {}_j \mathbf{\Lambda}_k^i \otimes \cdot_{i \cdot j \cdot k} + \mathbf{1}_k^{i, j} \otimes \mathbf{1}_j^i \cdot k + \mathbf{1}_j^{i, k} \otimes \mathbf{1}_k^i \cdot j + \cdot_{i, j, k} \otimes {}_j \mathbf{\Lambda}_k^i, \\ \delta(\mathbf{!}_i^k) &= \mathbf{!}_i^k \otimes \cdot_{i \cdot j \cdot k} + \mathbf{1}_{i, j}^k \otimes \mathbf{!}_i^j \cdot k + \mathbf{!}_i^{j, k} \otimes \cdot_i \mathbf{!}_j^k + \cdot_{i, j, k} \otimes \mathbf{!}_i^k. \end{aligned}$$

Remarks.

1. δ is the internal coproduct of [8].
2. $(\mathcal{H}_{\mathbf{QP}}, m, \delta)$ is not a Hopf algebra: for all $n \geq 1$, $\delta(\cdot_n) = \cdot_n \otimes \cdot_n$, and \cdot_n has no inverse in $\mathcal{H}_{\mathbf{QP}}$.
3. This coproduct is also compatible with the map $\llbracket \cdot \rrbracket$, so we obtain a bialgebra structure on $\mathcal{H}_{\mathbf{qp}}$ with the coproduct defined by:

$$\delta(\llbracket P \rrbracket) = \sum_{\sim \triangleleft P} \llbracket P/ \sim \rrbracket \otimes \llbracket P| \sim \rrbracket.$$

4. $\mathcal{H}_{\mathbf{P}}$ and $\mathcal{H}_{\mathbf{p}}$ are not stable under δ , as if P is a poset and $\sim \triangleleft P$, P/ \sim is not necessarily a poset (although $P| \sim$ is). However, there is a way to define a coproduct $\bar{\delta} = (\xi \otimes Id) \circ \delta$ on $\mathcal{H}_{\mathbf{p}}$:

$$\forall P \in \mathbf{P}(n), \bar{\delta}(\llbracket P \rrbracket) = \sum_{\sim \triangleleft P} \overline{\llbracket P/ \sim \rrbracket} \otimes \llbracket P| \sim \rrbracket.$$

$(\mathcal{H}_{\mathbf{p}}, m, \bar{\delta})$ is a quotient of $(\mathcal{H}_{\mathbf{qp}}, m, \delta)$ through the map ξ .

2.4 Cointeractions

Theorem 18 *We consider the map:*

$$\rho = (Id \otimes \llbracket \cdot \rrbracket) \circ \delta : \begin{cases} \mathcal{H}_{\mathbf{QP}} & \longrightarrow \mathcal{H}_{\mathbf{QP}} \otimes \mathcal{H}_{\mathbf{qp}} \\ P \in \mathbf{QP} & \longrightarrow \sum_{\sim \triangleleft P} (P/ \sim) \otimes \llbracket P| \sim \rrbracket. \end{cases}$$

The bialgebras $(\mathcal{H}_{\mathbf{QP}}, m, \Delta)$ and $(\mathcal{H}_{\mathbf{qp}}, m, \delta)$ are in cointeraction via ρ .

Proof. By composition, ρ is an algebra morphism. Let us take $P \in \mathbf{QP}(n)$. We put:

$$A = \{(r, O) \mid r \triangleleft P, O \in \text{top}(P/r)\}, \quad B = \{(O, s, s') \mid O \in \text{top}(P), s \triangleleft P_{[n] \setminus O}, s' \triangleleft P_{|O}\}.$$

First step. We define a map $F : A \longrightarrow B$, sending (r, O) to (O, s, s') , by:

- xy if xry and x, y are in the same connected component of $P_{|[n] \setminus O}$.
- $xs'y$ if xry and x, y are in the same connected component of $P_{|O}$.

Let us prove that F is well-defined. Let us take $x, y \in [n]$, with $x \in O$ and $x \leq_P y$. Then $x \leq_{P/r} y$. as O is an open set of P/r , $y \in O$: O is an open set of P . By definition, the equivalence classes of s' are the intersection of the equivalence classes of r and of the connected components of O . As O is a union of equivalence classes of r , they are $P_{|O}$ -connected. If $x \sim_{P_{|O}/s'} y$, then $x \sim_{P/r} y$ and x and y are in the same connected component of O . As $r \triangleleft r$, xry , so $xs'y$: $s' \triangleleft P_{|O}$. Similarly, $s \triangleleft P_{|[n] \setminus O}$.

Second step. We define a map $G : B \longrightarrow A$, sending (O, s, s') to (O, r, r') , by:

$$xry \text{ if } (x, y \notin O \text{ and } xsy) \text{ or } (x, y \in O \text{ and } xs'y).$$

Let us prove that G is well-defined. Let $x, y \in [N]$, with $x \in O$ and $x \leq_{P/r} y$. There exists $x_1, x'_1, \dots, x_k, x'_k$ such that:

$$x \leq_P x_1 r x'_1 \leq_P \dots \leq_P x_k r x'_k \leq_P y.$$

As O is an open set of P , $x_1 \in O$; by definition of r , $x'_1 \in O$. Iterating, we obtain that $x_2, x'_2, \dots, x_k, x'_k, y \in O$. So O is open in P/r .

Let us assume that xry . Then $x, y \in O$ or $x, y \notin O$. As $s \triangleleft P_{|[n] \setminus O}$ and $P_{|O}$, there exists a path from x to y in the Hasse graph of P formed by elements s - or s' - equivalent to x and y , so the equivalence classes of r are P -connected.

Let us assume that $x \sim_{P/r} y$. here exists $x_1, x'_1, \dots, x_k, s'_k$ and $y_1, y'_1, \dots, y_l, y_l$ such that:

$$x \leq_P x_1 r x'_1 \leq_P \dots \leq_P x_k r x'_k \leq_P y, \quad y \leq_P y_1 r y'_1 \leq_P \dots \leq_P y_l r y'_l \leq_P x.$$

If $x, y \in O$, then all these elements are in O , so $x \sim_{P_{|O}/s'} y$, and then $xs'y$, so xry . If $x, y \notin O$, as O is an open set, none of these elements is in O , so $x \sim_{P_{|[n] \setminus O}/s} y$, so xsy and finally xry : $r \triangleleft P$.

Third step. Let $(r, O) \in A$. We put $F(r, O) = (O, s, s')$ and $G(O, s, s') = (\tilde{r}, O)$. If xry , as O is an open set of P/r , both x and y are in O or both are not in O . Hence, xsy or $xs'y$, so $x\tilde{r}y$.

If $x\tilde{r}y$, then xsy or $xs'y$, so xry : $\tilde{r} = r$ and $G \circ F = Id_A$.

Let $(O, s, s') \in B$. We put $G(O, s, s') = (r, O)$ and $F(r, O) = (O, \tilde{s}, \tilde{s}')$. If xsy , then x and y are in the same connected component of $[n] \setminus O$ as $s \triangleleft P_{|[n] \setminus O}$ and xry , so $x\tilde{s}y$. If $x\tilde{s}y$, then xry , so xsy : we obtain that $\tilde{s} = s$. Similarly, $\tilde{s}' = s'$, which proves that $F \circ G = Id_B$.

We proved that F and G are inverse bijections. Let $(r, O) \in A$ and $(O, s, s') = F(O, r)$.

$$\begin{aligned} \leq_{(P/r)_{|[n] \setminus O}} &= \text{transitive closure of } (xry \text{ and } x \leq_P y) \text{ restricted to } [n] \setminus O \\ &= \text{transitive closure of } (xry \text{ and } x \leq_{P_{|[n] \setminus O}} y) \\ &= \text{transitive closure of } (xsy \text{ and } x \leq_{P_{|[n] \setminus O}} y) \\ &= \leq_{P_{|[n] \setminus [n]}/s}. \end{aligned}$$

So $(P/r)_{|[n] \setminus O} = P_{|[n] \setminus O}/s$. Similarly, $(P/r)_{|O} = P_{|O}/s'$.

Let us now consider $P|_R$. Its connected components are the equivalence classes of r , that is to say the equivalence classes of s and s' ; for any such equivalence class I , $(P|_R)|_I = P|_I$. So $P|_R$ is the disjoint union of $(P|_{[n]\setminus O})|_s$ and $(P|_O)|_{s'}$, and therefore is isomorphic to $Std(P|_{[n]\setminus O})|_s Std((P|_O)|_{s'})$, but not equal, because of the reindexation induced by the standardization. Hence, $[P|_R] = [(P|_{[n]\setminus O})|_s] [(P|_O)|_{s'}]$.

Finally:

$$\begin{aligned} (\Delta \otimes Id) \circ \rho(P) &= \sum_{(r,O) \in A} (G/r)|_{[n]\setminus O} \otimes (G/r)|_O \otimes [G|_r] \\ &= \sum_{(O,s,s') \in B} (P|_{[n]\setminus O})|_s \otimes (P|_O)|_{s'} \otimes [(P|_{[n]\setminus O})|_s] [(P|_O)|_{s'}] \\ &= m_{2,4}^3 \circ (\rho \otimes \rho) \circ \Delta(P). \end{aligned}$$

Moreover, $(\varepsilon \otimes Id) \circ \rho(P) = \delta_{P,1} 1 \otimes 1 = \varepsilon(P) 1 \otimes 1$. \square

Remark. As noticed in [8], $(\mathcal{H}_{\mathbf{QP}}, m, \Delta)$ and $(\mathcal{H}_{\mathbf{QP}}, m, \delta)$ are not in cointeraction through δ .

Taking the quotient through $[\]$:

Corollary 19 *The bialgebras $(\mathcal{H}_{\mathbf{qp}}, m, \Delta)$ and $(\mathcal{H}_{\mathbf{qp}}, m, \delta)$ are in cointeraction via δ . Moreover, $\mathcal{H}_{\mathbf{qp}}$ is given a graduation by:*

$$\forall n \geq 0, (\mathcal{H}_{\mathbf{qp}})_n = Vect(P \in \mathbf{qp} \mid cl(P) = n).$$

With this graduation, hypotheses 1–5 of section 1.3 are satisfied.

Proof. Here, V is the space generated by the set of connected quasi-posets; the basis $(g_i)_{i \in I}$ of group-like elements of $(\mathcal{H}_{\mathbf{qp}})_1$ is $(\bullet_n)_{n \geq 1}$. \square

We denote by $M_{\mathbf{qp}}$ the monoid of characters of $(\mathcal{H}_{\mathbf{qp}}, m, \delta)$. Its product is denoted by $*$. Using proposition 4 on $\mathcal{H}_{\mathbf{QP}}$:

Corollary 20 *Let $\lambda \in M_{\mathbf{qp}}$. The following map is a Hopf algebra endomorphism:*

$$\phi_\lambda : \begin{cases} (\mathcal{H}_{\mathbf{QP}}, m, \Delta) & \longrightarrow (\mathcal{H}_{\mathbf{QP}}, m, \Delta) \\ P \in \mathbf{QP} & \longrightarrow \sum_{\sim \triangleleft P} \lambda_{[P|\sim]} P / \sim. \end{cases}$$

It is bijective if, and only if, for all $n \geq 1$, $\lambda_{\bullet_n} \neq 0$. If this holds, $\phi_\lambda^{-1} = \phi_{\lambda^{-1}}$.

Proof. $\phi_\lambda = Id \leftarrow \lambda$, so is an element of $E_{\mathcal{H}_{\mathbf{QP}} \rightarrow \mathcal{H}_{\mathbf{QP}}}$.

\implies . For all $n \geq 0$, $\phi_\lambda(\bullet_n) = \lambda_{\bullet_n} \bullet_n$. As ϕ_λ is injective, $\lambda_{\bullet_n} \neq 0$.

\impliedby . By proposition 5, λ is invertible in $M_{\mathbf{qp}}$: let us denote its inverse by μ . Then, by proposition 4:

$$\phi_\lambda \circ \phi_\mu = Id \leftarrow (\lambda * \mu) = Id \leftarrow \varepsilon' = Id.$$

Similarly, $\phi_\mu \circ \phi_\lambda = Id$. \square

3 Ehrhart polynomials

Notations. For all $k \geq 0$, we denote by H_k the k -th Hilbert polynomial:

$$H_k(X) = \frac{X(X-1)\dots(X-k+1)}{k!}.$$

3.1 Definition

Definition 21 Let $P \in \mathbf{QP}(n)$ and $k \geq 1$. We put:

$$\begin{aligned} L_P(k) &= \{f : [n] \rightarrow [k] \mid \forall i, j \in [n], i \leq_P j \implies f(i) \leq f(j)\}, \\ L_P^{str}(k) &= \{f \in L_P(k) \mid \forall i, j \in [n], (i \leq_P j \text{ and } f(i) = f(j)) \implies i \sim_P j\}, \\ W_P(k) &= \{w \in L_P(k) \mid w([n]) = [k]\}, \\ W_P^{str}(k) &= \{w \in L_P^{str}(k) \mid w([n]) = [k]\}. \end{aligned}$$

By convention:

$$L_P(0) = L_P^{str}(0) = W_P(0) = W_P^{str}(0) = \begin{cases} \emptyset & \text{if } P \neq 1, \\ \{1\} & \text{if } P = 1. \end{cases}$$

We also put:

$$L_P = \bigcup_{k \geq 0} L_P(k), \quad L_P^{str} = \bigcup_{k \geq 0} L_P^{str}(k), \quad W_P = \bigsqcup_{k \geq 0} W_P(k), \quad W_P^{str} = \bigsqcup_{k \geq 0} W_P^{str}(k).$$

Note that the elements of W_P and W_P^{str} are packed words (see definition 40).

Proposition 22 Let $P \in \mathbf{QP}$. There exist unique polynomials ehr_P and $ehr_P^{str} \in \mathbb{Q}[X]$, such that for $k \geq 0$:

$$ehr_P(k) = \#L_P(k), \quad ehr_P^{str}(k) = \#L_P^{str}(k).$$

Proof. This is obvious if $P = 1$, with $ehr_1(X) = ehr_1^{str}(X) = 1$. Let us assume that $P \in \mathbf{QP}(n)$, $n \geq 1$. Note that if $i > n$, $W_P(i) = 0$. For all $k \geq 1$:

$$\#L_P(k) = \sum_{i=1}^k \#W_P(i) \binom{k}{i} = \sum_{i=1}^k \#W_P(i) H_i(k) = \sum_{i=1}^n \#W_P(i) H_i(k).$$

So:

$$ehr_P(X) = \sum_{i=1}^n \#W_P(i) H_i(X).$$

Moreover, if $k = 0$:

$$ehr_P(0) = \sum_{i=1}^n \#W_P(i) H_i(0) = \#L_P(0).$$

In the same way:

$$ehr_P^{str}(X) = \sum_{i=1}^n \#W_P^{str}(i) H_i(X).$$

These are indeed elements of $\mathbb{Q}[X]$. □

Remarks.

1. Let $P, Q \in \mathbf{QP}(n)$.

- If they are isomorphic, then $ehr_P(k) = ehr_Q(k)$ for all $k \geq 1$, so $ehr_P = ehr_Q$.
- If $w \in L_P$, for all $x, y \in P$ such that $x \sim_P y$, then $w(x) \leq w(y)$ and $w(y) \leq w(x)$, so $w(x) = w(y)$: w goes through the quotient by \sim_P . We obtain in this way a bijection from $L_P(k)$ to $L_{\overline{P}}(k)$ for all k , so $ehr_P = ehr_{\overline{P}}$. Similarly, $ehr_P^{str} = ehr_{\overline{P}}^{str}$.

So $f \in L_P(k+l)$, and $v(f) = (f_1, f_2)$: v is surjective, and finally bijective. Moreover, if $f_1 \in L_{[n]\setminus O}^{str}(k)$ and $f_2 \in L_{P|_O}^{str}(l)$, then $f = v^{-1}(f_1, f_2) \in L_P^{str}(k+l)$. Finally:

$$\begin{aligned} f(L_P(k+l)) &= \bigsqcup_{O \in Top(P)} L_{P|_{[n]\setminus O}}(k) \times L_{P|_O}(l), \\ f(L_P^{str}(k+l)) &= \bigsqcup_{O \in Top(P)} L_{P|_{[n]\setminus O}}^{str}(k) \times L_{P|_O}^{str}(l). \end{aligned}$$

Taking the cardinals, we obtain the announced result.

Second step. Let $P \in \mathbf{P}(m)$, $Q \in \mathbf{P}(n)$, and $f : [m+n] \rightarrow [k]$. Then $f \in L_{PQ}(k)$ if, and only if, $f|_{[m]} \in L_P(k)$ and $Std(f|_{[m+n]\setminus[m]}) \in L_Q(k)$. So $ehr_{PQ}(k) = ehr_P(k)ehr_Q(k)$, and then $ehr_{PQ}(X) = ehr_P(X)ehr_Q(X)$: ehr is an algebra morphism.

Let P be a finite poset, and $k, l \geq 0$. By the first step:

$$\begin{aligned} (ehr \otimes ehr) \circ \Delta(P)(k, l) &= \sum_{O \in Top(P)} ehr_{P|_{[m]\setminus O}}(k)ehr_{P|_O}(l) \\ &= ehr_P(k+l) \\ &= \Delta \circ ehr(P)(k, l). \end{aligned}$$

As this is true for all $k, l \geq 1$, $(ehr \otimes ehr) \circ \Delta(P) = \Delta \circ ehr(P)$. Moreover:

$$\varepsilon \circ ehr(P) = ehr_P(0) = \begin{cases} 1 & \text{if } P = 1, \\ 0 & \text{otherwise,} \end{cases}$$

so $\varepsilon \circ ehr = \varepsilon$.

The proof is similar for ehr^{str} . □

3.2 Recursive computation of ehr and ehr^{str}

Let us recall this classical result:

Lemma 24 *We consider the following maps:*

$$L : \begin{cases} \mathbb{K}[X] & \longrightarrow \mathbb{K}[X] \\ H_k(X) & \longrightarrow H_{k+1}(X). \end{cases}$$

The map L is injective, and $L(\mathbb{K}[X]) = \mathbb{K}[X]_+$. Moreover, for all $P \in \mathbb{K}[X]$, for all $n \geq 0$:

$$L(P)(n+1) = P(0) + \dots + P(n).$$

Proof. Let us consider $P = H_k(X)$. For all $n \geq 0$:

$$\begin{aligned} H_k(0) + \dots + H_k(n) &= \binom{0}{k} + \dots + \binom{n}{k} \\ &= \binom{k}{k} + \dots + \binom{n}{k} \\ &= \binom{n+1}{k+1} \\ &= H_{k+1}(n+1) \\ &= L(H_k)(n+1). \end{aligned}$$

By linearity, for any $P \in \mathbb{K}[X]$, $L(P)(n+1) = P(0) + \dots + P(n)$ for all $n \geq 1$. □

Proposition 25 Let $P \in \mathbf{P}(n)$.

$$\begin{aligned} ehr_P(X) &= L \left(\sum_{\emptyset \neq O \in Top(P)} ehr_{P_{|[n] \setminus O}}(X) \right), \\ ehr_P^{str}(X) &= L \left(\sum_{\emptyset \neq O \in Top(P), \text{ discrete}} ehr_{P_{|[n] \setminus O}}^{str}(X) \right). \end{aligned}$$

Proof. Let $n \geq 1$. As $L_Q(1)$ is reduced to a singleton for all finite poset Q :

$$\begin{aligned} ehr_P(n+1) &= \sum_{O \in Top(P)} ehr_{P_{|[n] \setminus O}}(n) ehr_{P_O}(1) \\ &= \sum_{\emptyset \neq O \in Top(P)} ehr_{P_{|[n] \setminus O}}(n) + ehr_P(n). \end{aligned}$$

We put:

$$Q(X) = \sum_{\emptyset \neq O \in Top(P)} ehr_{P_{|[n] \setminus O}}(X).$$

In particular:

$$Q(0) = \sum_{\emptyset \neq O \in Top(P)} ehr_{P_{|[n] \setminus O}}(0) = ehr_{\emptyset}(0) + 0 = 1 = ehr_P(1).$$

Then:

$$\begin{aligned} ehr_P(n+1) &= Q(n) + ehr_P(n) \\ &= Q(n) + Q(n-1) + ehr_P(n-1) \\ &\vdots \\ &= Q(n) + Q(n-1) + \dots + Q(1) + ehr_P(1) \\ &= Q(n) + \dots + Q(1) + Q(0) \\ &= L(Q)(n+1). \end{aligned}$$

So $ehr_P = L(Q)$.

For ehr_P^{str} , observe that $ehr_Q^{str}(1) = 1$ if Q is discrete, and 0 otherwise, which implies:

$$ehr_P^{str}(n+1) = \sum_{\emptyset \neq O \in Top(P), \text{ discrete}} ehr_{P_{|[n] \setminus O}}^{str}(n) + ehr_P^{str}(n).$$

The end of the proof is similar. □

Examples.

$$\begin{aligned}
ehr_{\bullet}(X) &= H_1(X) = X, \\
ehr_{\mathbf{1}}(X) &= H_1(X) + H_2(X) = \frac{X(X+1)}{2}, \\
ehr_{\mathbf{V}}(X) &= ehr_{\mathbf{\wedge}}(X) = H_1(X) + 3H_2(X) + 2H_3(X) = \frac{X(X+1)(2X+1)}{6}, \\
ehr_{\mathbf{\dagger}}(X) &= H_1(X) + 2H_2(X) + H_3(X) = \frac{X(X+1)(X+2)}{6}; \\
\\
ehr_{\bullet}^{str}(X) &= H_1(X) = X, \\
ehr_{\mathbf{\dagger}}^{str}(X) &= H_2(X) = \frac{X(X-1)}{2}, \\
ehr_{\mathbf{V}}^{str}(X) &= ehr_{\mathbf{\wedge}}^{str}(X) = H_2(X) + 2H_3(X) = \frac{X(X-1)(2X-1)}{6}, \\
ehr_{\mathbf{\dagger}}^{str}(X) &= H_3(X) = \frac{X(X-1)(X-2)}{6}.
\end{aligned}$$

3.3 Characterization of quasi-posets by packed words

Lemma 26 *Let $P \in \mathbf{QP}(n)$ and let I_1, \dots, I_k be distinct minimal classes of the poset \overline{P} ; let $w' \in W_{P_{[n] \setminus (I_1 \sqcup \dots \sqcup I_k)}}^{str}$. The following map belongs to W_P^{str} :*

$$w : \begin{cases} [n] & \longrightarrow \mathbb{N}^* \\ x \in I_p, 1 \leq p \leq k & \longrightarrow p \\ x \notin I_1 \sqcup \dots \sqcup I_k & \longrightarrow w'(x). \end{cases}$$

Proof. Let us assume that $i \leq_P j$.

- If $i \in I_p$, as I_p is a minimal class of \overline{P} , $j \in I_p$ or $j \notin I_1 \sqcup \dots \sqcup I_k$. In the first case, $w(i) = w(j)$; in the second case, $w(i) \leq k < w(j)$. If moreover $w(i) = w(j)$, then necessarily $j \in I_p$, so $i \sim_P j$.
- If $i \notin I_1 \sqcup \dots \sqcup I_k$, as $i \leq_P j$, $j \notin I_1 \sqcup \dots \sqcup I_k$, so $i \leq_{P_{[n] \setminus (I_1 \sqcup \dots \sqcup I_k)}} j$ and $w'(i) \leq w'(j)$, so $w'(i) \leq w'(j)$. If moreover $w(i) = w(j)$, then $w'(i) = w'(j)$, so $i \sim_{P_{[n] \setminus (I_1 \sqcup \dots \sqcup I_k)}} j$ and finally $i \sim_P j$.

As a conclusion, $w \in W_P^{str}$. □

Remark. This lemma implies that W_P^{str} is non-empty for any non-empty quasi-poset P .

Proposition 27 *Let $P = ([n], \leq_P)$ be a quasi-poset and let $i, j \in [n]$. The following assertions are equivalent:*

1. $i \leq_P j$.
2. For all $w \in L_P$, $w(i) \leq w(j)$.
3. For all $w \in L_P^{str}$, $w(i) \leq w(j)$.
4. For all $w \in W_P$, $w(i) \leq w(j)$.
5. For all $w \in W_P^{str}$, $w(i) \leq w(j)$.

Proof. Obviously:

$$\begin{array}{ccccc} 1. & \Longrightarrow & 2. & \Longrightarrow & 3. \\ & & \Downarrow & & \Downarrow \\ & & 4. & \Longrightarrow & 5. \end{array}$$

It is enough to prove that $5. \implies 1.$ We proceed by induction on n . If $n = 1$, there is nothing to prove. Let us assume the result at all ranks $< n$. Let $i, j \in [n]$, such that we do not have $i \leq_P j$. Let us prove that there exists $w \in W_P^{str}$, such that $w(i) > w(j)$. There exists a minimal element $k \in [n]$, such that $k \leq_P j$; let I be the class of k in \bar{P} . By hypothesis on i , i and k are not equivalent for \sim_P , so $i \notin I$. If $j \in I$, let us choose an element $w' \in W_{P_{[n] \setminus I}}^{str}$; if $j \notin I$, then by the induction hypothesis, there exists $w' \in W_{P_{[n] \setminus I}}^{str}$, such that $w'(i) > w'(j)$. By lemma 26, the following map is an element of W_P^{str} :

$$w : \begin{cases} [n] & \longrightarrow \mathbb{N} \\ x \in I & \longrightarrow 1 \\ x \notin I & \longrightarrow w'(x) + 1 \end{cases}$$

If $j \in I$, then $w(j) = 1 < w(i)$; if $j \notin I$, $w(i) = w'(i) + 1 > w'(j) + 1 = w(j)$. In both cases, $w(i) > w(j)$. \square

3.4 Link with linear extensions

Let $P \in \mathbf{QP}(n)$. Linear extensions, as defined in [10], are elements of W_P^{str} : they are the elements $f \in W_P^{str}$ such that

$$\forall i, j \in [n], f(i) = f(j) \iff i \sim_P j.$$

It may happens that not all elements of W_P^{str} are linear extensions. For example, if $P = {}^2\mathbf{V}_1^3$, $W_P^{str}(3) = \{(123), (132), (122)\}$, and (122) is not a linear extension of P . The set of linear extensions of P will be denoted by E_P .

Definition 28 Let w and w' be two packed words of the same length n . We shall say that $w \leq w'$ if:

$$\forall i, j \in [n], w(i) < w(j) \implies w'(i) < w'(j).$$

Proposition 29 Let $P \in \mathbf{QP}(n)$. Then:

$$W_P = \bigcup_{w \in E_P} \{w' \mid w' \leq w\}.$$

This union may be not disjoint. Moreover, the maximal elements of W_P for the order of definition 28 are the elements of E_P .

Proof. \subseteq . Let $w \in W_P$. For all $1 \leq p \leq \max(w)$, we put $I_p = w^{-1}(p)$. Let f_p be a linear extension of $P_{|I_p}$. Let us consider:

$$f : \begin{cases} [n] & \longrightarrow \mathbb{N} \\ i & \longrightarrow \max(f_1) + \dots + \max(f_{p-1}) + f_p(i) \text{ if } i \in I_p. \end{cases}$$

By construction, if $w(i) < w(j)$, then $f(i) < f(j)$: $w \leq f$. Let us prove that $f \in E_P$.

If $i \leq_P j$, then as $w \in W_P$, $w(i) \leq w(j)$. If $w(i) = w(j) = p$, then $i \leq_{P_{|I_p}} j$, so $f_p(i) \leq f_p(j)$, and $f(i) \leq f(j)$. If $w(i) < w(j)$, then $f(i) < f(j)$.

If $f(i) = f(j)$, then $w(i) = w(j) = p$, and $f_p(i) = f_p(j)$. As $f_p \in E_{P_{|I_p}}$, $i \sim_{P_{|I_p}} j$, so $i \sim_P j$.

\supseteq . Let $w \in E_P$ and $w' \leq w$. If $i \leq_P j$, then $w(i) \leq w(j)$ as w is a linear extension of P . As $w' \leq w$, $w'(i) \leq w'(j)$, so $w' \in W_P$.

Let w be a maximal element of W_P . There exists a linear extension w' of P , such that $w \leq w'$. As w is maximal, $w = w'$ is a linear extension of P . Conversely, if w is a linear extension of P and $w \leq w'$ in W_P , then as w is a linear extension of P , $\max(w) = cl(P)$. Moreover, as $w \leq w'$, $\max(w) \leq \max(w')$. As $w' \in W_P$, $\max(w') \leq cl(P)$, which implies that $\max(w) = \max(w') = cl(P)$, and finally $w = w'$: w is a maximal element of W_P . \square

Example. For $P = {}^2\mathbf{V}_1^3$:

$$\begin{aligned} E_P &= \{(123), (132)\}; \\ W_P &= \{(123), (122), (112), (111)\} \cup \{(132), (122), (121), (111)\} \\ &= \{(123), (132), (122), (112), (121), (111)\}. \end{aligned}$$

Note that the two components of W_P are not disjoint.

Remark. A similar result is proved in [10] for T -partitions of a quasi-poset, generalizing Stanley's result [17] for P -partitions of posets; nevertheless, this is different here, as the union is not a disjoint one.

4 Characters associated to ehr and ehr^{str}

Recall that $(M_{\mathbf{qp}}, *)$ the monoid of characters of $(\mathcal{H}_{\mathbf{qp}}, m, \delta)$.

By theorems 7 and 23:

Proposition 30 *1. There exists a unique homogeneous Hopf algebra morphism ϕ_0 from $(\mathcal{H}_{\mathbf{qp}}, m, \Delta)$ to $(\mathbb{K}[X], m, \Delta)$ such that:*

$$\forall n \geq 1, \phi_0(\bullet_n) = X.$$

There exists a unique character $\lambda \in M_{\mathbf{qp}}$ such that for all $P \in \mathbf{qp}$,

$$\phi_0(P) = \lambda_P X^{cl(P)}.$$

2. There exist unique characters $\alpha, \alpha^{str} \in M_{\mathbf{qp}}$, such that:

$$ehr = \phi_0 \leftarrow \alpha, \quad ehr^{str} = \phi_0 \leftarrow \alpha^{str}.$$

Remarks.

1. Let $P \in \mathbf{qp}$. Then $ehr_P = ehr_{\overline{P}}$ and $ehr_P^{str} = ehr_{\overline{P}}^{str}$, so $\alpha_P = \alpha_{\overline{P}}$ and $\alpha_P^{str} = \alpha_{\overline{P}}^{str}$.
2. Still by theorem 7, there exists a unique homogeneous Hopf algebra morphism ϕ'_0 from $\mathcal{H}_{\mathbf{p}}$ to $\mathbb{K}[X]$ such that $\phi'_0(\bullet) = X$. By unicity, $\phi_0 = \phi'_0 \circ \Xi$, so for any $P \in \mathbf{qp}$, $\lambda_P = \lambda_{\overline{P}}$.

4.1 The character λ

Lemma 31 *For all $P \in \mathbf{P}(n)$, $n \geq 0$:*

$$\lambda_{[P]} = \begin{cases} 1 & \text{if } P = 1, \\ \frac{1}{n} \sum_{M \in \max(P)} \lambda_{[P]_{[n] \setminus \{M\}}} & \text{otherwise.} \end{cases} = \frac{1}{n} \sum_{m \in \min(P)} \lambda_{[P]_{[n] \setminus \{m\}}}$$

Proof. Let $P \in \mathbf{P}(n)$, with $n \geq 0$. Recall that $\pi : \mathbb{K}[X] \rightarrow \text{Vect}(X)$ is the canonical projection.

$$\begin{aligned}
(Id \otimes \pi) \circ \Delta \circ \phi_0([P]) &= \lambda_{[P]}(Id \otimes \pi) \circ \Delta(X^n) \\
&= \lambda_{[P]} n X^{n-1}; \\
= (Id \otimes \pi) \circ (\phi_0 \otimes \phi_0) \circ \Delta([P]) &= \sum_{O \in \text{Top}(P)} \lambda_{[P|_O]} \lambda_{[P_{[n] \setminus O}]} X^{|[n] \setminus O|} \pi(X^{|O|}) \\
&= \sum_{O \in \text{Top}(P), |O|=1} \lambda_{[P|_O]} \lambda_{[P_{[n] \setminus O}]} X^{n-1} \\
&= \sum_{M \in \text{max}(P)} \lambda_{[P_{[n] \setminus \{M\}}]} X^{n-1}.
\end{aligned}$$

This implies the first equality. The second is proved by considering $(\pi \otimes Id) \circ \Delta \circ \phi_0([P])$. \square

Remarks.

1. This lemma allows to inductively compute λ_P . This gives:

P	\cdot	\dagger	\vee	\wedge	\ddagger	Ψ	\mathfrak{A}	\Downarrow	\Uparrow	Υ	\mathfrak{A}	\ddagger	\mathbb{N}	\mathfrak{M}	\diamond
λ_P	1	$\frac{1}{2}$	$\frac{1}{3}$	$\frac{1}{3}$	$\frac{1}{6}$	$\frac{1}{4}$	$\frac{1}{4}$	$\frac{1}{8}$	$\frac{1}{8}$	$\frac{1}{12}$	$\frac{1}{12}$	$\frac{1}{24}$	$\frac{5}{24}$	$\frac{1}{6}$	$\frac{1}{12}$

2. If $P = (P, \leq)$ is a finite poset, we denote by P^{op} the opposite poset (P, \geq) . It is not difficult to deduce from this lemma that $\lambda_P = \lambda_{P^{op}}$.

Proposition 32 *Let $P \in \mathbf{P}(n)$. The number of elements of $W_P(n)$ of P is denoted by μ_P : in other words, μ_P is the number of bijections f from $[n]$ to $[n]$ such that for all $x, y \in [n]$,*

$$x \leq_P y \implies f(x) \leq f(y).$$

These bijections are called heap-orderings of P . Then, for any finite poset P , $\lambda_P = \frac{\mu_P}{n!}$.

Proof. Let us fix a non-empty finite poset $P \in \mathbf{P}(n)$. The set of heap-orderings of P is $HO(P) = W_P(n)$. We consider the map:

$$\begin{cases} HO(P) & \rightarrow \bigsqcup_{M \in \text{max}(P)} HO(P \setminus \{M\}) \\ f & \rightarrow f_{[n-1]} \in HO(P \setminus \{f^{-1}(n)\}). \end{cases}$$

It is not difficult to prove that this is a bijection. So:

$$\mu_P = \sum_{M \in \text{max}(P)} \mu_{P \setminus \{M\}}; \quad \frac{\mu_P}{n!} = \frac{1}{n} \sum_{M \in \text{max}(P)} \frac{\mu_{P \setminus \{M\}}}{|P \setminus \{M\}|!}.$$

An easy induction on $|P|$ then proves that $\lambda_P = \frac{\mu_P}{n!}$ for all P . \square

This formula can be simplified for rooted forests.

Definition 33 *Let P be a non-empty finite poset.*

1. We put:

$$P! = \prod_{i \in V(P)} \#\{j \in V(P) \mid i \leq_P j\}.$$

By convention, $1! = 1$.

2. We shall say that P is a rooted forest if P does not contain any subposet isomorphic to Λ .

For example, here are isoclasses of rooted forests of cardinality ≤ 4 :

$$1; \quad \cdot; \quad \cdot, \cdot; \quad \vee, \cdot, \cdot, \cdot, \dots; \quad \Psi, \downarrow, \Upsilon, \uparrow, \vee, \cdot, \cdot, \cdot, \cdot, \dots, \dots$$

Examples.

P	\cdot	\cdot, \cdot	\vee	Λ	\uparrow	Ψ	\downarrow	\downarrow	\uparrow	Υ	\downarrow	\uparrow	\mathbb{N}	\mathbb{N}	\diamond
$P!$	1	2	3	4	6	4	8	8	12	12	18	24	6	9	16

Proposition 34 For all finite poset P , $\lambda_P \geq \frac{1}{P!}$, with equality if, and only if, P is a rooted forest.

Proof. We proceed by induction on $n = |P|$. It is obvious if $n = 0$. Let us assume the result at all ranks $< n$.

$$\begin{aligned} \lambda_P &= \frac{1}{|P|} \sum_{m \in \min(P)} \lambda_{P \setminus \{m\}} \\ &\geq \frac{1}{|P|} \sum_{m \in \min(P)} \prod_{i \in V(P), i \neq m} \frac{1}{\#\{j \in V(P) \mid j \neq m, i \leq_P j\}} \\ &= \frac{1}{|P|} \sum_{m \in \min(P)} \prod_{i \in V(P), i \neq m} \frac{1}{\#\{j \in V(P) \mid i \leq_P j\}} \\ &= \frac{1}{P!} \frac{1}{|P|} \sum_{m \in \min(P)} \#\{j \in V(P) \mid m \leq_P j\}. \end{aligned}$$

For any $j \in A$, there exists $m \in \min(P)$ such that $m \leq_P j$, so:

$$\sum_{m \in \min(P)} \#\{j \in V(P) \mid m \leq_P j\} \geq |P|.$$

Consequently, $\lambda_P \geq \frac{1}{P!}$.

Let us assume that this is an equality. Then:

$$\sum_{m \in \min(P)} \#\{j \in V(P) \mid m \leq_P j\} = |P|.$$

Consequently, for all $j \in \min(P)$, there exists a unique $m \in \min(P)$ such that $m \leq_P j$. Moreover, for all $m \in \min(P)$, $\lambda_{P \setminus \{m\}} = \frac{1}{P \setminus \{m\}!}$. By the induction hypothesis, $P \setminus \{m\}$ is a rooted forest; this implies that P is also a rooted forest.

Let us assume that P is a rooted forest. For any $j \in V(P)$, there exists a unique $m \in \min(P)$ such that $m \leq_P j$, so:

$$\sum_{m \in \min(P)} \#\{j \in V(P) \mid m \leq_P j\} = |P|.$$

Moreover, for all $m \in \min(P)$, $P \setminus \{m\}$ is also a rooted forest. By the induction hypothesis, $\lambda_{P \setminus \{m\}} = \frac{1}{P \setminus \{m\}!}$. Hence, $\lambda_P = \frac{1}{P!}$. \square

4.2 The character α^{str}

Let us now apply theorem 8 to ehr and ehr^{str} :

Theorem 35 *For all finite connected quasi-poset P , we have:*

$$\alpha_P = \frac{d \, ehr_P}{dX}(0), \quad \alpha_P^{str} = \frac{d \, ehr_P^{str}}{dX}(0).$$

For any quasi-poset P :

$$ehr_P(X) = \sum_{\sim \triangleleft P} \frac{\mu_{P/\sim}}{cl(\sim)!} \alpha_{P|\sim} X^{cl(\sim)}, \quad ehr_P^{str}(X) = \sum_{\sim \triangleleft P} \frac{\mu_{P/\sim}}{cl(\sim)!} \alpha_{P|\sim}^{str} X^{cl(\sim)},$$

where $cl(\sim)$ is the number of equivalence classes of \sim .

Let us give a few values of α :

P	\cdot	\mathfrak{I}	\mathfrak{V}	\mathfrak{A}	\mathfrak{I}	\mathfrak{V}	\mathfrak{A}	\mathfrak{V}	\mathfrak{I}	\mathfrak{Y}	\mathfrak{A}	\mathfrak{I}	\mathfrak{N}	\mathfrak{M}	\mathfrak{D}
α_P	1	$\frac{1}{2}$	$\frac{1}{6}$	$\frac{1}{6}$	$\frac{1}{3}$	0	0	$\frac{1}{12}$	$\frac{1}{12}$	$\frac{1}{6}$	$\frac{1}{6}$	$\frac{1}{4}$	$\frac{1}{12}$	$\frac{1}{6}$	$\frac{1}{6}$

Lemma 36 *Let $P \in \mathbf{QP}$, not discrete. Then:*

$$(-1)^{cl(P)} ehr_P(-1) = ehr_P^{str}(1) = \begin{cases} 1 & \text{if } P \text{ is discrete,} \\ 0 & \text{otherwise.} \end{cases}$$

Proof. If P is discrete, then $ehr_P(X) = ehr_P^{str}(X) = X^{cl(P)}$ and the result is obvious. Let us assume that P is not discrete. There exists a unique map f from $[n]$ to $[1]$; as P is not discrete, $f \notin L_P^{str}(1)$, so $ehr_P^{str}(1) = 0$. We now proceed with $ehr_P(-1)$.

First step. Let us prove that $L(H_k(-X)) = -H_{k+1}(-X)$ for all $k \geq 0$. For all $l, n \geq 0$:

$$H_l(-n) = (-1)^l \binom{n+l-1}{l}.$$

For all $k, n \geq 0$:

$$\begin{aligned} L(H_k(-X))(n+1) &= H_k(0) + \dots + H_k(-n) \\ &= (-1)^k \sum_{i=0}^n \binom{i+k-1}{k} \\ &= (-1)^k \sum_{j=k}^{n+k-1} \binom{j}{k} \\ &= (-1)^k \binom{n+k}{k+1} \\ &= -H_{k+1}(-(n+1)). \end{aligned}$$

Second step. Let us prove that $L((X+1)\mathbb{K}[X]) \subseteq (X+1)\mathbb{K}[X]$. For all $k \geq 2$, let us put $H_k(-X) = X(X+1)L_k(X)$; $(L_k(X))_{k \geq 2}$ is a basis of $\mathbb{K}[X]$, which implies that $(H_k(-X))_{k \geq 2}$ is a basis of $X(X+1)\mathbb{K}[X]$, and that $(X+1) \sqcup (H_k(-X))_{k \geq 2}$ is a basis of $(X+1)\mathbb{K}[X]$. First:

$$L(X+1) = L(H_1(X) + H_0(X)) = H_2(X) + H_1(X) = \frac{X(X-1)}{2} + X = \frac{X(X+1)}{2} \in (X+1)\mathbb{K}[X];$$

if $k \geq 2$, by the first step, $L(H_k(-X)) = -H_{k+1}(-X) \in (X+1)\mathbb{K}[X]$.

Last step. We can replace P by \overline{P} , and we now assume that $P \in \mathbf{P}(n)$. There is nothing to prove if $n = 0, 1$. Let us assume the result at all rank $< n$. Then, by the second step and the induction hypothesis:

$$\begin{aligned}
ehr_{[P]}(-1) &= L \left(\sum_{\emptyset \neq O \in Top(P)} ehr_{[P_{[n] \setminus O}]}(X) \right)_{|X=-1} \\
&= L \left(\sum_{\substack{\emptyset \neq O \in Top(P) \\ P_{[n] \setminus O} \text{ discrete}}} ehr_{[P_{[n] \setminus O}]}(X) \right)_{|X=-1} \\
&= L \left(\sum_{[n] \neq J \subseteq min(P)} ehr_{[P_{[J]}]}(X) \right)_{|X=-1} \\
&= L \left(\sum_{J \subseteq min(P)} ehr_{[P_{[J]}]} J(X) \right)_{|X=-1} \\
&= L \left(\sum_{J \subseteq min(P)} X^{|J|} \right)_{|X=-1} \\
&= L \left(\underbrace{(1+X)^{|min(P)|}}_{\in (X+1)\mathbb{K}[X]} \right)_{|X=-1} \\
&= 0.
\end{aligned}$$

For the fourth equality, note that P is not discrete, so $min(P) \neq P$. □

In other words, for any $P \in \mathbf{qp}$, $(-1)^{cl(P)} ehr_P(-1) = ehr_P^{str}(1) = \varepsilon'(P)$. By corollary 10:

Theorem 37 *ehr^{str} is the unique morphism from $\mathcal{H}_{\mathbf{qp}}$ to $\mathbb{K}[X]$ such that:*

1. *ehr^{str} is a Hopf algebra morphism from $(\mathcal{H}_{\mathbf{qp}}, m, \Delta)$ to $(\mathbb{K}[X], m, \Delta)$.*
2. *ehr^{str} is a bialgebra morphism from $(\mathcal{H}_{\mathbf{qp}}, m, \delta)$ to $(\mathbb{K}[X], m, \delta)$.*

Moreover, the character α^{str} is the inverse of λ in M_b .

4.3 The character α and the duality principle

Theorem 38 1. *(Duality principle). For any quasi-poset P :*

$$ehr_P^{str}(X) = (-1)^{cl(P)} ehr_P(-X).$$

2. *For any quasi-poset P , $\alpha_P = (-1)^{cl(P)+cc(P)} \alpha_P^{str}$.*

3. *α is invertible in $M_{\mathbf{qp}}$. We denote by β its inverse. For any quasi-poset P :*

$$\beta_P = (-1)^{cl(P)+cc(P)} \lambda_P = (-1)^{cl(P)+cc(P)} \frac{\mu_P}{cl(P)!}.$$

Proof. 1. We consider the morphism:

$$\phi : \begin{cases} \mathcal{H}_{\mathbf{qp}} & \longrightarrow \mathbb{K}[X] \\ P \in \mathbf{qp} & \longrightarrow (-1)^{cl(P)} ehr_P(-X). \end{cases}$$

We put:

$$\theta_1 : \begin{cases} \mathcal{H}_{\mathbf{qp}} & \longrightarrow \mathcal{H}_{\mathbf{qp}} \\ P \in \mathbf{qp} & \longrightarrow (-1)^{cl(P)} P, \end{cases} \quad \theta_2 : \begin{cases} \mathbb{K}[X] & \longrightarrow \mathbb{K}[X] \\ P(X) & \longrightarrow P(-X). \end{cases}$$

Both are Hopf algebra morphisms, and $\phi = \theta_2 \circ ehr \circ \phi_1$, so ϕ is a Hopf algebra morphism. If P is a non discrete quasi-poset, then $\phi(P)(1) = (-1)^{cl(P)} ehr_P(-1) = 0 = \varepsilon'(x)$. If P is a discrete quasi-poset, then $ehr_P(X) = X^{cl(P)}$, so $\phi(P)(1) = 1 = \varepsilon'(x)$. By corollary 10, $\phi = \phi_1 = ehr^{str}$.

2. and 3. Immediate consequences of the first point. \square

Proposition 39 *The following map is a Hopf algebra automorphism:*

$$\theta : \begin{cases} (\mathcal{H}_{\mathbf{qp}}, m, \Delta) & \longrightarrow (\mathcal{H}_{\mathbf{qp}}, m, \Delta) \\ P & \longrightarrow \sum_{\sim \triangleleft P} P / \sim. \end{cases}$$

Its inverse is:

$$\theta^{-1} : \begin{cases} (\mathcal{H}_{\mathbf{qp}}, m, \Delta) & \longrightarrow (\mathcal{H}_{\mathbf{qp}}, m, \Delta) \\ P & \longrightarrow \sum_{\sim \triangleleft P} (-1)^{cl(\sim)+cl(P)} P / \sim. \end{cases}$$

Moreover:

$$ehr^{str} \circ \theta = ehr, \quad ehr \circ \theta^{-1} = ehr^{str}.$$

Proof. Let ι be the character of $\mathcal{H}_{\mathbf{qp}}$ which sends any quasi-poset to 1. By corollary 20, θ is an automorphism and $\theta^{-1} = \phi_{\iota^* - 1}$. For any quasi-poset P :

$$\iota(1) = 1 = ehr_P(1) = \sum_{\sim \triangleleft P} \frac{\mu_{P/\sim}}{cl(P/\sim)!} \alpha_{P/\sim} = \sum_{\sim \triangleleft P} \lambda_{P/\sim} \alpha_{P/\sim} = \lambda * \alpha(P),$$

so $\iota = \lambda * \alpha$; hence, its inverse is $\beta * \alpha^{str}$, and for any quasi-poset P , as $cl(P/\sim) = cl(\sim)$ and $cc(P/\sim) = cc(P)$ for any $\sim \triangleleft P$:

$$\begin{aligned} \beta * \alpha^{str}(P) &= \sum_{\sim \triangleleft P} (-1)^{cl(\sim)+cc(P)} \frac{\mu_{P/\sim}}{cl(P/\sim)!} \alpha_{P/\sim}^{str} \\ &= (-1)^{cc(P)} ehr_P^{str}(-1) \\ &= (-1)^{cc(P)+cl(P)} ehr^{str}(1) \\ &= (-1)^{cc(P)+cl(P)}. \end{aligned}$$

Hence:

$$\theta^{-1}(P) = Id \leftarrow (\beta * \alpha^{str})(P) = \sum_{\sim \triangleleft P} (-1)^{cc(P/\sim)+cl(P/\sim)} P / \sim = \sum_{\sim \triangleleft P} (-1)^{cl(\sim)+cl(P)} P / \sim.$$

Moreover:

$$\begin{aligned} ehr^{str} \circ \theta &= (\phi_0 \leftarrow \alpha^{str}) \circ (Id \leftarrow \iota) \\ &= ((\phi_0 \leftarrow \alpha^{str}) \circ Id) \leftarrow \iota \\ &= (\phi_0 \leftarrow \alpha^{str}) \leftarrow \iota \\ &= \phi_0 \leftarrow (\alpha^{str} * \iota) \\ &= \phi_0 \leftarrow (\alpha^{str} * \lambda * \alpha) \\ &= \phi_0 \leftarrow \alpha \\ &= ehr. \end{aligned}$$

\square

4.4 A link with Bernoulli numbers

For any $k \in \mathbb{N}$, let c_k be the corolla quasi-poset with k leaves: $c_k = ([k+1], \leq_{c_k})$, with $1 \leq_{c_k} 2, \dots, k+1$:

$$c_0 = \bullet_1, \quad c_1 = \mathbf{1}_1^2, \quad c_2 = {}^2\mathbf{V}_1^3, \quad c_3 = {}^2\mathbf{V}_1^4 \dots$$

By proposition 34, $\lambda_{c_k} = \frac{1}{k+1}$. Moreover:

$$\begin{aligned} L_{c_k} &= \{f : [k+1] \longrightarrow \mathbb{N}^* \mid f(1) \leq f(2), \dots, f(k+1)\}, \\ L_{c_k}^{str} &= \{f : [k+1] \longrightarrow \mathbb{N}^* \mid f(1) < f(2), \dots, f(k+1)\}, \end{aligned}$$

so, for all $n \geq 1$:

$$Ehr_{c_k}^{str}(n) = (n-1)^k + \dots + 1^k = S_k(n),$$

where $S_k(X)$ is the unique polynomial such that for all $n \geq 1$, $S_k(n) = 1^k + \dots + (n-1)^k$. As a consequence, $\alpha_{c_k}^{str}$ is equal to the k -th Bernoulli number b_k .

Let $\sim \triangleleft c_k$. As the equivalence classes of \sim are connected:

- The equivalence class of the minimal element 1 of c_k contains i leaves, $0 \leq i \leq k$.
- The other equivalence classes are formed by a unique leaf.

Hence:

$$\delta([c_k]) = \sum_{i=0}^k \binom{k}{i} [c'_{i,k-i}] \otimes [c_i] \bullet^{k-i},$$

where $c'_{i,k-i}$ is the quasi poset on $[k+1]$ such that:

$$1 \sim_{c'_{i,k-i}} \dots \sim_{c'_{i,k-i}} i+1 \leq_{c'_{i,k-i}} i+2, \dots, k+1.$$

Hence, by theorem 35:

$$\begin{aligned} S_k(X) &= \sum_{i=0}^k \binom{k}{i} \lambda_{c'_{i,k-i}} b_i X^{k-i+1} \\ &= \sum_{i=0}^k \binom{k}{i} \lambda_{c'_{i,k-i}} b_i X^{k-i+1} \\ &= \sum_{i=0}^k \binom{k}{i} \lambda_{c_{k-i}} b_i X^{k-i+1} \\ &= \sum_{i=0}^k \binom{k}{i} \frac{b_i}{k-i+1} X^{k-i+1}. \end{aligned}$$

We recover in this way Faulhaber's formula. For all $n \geq 1$, $ehr_{c_k}(n) = n^k + \dots + 1^k$, and the duality principle gives, for all $n \geq 1$:

$$(-1)^{k+1} S_k(-n) = 1^k + \dots + n^k = S_k(n) + n^k.$$

5 Noncommutative version

5.1 Reminders on packed words

Let us recall the construction of the Hopf algebra of packed words \mathbf{WQSym} [15, 16].

Definition 40 *Let $w = x_1 \dots x_n$ be a word whose letters are positive integers.*

1. *We shall say that w is a packed word if there exists an integer k such that $\{x_1, \dots, x_n\} = [k]$. The set of packed words of length n is denoted by $\mathbf{PW}(n)$; the set of all packed words is denoted by \mathbf{PW} .*
2. *There exists a unique increasing bijection $f : \{x_1, \dots, x_n\} \rightarrow [k]$ for a well-chosen k . We denote by $\text{Pack}(w)$ the packed word $f(x_1) \dots f(x_n)$. Note that w is packed if, and only if, $w = \text{Pack}(w)$.*
3. *Let $I \subseteq \mathbb{N}$. Let $i_1 < \dots < i_p$ be the indices i such that $x_i \in I$. We denote by w_I the word $x_{i_1} \dots x_{i_p}$.*

As a vector space, \mathbf{WQSym} is generated by the set \mathbf{PW} . The product is given by:

$$\forall u \in \mathbf{PW}(k), \forall v \in \mathbf{PW}(l), u.v = \sum_{\substack{w=x_1 \dots x_{k+l} \in \mathbf{PW}(k+l), \\ \text{Pack}(x_1 \dots x_k)=u, \\ \text{Pack}(x_{k+1} \dots x_{k+l})=v}} w.$$

The unit is the empty word 1. The coproduct is given by:

$$\forall w \in \mathbf{PW}, \Delta(w) = \sum_{k=0}^{\max(w)} w_{\{1, \dots, k\}} \otimes \text{Pack}(w_{\{k+1, \dots, \max(w)\}}).$$

For example:

$$\begin{aligned} (11).(11) &= (1111) + (1122) + (2211), \\ (11).(12) &= (1112) + (1123) + (2212) + (2213) + (3312), \\ (11).(21) &= (1121) + (1132) + (2231) + (3321), \\ (12).(11) &= (1211) + (1222) + (1233) + (1322) + (2311), \\ (12).(12) &= (1212) + (1213) + (1223) + (1234) + (1323) + (1324) \\ &\quad + (1423) + (2312) + (2313) + (2314) + (2413) + (3412), \\ (12).(21) &= (1221) + (1231) + (1232) + (1243) + (1332) + (1342) \\ &\quad + (1432) + (2321) + (2331) + (2341) + (2431) + (3421), \end{aligned}$$

$$\begin{aligned} \Delta(111) &= (111) \otimes 1 + (111) \otimes 1, \\ \Delta(212) &= (212) \otimes 1 + (1) \otimes (11) + 1 \otimes (212), \\ \Delta(312) &= (312) \otimes 1 + (1) \otimes (21) + (12) \otimes (1) + 1 \otimes (312). \end{aligned}$$

5.2 Hopf algebra morphisms in \mathbf{WQSym}

Proposition 41 *The two following maps are surjective Hopf algebra morphisms:*

$$\begin{aligned} EHR : & \begin{cases} (\mathcal{H}_{\mathbf{QP}}, m, \Delta) & \longrightarrow & \mathbf{WQSym} \\ P & \longrightarrow & \sum_{w \in W_P} w, \end{cases} \\ EHR^{str} : & \begin{cases} (\mathcal{H}_{\mathbf{QP}}, m, \Delta) & \longrightarrow & \mathbf{WQSym} \\ P & \longrightarrow & \sum_{w \in W_P^{str}} w. \end{cases} \end{aligned}$$

Proof. Let $P \in \mathbf{QP}(k)$, $Q \in \mathbf{QP}(l)$, and w be a packed word of length $k + l$. Then:

- $w \in W_{PQ}$ if, and only if, $Pack(w_1 \dots w_k) \in W_P$ and $Pack(w_{k+1} \dots w_{k+l}) \in W_Q$.
- $w \in W_{PQ}^{str}$ if, and only if, $Pack(w_1 \dots w_k) \in W_P^{str}$ and $Pack(w_{k+1} \dots w_{k+l}) \in W_Q^{str}$.

This implies that :

$$EHR(PQ) = EHR(P)EHR(Q), \quad EHR^{str}(PQ) = EHR^{str}(P)EHR^{str}(Q).$$

Let $P \in \mathbf{QP}(n)$. We consider the two sets:

$$\begin{aligned} A &= \{(w, k) \mid w \in W_P, 0 \leq k \leq \max(w)\}, \\ B &= \{(O, w_1, w_2) \mid O \in Top(P), w_1 \in W_{Pack(P_{[n] \setminus O})}, w_2 \in W_{Pack(P|_O)}\}. \end{aligned}$$

We define a bijection between A and B by $F(w, k) = (O, w_1, w_2)$, where:

- $O = w^{-1}(\{k + 1, \dots, \max(w)\})$.
- $w_1 = Pack(w_{\{1, \dots, k\}})$.
- $w_2 = Pack(w_{\{k+1, \dots, \max(w)\}})$.

Then:

$$\begin{aligned} \Delta \circ EHR(P) &= \sum_{(w,k) \in A} Pack(w_{\{1, \dots, k\}}) \otimes Pack(w_{\{k+1, \dots, \max(w)\}}) \\ &= \sum_{(O, w_1, w_2) \in B} w_1 \otimes w_2 \\ &= (EHR \otimes EHR) \circ \Delta(P). \end{aligned}$$

So EHR is a Hopf algebra morphism. In the same way, EHR^{str} is a Hopf algebra morphism.

Let w be a packed word of length n . We define a quasi-poset structure on $[n]$ by $i \leq_P j$ if, and only if, $w_i \leq w_j$. Then $W_P^{str} = \{w\}$, so $EHR^{str}(P) = w$: EHR^{str} is surjective. If $w' \in W_P$, then $\max(w') \leq \max(w)$ with equality if, and only if, $w = w'$. Hence:

$$EHR(P) = w + \text{words } w' \text{ with } \max(w') < \max(w).$$

By a triangular argument, EHR is surjective. □

Examples.

$$\begin{array}{ll} EHR(\bullet_1) = (1), & EHR^{str}(\bullet_1) = (1), \\ EHR(\mathbf{1}_1^2) = (12) + (11), & EHR^{str}(\mathbf{1}_1^2) = (12), \\ EHR(\mathbf{1}_2^1) = (21) + (11), & EHR^{str}(\mathbf{1}_2^1) = (21), \\ EHR(\bullet_{1 \bullet 2}) = (12) + (21) + (11), & EHR^{str}(\bullet_{1 \bullet 2}) = (12) + (21) + (11), \\ EHR(\bullet_{\bullet 1, 2}) = (11), & EHR^{str}(\bullet_{\bullet 1, 2}) = (11). \end{array}$$

Remark. The Hopf algebra \mathbf{WQSymb} has a polynomial representation [16]. Let $X = \{x_1, x_2, \dots\}$ be an infinite, totally ordered alphabet; then, for any packed word w :

$$P_w(X) = \sum_{w' \in X^*, Pack(w')=w} w'.$$

With this polynomial representation, for any $P \in \mathbf{QP}(n)$:

$$EHR(P) = \sum_{f \in L_P} x_{f(1)} \dots x_{f(n)}, \quad EHR^{str}(P) = \sum_{f \in L_P^{str}} x_{f(1)} \dots x_{f(n)}.$$

Proposition 42 *The following map is a Hopf algebra automorphism:*

$$\Theta : \begin{cases} (\mathcal{H}_{\mathbf{QP}}, m, \Delta) & \longrightarrow (\mathcal{H}_{\mathbf{QP}}, m, \Delta) \\ P & \longrightarrow \sum_{\sim \triangleleft P} P / \sim . \end{cases}$$

Its inverse is:

$$\Theta^{-1} : \begin{cases} (\mathcal{H}_{\mathbf{QP}}, m, \Delta) & \longrightarrow (\mathcal{H}_{\mathbf{QP}}, m, \Delta) \\ P & \longrightarrow \sum_{\sim \triangleleft P} (-1)^{cl(P)+cl(\sim)} P / \sim . \end{cases}$$

Moreover, $EHR^{str} \circ \Theta = EHR$ and $EHR \circ \Theta^{-1} = EHR^{str}$.

Proof. By corollary 20, $\Theta = \phi_\iota$ is a Hopf algebra automorphism, where ι is defined in the proof of proposition 39. Its inverse is $\phi_{\iota^{-1}}$.

Let us prove that:

$$W_G = \sum_{\sim \triangleleft P} W_{G/\sim}^{str}.$$

Let $w \in W_G$; we define an equivalence \sim_w by $x \sim_w y$ if $w(x) = w(y)$ and x and y are in the same connected component of $w^{-1}(w(x))$. By definition, the equivalence classes of \sim_w are connected. If $x \sim_{P/\sim_w} y$, there exists $x_1, x'_1, \dots, x_k, x'_k, y_1, y'_1, \dots, y_l, y'_l$ such that:

$$\begin{aligned} x \leq_P x_1 \sim_w x'_1 \leq_P \dots \leq_P x_k \sim_w x'_k \leq_P y, \\ y \leq_P y_1 \sim_w y'_1 \leq_P \dots \leq_P y_l \sim_w y'_l \leq_P x. \end{aligned}$$

As $w \in W_P$, $w(x) \leq w(x_1) = w(x'_1) \leq \dots \leq w(x_k) = w(x'_k) \leq w(y)$; by symmetry, $w(x) = w(x_1) = \dots = w(x'_k) = w(y) = i$. Moreover, as the equivalence classes of \sim_w are connected, x and y are in the same connected component of $w^{-1}(i)$, so $x \sim_w y$: $\sim_w \triangleleft P$.

If $x \leq_P y$ or $x \sim_w y$, then $w(x) \leq w(y)$. By transitive closure, if $x \leq_{P/\sim_w} y$, then $w(x) \leq w(y)$, so $w \in W_{P/\sim_w}$. Moreover, if $w(i) \neq w(j)$, we do not have $x \sim_w y$, so $w \in W_{P/\sim_w}^{str}$.

Let us assume that $\sim \triangleleft P$ and let $w \in W_{P/\sim}^{str}$. If $x \leq_P y$, then $x \leq_{P/\sim} y$, so $w(x) \leq w(y)$: $W_{P/\sim}^{str} \subseteq W_P$.

Let us assume that $w \in W_{P/\sim}^{str}$, with $\sim \triangleleft P$. If $x \sim y$, then $w(x) = w(y) = i$ and x and y are in the same connected component of P/\sim , so are in the the same connected component of $w^{-1}(i)$: $x \sim_w y$. If $x \sim_w y$, then $w(x) = w(y) = i$ and there exists $x_1, x'_1, \dots, x_k, x'_k$ with $w(x_1) = w(x'_1) = \dots = w(x_k) = w(x'_k) = i$ such that:

$$x \leq_P x_1 \geq_P x'_1 \leq_P \dots \geq_P x'_k \leq_P y.$$

As $w \in W_{P/\sim}^{str}$, $x \sim_{P/\sim} x_1, x_1 \sim_{P/\sim} x'_1, \dots, x'_k \sim_{P/\sim} y$. So $x \sim_{P/\sim} y$; as $\sim \triangleleft P$, $x \sim y$. Finally, $\sim = \sim_w$.

We obtain that:

$$EHR(P) = \sum_{w \in W_P} w = \sum_{\sim \triangleleft P} \sum_{w \in W_{P/\sim}^{str}} w = \sum_{\sim \triangleleft P} EHR^{str}(P/\sim) = EHR^{str}(\Theta(P)).$$

So $Ehr^{str} \circ \Theta = EHR$. □

Proposition 43 *Let us consider the following map:*

$$H : \begin{cases} \mathbf{WQSym} & \longrightarrow \mathbb{K}[X] \\ w \in \mathbf{PW} & \longrightarrow H_{\max(w)}(X). \end{cases}$$

This is a surjective Hopf algebra morphism, making the following diagram commuting:

$$\begin{array}{ccccc}
& & \mathcal{H}_{\mathbf{QP}} & & \\
& \swarrow \sqcup & \downarrow \theta & \searrow EHR & \\
\mathcal{H}_{\mathbf{QP}} & & \mathcal{H}_{\mathbf{QP}} & \xrightarrow{EHR^{str}} & \mathbf{WQSym} \\
\downarrow \theta & \swarrow \sqcup & \downarrow \theta & \searrow H & \\
\mathcal{H}_{\mathbf{QP}} & \xrightarrow{ehr^{str}} & \mathbb{K}[X] & &
\end{array}$$

Proof. Let $P \in \mathbf{QP}$. Then:

$$ehr(\lfloor P \rfloor) = \sharp W_P(k) H_k(X) = \sum_{w \in W_P} H_{\max(w)}(X) = \sum_{w \in W_P} H(w) = H \circ EHR(P).$$

So $ehr \circ \lfloor \rfloor = H \circ EHR$. Similarly, $ehr^{str} \circ \lfloor \rfloor = H \circ EHR^{str}$.

Let us prove that H is a Hopf algebra morphism. Let $w_1, w_2 \in \mathbf{WQSym}$. There exist $x_1, x_2 \in \mathcal{H}_{\mathbf{QP}}$, such that $w_1 = EHR(x_1)$ and $w_2 = EHR(x_2)$. Then:

$$\begin{aligned}
H(w_1 w_2) &= H(EHR(x_1) EHR(x_2)) \\
&= H \circ EHR(x_1 x_2) \\
&= ehr(\lfloor x_1 x_2 \rfloor) \\
&= ehr(\lfloor x_1 \rfloor) ehr(\lfloor x_2 \rfloor) \\
&= H \circ EHR(x_1) H \circ EHR(x_2) \\
&= H(w_1) H(w_2).
\end{aligned}$$

Let $w \in \mathbf{WQSym}$. There exists $x \in \mathcal{H}_{\mathbf{QP}}$ such that $w = EHR(x)$.

$$\begin{aligned}
\Delta \circ H(w) &= \Delta \circ H \circ EHR(x) \\
&= (H \otimes H) \circ (EHR \otimes EHR) \circ \Delta(x) \\
&= (H \otimes H) \circ \Delta \circ EHR(x) \\
&= (H \otimes H) \circ \Delta(w).
\end{aligned}$$

So H is a Hopf algebra morphism. □

5.3 The non-commutative duality principle

Lemma 44 *The following map is an involution and a Hopf algebra automorphism:*

$$\Phi_{-1} : \begin{cases} \mathbf{WQSym} & \longrightarrow & \mathbf{WQSym} \\ w & \longrightarrow & (-1)^{\max(w)} \sum_{\sigma : [\max(w)] \rightarrow [l], \text{ non-decreasing}} \sigma \circ w. \end{cases}$$

Proof. Using the surjective morphisms EHR^{str} and ehr^{str} , taking the quotients of the cointeracting bialgebras $(\mathcal{H}_{\mathbf{QP}}, m, \Delta)$ and $(\mathcal{H}_{\mathbf{QP}}, m, \delta)$, we obtain that $(\mathbf{WQSym}, m, \Delta)$ and $(\mathbb{K}[X], m, \delta)$ are cointeracting bialgebras, with the coaction defined by:

$$\rho = (Id \otimes H) \circ \delta : \mathbf{WQSym} \longrightarrow \mathbf{WQSym} \otimes \mathbb{K}[X]$$

For any packed word w :

$$\rho(w) = \sum_{\sigma : [k] \rightarrow [l], \text{ non-decreasing}} \sigma \circ w \otimes H_{\max(\text{Pack}(w_{|(\sigma \circ w)^{-1}(1)}))}(X) \dots H_{\max(\text{Pack}(w_{|(\sigma \circ w)^{-1}(l)}))}(X).$$

Using proposition 4, for any $\lambda \in \mathbb{K}$, considering the character:

$$ev_\lambda : \begin{cases} \mathbb{K}[X] & \longrightarrow \mathbb{K} \\ P & \longrightarrow P(\lambda), \end{cases}$$

we obtain an endomorphism Φ_λ of $(\mathbf{WQSym}, m, \Delta)$ defined by $\Phi_\lambda = Id \leftarrow ev_\lambda$. if $\lambda \neq 0$, Φ_λ is invertible, of inverse $\Phi_{\lambda^{-1}}$. For any packed word w , denoting by k its maximum:

$$\Phi_\lambda(w) = \sum_{\sigma : [k] \rightarrow [l], \text{ non-decreasing}} H_{\max(\text{Pack}(w_{|(\sigma \circ w)^{-1}(1)}))}(\lambda) \cdots H_{\max(\text{Pack}(w_{|(\sigma \circ w)^{-1}(l)}))}(\lambda) \sigma \circ w.$$

In particular, for $\lambda = -1$, for any $p \in \mathbb{N}$:

$$H_p(-1) = \frac{(-1)(-2) \cdots (-k)}{k!} = (-1)^k.$$

Hence:

$$\begin{aligned} \Phi_{-1}(w) &= \sum_{\sigma : [k] \rightarrow [l], \text{ non-decreasing}} (-1)^{\max(\text{Pack}(w_{|(\sigma \circ w)^{-1}(1)})) + \dots + \max(\text{Pack}(w_{|(\sigma \circ w)^{-1}(l)}))} \sigma \circ w \\ &= (-1)^k \sum_{\sigma : [k] \rightarrow [l], \text{ non-decreasing}} \sigma \circ w. \end{aligned}$$

Indeed, if $x \in (\sigma \circ w)^{-1}(p)$ and $y \in (\sigma \circ w)^{-1}(q)$, with $p < q$, then $\sigma \circ w(x) < \sigma \circ w(y)$; as σ is non-decreasing, $x < y$. So there exists $n_1 < n_2 < \dots < n_l = k$ such that for all p , the values taken by w on $(\sigma \circ w)^{-1}(p)$ are $n_{p-1} + 1, \dots, n_p$. Hence, the values taken by $\text{Pack}(w_{|(\sigma \circ w)^{-1}(p)})$ are $1, \dots, n_p - n_{p-1}$, so:

$$\max(\text{Pack}(w_{|(\sigma \circ w)^{-1}(1)})) + \dots + \max(\text{Pack}(w_{|(\sigma \circ w)^{-1}(l)})) = n_1 + n_2 - n_1 + \dots + n_l - n_{l-1} = n_l = k.$$

In particular, Φ_{-1} is an involution and a Hopf algebra automorphism of $(\mathbf{WQSym}, m, \Delta)$. \square

Theorem 45 (Non commutative duality principle) For any quasi-poset $P \in \mathbf{QP}$:

$$EHR(P) = (-1)^{cl(P)} \Phi_{-1} \circ ERH^{str}(P), \quad EHR^{str}(P) = (-1)^{cl(P)} \Phi_{-1} \circ ERH(P).$$

Proof. We shall use the following involution and Hopf algebra automorphism:

$$\Psi : \begin{cases} \mathcal{H}_{\mathbf{QP}} & \longrightarrow \mathcal{H}_{\mathbf{QP}} \\ p \in \mathbf{QP} & \longrightarrow (-1)^{cl(P)} P. \end{cases}$$

Recall that the character ι of $\mathcal{H}_{\mathbf{QP}}$ sends any $P \in \mathbf{QP}$ to 1. By the duality principle:

$$\iota \circ \Psi(P) = (-1)^{cl(P)} = (-1)^{cl(P)} ehr(P)(1) = ehr^{str}(-1) = ev_{-1} \circ ehr^{str}(P).$$

So $\iota \circ \Psi = ev_{-1} \circ ehr^{str}$.

Let $P \in \mathbf{QP}$. Recalling that if $\sim \triangleleft P$, $cl(P| \sim) = cl(P)$:

$$\delta \circ \Psi(P) = (-1)^{cl(P)} \sum_{\sim \triangleleft P} P/ \sim \otimes P| \sim = \sum_{\sim \triangleleft P} P/ \sim \otimes (-1)^{cl(P| \sim)} P| \sim = (Id \otimes \Psi) \circ \delta(P).$$

So $\delta \circ \Psi = (Id \otimes \Psi) \circ \delta$. Hence, for any $x \in \mathcal{H}_{\mathbf{QP}}$:

$$\begin{aligned} EHR \circ \Psi(x) &= EHR^{str} \circ (Id \leftarrow \iota) \circ \Psi(x) \\ &= EHR^{str}(\Psi(x)_0) \iota \circ \Psi(x)_1 \\ &= EHR^{str}(x_0) \iota \circ \Psi(x_1) \\ &= EHR^{str}(x_0) ev_{-1} \circ ehr^{str}(x_1) \\ &= EHR^{str}(x^{(1)}) ev_{-1} \circ EHR^{str}(x^{(2)}) \\ &= EHR \leftarrow ev_{-1}(x) \\ &= (Id \leftarrow ev_{-1}) \circ EHR^{str}(x) \\ &= \Phi_{-1} \circ EHR^{str}(x), \end{aligned}$$

where we denote $\delta(x) = x^{(1)} \otimes x^{(2)}$ and $\rho(x) = x_0 \otimes x_1$. As Φ_{-1} and Ψ are involutions, $EHR^{str} \circ \Psi = \Phi_{-1} \circ EHR$. \square

In $E_{\mathbb{K}[X] \rightarrow \mathbb{K}[X]}$, putting $\phi_\lambda = Id \leftarrow ev_\lambda$, for any $P \in \mathbb{K}[X]$, $\phi_\lambda(P) = P(\lambda X)$. Moreover, as H is compatible with the coactions:

$$H \circ \Phi_\lambda = H \circ (Id \leftarrow ev_\lambda) = H \leftarrow ev_\lambda = (Id \leftarrow ev_\lambda) \circ H = \phi_\lambda \circ H,$$

so:

$$ehr \circ \Psi = H \circ EHR \circ \Psi = H \circ \Phi_{-1} \circ EHR^{str} = \phi_{-1} \circ H \circ EHR^{str} = \phi_{-1} \circ ehr^{str}.$$

In other words, for any $P \in \mathbf{QP}$, $(-1)^{cl(P)} ehr_P(X) = ehr_P^{str}(-X)$: we recover the duality principle.

We obtain the commutative diagram of Hopf algebra morphisms:

$$\begin{array}{ccccc}
\mathcal{H}_{\mathbf{QP}} & & & & \\
\downarrow \ominus & \searrow EHR & & & \\
\mathcal{H}_{\mathbf{QP}} & \xrightarrow{EHR^{str}} & \mathbf{WQSym} & \sqcup & \\
\downarrow \Psi & & \downarrow \Phi_{-1} & & \\
\mathcal{H}_{\mathbf{QP}} & \xrightarrow{EHR} & \mathbf{WQSym} & \sqcup & \\
& & & \sqcup & \\
& & & & \mathcal{H}_{\mathbf{qp}} \\
& & & \searrow H & \downarrow \theta \\
& & & & \mathcal{H}_{\mathbf{qp}} \xrightarrow{ehr} \mathbb{K}[X] \\
& & & \searrow H & \downarrow \psi \\
& & & & \mathcal{H}_{\mathbf{qp}} \xrightarrow{ehr^{str}} \mathbb{K}[X] \\
& & & & \downarrow \phi_{-1} \\
& & & & \mathcal{H}_{\mathbf{qp}} \xrightarrow{chr} \mathbb{K}[X]
\end{array}$$

5.4 Compatibility with the other product and coproduct

Theorem 46 We define a second coproduct δ on \mathbf{WQSym} :

$$\forall w \in \mathbf{PW}, \delta(w) = \sum_{(\sigma, \tau) \in A_w} (\sigma \circ w) \otimes (\tau \circ w),$$

where A_w is the set of pairs of packed words (σ, τ) of length $\max(w)$ such that:

- σ is non-decreasing.
- If $1 \leq i < j \leq \max(w)$ and $\sigma(i) = \sigma(j)$, then $\tau(i) < \tau(j)$.

Then $(\mathbf{WQSym}, m, \delta)$ is a bialgebra and EHR^{str} is a bialgebra morphism from $(\mathcal{H}_{\mathbf{QP}}, m, \delta)$ to $(\mathbf{WQSym}, m, \delta)$.

Proof. Let us prove that $\delta \circ EHR^{str} = (EHR^{str} \otimes EHR^{str}) \circ \delta$. Let $P \in \mathbf{QP}$. We consider the two following sets:

$$\begin{aligned}
A &= \{(\sim, w_1, w_2) \mid \sim \triangleleft P, w_1 \in W_{P/\sim}^{str}, w_2 \in W_{P|\sim}^{str}\}, \\
B &= \{(w, \sigma, \tau) \mid w \in W_P^{str}, (\sigma, \tau) \in A_w\}.
\end{aligned}$$

Let $(\sim, w_1, w_2) \in A$. We put $I_p = w_1^{-1}(p)$ for all $1 \leq p \leq \max(w_1)$, and $w_2^{(p)}$ the standardization of the restriction of w_2 to I_p . We define w by:

$$w(i) = w_2^{(p)}(i) + \max w_1^{(2)} + \dots + \max w_{p-1}^{(2)} \text{ if } i \in I_p.$$

Let us prove that $w \in W_P^{str}$. If $x \leq_P y$, then $x \leq_{P/\sim} y$, so $p = w_1(x) \leq w_2(y) = q$.

- If $p < q$, then $w(x) < w(y)$.
- If $p = q$, then $w_1(x) = w_2(y)$ and, as $x \leq_P y$, x and y are in the same connected component of $w^{-1}(p)$. So $x \sim_{w_1} y$, that is to say $x \sim y$ as $w_1 \in W_{P/\sim}^{str}$, and $x \leq_{P|\sim} y$, which implies that $w_2(x) \leq w_2(y)$ and finally $w(x) \leq w(y)$.

Let us assume that moreover $w(x) = w(y)$. Then $p = q$ and necessarily, $w_2(x) = w_2(y)$. As $w_2 \in W_{P|\sim}^{str}$, $x \sim_{P|\sim} y$, so $x \sim_P y$.

If $w(x) = w(y)$, then by definition of w , $w_1(x) = w_1(y)$. So there exists a unique $\sigma : [\max(w)] \rightarrow [\max(w_1)]$, such that $w_1 = \sigma \circ w$. If $w(x) < w(y)$, then, by construction of w , $w_1(x) \leq w_1(y)$: σ is non-decreasing.

There exists a unique $\tau : [\max(w)] \rightarrow [\max(w_2)]$, such that $w_2 = \tau \circ \sigma$. As $Pack(w|_{I_p}) = Pack((w_2)|_{I_p})$ for all p , τ is increasing on I_p .

To any $(\sim, w_1, w_2) \in A$, we associate $(w, \sigma, \tau) = F(\sim, w_1, w_2) \in B$, such that $w_1 = \sigma \circ w$, $w_2 = \tau \circ \sigma$, and $\sim = \sim_{\sigma \circ \tau}$. This defines a map $F : A \rightarrow B$.

Let $(w, \sigma, \tau) \in B$. We put $G(w, \sigma, \tau) = (\sim, \sigma, \tau) = (\sim_{\sigma \circ w}, \sigma \circ w, \tau \circ w)$. If $x \leq_P y$, then $w(x) \leq w(y)$, so $w_1(x) = \sigma \circ w(x) \leq \sigma \circ w(y) = w_1(y)$. If moreover $w_1(x) = w_1(y)$, then as $x \leq_P y$, x and y are in the same connected component of $w_1^{-1}(w_1(x))$, so $x \sim_{w_1} y$: $w_1 \in W_{P/\sim}^{str}$.

If $x \leq_{P|\sim} y$, then $x \sim_{w_1} y$ and $x \leq_P y$, so $w_1(x) = w_1(y)$ and $w(x) \leq w(y)$. By hypothesis on τ , $\tau \circ w(x) \leq \tau \circ w(y)$, so $w_2(x) \leq w_2(y)$. If moreover $w_2(x) = w_2(y)$, by hypothesis on τ , $w(x) = w(y)$. As $w \in W_P^{str}$, $x \sim_P y$, so $x \sim_{P|\sim} y$: $w_2 \in W_{P|\sim}^{str}$.

We defined in this way a map $G : B \rightarrow A$. If $(\sim, w_1, w_2) \in A$:

$$G \circ F(\sim, w_1, w_2) = G(w, \sigma, \tau) = (\sim_{\sigma \circ w}, \sigma \circ w, \tau \circ w) = (\sim_{w_1}, w_1, w_2) = (\sim, w_1, w_2).$$

So $G \circ F = Id_A$. If $(w, \sigma, \tau) \in B$:

$$F \circ G(w, \sigma, \tau) = F(\sim_{\sigma \circ w}, \sigma \circ w, \tau \circ w) = (w, \sigma, \tau).$$

So $F \circ G = Id_B$: F and G are inverse bijections.

We obtain:

$$\begin{aligned} (EHR^{str} \otimes EHR^{str}) \circ \delta(P) &= \sum_{(\sim, w_1, w_2) \in A} w_1 \otimes w_2 \\ &= \sum_{(w, \sigma, \tau) \in B} \sigma \circ w \otimes \tau \circ w \\ &= \sum_{w \in W_P^{str}} \delta(w) \\ &= \delta \circ EHR^{str}(P). \end{aligned}$$

So EHR^{str} is compatible with δ .

As EHR^{str} is compatible with the product m and the coproduct δ , $Ker(EHR^{str})$ is a biideal of $(\mathcal{H}_{\mathbf{QP}}, m, \delta)$, and $(\mathbf{WQSym}, m, \delta)$ is identified with the quotient $\mathcal{H}_{\mathbf{QP}}/Ker(EHR^{str})$, so is a bialgebra. \square

Examples.

$$\begin{aligned} \delta(11) &= (11) \otimes (11), \\ \delta(12) &= (12) \otimes ((11) + (12) + (21)) + (11) \otimes (12), \\ \delta(21) &= (21) \otimes ((11) + (12) + (21)) + (11) \otimes (21). \end{aligned}$$

This coproduct δ on **WQSym** is the internal coproduct of [16], dual to the product of the Solomon-Tits algebra.

Remarks.

1. The count of $(\mathbf{WQSym}, m, \delta)$ is given by:

$$\varepsilon_B(w) = \begin{cases} 1 & \text{if } w = (1 \dots 1), \\ 0 & \text{otherwise.} \end{cases}$$

2. There is no coproduct δ' on **WQSym** such that $(EHR \otimes EHR) \circ \delta = \delta' \circ EHR$. Indeed, if δ' is any coproduct on **WQSym**, for $x = \mathbf{1}_1^2 + \mathbf{1}_2^1 - \bullet_{1,2} - \bullet_{1,2}$:

$$\delta' \circ EHR(x) = \delta'(0) = 0,$$

but:

$$\begin{aligned} & (EHR \otimes EHR) \circ \delta(x) \\ &= (EHR \otimes EHR)((\mathbf{1}_1^2 + \mathbf{1}_2^1 - \bullet_{1,2}) \otimes \bullet_{1,2} + \bullet_{1,2} \otimes (\mathbf{1}_1^2 + \mathbf{1}_2^1 - \bullet_{1,2} - \bullet_{1,2})) \\ &= (11) \otimes (11). \end{aligned}$$

Proposition 47 $H : (\mathbf{WQSym}, m, \delta) \longrightarrow (\mathbb{K}[X], m, \delta)$ is a bialgebra morphism.

Proof. Let w be a packed word. We denote $k = \max(w)$. Let $a, b \in \mathbb{N}$.

$$\begin{aligned} (H \otimes H) \circ \delta(w)(a, b) &= \sum_{(\sigma, \tau) \in A_w} H_{\max(\sigma \circ w)}(a) H_{\max(\tau \circ w)}(b) \\ &= \sum_{\sigma : [k] \rightarrow [l], \text{ non-decreasing}} \binom{a}{l} \binom{b}{|\sigma^{-1}(1)|} \cdots \binom{b}{|\sigma^{-1}(l)|} \\ &= \sum_{\substack{1 \leq l \leq k, \\ i_1 + \dots + i_l = k}} \binom{a}{l} \binom{b}{i_1} \cdots \binom{b}{i_l} \\ &= \binom{ab}{k} \\ &= H_k(ab) \\ &= \delta(H(w))(a, b). \end{aligned}$$

As this is true for any $a, b \in \mathbb{N}$, $(H \otimes H) \circ \delta(w) = \delta \circ H$. □

Definition 48 Let $w = w_1 \dots w_k$ and $w' = w'_1 \dots w'_l$ be two packed words. We put:

$$\begin{aligned} w \downarrow w' &= w_1 \dots w_k (w'_1 + \max(w)) \dots (w'_l + \max(w)), \\ w \circledast w' &= w_1 \dots w_k (w'_1 + \max(w) - 1) \dots (w'_l + \max(w) - 1), \\ w \downarrow\! \! \! \downarrow w' &= w \downarrow w' + w \circledast w'. \end{aligned}$$

These three products are extended to **WQSym** by bilinearity.

Proposition 49 For all $x, y \in \mathcal{H}_{\mathbf{QP}}$:

$$EHR^{str}(x \downarrow y) = EHR^{str}(x) \downarrow EHR^{str}(y), \quad EHR(x \downarrow y) = EHR(x) \downarrow\! \! \! \downarrow EHR(y).$$

Proof. Let $P \in \mathbf{QP}(k)$ and $Q \in \mathbf{QP}(l)$. If $w = w_1 \dots w_{k+l}$ is a packed word of length $k+l$:

$$\begin{aligned} w \in W_{P \downarrow Q}^{str} &\iff w_1 \dots w_k \in L_P^{str}, w_{k+1} \dots w_{k+l} \in L_Q^{str}, w_1, \dots, w_k < w_{k+1}, \dots, w_{k+l} \\ &\iff w = w_P \downarrow w_Q, \text{ with } w_P \in W_P^{str}, w_Q \in W_Q^{str}. \end{aligned}$$

So $W_{P \downarrow Q}^{str} = W_P^{str} \downarrow W_Q^{str}$, and:

$$EHR^{str}(P \downarrow Q) = \sum_{w_P \in W_P^{str}, w_Q \in W_Q^{str}} w_P \downarrow w_Q = EHR^{str}(P) \downarrow EHR^{str}(Q).$$

If $w = w_1 \dots w_{k+l}$ is a packed word of length $k+l$:

$$\begin{aligned} w \in W_{P \downarrow Q} &\iff w_1 \dots w_k \in L_P, w_{k+1} \dots w_{k+l} \in L_Q, w_1, \dots, w_k \leq w_{k+1}, \dots, w_{k+l} \\ &\iff w = (w_P \downarrow w_Q, \text{ with } w_P \in W_P, w_Q \in W_Q) \\ &\text{or } w = (w_P \otimes w_Q, \text{ with } w_P \in W_P, w_Q \in W_Q). \end{aligned}$$

Note that these two conditions are incompatible:

- in the first case, $\max(w_1 \dots w_k) = \min(w_{k+1} \dots w_{k+l}) - 1$;
- in the second case, $\max(w_1 \dots w_k) = \min(w_{k+1} \dots w_{k+l})$.

So $W_{P \downarrow Q} = (W_P \downarrow W_Q) \sqcup (W_P \otimes W_Q)$, and:

$$\begin{aligned} EHR(P \downarrow Q) &= \sum_{w_P \in W_P, w_Q \in W_Q} w_P \downarrow w_Q + w_P \otimes w_Q \\ &= EHR(P) \downarrow EHR(Q) + EHR(P) \otimes EHR(Q), \end{aligned}$$

so $EHR(P \downarrow Q) = EHR(P) \not\downarrow EHR(Q)$. □

Remark. As a consequence, $(\mathbf{WQSym}, \downarrow, \Delta)$ and $(\mathbf{WQSym}, \not\downarrow, \Delta)$ are infinitesimal bialgebras [12], as $(\mathcal{H}_{\mathbf{QP}}, \downarrow, \Delta)$ is [11, 10].

Corollary 50 For all $x, y \in \mathbf{WQSym}$:

$$\Phi_{-1}(x \downarrow y) = \Phi_{-1}(x) \not\downarrow \Phi_{-1}(y) \qquad \Phi_{-1}(x \not\downarrow y) = \Phi_{-1}(x) \downarrow \Phi_{-1}(y).$$

Proof. If $P, Q \in \mathbf{QP}$, then $cl(P \downarrow Q) = cl(P) + cl(Q)$, so:

$$\Psi(P \downarrow Q) = (-1)^{cl(P)+cl(Q)} P \downarrow Q = \Psi(P) \downarrow \Psi(Q).$$

Let $x, y \in \mathbf{WQSym}$. There exist $x', y' \in \mathcal{H}_{\mathbf{QP}}$, such that $EHR^{str}(x') = x$ and $EHR^{str}(y') = y$. Hence, using the non-commutative duality principle:

$$\begin{aligned} \Phi_{-1}(x \downarrow y) &= \Phi_{-1}(EHR^{str}(x') \downarrow EHR^{str}(y')) \\ &= \Phi_{-1} \circ EHR^{str}(x' \downarrow y') \\ &= \Phi_{-1} \circ EHR^{str} \circ \Psi(x') \downarrow \Psi(y') \\ &= EHR(\Psi(x') \downarrow \Psi(y')) \\ &= EHR \circ \Psi(x') \not\downarrow EHR \circ \Psi(y') \\ &= \Phi_{-1}(\Phi_{-1} \circ EHR \circ \Psi(x')) \not\downarrow \Phi_{-1}(\Phi_{-1} \circ EHR \circ \Psi(y')) \\ &= \Phi_{-1}(EHR^{str}(x')) \not\downarrow \Phi_{-1}(EHR^{str}(y')) \\ &= \Phi_{-1}(x) \not\downarrow \Phi_{-1}(y). \end{aligned}$$

As Φ_{-1} is an involution, we obtain the second point. □

5.5 Restriction to posets

In [10], the image of the restriction to $\mathcal{H}_{\mathbf{P}}$ of the map from $\mathcal{H}_{\mathbf{QP}}$ to \mathbf{WQSymb} defined by T -partitions is a Hopf subalgebra, isomorphic to the Hopf algebra of permutations \mathbf{FQSymb} [13, 7]. This is not the case here:

Proposition 51 $EHR(\mathcal{H}_{\mathbf{P}}) = EHR^{str}(\mathcal{H}_{\mathbf{P}}) = \mathbf{WQSymb}$.

Proof. Let w be a packed word of length n . We define a poset P on $[n]$ by:

$$\forall i, j \in [n], i \leq_P j \text{ if } (i = j) \text{ or } (w(i) < w(j)).$$

Note that if $i \leq_P j$, then $w(i) \leq w(j)$. If $i \leq_P j$ and $j \leq_P k$, then:

- if $i = j$ or $j = k$, then obviously $i \leq_P k$.
- Otherwise, $w(i) < w(j)$ and $w(j) < w(k)$, so $w(i) < w(k)$ and $i \leq_P k$.

Let us assume that $i \leq_P j$ and $j \leq_P i$. Then $w(i) \leq w(j)$ and $w(j) \leq w(i)$, so $w(i) = w(j)$. As $i \leq_P j$, $i = j$. So P is indeed a poset, and we observed that $w \in W_P$.

Let w' be a packed word of length n . Let us prove that $w' \in W_P^{str}$ if, and only if, $w \leq w'$, where \leq is the order on packed words defined in definition 28.

\implies . Let us assume that $w' \in W_P^{str}$. If $w(i) < w(j)$, then $i \leq_P j$, so $w'(i) \leq w'(j)$. Moreover, if $w'(i) = w'(j)$, then $i \leq_P j$, so $i = j$ as P is a poset, and finally $w(i) = w(j)$: contradiction. So $w'(i) < w'(j)$, we shows that $w \leq w'$.

\impliedby . Let us assume that $w' \leq w$. If $i \leq_P j$, then $i = j$ or $w(i) < w(j)$, so $w'(i) = w'(j)$ or $w'(i) < w'(j)$. If, moreover, $w'(i) = w'(j)$, then $i = j$; so $w' \in W_P^{str}$.

We obtain an element $P \in \mathcal{H}_{\mathbf{P}}$ such that:

$$EHR^{str}(P) = \sum_{w \leq w'} w'.$$

As this holds for any w , by a triangularity argument, $EHR^{str}(\mathcal{H}_{\mathbf{P}}) = \mathbf{WQSymb}$. By the non-commutative duality principle:

$$EHR(\mathcal{H}_{\mathbf{P}}) = \Phi_{-1} \circ EHR^{str} \circ \Psi(\mathcal{H}_{\mathbf{P}}) = \Phi_{-1} \circ EHR^{str}(\mathcal{H}_{\mathbf{P}}) = \Phi_{-1}(\mathbf{WQSymb}) = \mathbf{WQSymb},$$

as Φ_{-1} is an automorphism of \mathbf{WQSymb} . □

References

- [1] P. Alexandroff, *Diskrete Räume.*, Rec. Math. Moscou, n. Ser. **2** (1937), 501–519 (German).
- [2] Matthias Beck and Sinai Robins, *Computing the continuous discretely*, second ed., Undergraduate Texts in Mathematics, Springer, New York, 2015, Integer-point enumeration in polyhedra, With illustrations by David Austin.
- [3] Ch. Brouder, *Trees, renormalization and differential equations*, BIT **44** (2004), no. 3, 425–438.
- [4] J. C. Butcher, *An algebraic theory of integration methods*, Math. Comp. **26** (1972), 79–106.
- [5] Damien Calaque, Kurusch Ebrahimi-Fard, and Dominique Manchon, *Two interacting Hopf algebras of trees: a Hopf-algebraic approach to composition and substitution of B-series*, Adv. in Appl. Math. **47** (2011), no. 2, 282–308, arXiv:0806.2238.

- [6] F. Chapoton, *Sur une série en arbres à deux paramètres*, Sém. Lothar. Combin. **70** (2013), Art. B70a, 20.
- [7] Gérard Duchamp, Florent Hivert, and Jean-Yves Thibon, *Noncommutative symmetric functions. VI. Free quasi-symmetric functions and related algebras*, Internat. J. Algebra Comput. **12** (2002), no. 5, 671–717.
- [8] Frédéric Fauvet, Loïc Foissy, and Dominique Manchon, *The Hopf algebra of finite topologies and mould composition*, arXiv:1503.03820, 2015.
- [9] Loïc Foissy, *Chromatic polynomials and bialgebras of graphs*, in preparation.
- [10] Loïc Foissy and Claudia Malvenuto, *The Hopf algebra of finite topologies and T -partitions*, J. Algebra **438** (2015), 130–169, arXiv:1407.0476.
- [11] Loïc Foissy, Claudia Malvenuto, and Frédéric Patras, *Infinitesimal and B_∞ -algebras, finite spaces, and quasi-symmetric functions*, J. Pure Appl. Algebra **220** (2016), no. 6, 2434–2458, arXiv:1403.7488.
- [12] Jean-Louis Loday and María Ronco, *On the structure of cofree Hopf algebras*, J. Reine Angew. Math. **592** (2006), 123–155, arXiv:math/0405330.
- [13] Claudia Malvenuto and Christophe Reutenauer, *A self paired Hopf algebra on double posets and a Littlewood-Richardson rule*, J. Combin. Theory Ser. A **118** (2011), no. 4, 1322–1333, arXiv:0905.3508.
- [14] Dominique Manchon, *On bialgebras and Hopf algebras or oriented graphs*, Confluentes Math. **4** (2012), no. 1, 1240003, 10, arXiv:1011.3032.
- [15] J.-C. Novelli and J.-Y. Thibon, *Construction of dendriform trialgebras*, C. R. Acad. Sci. Paris **342** (2006), no. 6, 365–446, arXiv:math/0605061.
- [16] ———, *Polynomial realization of some trialgebras*, FPSAC’06 (San Diego) (2006), arXiv:math/0605061v1.
- [17] Richard P. Stanley, *Ordered structures and partitions*, American Mathematical Society, Providence, R.I., 1972, Memoirs of the American Mathematical Society, No. 119.
- [18] R. E. Stong, *Finite topological spaces*, Trans. Amer. Math. Soc. **123** (1966), 325–340.
- [19] David Wright and Wenhua Zhao, *D -log and formal flow for analytic isomorphisms of n -space*, Trans. Amer. Math. Soc. **355** (2003), no. 8, 3117–3141 (electronic), arXiv:math/0209274.