



**HAL**  
open science

# Necessary and sufficient conditions for the convergence of the consistent maximal displacement of the branching random walk

Bastien Mallein

► **To cite this version:**

Bastien Mallein. Necessary and sufficient conditions for the convergence of the consistent maximal displacement of the branching random walk. *Brazilian Journal of Probability and Statistics*, 2019, 33 (2). hal-01322467

**HAL Id: hal-01322467**

**<https://hal.science/hal-01322467>**

Submitted on 27 May 2016

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Convergence of the consistent maximal displacement of the branching random walk

Bastien Mallein\*

May 27, 2016

## Abstract

We consider a supercritical branching random walk on  $\mathbb{R}$ . The consistent maximal displacement is the smallest of the distances between the trajectories of individuals at the  $n$ th generation and the boundary of the process. It has been proved by Fang and Zeitouni [7] and by Faraud, Hu and Shi [8] that the consistent maximal displacement grows almost surely at rate  $\lambda^* n^{1/3}$  for an explicit  $\lambda^*$ . We obtain here a necessary and sufficient condition for this asymptotic behaviour to hold.

## 1 Introduction

A branching random walk on  $\mathbb{R}$  is a process defined as follows. It starts with one individual located at 0 at time 0. Its children are positioned on  $\mathbb{R}$  according to the law  $\mathcal{L}$  of a point process, and form the first generation of the process. Then for any  $n \in \mathbb{N}$ , each individual in the  $n$ -th generation makes children around its current position according to an independent point process with law  $\mathcal{L}$ . We write  $\mathbf{T}$  for the genealogical tree of the population. For any  $u \in \mathbf{T}$  we denote by  $V(u)$  the position of the individual  $u$  and by  $|u|$  the generation to which  $u$  belongs. The random marked tree  $(\mathbf{T}, V)$  is the branching random walk with reproduction law  $\mathcal{L}$ . We assume the Galton-Watson tree  $\mathbf{T}$  is supercritical:

$$\mathbb{E} \left( \sum_{|u|=1} 1 \right) > 1, \quad (1.1)$$

and we write  $S = \{\#\mathbf{T} = +\infty\}$  the survival event. We also assume the branching random walk  $(\mathbf{T}, V)$  is in the boundary case (in the sense of [5])

$$\mathbb{E} \left( \sum_{|u|=1} e^{-V(u)} \right) = 1 \quad \text{and} \quad \mathbb{E} \left( \sum_{|u|=1} V(u) e^{-V(u)} \right) = 0. \quad (1.2)$$

Under these assumptions, Biggins [4] proved that  $\frac{1}{n} \max_{|u|=n} V(u)$  converges to 0 almost surely on  $S$ . Any branching random walk with mild integrability

---

\*DMA, École Normale Supérieure (Paris).

assumption can be normalized to be in the boundary case, see e.g. Bérard and Gouéré [3]. We also assume that

$$\sigma^2 := \mathbb{E} \left( \sum_{|u|=1} V(u)^2 e^{-V(u)} \right) < +\infty. \quad (1.3)$$

Let  $n \geq 0$ . For any  $u \in \mathbf{T}$  such that  $|u| = n$  and  $k \leq n$  we denote by  $u_k$  the ancestor of  $u$  alive at generation  $k$ . The consistent maximal displacement of the branching random walk is the quantity defined as

$$L_n := \min_{|u|=n} \max_{k \leq n} V(u_k).$$

It correspond to the distance between the boundary of the branching random walk and the individual that stayed as close as possible to it. The asymptotic behaviour of  $L_n$  has been studied by Fang and Zeitouni [7] and by Fauraud, Hu and Shi [8]. Under stronger integrability assumptions, they proved that  $L_n$  behaves as  $\lambda^* n^{1/3}$  almost surely for some explicit  $\lambda^*$ . The main result of this article is a necessary and sufficient condition for this asymptotic behaviour to hold. Roberts [15] computed the second order of the asymptotic behaviour of  $L_n$  for the branching Brownian motion.

We now introduce the integrability assumption

$$\lim_{x \rightarrow +\infty} x^2 \mathbb{E} \left( \sum_{|u|=1} e^{-V(u)} \mathbf{1}_{\left\{ \log \left( \sum_{|v|=1} e^{-V(v)} \right) \geq x \right\}} \right) = 0. \quad (1.4)$$

Observe that (1.4) is strictly weaker than the classical integrability assumption that is [1, Assumption (1.4)]. This stronger assumption is necessary and sufficient to obtain the asymptotic behaviour of many quantities associated to the branching random walk, such as the minimal displacement, or the derivative martingale (see [1, 6]).

**Theorem 1.1.** *We assume (1.1), (1.2) and (1.3) hold. Then (1.4) is a necessary and sufficient condition for*

$$\lim_{n \rightarrow +\infty} \frac{L_n}{n^{1/3}} = \left( \frac{3\pi^2 \sigma^2}{2} \right)^{1/3} \quad \text{a.s. on } S.$$

If (1.4) is changed into

$$\lim_{x \rightarrow +\infty} x^2 \mathbb{E} \left( \sum_{|u|=1} e^{-V(u)} \mathbf{1}_{\left\{ \log \left( \sum_{|v|=1} e^{-V(v)} \right) \geq x \right\}} \right) = \rho \in (0, +\infty),$$

we were not able to compute the precise asymptotic behaviour of  $\frac{L_n}{n^{1/3}}$ . Using the methods developed in this article, it can be proved that a.s. on  $S$ , for  $n$  large enough we have

$$\left( \frac{3\pi^2 \sigma^2}{2} + 3\rho \right)^{1/3} \leq \frac{L_n}{n^{1/3}} \leq \left( \frac{3\pi^2 \sigma^2}{2} \right)^{1/3} + (3\rho)^{1/3}.$$

The rest of the article is organised as follows. In Section 2 we introduce the spinal decomposition of the branching random walk and the Mogul'skiĭ's small deviations estimate. These results are used to bound the left tail of  $L_n$  in Section 3, yielding to its a.s. asymptotic behaviour.

## 2 Preliminary results

### 2.1 Spinal decomposition of the branching random walk

For  $n \in \mathbb{N}$ , we write  $W_n = \sum_{|u|=n} e^{-V(u)}$  and  $\mathcal{F}_n = \sigma(u, V(u), |u| \leq n)$ . Under assumption (1.2),  $(W_n)$  is a non-negative  $(\mathcal{F}_n)$ -martingale. We introduce the probability  $\overline{\mathbb{P}}$  such that for any  $n \in \mathbb{N}$ ,  $\overline{\mathbb{P}}|_{\mathcal{F}_n} = W_n \cdot \mathbb{P}|_{\mathcal{F}_n}$ .

The spinal decomposition consists in an alternative description of  $\overline{\mathbb{P}}$  as a branching random walk with a distinguished individual with a different reproduction law. It generalizes a similar construction for Galton-Watson processes, that can be found in [11]. This result has been proved by Lyons in [10]. Let  $\mathbf{T}$  be a tree, a spine of  $\mathbf{T}$  is a sequence  $w = (w_n) \in \mathbf{T}^{\mathbb{N}}$  such that  $|w_n| = n$  and for any  $k \leq n$ ,  $(w_n)_k = w_k$ . We write  $\widehat{\mathcal{L}}$  for the law of the point process  $(V(u), |u| = 1)$  under the law  $\overline{\mathbb{P}}$ .

We now define the law  $\widehat{\mathbb{P}}$  of a branching random walk with spine  $(\mathbf{T}, V, w)$ . It starts with a unique individual  $w_0$  located at 0 at time 0. Its children are positioned according to a point process of law  $\widehat{\mathcal{L}}$ . The individual  $w_1$  is then chosen at random among these children  $u$  with probability proportional to  $e^{-V(u)}$ . Similarly at each generation  $n$ , every individual  $u$  makes children independently, according to law  $\widehat{\mathcal{L}}$  if  $u \neq w_n$  and  $\widehat{\mathcal{L}}$  otherwise; and  $w_{n+1}$  is chosen at random among the children  $v$  of  $w_n$  with probability proportional to  $e^{-V(v)}$ .

**Proposition 2.1** (Spinal decomposition, Lyons [10]). *Assuming (1.2) and (1.3), for any  $n \in \mathbb{N}$ , we have  $\widehat{\mathbb{P}}|_{\mathcal{F}_n} = \overline{\mathbb{P}}|_{\mathcal{F}_n}$ , and for any  $|u| = n$ ,*

$$\widehat{\mathbb{P}}(w_n = u | \mathcal{F}_n) = e^{-V(u)} / W_n,$$

and  $(V(w_n), n \geq 0)$  is a centred random walk with variance  $\sigma^2$ .

### 2.2 Small deviations estimate for enriched random walk

Let  $(X_n, \xi_n)_{n \in \mathbb{N}}$  be a sequence of i.i.d. vectors in  $\mathbb{R}^2$  such that

$$\mathbb{E}(X_n) = 0 \quad \text{and} \quad \mathbb{E}(X_n^2) = \sigma^2 \in (0, +\infty). \quad (2.1)$$

We set  $\rho_+ = \limsup_{x \rightarrow +\infty} x^2 \mathbb{P}(\xi_1 \geq x)$  and  $\rho_- = \liminf_{x \rightarrow +\infty} x^2 \mathbb{P}(\xi_1 \geq x)$ .

We write  $T_n = T_0 + X_1 + \dots + X_n$  and we call  $((T_n, \xi_n), n \geq 0)$  an enriched random walk. For any  $z \in \mathbb{R}$ ,  $\mathbb{P}_z$  is the probability such that  $\mathbb{P}_z(T_0 = z) = 1$ . We simply write  $\mathbb{P}$  for  $\mathbb{P}_0$ . We study in this section the probability that an enriched random walk stays during  $n$  unites of time in an interval of width  $o(n^{1/2})$ , generalizing the Mogul'skiĭ small deviations estimate [14].

**Theorem 2.2.** *Let  $(a_n)$  be a sequence of real non-negative numbers such that  $\lim_{n \rightarrow +\infty} a_n = +\infty$  and  $\lim_{n \rightarrow +\infty} \frac{a_n^2}{n} = 0$ . Assuming (2.1), for any continuous functions  $f < g$  and  $h > 0$ , for any  $x, x'$  such that  $f(0) < x \leq 0 \leq x' < g(0)$ , we have*

$$\begin{aligned} \limsup_{n \rightarrow +\infty} \frac{a_n^2}{n} \log \sup_{z \in [x, x']} \mathbb{P}_{z a_n} \left( \frac{T_j}{a_n} \in [f(j/n), g(j/n)], \frac{\xi_j}{a_n} \leq h(j/n), j \leq n \right) \\ = - \int_0^1 \frac{\pi^2 \sigma^2}{2(g(s) - f(s))^2} + \frac{\rho_-}{h(s)^2} ds, \quad (2.2) \end{aligned}$$

$$\begin{aligned} \liminf_{n \rightarrow +\infty} \frac{a_n^2}{n} \log \inf_{z \in [x, x']} \mathbb{P}_{za_n} \left( \frac{T_j}{a_n} \in [f(j/n), g(j/n)], \frac{\xi_j}{a_n} \leq h(j/n), j \leq n \right) \\ = - \int_0^1 \frac{\pi^2 \sigma^2}{2(g(s) - f(s))^2} + \frac{\rho_+}{h(s)^2} ds. \end{aligned} \quad (2.3)$$

We prove this result using the same techniques as in [13, Lemma 2.6]. We prove in a first time the following result.

**Lemma 2.3.** *Under the assumptions of Theorem 2.2, if  $\rho_+ = \rho_- =: \rho \in [0, +\infty]$  then for any  $a < 0 < b$ ,  $\lambda > 0$  and  $a \leq c < d \leq b$ , we have*

$$\lim_{n \rightarrow +\infty} \frac{a_n^2}{n} \log \mathbb{P} \left( \frac{T_n}{a_n} \in [c, d], \frac{T_j}{a_n} \in [a, b], \frac{\xi_j}{a_n} \leq \lambda, j \leq n \right) = - \frac{\pi^2 \sigma^2}{2(b-a)^2} - \frac{\rho}{\lambda^2}.$$

*Proof.* Note that

$$\mathbb{P}(|T_j| < ha_n, \xi_j \leq \lambda a_n, j \leq n) \leq \mathbb{P}(\xi_1 \leq \lambda a_n)^n \leq \exp(-n\mathbb{P}(\xi_1 > \lambda a_n)).$$

Consequently, if  $\rho = +\infty$ , the proof is immediate. We assume in the rest of the proof that  $\rho < +\infty$ .

Let  $B$  be a Brownian motion and  $P$  an independent Poisson process with intensity 1. For any  $n \in \mathbb{N}$  and  $k \leq n$ , we denote by  $P_k^{(n)} = \sum_{j=1}^k \mathbf{1}_{\{\xi_j \geq \lambda a_n\}}$ . By [16, Theorem 5.1], for any  $T > 0$  we have

$$\lim_{n \rightarrow +\infty} \left( \frac{T_{ia_n^2}}{a_n}, P_{ia_n^2}^{(n)}, t \leq T \right) = (B_{\sigma^2 t}, P_{\rho t / \lambda^2}, t \leq T) \quad (2.4)$$

in the Skorohod sense.

Let  $a < 0 < b$ ,  $a \leq c < d \leq b$  and  $T > 0$ . We set  $r_n = \lfloor Ta_n^2 \rfloor$ . Applying the Markov property, for any  $\epsilon > 0$  small enough, we have

$$\begin{aligned} \left( \inf_{\substack{h \leq b-a-4\epsilon \\ |x| \leq \epsilon}} \mathbb{P}_{xa_n} \left( \left| \frac{T_{r_n}}{a_n} - h \right| \leq \epsilon, \frac{T_j}{a_n} \in [a, b], \frac{\xi_j}{a_n} \leq \lambda, j \leq r_n \right) \right)^{\lfloor n/r_n \rfloor} \\ \leq \mathbb{P} \left( \frac{T_n}{a_n} \in [c, d], \frac{T_j}{a_n} \in [a, b], \frac{\xi_j}{a_n} \leq \lambda, j \leq n \right) \\ \leq \left( \sup_{x \in [a, b]} \mathbb{P}_{xa_n} \left( \frac{T_j}{a_n} \in [a, b], \frac{\xi_j}{a_n} \leq \lambda, j \leq r_n \right) \right)^{\lfloor n/r_n \rfloor}. \end{aligned}$$

Consequently, letting  $n \rightarrow +\infty$  and using (2.4), we obtain:

$$\begin{aligned} \liminf_{n \rightarrow +\infty} \frac{a_n^2}{n} \log \mathbb{P} \left( \frac{T_n}{a_n} \in [c, d], \frac{T_j}{a_n} \in [a, b], \frac{\xi_j}{a_n} \leq \lambda, j \leq n \right) \\ \geq \frac{1}{T} \log \inf_{\substack{h \leq b-a-4\epsilon \\ |x| \leq \epsilon}} \mathbb{P}_x (|B_{\sigma^2 T} - h| \leq \epsilon, P_{\rho T / \lambda^2} = 0, B_{\sigma^2 s} \in [a, b], s \leq T) \\ \geq - \frac{\rho}{\lambda^2} + \frac{1}{T} \log \inf_{\substack{h \leq b-a-4\epsilon \\ |x| \leq \epsilon}} \mathbb{P}_x (|B_{\sigma^2 T} - h| \leq \epsilon, B_{\sigma^2 s} \in [a, b], s \leq T), \end{aligned}$$

$$\begin{aligned}
& \limsup_{n \rightarrow +\infty} \frac{a_n^2}{n} \log \mathbb{P} \left( \frac{T_j}{a_n} \in [a, b], \frac{\xi_j}{a_n} \leq \lambda, j \leq n \right) \\
& \leq \frac{1}{T} \log \sup_{x \in [a, b]} \mathbb{P} (P_{\rho T / \lambda^2} = 0, B_{\sigma^2 s} \in [a, b], s \leq T) \\
& \leq -\frac{\rho}{\lambda^2} + \frac{1}{T} \log \sup_{x \in [a, b]} \mathbb{P} (B_{\sigma^2 s} \in [a, b], s \leq T).
\end{aligned}$$

Letting  $T \rightarrow +\infty$ , using e.g. [9, Chapter 1.7, Problem 8], we conclude that

$$\lim_{n \rightarrow +\infty} \frac{a_n^2}{n} \log \mathbb{P} \left( \frac{T_j}{a_n} \in [a, b], \frac{\xi_j}{a_n} \leq \lambda, j \leq n \right) = -\frac{\pi^2 \sigma^2}{2(b-a)^2} - \frac{\rho}{\lambda^2}.$$

□

**Corollary 2.4.** *Under the assumptions of Theorem 2.2, for any  $a < 0 < b$ ,  $\lambda > 0$  and  $a \leq c < d \leq b$ , we have*

$$\begin{aligned}
& \limsup_{n \rightarrow +\infty} \frac{a_n^2}{n} \log \mathbb{P} \left( \frac{T_j}{a_n} \in [a, b], \frac{\xi_j}{a_n} \leq \lambda, j \leq n \right) = -\frac{\pi^2 \sigma^2}{2(b-a)^2} - \frac{\rho_-}{\lambda^2}, \\
& \liminf_{n \rightarrow +\infty} \frac{a_n^2}{n} \log \mathbb{P} \left( \frac{T_n}{a_n} \in [c, d], \frac{T_j}{a_n} \in [a, b], \frac{\xi_j}{a_n} \leq \lambda, j \leq n \right) = -\frac{\pi^2 \sigma^2}{2(b-a)^2} - \frac{\rho_+}{\lambda^2}.
\end{aligned}$$

*Proof.* The two equations being proved the same way, we only prove the first one. As  $\mathbb{P}(|T_j| < ha_n, \xi_j \leq n, j \leq n) \leq \mathbb{P}(\xi_1 \leq n)^n \leq \exp(-n\mathbb{P}(\xi_1 > n))$ , we only consider the case  $\rho_- < +\infty$ .

There exists a subsequence  $(n_k)$  such that  $a_{n_k}^2 \mathbb{P}(\xi_1 > \lambda a_{n_k})$  converges to  $\rho_- / \lambda^2$ , therefore by Lemma 2.3,

$$\begin{aligned}
& \limsup_{n \rightarrow +\infty} \frac{a_n^2}{n} \log \mathbb{P} \left( \frac{T_j}{a_n} \in [a, b], \frac{\xi_j}{a_n} \leq \lambda, j \leq n \right) \\
& \geq \lim_{k \rightarrow +\infty} \frac{a_{n_k}^2}{n_k} \log \mathbb{P} \left( \frac{T_j}{a_{n_k}} \in [a, b], \frac{\xi_j}{a_{n_k}} \leq \lambda, j \leq n_k \right) = -\frac{\pi^2 \sigma^2}{2(b-a)^2} - \frac{\rho_-}{\lambda^2}.
\end{aligned}$$

Moreover, we may couple the random variable  $\xi_j$  with a random variable  $\tilde{\xi}_j$  such that  $\tilde{\xi}_j \leq \xi_j$  and  $\lim_{n \rightarrow +\infty} x^2 \mathbb{P}(\tilde{\xi}_1 \geq x) = \rho_-$ . By Lemma 2.3 again,

$$\begin{aligned}
& \limsup_{n \rightarrow +\infty} \frac{a_n^2}{n} \log \mathbb{P} \left( \frac{T_j}{a_n} \in [a, b], \frac{\xi_j}{a_n} \leq \lambda, j \leq n \right) \\
& \leq \limsup_{n \rightarrow +\infty} \frac{a_n^2}{n} \log \mathbb{P} \left( \frac{T_j}{a_n} \in [a, b], \frac{\tilde{\xi}_j}{a_n} \leq \lambda, j \leq n \right) = -\frac{\pi^2 \sigma^2}{2(b-a)^2} - \frac{\rho_-}{\lambda^2},
\end{aligned}$$

concluding the proof. □

*Proof of Theorem 2.2.* We first prove that for any  $a < b$  and  $\lambda > 0$ ,

$$\limsup_{n \rightarrow +\infty} \frac{a_n^2}{n} \log \sup_{x \in [a, b]} \mathbb{P}_{x a_n} \left( \frac{T_j}{a_n} \in [a, b], \frac{\xi_j}{a_n} \leq \lambda, j \leq n \right) = -\frac{\pi^2 \sigma^2}{2(b-a)^2} - \frac{\rho_-}{\lambda^2}. \tag{2.5}$$

Let  $\delta > 0$ , we write  $M = \lceil \frac{b-a}{\delta} \rceil$ . We observe that

$$\begin{aligned} & \sup_{x \in [a, b]} \mathbb{P}_{xa_n} \left( \frac{T_j}{a_n} \in [a, b], \frac{\xi_j}{a_n} \leq \lambda, j \leq n \right) \\ & \leq \max_{0 \leq m < M} \sup_{x \in [a+m\delta, a+(m+1)\delta]} \mathbb{P}_{xa_n} \left( \frac{T_j}{a_n} \in [a, b], \frac{\xi_j}{a_n} \leq \lambda, j \leq n \right) \\ & \leq \max_{0 \leq m < M} \mathbb{P} \left( \frac{T_j}{a_n} \in [-(m+1)\delta, b-a-m\delta], \frac{\xi_j}{a_n} \leq \lambda, j \leq n \right). \end{aligned}$$

We apply Corollary 2.4 and let  $\delta \rightarrow 0$  to obtain (2.5). Then we approach functions  $f$ ,  $g$  and  $h$  by staircase functions, using (2.5) we obtain (2.2).

For any  $a < 0 < b$ ,  $\lambda > 0$ ,  $y \in [a, b]$  and  $\epsilon > 0$  small enough, we prove in a second time that

$$\begin{aligned} \liminf_{n \rightarrow +\infty} \frac{a_n^2}{n} \log \inf_{|x| \leq \epsilon} \mathbb{P}_{xa_n} \left( \left| \frac{T_n}{a_n} - y \right| \leq \epsilon, \frac{T_j}{a_n} \in [a, b], \frac{\xi_j}{a_n} \leq \lambda, j \leq n \right) \\ = -\frac{\pi^2 \sigma^2}{2(b-a)^2} - \frac{\rho_+}{\lambda^2}. \end{aligned} \quad (2.6)$$

Let  $\epsilon > 0$  small enough such that  $[-2\epsilon, 2\epsilon] \subset [a, b]$ , let  $0 < \delta < \epsilon$ , we set  $M = \lceil \frac{2\delta}{\epsilon} \rceil$ . We have again

$$\begin{aligned} & \inf_{|x| \leq \epsilon} \mathbb{P}_{xa_n} \left( \left| \frac{T_n}{a_n} - y \right| \leq \epsilon, \frac{T_j}{a_n} \in [a, b], \xi_j \leq n, j \leq n \right) \\ & \geq \min_{0 \leq m < M} \inf_{x \in [-\epsilon+m\delta, -\epsilon+(m+1)\delta]} \mathbb{P}_{xa_n} \left( \left| \frac{T_n}{a_n} - y \right| \leq \epsilon, \frac{T_j}{a_n} \in [a, b], \xi_j \leq n, j \leq n \right) \\ & \geq \min_{0 \leq m < M} \mathbb{P} \left( \begin{array}{l} \frac{T_j}{a_n} \in [a + \epsilon - m\delta, b + \epsilon - (m+1)\delta], \xi_j \leq n, j \leq n \\ \frac{T_n}{a_n} \in [y - m\delta, y + 2\epsilon - (m+1)\delta] \end{array} \right). \end{aligned}$$

Applying again Corollary 2.4, (2.6) holds. We finally approach functions  $f$ ,  $g$  and  $h$  by staircase functions and use the Markov property, to obtain the lower bound (2.3).  $\square$

### 3 Tail of the consistent maximal displacement

For any  $u \in \mathbf{T}$ , we write  $\pi u$  for the parent of  $u$ ,  $\Omega(u)$  for the set of children of  $u$ ,

$$\tilde{\xi}(u) = \log \sum_{v \in \Omega(u)} e^{V(u) - V(v)} \quad \text{and} \quad \xi(u) = \tilde{\xi}(\pi u).$$

Note that by (1.2), we have  $\mathbb{P}(\tilde{\xi}(u) \geq x) \leq e^{-x}$  for any  $x \geq 0$ . We introduce

$$\rho_+ = \limsup_{x \rightarrow +\infty} x^2 \widehat{\mathbb{P}} \left( \tilde{\xi}(w_0) \geq x \right) \quad \text{and} \quad \rho_- = \liminf_{x \rightarrow +\infty} x^2 \widehat{\mathbb{P}} \left( \tilde{\xi}(w_0) \geq x \right) \quad (3.1)$$

$$\lambda_+^* = \left( \frac{3\pi^2 \sigma^2}{2} + 3\rho_+ \right)^{1/3}, \quad \lambda_-^* = \left( \frac{3\pi^2 \sigma^2}{2} + 3\rho_- \right)^{1/3} \quad \text{and} \quad \lambda^* = \left( \frac{3\pi^2 \sigma^2}{2} \right)^{1/3}. \quad (3.2)$$

We use here the results of the previous section to obtain upper and lower bounds for the left tail of  $L_n$ . We first provide an upper bound.

**Lemma 3.1.** *We assume (1.1), (1.2) and (1.3). For any  $\lambda \in (0, \lambda^*)$ , we have*

$$\limsup_{n \rightarrow +\infty} \frac{1}{n^{1/3}} \log \mathbb{P} \left( L_n \leq \lambda n^{1/3} \right) \leq \lambda - \lambda_-^*, \quad (3.3)$$

and for any  $\lambda \in (0, \lambda_+^*)$ , we have

$$\liminf_{n \rightarrow +\infty} \frac{1}{n^{1/3}} \log \mathbb{P} \left( L_n \leq \lambda n^{1/3} \right) \leq \lambda - \lambda_+^*. \quad (3.4)$$

*Proof.* Let  $\lambda > 0$ ,  $f$  be a continuous increasing function such that  $f(0) < 0$  and  $f(1) = \lambda$ , and  $h = \lambda - f$ . We set  $I_k^{(n)} = [f(k/n)n^{1/3}, \lambda n^{1/3}]$  for  $k \leq n$  and we denote by

$$G_n = \left\{ u \in \mathbf{T} : |u| \leq n, V(u_j) \in I_j^{(n)}, \xi(u_j) \leq h(j/n)n^{1/3}, j \leq |u| \right\}.$$

We introduce the quantities

$$\begin{aligned} X_k^{(n)} &= \sum_{|u|=k} \mathbf{1}_{\{V(u) < f(k/n)n^{1/3}\}} \mathbf{1}_{\{\pi u \in G_n\}} \quad \text{and} \\ Y_k^{(n)} &= \sum_{|u|=k-1} \mathbf{1}_{\{\tilde{\xi}(u) > h(k/n)n^{1/3}\}} \mathbf{1}_{\{u \in G_n\}}. \end{aligned}$$

We observe that

$$\begin{aligned} \mathbb{P} \left( L_n \leq \lambda n^{1/3} \right) &= \mathbb{P} \left( \exists |u| = n : V(u_j) \leq \lambda n^{1/3}, j \leq n \right) \\ &\leq \mathbb{P} \left( \sum_{j=1}^n X_j^{(n)} + Y_j^{(n)} \geq 1 \right) \leq \sum_{j=1}^n \mathbb{E} \left( X_j^{(n)} + Y_j^{(n)} \right). \end{aligned} \quad (3.5)$$

For any  $k \leq n$ , using the spinal decomposition we have

$$\begin{aligned} \mathbb{E} \left( X_k^{(n)} \right) &= \widehat{\mathbb{E}} \left( \sum_{|u|=k} \frac{e^{-V(u)}}{W_k} e^{V(u)} \mathbf{1}_{\{V(u) < f(k/n)n^{1/3}\}} \mathbf{1}_{\{\pi u \in G_n\}} \right) \\ &= \widehat{\mathbb{E}} \left( e^{V(w_k)} \mathbf{1}_{\{V(w_k) < f(k/n)n^{1/3}\}} \mathbf{1}_{\{w_{k-1} \in G_n\}} \right) \\ &\leq e^{f(k/n)n^{1/3}} \widehat{\mathbb{P}}(w_{k-1} \in G_n). \end{aligned}$$

Moreover, as  $\xi(u)$  is independent of  $\mathcal{F}_{|u|}$ , we also note that

$$\begin{aligned} \mathbb{E} \left( Y_k^{(n)} \right) &= \mathbb{E} \left( \sum_{|u|=k-1} \mathbf{1}_{\{u \in G_n\}} \right) \mathbb{P} \left( \tilde{\xi}(u) > h(k/n)n^{1/3} \right) \\ &\leq e^{-h(k/n)n^{1/3}} \widehat{\mathbb{E}} \left( e^{V(w_{k-1})} \mathbf{1}_{\{w_{k-1} \in G_n\}} \right) \leq e^{f(k/n)n^{1/3}} \widehat{\mathbb{P}}(w_{k-1} \in G_n). \end{aligned}$$

Consequently, (3.5) becomes

$$\mathbb{P} \left( L_n \leq \lambda n^{1/3} \right) \leq 2 \sum_{k=1}^n e^{f(k/n)n^{1/3}} \widehat{\mathbb{P}}(w_{k-1} \in G_n).$$



We set  $A > 0$ , for any  $a \leq A$ , we write  $m_a = \lfloor na/A \rfloor$ . As  $f$  is increasing, for any  $k \in (m_a, m_{a+1}]$ , we have

$$e^{f(k/n)n^{1/3}} \widehat{\mathbb{P}}(w_{k-1} \in G_n) \leq e^{f((a+1)/A)n^{1/3}} \widehat{\mathbb{P}}(w_{m_a} \in G_n).$$

Moreover, by the spinal decomposition,  $(V(w_j), \xi(w_j))$  is an enriched random walk under law  $\widehat{\mathbb{P}}$ . We apply Theorem 2.2 to obtain for any  $a \leq A$ ,

$$\begin{aligned} \limsup_{n \rightarrow +\infty} \frac{1}{n^{1/3}} \log \widehat{\mathbb{P}}(w_{m_a} \in G_n) &= - \int_0^{a/A} \frac{3(\lambda_-^*)^3}{(\lambda - f(s))^2} ds, \\ \liminf_{n \rightarrow +\infty} \frac{1}{n^{1/3}} \log \widehat{\mathbb{P}}(w_{m_a} \in G_n) &= - \int_0^{a/A} \frac{3(\lambda_+^*)^3}{(\lambda - f(s))^2} ds. \end{aligned}$$

We conclude that, letting  $n \rightarrow +\infty$  then  $A \rightarrow +\infty$  we have

$$\limsup_{n \rightarrow +\infty} \frac{1}{n^{1/3}} \log \mathbb{P}(L_n \leq \lambda n^{1/3}) \leq \sup_{t \in [0,1]} f(t) - \int_0^t \frac{3(\lambda_-^*)^3}{(\lambda - f(s))^2} ds, \quad (3.6)$$

$$\liminf_{n \rightarrow +\infty} \frac{1}{n^{1/3}} \log \mathbb{P}(L_n \leq \lambda n^{1/3}) \leq \sup_{t \in [0,1]} f(t) - \int_0^t \frac{3(\lambda_+^*)^3}{(\lambda - f(s))^2} ds. \quad (3.7)$$

Optimizing these equations in  $f$  concludes the proof.

Indeed, to prove (3.3), we consider the following two cases. If  $\rho_- = +\infty$  then  $\lambda_-^* = +\infty$ . Choosing a sequence of functions  $f_M$  such that  $f_M(0) < -M$ , and letting  $M \rightarrow +\infty$ , (3.6) yields

$$\limsup_{n \rightarrow +\infty} \frac{1}{n^{1/3}} \log \mathbb{P}(L_n \leq \lambda n^{1/3}) = -\infty.$$

If  $\rho_- < +\infty$ , then setting  $f(t) = \lambda - \lambda_-^*(1-t)^{1/3}$  we have

$$\begin{aligned} \limsup_{n \rightarrow +\infty} \frac{1}{n^{1/3}} \log \mathbb{P}(L_n \leq \lambda n^{1/3}) &\leq \sup_{t \in [0,1]} f(t) - \int_0^t \frac{3(\lambda_-^*)^3}{(\lambda - f(s))^2} ds \\ &\leq \sup_{t \in [0,1]} f(t) - \int_0^t \frac{3\lambda_-^*}{(1-s)^{2/3}} ds = \lambda - \lambda_-^*. \end{aligned}$$

A similar reasoning leads to (3.4).  $\square$

**Corollary 3.2.** *Under assumptions (1.1), (1.2) and (1.3), we have*

$$\liminf_{n \rightarrow +\infty} \frac{L_n}{n^{1/3}} \geq \lambda_-^* \quad \text{and} \quad \limsup_{n \rightarrow +\infty} \frac{L_n}{n^{1/3}} \geq \lambda_+^* \quad a.s.$$

*In particular, if  $\rho_+ > 0$  then  $\limsup_{n \rightarrow +\infty} \frac{L_n}{n^{1/3}} > \lambda^*$  a.s.*

*Proof.* By Lemma 3.1, for any  $\epsilon > 0$ ,  $\sum_{n=1}^{+\infty} \mathbb{P}(\frac{L_n}{n^{1/3}} \geq (\lambda_-^* + \epsilon)) < +\infty$ , and there exists an increasing sequence  $(n_k)$  such that  $\sum_{k=1}^{+\infty} \mathbb{P}(\frac{L_{n_k}}{n_k^{1/3}} \geq (\lambda_+^* + \epsilon)) < +\infty$ .

We conclude the proof by Borel-Cantelli lemma.  $\square$

In a second time, to bound from above the consistent maximal displacement, we prove that with high probability there exists an individual staying below  $\lambda n^{1/3}$  for  $n$  units of time, as soon as  $\lambda$  is large enough. To do so, we compute the first two moments of the number of individuals staying in two well-chosen lines, while making “not too many children”.

**Lemma 3.3.** *Assuming (1.1), (1.2), (1.3) and (1.4) (i.e.  $\rho_+ = \rho_- = 0$ ), for any  $0 < \lambda < \lambda^*$ , we have  $\liminf_{n \rightarrow +\infty} \frac{1}{n^{1/3}} \log \mathbb{P}(L_n \leq \lambda n^{1/3}) \geq \lambda - \lambda^*$ .*

*Proof.* Let  $\lambda \in (0, \lambda^*)$ ,  $\delta > 0$ , and  $f : t \in [0, 1] \mapsto \lambda - \lambda^*(1 + \delta - t)^{1/3}$ . We denote by  $I_j^{(n)} = [f(j/n)n^{1/3}, \lambda n^{1/3}]$  for  $j \leq n$ . We set

$$Z_n = \sum_{|u|=n} \mathbf{1}_{\{V(u_j) \in I_j^{(n)}, \xi(u_j) \leq \delta n^{1/3}, j \leq n\}}.$$

We compute the first two moments of  $Z_n$  to bound from below  $\mathbb{P}(Z_n > 0)$ .

By spinal decomposition, we have

$$\begin{aligned} \mathbb{E}(Z_n) &= \widehat{\mathbb{E}} \left( e^{V(w_n)} \mathbf{1}_{\{V(w_j) \in I_j^{(n)}, \xi(w_j) \leq \delta n^{1/3}, j \leq n\}} \right) \\ &\geq e^{f(1)n^{1/3}} \widehat{\mathbb{P}} \left( V(w_j) \in I_j^{(n)}, \xi(w_j) \leq \delta n^{1/3}, j \leq n \right). \end{aligned}$$

As  $\lim_{n \rightarrow +\infty} n^{2/3} \widehat{\mathbb{P}}(\xi(w_1) \geq \delta n^{1/3}) = 0$  by (1.4), Theorem 2.2 yields

$$\liminf_{n \rightarrow +\infty} \frac{1}{n^{1/3}} \log \mathbb{E}(Z_n) \geq f(1) - \frac{\pi^2 \sigma^2}{2} \int_0^1 \frac{ds}{(\lambda - f(s))^2} \geq \lambda - \lambda^*(1 + \delta)^{1/3}.$$

Similarly, to compute the second moment we observe that

$$\begin{aligned} \mathbb{E}(Z_n^2) &= \widehat{\mathbb{E}} \left( Z_n \sum_{|u|=n} \frac{e^{-V(u)}}{W_n} e^{V(u)} \mathbf{1}_{\{V(u_j) \in I_j^{(n)}, \xi(u_j) \leq \delta n^{1/3}, j \leq n\}} \right) \\ &= \widehat{\mathbb{E}} \left( Z_n e^{V(w_n)} \mathbf{1}_{\{V(w_j) \in I_j^{(n)}, \xi(w_j) \leq \delta n^{1/3}, j \leq n\}} \right) \\ &\leq e^{\lambda n^{1/3}} \widehat{\mathbb{E}} \left( Z_n \mathbf{1}_{\{V(w_j) \in I_j^{(n)}, \xi(w_j) \leq \delta n^{1/3}, j \leq n\}} \right). \end{aligned}$$

Under the law  $\widehat{\mathbb{P}}$ ,  $Z_n$  can be decomposed as follows

$$Z_n = \mathbf{1}_{\{V(w_j) \in I_j^{(n)}, \xi(w_j) \leq \delta n^{1/3}, j \leq n\}} + \sum_{k=0}^{n-1} \sum_{\substack{u \in \Omega(w_k) \\ u \neq w_{k+1}}} Z_n(u),$$

where  $Z_n(u) = \sum_{|v|=n, v > u} \mathbf{1}_{\{V(u_j) \in I_j^{(n)}, \xi(u_j) \leq \delta n^{1/3}, j \leq n\}}$ . We denote by

$$\mathcal{G} = \sigma(w_n, \Omega(w_n), V(u), u \in \Omega(w_n), n \geq 0).$$

Observe that conditionally on  $\mathcal{G}$ , for any  $u \in \Omega(w_k)$  such that  $u \neq w_{k+1}$ , the subtree of the descendants of  $u$  has the law of a branching random walk starting

from  $V(u)$ . Therefore, writing  $\mathbb{P}_x$  for the law of  $(\mathbf{T}, V + x)$ , for any  $k < n$  and  $u \in \Omega(w_k)$  such that  $u \neq w_{k+1}$ , we have

$$\begin{aligned}\widehat{\mathbb{E}}(Z_n(u)|\mathcal{G}) &\leq \mathbb{E}_{V(u)} \left( \sum_{|v|=n-k-1} \mathbf{1}_{\{V(v_j) \in I_{k+j+1}^{(n)}, \xi(v_j) \leq \delta n^{1/3}, j \leq n-k-1\}} \right) \\ &\leq \mathbb{E}_{V(u)} \left( \sum_{|v|=n-k-1} \mathbf{1}_{\{V(v_j) \in I_{k+j+1}^{(n)}, j \leq n-k-1\}} \right).\end{aligned}$$

Applying spinal decomposition, for any  $x \in \mathbb{R}$  and  $p \leq n$ ,

$$\begin{aligned}\mathbb{E}_x \left( \sum_{|v|=n-p} \mathbf{1}_{\{V(v_j) \in I_{p+j}^{(n)}, j \leq n-p\}} \right) &= \widehat{\mathbb{E}} \left( e^{V(w_{n-p}+x)} \mathbf{1}_{\{V(w_j)+x \in I_{p+j}^{(n)}, j \leq n-p\}} \right) \\ &\leq e^{\lambda n^{1/3}-x} \widehat{\mathbb{P}} \left( V(w_j) + x \in I_{p+j}^{(n)}, j \leq n-p \right).\end{aligned}$$

Let  $A > 0$ , for any  $a \leq A$  we set  $m_a = \lfloor na/A \rfloor$  and

$$\Psi_{a,A}^{(n)} = \sup_{y \in I_{m_a}^{(n)}} \widehat{\mathbb{P}} \left( V(w_j) + y \in I_{m_a+j}^{(n)}, j \leq n - m_a \right).$$

Using the previous equation, for any  $m_a \leq k < m_{a+1}$ , we have

$$\begin{aligned}\sum_{\substack{u \in \Omega(w_k) \\ u \neq w_{k+1}}} \widehat{\mathbb{E}}(Z_n(u)|\mathcal{G}) &\leq e^{\lambda n^{1/3}} \Psi_{a+1,A}^{(n)} \sum_{u \in \Omega(w_k)} e^{V(u)} \\ &\leq e^{\lambda n^{1/3}+V(w_k)+\xi(w_{k+1})} \Psi_{a+1,A}^{(n)}.\end{aligned}$$

As  $\xi(w_{k+1}) \leq \delta n^{1/3}$ , we obtain

$$\begin{aligned}\mathbb{E}(Z_n^2) &\leq e^{\lambda n^{1/3}} \widehat{\mathbb{P}} \left( V(w_j) \in I_j^{(n)}, j \leq n \right) \\ &\quad + e^{(2\lambda+\delta)n^{1/3}} \sum_{a=0}^{A-1} n \Psi_{a+1,A}^{(n)} e^{-f(a/A)n^{1/3}} \widehat{\mathbb{P}} \left( V(w_j) \in I_j^{(n)}, j \leq n \right).\end{aligned}$$

Therefore, applying Theorem 2.2, we have

$$\begin{aligned}\limsup_{n \rightarrow +\infty} \frac{1}{n^{1/3}} \log \mathbb{E}(Z_n^2) &\leq 2\lambda + \delta - \frac{\pi^2 \sigma^2}{2} \int_0^1 \frac{ds}{(\lambda - f(s))^2} \\ &\quad - \min_{a < A} f\left(\frac{a}{A}\right) - \frac{\pi^2 \sigma^2}{2} \int_{(a+1)/A}^1 \frac{ds}{(\lambda - f(s))^2}.\end{aligned}$$

Letting  $A \rightarrow +\infty$ , we obtain

$$\limsup_{n \rightarrow +\infty} \frac{1}{n^{1/3}} \log \mathbb{E}(Z_n^2) \leq \lambda + \delta - \lambda^*(1 + \delta)^{1/3}.$$

By Cauchy-Schwarz inequality,  $\mathbb{P}(L_n \leq \lambda n^{1/3}) \geq \mathbb{P}(Z_n > 0) \geq \frac{\mathbb{E}(Z_n)^2}{\mathbb{E}(Z_n^2)}$ , thus

$$\liminf_{n \rightarrow +\infty} \frac{1}{n^{1/3}} \log \mathbb{P}(L_n \leq \lambda n^{1/3}) \geq \lambda - \lambda^*(1 + \delta)^{1/3} - \delta.$$

Letting  $\delta \rightarrow 0$ , we conclude the proof.  $\square$

Note that Lemmas 3.1 and 3.3 imply that, assuming (1.4)

$$\lim_{n \rightarrow +\infty} \frac{1}{n^{1/3}} \log \mathbb{P}(L_n \leq \lambda n^{1/3}) = \lambda - \lambda^*. \quad (3.8)$$

We use Lemma 3.3 to obtain an a.s. behaviour of  $L_n$ .

**Lemma 3.4.** *Assuming (1.1), (1.2), (1.3) and (1.4), we have  $\lim_{n \rightarrow +\infty} \frac{L_n}{n^{1/3}} = \lambda^*$  almost surely on  $S$ .*

*Proof.* We have  $\liminf_{n \rightarrow +\infty} \frac{L_n}{n^{1/3}} \geq \lambda^*$  by Corollary 3.2. We now turn to the upper bound of  $L_n$ . By Lemma 3.3, for any  $\delta > 0$ ,

$$\liminf_{n \rightarrow +\infty} \frac{1}{n^{1/3}} \log \mathbb{P}(L_n \leq \lambda^* n^{1/3}) > -\delta.$$

We work in the rest of the proof conditionally on the survival event  $S$ . We write  $\widehat{\mathbf{T}}$  for the subtree of  $\mathbf{T}$  consisting of individuals having an infinite line of descent. By [2, Chapter 1, Theorem 12.1],  $\widehat{\mathbf{T}}$  is a supercritical Galton-Watson process that never dies out. Applying [12, Lemma 2.4] to the branching random walk  $(\widehat{\mathbf{T}}, V)$ , there exists  $a > 0$  and  $\rho > 1$  such that the event

$$\mathcal{A}(p) = \{\#\{|u| = p : \forall j \leq p, V(u_j) \geq -pa\} \geq \rho^p\}$$

is verified a.s. for  $p \geq 1$  large enough. Let  $\eta > 0$ , we set  $p = \lfloor \eta n^{1/3} \rfloor$ . Applying the Markov property at time  $p$ , we have

$$\mathbb{P}\left(L_{n+p} \geq (\lambda^* + A\eta)n^{1/3} | \mathcal{A}(p)\right) \leq \left(1 - \mathbb{P}\left(L_n \leq \lambda^* n^{1/3}\right)\right)^{\rho^p}.$$

Therefore by Borel-Cantelli lemma,  $\limsup_{n \rightarrow +\infty} \frac{L_n}{n^{1/3}} \leq \lambda^* + A\eta$  a.s. on  $S$ . We let  $\eta \rightarrow 0$  to conclude the proof.  $\square$

Theorem 1.1 is a consequence of Lemma 3.4 and Corollary 3.2: if (1.4) does not hold, then  $\rho_+ > 0$  and  $\frac{L_n}{n^{1/3}}$  does not converge toward  $\lambda^*$ .

## References

- [1] Elie Aïdékon. Convergence in law of the minimum of a branching random walk. *Ann. Probab.*, 41(3A):1362–1426, 2013.
- [2] K. B. Athreya and P. E. Ney. *Branching processes*. Dover Publications, Inc., Mineola, NY, 2004. Reprint of the 1972 original [Springer, New York; MR0373040].
- [3] J. Bérard and J.-B. Gouéré. Survival probability of the branching random walk killed below a linear boundary. *Electron. J. Probab.*, 16:no. 14, 396–418, 2011.
- [4] J. D. Biggins. The first- and last-birth problems for a multitype age-dependent branching process. *Advances in Appl. Probability*, 8(3):446–459, 1976.

- [5] J. D. Biggins and A. E. Kyprianou. Fixed points of the smoothing transform: the boundary case. *Electron. J. Probab.*, 10:no. 17, 609–631, 2005.
- [6] X. Chen. A necessary and sufficient condition for the non-trivial limit of the derivative martingale in a branching random walk. *Adv. in Appl. Probab.*, 47:no. 3, 741–760, 2015.
- [7] M. Fang and O. Zeitouni. Consistent minimal displacement of branching random walks. *Electron. Commun. Probab.*, 15:106–118, 2010.
- [8] G. Faraud, Y. Hu, and Z. Shi. Almost sure convergence for stochastically biased random walks on trees. *Probab. Theory Related Fields*, 154(3-4):621–660, 2012.
- [9] Kiyosi Itô and Henry P. McKean, Jr. *Diffusion processes and their sample paths*. Springer-Verlag, Berlin-New York, 1974. Second printing, corrected, Die Grundlehren der mathematischen Wissenschaften, Band 125.
- [10] R. Lyons. A simple path to Biggins’ martingale convergence for branching random walk. In *Classical and modern branching processes (Minneapolis, MN, 1994)*, volume 84 of *IMA Vol. Math. Appl.*, pages 217–221. Springer, New York, 1997.
- [11] R. Lyons, R. Pemantle, and Y. Peres. Conceptual proofs of  $L \log L$  criteria for mean behavior of branching processes. *Ann. Probab.*, 23(3):1125–1138, 1995.
- [12] B. Mallein. Branching random walk with selection at critical rate. *arXiv preprint*, 2015. arXiv:1502.07390.
- [13] B. Mallein.  $N$ -branching random walk with  $\alpha$ -stable spine. *arXiv preprint*, 2015. arXiv:1503.03762.
- [14] A. A. Mogul’skiĭ. Small deviations in the space of trajectories. *Teor. Veroyatnost. i Primenen.*, 19:755–765, 1974.
- [15] M. I. Roberts. Fine asymptotics for the consistent maximal displacement of branching Brownian motion. *Electron. J. Probab.*, 20:no. 28, 26, 2015.
- [16] A. V. Skorohod. Limit theorems for stochastic processes with independent increments. *Teor. Veroyatnost. i Primenen.*, 2:145–177, 1957.