



**HAL**  
open science

## **Drying of Indian subcontinent by rapid Indian Ocean warming and a weakening land-sea thermal gradient**

Mathew Koll Roxy, Kapoor Ritika, Pascal Terray, Raghu Murtugudde, Karumuri Ashok, B. N. Goswami

### ► **To cite this version:**

Mathew Koll Roxy, Kapoor Ritika, Pascal Terray, Raghu Murtugudde, Karumuri Ashok, et al.. Drying of Indian subcontinent by rapid Indian Ocean warming and a weakening land-sea thermal gradient. *Nature Communications*, 2015, 6, pp.7423. <10.1038/ncomms8423>. <hal-01322870>

**HAL Id: hal-01322870**

**<https://hal.science/hal-01322870v1>**

Submitted on 30 May 2016

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

# **Drying of Indian subcontinent by rapid Indian Ocean warming and a weakening land-sea thermal gradient.**

**Mathew Koll ROXY<sup>1</sup>, Kapoor RITIKA<sup>1,2</sup>, Pascal TERRAY<sup>3,4</sup>, Raghu MURTUGUDDE<sup>5</sup>,  
Karumuri ASHOK<sup>1</sup> and B N GOSWAMI<sup>1</sup>**

<sup>1</sup>*Centre for Climate Change Research, Indian Institute of Tropical Meteorology, Pune, India*

<sup>2</sup>*Department of Environmental Sciences, Fergusson College, University of Pune, Pune, India*

<sup>3</sup>*Sorbonne Universites (UPMC, Univ Paris 06)-CNRS-IRD-MNHN, LOCEAN Laboratory, 4  
place Jussieu, F-75005 Paris, France*

<sup>4</sup>*Indo-French Cell for Water Sciences, IISc-IITM-NIO-IRD Joint International Laboratory,  
IITM, Pune, India*

<sup>5</sup>*ESSIC, University of Maryland, College Park, Maryland, USA*

*Nature Communications*, 13<sup>th</sup> March 2015

---

Corresponding author address: Mathew Koll Roxy, Indian Institute of Tropical Meteorology,  
Pune 411008, India. E-mail: roxy@tropmet.res.in

1 **Abstract**

2 There are large uncertainties looming over the status and fate of the South Asian summer  
3 monsoon, with several studies debating whether the monsoon is weakening or strengthening in  
4 a changing climate. Our analysis using multiple observed datasets demonstrates a significant  
5 weakening trend in summer rainfall during 1901-2012 over central South Asia, extending from  
6 south of Pakistan through central India to Bangladesh, with up to 10-20% reduction over  
7 central-eastern India. Earlier studies have suggested an increase in moisture availability and  
8 land-sea thermal contrast in the tropics due anthropogenic warming, favoring an increase in  
9 tropical rainfall. In contrast, we notice a weakened land-sea thermal contrast due to rapid  
10 warming in the Indian Ocean, along with a relatively subdued warming over the subcontinent.  
11 Using long-term observations and coupled model experiments, we provide compelling  
12 evidence that the enhanced Indian Ocean warming, especially that over the western region,  
13 potentially weakens the land-sea thermal contrast, dampens the summer monsoon Hadley  
14 circulation, and thereby reduces the rainfall over South Asia.

15

16 **Introduction**

17 The South Asian summer monsoon rainfall during June to September plays a significantly large  
18 role in the *roti and dal*\* of roughly one-half of the world population<sup>1</sup>. The variability of the  
19 monsoon makes the region one of the most susceptible areas around the world to the impacts  
20 of climate-related natural disasters such as droughts and floods. Through observations and  
21 modeling, the scientific community has advanced its understanding of the past and future  
22 changes in the monsoons. With regard to the northern summer monsoon, some studies suggest  
23 that the monsoon circulation and rainfall have intensified in a changing climate<sup>2</sup>. Wang et al.<sup>2</sup>  
24 show that the rainfall over the northern hemisphere summer monsoon domain, as well as the

---

\* *bread and butter* in south Asian nomenclature

25 Hadley and Walker circulations have all undergone substantial intensification during the last  
26 three decades (1979-2011), with a striking increase of monsoon rainfall by 9.5% per degree of  
27 global warming. A few recent studies have also pointed out the increasing frequency of extreme  
28 rainfall events over the south Asian monsoon domain, due to a warming climate<sup>3</sup>. However,  
29 quite a few other studies indicate that the monsoon rainfall<sup>4, 5, 6</sup> and the Hadley and Walker  
30 circulations<sup>7, 8, 9</sup> are weakening over the South Asian domain during the past half century (since  
31 1950s). Some of these studies suggest that, though the extreme rainfall events have increased  
32 over some regions<sup>3</sup>, the frequency of moderate-to-heavy rainfall events has decreased over the  
33 subcontinent<sup>8</sup>. In addition, several recent studies based on the Coupled Model Inter-comparison  
34 Project phase 5 (CMIP5) indicate a significant weakening of the large-scale Asian summer  
35 monsoon circulation, especially in the middle to upper atmosphere<sup>10, 11, 12</sup>. Monsoonal changes  
36 are found to be different when the region and time-periods considered are different—and so are  
37 the implications<sup>13</sup>. In fact, a recent study<sup>14</sup> using proxy records for the Asian monsoon over the  
38 past two millennia suggests that the monsoon is currently in the decreasing phase of a multi-  
39 decadal oscillation. This study also cautions that monsoonal changes due to anthropogenic  
40 forcing will be difficult to detect against a backdrop of large natural variability. The  
41 Intergovernmental Panel on Climate Change (IPCC) Fifth Assessment Report (AR5) is also  
42 suggestive of the uncertainties looming over the status and fate of the monsoons<sup>15, 16</sup>.  
43 Considering these uncertainties, there is an urgent need to provide robust information through  
44 the use of long-term observed datasets and climate models that could resolve the spatial and  
45 temporal lacuna associated with the monsoon variability in a changing climate.

46 Climate models consistently show that, under increasing greenhouse gas forcing, the  
47 land-sea temperature contrast increases, possibly due to enhanced evaporation (latent heat loss)  
48 over the ocean<sup>17</sup>. An increase in the land-sea thermal contrast should ideally increase the  
49 strength of the monsoon<sup>18, 19, 20</sup>. Also, the sea surface temperatures (SST) over the Indian Ocean

50 have shown a warming trend over the last several decades<sup>21, 22, 23</sup>, potentially enhancing the  
51 supply of moisture to the monsoon regions and increasing the rainfall amounts<sup>24</sup>. In a recent  
52 review of climate change and the South Asian monsoon, Turner and Annamalai<sup>13</sup> point out that  
53 though the land-sea contrast and the increased moisture availability should favor an increased  
54 monsoon rainfall, the observations do not provide any convincing evidence for such a positive  
55 trend. On the contrary, the South Asian monsoon rainfall has decreased significantly during the  
56 past several decades and this time evolution is not well reproduced by the CMIP5 models used  
57 in the IPCC AR5<sup>4, 11, 12, 25, 26, 27</sup>. This is indeed intriguing, and escalates the uncertainty  
58 associated with the monsoon in a changing climate.

59 In this study, we find that the summer monsoon rainfall during 1901-2012 shows a  
60 significant weakening trend over central South Asia, especially over the central-eastern parts  
61 of India where agriculture is still mostly rain-fed. Using observations and climate model  
62 experiments, we demonstrate that this reduction in rainfall is linked to the rapid warming of  
63 the Indian Ocean, especially its western part, during the past century. The Indian Ocean  
64 warming, along with a relatively subdued warming of the Indian subcontinent, has played a  
65 key role in weakening the land-sea thermal contrast, which is a major driver of the South Asian  
66 monsoon. The weakened land-sea thermal contrast dampens the summer monsoon Hadley  
67 circulation, reducing the rainfall over South Asia.

68

## 69 **Results**

### 70 **Rainfall trends during the past century**

71 Figures 1a and 1b show the trend in precipitation over the South Asian region, during summer  
72 monsoon for the years 1901-2012, using gridded datasets from the India Meteorological  
73 Department (IMD) and the Climate Research Unit (CRU). The trends are similar in both the  
74 datasets, with a weakening trend over the central and northeast Indian subcontinent, and south

75 of the Western Ghats region. The CRU dataset shows that the negative trend over the central  
76 subcontinent extends west to east, from south of Pakistan to central-eastern India, with a  
77 prominent horseshoe pattern with one of its arms placed on the foothills of the Himalayas, and  
78 the other obliquely along the central east coast of India. The reduction in summer rainfall over  
79 central-eastern India during the past century is about 10-20%. This could have socio-economic  
80 consequences since the agriculture in this region is still largely rain-fed. Significant positive  
81 trends in precipitation are consistently observed to the north of the Western Ghats region, in  
82 both the datasets. However, the positive rainfall trends are confined to a small domain along  
83 the west coast.

84

#### 85 **Indian Ocean warming and its association with the rainfall trends**

86 Previous studies using atmospheric general circulation models (AGCM) forced by observed  
87 SST variations suggest that a decrease in global land monsoon rainfall can be caused by  
88 warming trends over the tropical oceans, especially the central-eastern Pacific and the western  
89 Indian Ocean<sup>28</sup>. Among these tropical oceans, the Pacific SST anomalies do not show any long-  
90 term significant trend, while the western Indian Ocean exhibited rapid warming throughout the  
91 past century<sup>29</sup> (Fig. 2). In comparison with the rest of the Indian Ocean, the WIO generally has  
92 cooler mean SSTs in summer, owing to the strong monsoon winds and the resultant upwelling  
93 over the west. Meanwhile, the central-east Indian Ocean is characterized by a warm pool with  
94 SSTs greater than 28.0°C. Though previous studies demonstrate a basin-wide warming in the  
95 Indian Ocean during the past half century<sup>22, 23</sup>, our analysis using long-term SST datasets show  
96 that over an extended period of 112 years, the western Indian Ocean (WIO, 50-65°E, 5°S-10°N)  
97 has experienced anomalous warming of 1.2°C (Fig. 2), which is 0.5°C greater than the warming  
98 over the warm pool region<sup>29</sup>. Even though a warming trend over the Indian Ocean may result  
99 in an increased supply of moisture locally over the ocean, it is not necessary that the surplus

100 moisture is carried on to the peninsula<sup>30</sup>. The sustained Indian Ocean warming in the west has  
101 led to a greater westward extension of this Indian Ocean warm pool in recent decades<sup>22, 29</sup>. The  
102 expansion of the Indian Ocean warm pool can in turn modify the land-sea thermal gradient,  
103 which could modulate the strength and flow of the monsoon circulation and the moisture laden  
104 winds towards the South Asian subcontinent. Hence, deciphering the relationship between the  
105 Indian Ocean warming and the monsoon precipitation might give a clue to the cause and effect  
106 of the observed rainfall trends over the region.

107         In order to evaluate the association of the nearly monotonic warming over the WIO on  
108 the observed monsoon trends over the subcontinent, we carried out a correlation analysis  
109 between SST and precipitation anomalies over the two regions. Figures 3a and 3b show the  
110 correlation between the WIO SST anomalies and precipitation anomalies over the South Asian  
111 region, using multiple datasets. The spatial distribution of the correlation coefficients is  
112 consistent among the different datasets. The fascinating aspect is that, the correlation of  
113 precipitation anomalies with the Indian Ocean SST anomalies, and the trend in these  
114 precipitation anomalies are strikingly similar. A pattern correlation computed between the  
115 trends in Fig.1b and the correlation coefficients in Fig.3b (for the region 60°-100°E, 5°-35°N  
116 at 0.5° horizontal grid resolution) indicates that they are strongly correlated ( $r=0.73$ , significant  
117 at 95% confidence level). The most robust common feature among the trend and correlation  
118 patterns is a horseshoe pattern over central India. Apart from this, a dipolar pattern in the  
119 rainfall trends, with positive in the north and negative in the south is also visible along the  
120 Western Ghats of the Indian subcontinent (Fig. 1). A recent study<sup>31</sup> suggests that enhanced  
121 warming to the north of WIO in recent decades has also led to a poleward shift of the core of  
122 low level monsoon winds, increasing the precipitation to the north of the Western Ghats, while  
123 weakening it over the south. This dipolar rainfall pattern between the south and north of the

124 Western Ghats is also found to be associated with the WIO warming at both the interannual  
125 and low frequency (e.g. trend) time scales (Fig.3).

126 For a time series analysis, we consider the precipitation over the central South Asia  
127 (60°-100°E, 20°-30°N, inset box in Fig. 3b) along with the WIO SST anomalies (Fig. 3c). The  
128 rainfall region under consideration is chosen because the decreasing trend in rainfall is  
129 significant over this region, and also consistent with other studies utilizing station-wise analysis  
130 of rain gauge data<sup>6</sup>. Indeed, the SST and precipitation time series in Figure 3c are negatively  
131 correlated ( $r = -0.35$  at  $df = 110$ , for the unsmoothed time series), significant at 95% confidence  
132 level. Analysis using detrended time series data also shows significant negative correlation  
133 over the same region (Supplementary Fig. 1). This means that similar processes may operate  
134 at interannual and lower frequency timescales to generate a secular downward monsoon rainfall  
135 trend in response to the warming of the WIO, or alternatively, that a part of the Indian Ocean  
136 warming is due to the weakening of the monsoon<sup>23</sup>. The negative correlation is not observed  
137 during the first half of the 20<sup>th</sup> century, but becomes statistically significant post-1950s,  
138 probably due to the accelerated warming observed during the latter half. This implies that the  
139 relationship between a warm ocean and a weak monsoon is not linear. Since the ocean warming  
140 is a slow process, the heat has to build up over several decades to make a substantial difference  
141 and reach the warm pool SST values (28.0°C)<sup>22, 29</sup>. The sustained WIO warming has led to a  
142 greater spatial extension of the Indian Ocean warm pool in the recent decades only<sup>22</sup>.

143

#### 144 **Changing land-sea thermal contrast**

145 As pointed out earlier, changes in the land-sea thermal contrast impact the strength of the  
146 monsoon<sup>17, 18, 19, 20</sup>. Studies show that over the northern hemisphere, the surface temperatures  
147 over land increase more rapidly than over sea, under increasing greenhouse gases<sup>20</sup>. Ideally,  
148 such a scenario should strengthen the South Asian monsoon<sup>17, 18, 19, 20</sup>. Figures 4a and 4b show

149 the climatological mean and trends of surface temperatures over the Indian Ocean and the south  
150 Asian monsoon domain, during summer. Obviously, the monotonic warming trend over the  
151 Indian Ocean is prominent, with the strongest warming over the west. At the same time the  
152 Indian subcontinent exhibits suppressed warming during the past century. In fact, the surface  
153 temperatures in the northern Indian peninsula show a cooling trend. This implies a weakening  
154 meridional thermal gradient over the South Asian domain, in contrast with earlier studies,  
155 which suggest a strengthening of the thermal gradient under a globally warming scenario. The  
156 weakening land-sea thermal gradient is reproduced in other datasets also, indicating the  
157 robustness of the trend (Supplementary Fig. 2).

158 The land-sea contrast in surface temperatures, though important, may not provide a  
159 comprehensive picture of the factors modulating the strength of the monsoon circulation. Once  
160 the monsoon rains set in on the South Asian subcontinent, the land surface cools down  
161 considerably. However, the troposphere above the land remains warm due to the latent heat  
162 release from the convective activity, keeping the thermal contrast functional<sup>32</sup>. This means that  
163 the tropospheric temperature is an ideal parameter for examining the thermodynamic forcing  
164 related to the monsoon<sup>32</sup>. However, unlike the surface temperature, robust records of air  
165 temperature at different levels of the troposphere are not available prior to the 1950s, which  
166 makes it difficult to examine the role of long-term tropospheric temperature trends on the  
167 monsoon. Nevertheless, to confirm that our results are valid for the recent decades also, we  
168 analyze the tropospheric temperatures over the monsoon domain. The trend in the thermal  
169 contrast is well reflected in the upper tropospheric (200 hPa) temperature anomalies also, as  
170 demonstrated in Figure 4c. Along with a cooling trend over the larger Indo-Tibetan landmass,  
171 increasing positive temperature anomalies are observed over the WIO, indicating its potential  
172 role in weakening the monsoon flow towards the Indian peninsula. The warming of the upper  
173 troposphere over the Indian Ocean (Fig. 4c) is co-located with that over the sea surface (Fig.

174 4b). This may be attributed to the fact that the tropical Indian Ocean is a highly convective  
175 region during the northern summer, as the mean SSTs are above the minimum convective  
176 threshold (26°C) and the southwesterly monsoon winds are conducive for enhanced moisture  
177 convergence<sup>33</sup>. Further warming of the Indian Ocean enhances the convection<sup>24</sup>, transferring  
178 surplus heat to the upper troposphere<sup>34, 35, 36</sup>.

179 To evaluate the trends observed in land-sea thermal contrast in Figs. 4b and 4c, we  
180 analyze the difference between the summer tropospheric temperatures over WIO and the Indian  
181 subcontinent, at 850 and 200 hPa levels (Fig. 4d). The surface temperature gradient ( $T_{surf}$ ) shows  
182 a sharp decline throughout the past century, largely contributed by an increased warming over  
183 the WIO. The tropospheric temperature gradient ( $T_{trop}$ ) post-1950s also indicates a trend similar  
184 to that in the surface temperatures. This is suggestive of a warm Indian Ocean having a strong  
185 hold on the whole troposphere, from the surface to the top, possibly due to enhanced convective  
186 mixing over the ocean<sup>34, 35, 36</sup>. Indeed, a separate analysis of the trend at each of these levels  
187 gave consistent results of a warming troposphere over the Indian Ocean, and a weakening  
188 thermal contrast between the subcontinent and the ocean. All these results indicate a reduced  
189 land-sea thermal contrast during the past century, partly due to increased warming over the  
190 Indian Ocean, especially over the western region, and partly due to a relatively weaker warming  
191 over the Indian peninsula.

192

### 193 **Weakening of the summer monsoon Hadley circulation**

194 Many studies have shown that warm SST anomalies are accompanied by large variations in the  
195 lower and upper troposphere due to enhanced latent heating aloft from convection over the  
196 ocean<sup>34, 35, 36</sup>. Furthermore, these changes are highly correlated with the strength of the  
197 monsoon circulation<sup>32, 35</sup>. A more recent study using a coupled model framework also points  
198 out that a spatial extension of Indian Ocean warm pool could enhance the convection over the

199 ocean while introducing a dry bias over land, by modulating the meridional Hadley  
200 circulation<sup>37</sup>. It is therefore likely that the impact of the Indian Ocean warming on the monsoon  
201 has become more prominent in recent decades. Consistently, an examination of the vertical  
202 wind velocity over the South Asian domain (50-100°E) during the years 1948-2012 indicates  
203 large scale upward motion over the equatorial ocean (10°S-10°N), extending up to the upper  
204 troposphere and favoring intense local convection (Fig. 5a). Furthermore, this enhanced  
205 upward motion over the ocean is compensated by subsidence of air over the subcontinent (10-  
206 20°N), inhibiting convection over the landmass and drying the region, through a modulation of  
207 the local Hadley cell (Fig. 5a). This suggests that though the warming ocean engenders  
208 enhanced local rainfall due to increased moisture availability, it weakens the monsoon Hadley  
209 circulation and reduces the rainfall over the land, ultimately building up a competition among  
210 the land and ocean rainfall in the south Asian monsoon domain.

211

### 212 **Climate model response to Indian Ocean warming**

213 Attribution of the decreasing monsoon rainfall to the Indian Ocean warming from observations  
214 alone is very difficult and probably impossible. Thus, to delineate and examine the role of the  
215 Indian Ocean warming on the monsoon circulation and rainfall, model sensitivity experiments  
216 using an ocean-atmosphere coupled climate model were carried out. In the sensitivity  
217 experiment, positive SST anomalies similar to those in the observations were added to the WIO  
218 region (Supplementary Fig. 3). Figure 5b shows the model response in the summer monsoon  
219 Hadley circulation, and features a weakening of the local Hadley cell, much similar to the  
220 observed trends. The fact that the warming has resulted in enhanced convection over the ocean  
221 and subsidence over the landmass serves as the evidence that the observed warming trend in  
222 the Indian Ocean has the potential to weaken the monsoon circulation.

223 Figure 6 compares the observed trends in precipitation and surface winds with the  
224 model-simulated anomalies in response to warming over the WIO. The interesting, robust  
225 element in the model simulations (Fig. 6b) is the horseshoe pattern in the negative precipitation  
226 anomalies, with one arm placed over the Himalayan foothills and the other over the central  
227 Indian landmass, similar to the one in the precipitation trends (Figs. 1, 6a) and the correlation  
228 pattern with WIO SST anomalies (Figs. 3a,3b). Consistent with the reduced precipitation, the  
229 surface winds also suggest weakened mean southwesterly winds. The trend analysis (Fig. 6a)  
230 shows a Rossby wave-like response<sup>38</sup> to the Indian Ocean warming, with northeasterly wind  
231 anomalies over most of the Arabian Sea, converging towards the central Indian Ocean. The  
232 model simulation (Fig. 6b) also shows similar results, with respect to the Rossby wave response  
233 near the subcontinent and over the Arabian Sea. However, the zonal direction of the wind  
234 anomalies near the equator appears different in the observations and the model. This is not an  
235 unexpected result as the periods under consideration and the model simulations are different,  
236 since the observed winds are available only for 1948-2012—a period during which the Indian  
237 Ocean experienced a basin-wide warming<sup>22, 23</sup>. In fact previous studies using AGCM  
238 experiments driven by recent basin-wide warming in the tropical Indian Ocean reproduced the  
239 observed zonal wind changes in the recent decades<sup>39</sup>. In the model sensitivity experiment,  
240 however, the positive SST offset is prescribed only over the WIO region, where the long-term  
241 warming trend is observed in the SST records during 1901-2012. This suggests that while WIO  
242 warming plays a significant role in the long-term change in winds, basin-wide warming over  
243 the tropical Indian Ocean has emerged as a co-conspirator in the change in land-sea thermal  
244 contrast and winds in recent decades.

245 A further comparison of the model response to the warming is provided in  
246 Supplementary Fig. 4. The thermal response (surface and upper troposphere) is similar to those  
247 in the observed trends over the ocean, but exhibit some discrepancies over land. More

248 specifically, for the model response in temperature at 200 hPa, the results reproduce the upper  
249 tropospheric warming over the ocean, but do not replicate the cooling over the Indo-Tibetan  
250 landmass. This is possibly because of the role of other processes on the upper tropospheric  
251 temperatures, such as aerosol forcing<sup>40, 41</sup> and stratosphere-troposphere interactions<sup>42</sup>, which  
252 are not included in the model experiment. It may also be noted that previous studies<sup>42</sup> have  
253 suggested that the NCEP upper tropospheric data overestimates the cooling trend over the Indo-  
254 Tibetan landmass. Nevertheless, the similarity between the rainfall and local Hadley responses  
255 between the model experiments and observed fields clearly demonstrates the importance of  
256 WIO warming in determining the observed monsoon response.

257

## 258 **Discussion**

259 The above analysis using observations and climate model sensitivity experiments indicates the  
260 potential role of the Indian Ocean warming in weakening the monsoon circulation and rainfall  
261 over central South Asia. The Indian Ocean and monsoon exhibit a coupled relationship<sup>33, 43, 44</sup>,  
262 and it is also possible that a feedback from a weak monsoon accelerates the warming in the  
263 Indian Ocean<sup>23</sup>. Some studies have suggested a role for anthropogenic aerosols in weakening  
264 the monsoon circulation during the recent decades<sup>41, 45, 46, 47, 48, 49</sup>. It is possible that such a  
265 weakened circulation amplifies the SST warming due to a decreased evaporation. In contrast,  
266 recent studies show that the anthropogenic aerosols tend to have a cooling effect and compete  
267 with the greenhouse warming over the Indian Ocean<sup>50, 51</sup>. Despite the fact that aerosols are a  
268 competing factor, the observed SST trends do not show any change during the post-1980s  
269 where aerosols have increased (Fig.3c). In fact, a study<sup>51</sup> points out that a large share of the  
270 CMIP5 models with aerosol forcing are not able to reproduce the observed SST warming over  
271 the Indian Ocean. They attribute this issue to the large uncertainties in representing the direct  
272 and indirect effects of aerosols in these models. Meanwhile, a recent study demonstrates that

273 apart from the warming due to greenhouse gas forcing<sup>52</sup>, the long-term warming trend in the  
274 western Indian Ocean is significantly associated with the asymmetry in the ENSO forcing<sup>29</sup>.  
275 That is, while the quasi-periodic El Niño events induce a warming over the Indian Ocean via  
276 the Walker circulation, the La Niña events fail to do the inverse. Along with this, an increase  
277 in the magnitude and frequency of El Niños in the recent decades has accelerated this  
278 warming<sup>29</sup>. The heat pile-up due to this ENSO forcing<sup>53</sup> persists in the land-locked Indian  
279 Ocean which has a restricted thermohaline circulation, sustained by a feedback from the local  
280 air-sea interaction<sup>54, 55</sup> and ocean dynamics<sup>56, 57</sup>. Considering the strong seasonal variability in  
281 this region and short-term persistence in atmospheric changes, it is rather self-evident that, on  
282 longer time scales, the warmer ocean plays a larger role in weakening the monsoon than vice-  
283 versa.

284         Though the observed increase in anthropogenic aerosols may not have a direct effect in  
285 warming the Indian Ocean, they may have a potential role in suppressing (or even cooling) the  
286 surface warming over the South Asian domain in the recent decades<sup>40, 41</sup>. Aerosols could be  
287 considered as the perfect partner in reducing the land-sea thermal contrast, along with a  
288 warming Indian Ocean. An evaluation of the CMIP5 models<sup>41</sup> suggest that aerosols could  
289 potentially reduce the incoming solar radiation over the Indo-Gangetic plain, thereby cooling  
290 the surface. However, it may be pointed out that majority of these models could not simulate  
291 the spatial distribution of aerosols or related processes over the South Asian subcontinent<sup>41</sup>.  
292 Also, ambiguity still remains on the extent of cooling (or warming) induced by different types  
293 of aerosols<sup>46, 47, 48</sup>. Meanwhile, other studies link the upper tropospheric cooling over the larger  
294 Indo-Tibetan landmass during summer to stratospheric temperature changes which are  
295 channeled through stratosphere-troposphere interactions<sup>42</sup>. Further research is required to  
296 differentiate the individual contributions of these components on the observed trends in surface  
297 and upper tropospheric temperatures over land.

298           The current study discusses the role of Indian Ocean warming in weakening of the  
299 monsoon, but does not rule out the possibility that similar impacts may also arise due to SST  
300 changes in the tropical Pacific and the Atlantic<sup>9, 55, 58, 59, 60, 61, 62</sup>. Recent studies show that the  
301 Pacific warming trend during the past century is not prominent as compared to the Indian Ocean  
302 warming while the tropical and extratropical Atlantic show strong warming trends<sup>29</sup>. However,  
303 Prodhomme et al.<sup>62</sup> pointed out that a warming in the tropical Atlantic does not change the  
304 tropospheric temperature gradient between the Indian Ocean and the subcontinent as the  
305 temperature changes due to the teleconnection are uniform over this domain. Meanwhile,  
306 Goswami et al.<sup>60</sup> suggest that a North Atlantic warming should ideally strengthen the Asian  
307 monsoon by increasing the tropospheric temperatures over northern India and southern Eurasia.  
308 However, temperature and rainfall trends during recent decades suggest that any such role of  
309 the Atlantic is offset by the tropospheric temperature trends induced by a warming Indian  
310 Ocean. Deconvolving the non-stationary relations between the monsoon and the warming over  
311 these remote regions is beyond the scope of this study. But considering the robustness of the  
312 Indian Ocean warming and its direct role in monsoon dynamics in comparison with the other  
313 tropical oceans, it is justified to say that the Indian Ocean SST trends play a relatively larger  
314 role on the observed rainfall trends in the South Asian domain.

315           A recent study comparing present observations and future regional climate model  
316 projections of the summer monsoon suggests a further weakening of the monsoon circulation  
317 and peninsular rainfall in the future<sup>63</sup>. Meanwhile, the CMIP5 future projections suggest an  
318 increase in rainfall over the central Indian subcontinent<sup>28</sup>, though one must note the ongoing  
319 debate about whether CMIP5 models are in fact capturing the historic monsoon trends  
320 accurately<sup>4, 11, 12, 25</sup> and whether the monsoon projections are reliable in these models<sup>26</sup>. The  
321 overall weakening trend of monsoon rainfall over South Asia is a matter of grave concern since  
322 the socio-economic livelihood in this region, including agriculture, water resources and power

323 generation are irrevocably dependent on it. The role of Indian Ocean warming might not be  
324 solely limited to weakening the Asian monsoon. Research based on climate proxy records and  
325 models point out that the climate signals forced by warm SST anomalies over the tropical  
326 Indian and western Pacific oceans synergistically contribute to widespread drying over south  
327 Asia and mid-latitudes, an ideal scenario for a large-scale drought<sup>64, 65</sup>. In addition, a recent  
328 study points out that the Indian Ocean warming may also be playing an important role in  
329 modulating the Pacific climate change in recent decades—by favoring stronger trade winds in  
330 the Pacific, thereby contributing to the recent global warming hiatus<sup>66</sup>. Hence, the critical role  
331 of the warm Indian Ocean deserves special attention not only for its decisive effect on the food  
332 security of a large fraction of the world's population, but also due to its disproportionately large  
333 role as a partner to the warm pool Pacific in inducing a drought, as well as its potential role in  
334 the global warming pause.

335

## 336 **Methods**

### 337 **Rainfall and SST historical data analysis**

338 Two long-term precipitation datasets for the northern summer during the period 1901-2012,  
339 obtained from the India Meteorological Department (IMD) and Climate Research Unit (CRU)  
340 are used in this study. The recent IMD precipitation data uses daily rainfall data from all the  
341 rain gauge stations over India, gridded at 0.25° horizontal resolution<sup>67</sup>. This includes rainfall  
342 records from 6,955 stations, which is the highest number of stations used by any gridded  
343 datasets currently available. The data density varies from year to year from about 1,500 in the  
344 first couple of years to about 4,000 by the end of the century. The trend in precipitation over  
345 the central Indian region under consideration, is compared with studies utilizing station wise  
346 analysis of rain gauge data<sup>6</sup>. SST data for the same period is obtained from the HadISST1  
347 dataset provided by the Met Office Hadley Centre, and also the Extended Reconstructed Sea

348 Surface Temperature (ERSST) provided by NOAA. In order to examine the role of land-sea  
349 temperature gradient, surface temperatures from HadCRUT (1901-2012) and tropospheric  
350 temperature values from NCEP reanalysis (1948-2012) are used. Observed trends in the  
351 monsoon southwesterlies are inferred from the wind data based on NCEP reanalysis during  
352 1948-2012. The significance of the trends and correlations is examined using standard two-  
353 tailed student's t-tests. The correlation analysis in Fig.3c has also been tested using Kendall's  
354 rank correlation which is non-parametric and therefore makes no assumptions about the  
355 distribution and at the same time determine the direction and significance of the relation  
356 between the two variables<sup>68</sup>. The correlated variables are said to be concordant if their ranks  
357 vary together (-1) and discordant if they vary differently (+1). The significance of the rainfall  
358 and SST linear trend values are further assessed with Mann-Kendall tests<sup>69</sup> with block bootstrap  
359 to improve the significance test when a time series shows auto-correlation.

360

### 361 **Model and experimental design**

362 For the numerical model experiments, we use the standard configuration of a global coupled  
363 ocean-atmosphere model, the Climate Forecast System version 2 (CFSv2). The CFSv2 model  
364 simulates the mean monsoon and its interannual variability reasonably well, and hence used in  
365 this study<sup>43</sup>. The oceanic component has a 0.25-0.5° horizontal resolution, 40 vertical levels  
366 and includes an ice model. The atmospheric component is at T126 (~0.9°) horizontal resolution,  
367 and 64 sigma-pressure hybrid levels. The atmosphere and ocean exchange quantities such as  
368 heat and momentum fluxes every half an hour, with no flux adjustment or correction. See Saha  
369 et al.<sup>70</sup> for further details on the model and its components. The coupled configuration of CFSv2  
370 is time integrated over a period of 100 years, and utilized as the reference run (CFSv2<sub>CTL</sub>). In  
371 order to examine the role of a warming Indian Ocean, a sensitivity experiment (CFSv2<sub>WIO</sub>)  
372 using ensembles (20 members) of short integrations for the summer monsoon season during

373 June-September were performed by adding temperature anomalies to the SSTs passed over the  
374 WIO, to the atmosphere. Positive anomalies of the order of 1.5°C were added over the region,  
375 in such a way that it tapers out by the limits of the domain (50-65°E, 5°S-10°N). The positive  
376 offset added to the simulated SSTs over WIO in the sensitivity experiments is similar to the  
377 amplitude of the observed trends (Fig. 2, Supplementary Fig. 3). The difference between  
378 CFSv2<sub>WIO</sub> and CFSv2<sub>CTL</sub> is taken as the model response to the summer warming over the WIO.

379 **Acknowledgments**

380 The authors gratefully acknowledge the financial support given by the Earth System Science  
381 Organization, Ministry of Earth Sciences, Government of India, to conduct this research under  
382 the National Monsoon Mission (Grant #MM/SERP/CNRS/2013/INT-10/002, Contribution  
383 #MM/PASCAL/RP/03).

384

385 **Author Contributions**

386 M.K.R. designed the study and the model experiment. M.K.R. and K.R. performed the model  
387 sensitivity experiments and the analysis. All authors contributed ideas in developing the  
388 research, discussed the results and wrote the paper.

389

390 **Competing financial interests**

391 The authors declare no competing financial interests.

392 **References**

- 393 1. Gadgil S, Rupa Kumar K. The Asian monsoon — agriculture and economy. *The Asian*  
394 *Monsoon*. Springer Berlin Heidelberg, 2006, pp 651-683.
- 395 2. Wang B, Liu J, Kim H-J, Webster PJ, Yim S-Y, Xiang B. Northern Hemisphere summer  
396 monsoon intensified by mega-El Niño/southern oscillation and Atlantic multidecadal  
397 oscillation. *Proceedings of the National Academy of Sciences* 2013, **110**(14): 5347-  
398 5352.
- 399 3. Goswami BN, Venugopal V, Sengupta D, Madhusoodanan M, Xavier PK. Increasing  
400 trend of extreme rain events over India in a warming environment. *Science* 2006,  
401 **314**(5804): 1442-1445.
- 402 4. Kitoh A, Endo H, Krishna Kumar K, Cavalcanti IF, Goswami P, Zhou T. Monsoons in  
403 a changing world: a regional perspective in a global context. *Journal of Geophysical*  
404 *Research: Atmospheres* 2013, **118**(8): 3053-3065.
- 405 5. Zhou T, Zhang L, Li H. Changes in global land monsoon area and total rainfall  
406 accumulation over the last half century. *Geophysical Research Letters* 2008, **35**(16):  
407 L16707.
- 408 6. Guhathakurta P, Rajeevan M. Trends in the rainfall pattern over India. *International*  
409 *Journal of Climatology* 2008, **28**(11): 1453-1469.
- 410 7. Tokinaga H, Xie S-P, Deser C, Kosaka Y, Okumura YM. Slowdown of the Walker  
411 circulation driven by tropical Indo-Pacific warming. *Nature* 2012, **491**(7424): 439-443.
- 412 8. Krishnan R, Sabin T, Ayantika D, Kitoh A, Sugi M, Murakami H, *et al.* Will the South  
413 Asian monsoon overturning circulation stabilize any further? *Climate Dynamics* 2013,  
414 **40**(1-2): 187-211.
- 415 9. Vecchi GA, Soden BJ, Wittenberg AT, Held IM, Leetmaa A, Harrison MJ. Weakening  
416 of tropical Pacific atmospheric circulation due to anthropogenic forcing. *Nature* 2006,  
417 **441**(7089): 73-76.

- 418 10. Ma J, Yu JY. Paradox in South Asian summer monsoon circulation change: Lower  
419 tropospheric strengthening and upper tropospheric weakening. *Geophysical Research*  
420 *Letters* 2014, **41**(8): 2934-2940.
- 421 11. Ogata T, Ueda H, Inoue T, Hayasaki M, Yoshida A, Watanabe S, *et al.* Projected Future  
422 Changes of the Asian Monsoon: A Comparison of CMIP3 and CMIP5 model results.  
423 *Journal of Meteorological Society of Japan* 2014.
- 424 12. Sooraj KP, Terray P, Mujumdar M. Global warming and the weakening of the Asian  
425 summer Monsoon circulation : assessments from the CMIP5 models. *Climate*  
426 *Dynamics* 2014.
- 427 13. Turner AG, Annamalai H. Climate change and the South Asian summer monsoon.  
428 *Nature Climate Change* 2012, **2**: 587–595.
- 429 14. Sinha A, Kathayat G, Cheng H, Breitenbach SF, Berkelhammer M, Mudelsee M, *et al.*  
430 Trends and oscillations in the Indian summer monsoon rainfall over the last two  
431 millennia. *Nature communications* 2015, **6**.
- 432 15. IPCC. IPCC WGI Fifth Assessment Report, Chapter 2 - Observations: Atmosphere and  
433 Surface; 2013.
- 434 16. Christensen JH, Kanikicharla KK, Marshall G, Turner J. Climate phenomena and their  
435 relevance for future regional climate change. 2013.
- 436 17. Sutton RT, Dong B, Gregory JM. Land/sea warming ratio in response to climate  
437 change: IPCC AR4 model results and comparison with observations. *Geophysical*  
438 *Research Letters* 2007, **34**(2).
- 439 18. Chou C. Land–sea heating contrast in an idealized Asian summer monsoon. *Climate*  
440 *Dynamics* 2003, **21**(1): 11-25.
- 441 19. Wu G, Liu Y, He B, Bao Q, Duan A, Jin F-F. Thermal controls on the Asian summer  
442 monsoon. *Scientific reports* 2012, **2**.
- 443 20. Kamae Y, Watanabe M, Kimoto M, Shiogama H. Summertime land–sea thermal  
444 contrast and atmospheric circulation over East Asia in a warming climate—Part I: Past  
445 changes and future projections. *Climate Dynamics* 2014: 1-16.

- 446 21. Alory G, Wijffels S, Meyers G. Observed temperature trends in the Indian Ocean over  
447 1960–1999 and associated mechanisms. *Geophysical Research Letters* 2007, **34**(2).
- 448 22. Rao SA, Dhakate AR, Saha SK, Mahapatra S, Chaudhari HS, Pokhrel S, *et al.* Why is  
449 Indian Ocean warming consistently? *Climatic change* 2012, **110**(3-4): 709-719.
- 450 23. Swapna P, Krishnan R, Wallace J. Indian Ocean and monsoon coupled interactions in  
451 a warming environment. *Climate Dynamics* 2014, **42**(9-10): 2439-2454.
- 452 24. Roxy M. Sensitivity of precipitation to sea surface temperature over the tropical  
453 summer monsoon region—and its quantification. *Climate Dynamics* 2013, **43**(5-6):  
454 1159-1169.
- 455 25. Saha A, Ghosh S, Sahana A, Rao E. Failure of CMIP5 climate models in simulating  
456 post-1950 decreasing trend of Indian monsoon. *Geophysical Research Letters* 2014.
- 457 26. Sabeerali C, Rao SA, Dhakate A, Salunke K, Goswami B. Why ensemble mean  
458 projection of south Asian monsoon rainfall by CMIP5 models is not reliable? *Climate*  
459 *Dynamics* 2014: 1-14.
- 460 27. Sharmila S, Joseph S, Sahai A, Abhilash S, Chattopadhyay R. Future projection of  
461 Indian summer monsoon variability under climate change scenario: An assessment  
462 from CMIP5 climate models. *Global and Planetary Change* 2015, **124**: 62-78.
- 463 28. Zhou T, Yu R, Li H, Wang B. Ocean forcing to changes in global monsoon precipitation  
464 over the recent half-century. *Journal of Climate* 2008, **21**(15): 3833-3852.
- 465 29. Roxy MK, Ritika K, Terray P, Masson S. The curious case of Indian Ocean warming.  
466 *Journal of Climate* 2014, **27**(22): 8501-8509.
- 467 30. Prodhomme C, Terray P, Masson S, Izumo T, Tozuka T, Yamagata T. Impacts of Indian  
468 Ocean SST biases on the Indian Monsoon: as simulated in a global coupled model.  
469 *Climate Dynamics* 2014, **42**(1-2): 271-290.
- 470 31. Sandeep S, Ajayamohan R. Poleward shift in Indian summer monsoon low level  
471 jetstream under global warming. *Climate Dynamics* 2014: 1-15.

- 472 32. Xavier PK, Marzin C, Goswami BN. An objective definition of the Indian summer  
473 monsoon season and a new perspective on the ENSO–monsoon relationship. *Quarterly*  
474 *Journal of the Royal Meteorological Society* 2007, **133**(624): 749-764.
- 475 33. Gadgil S, Joshi NV, Joseph PV. Ocean-atmosphere coupling over monsoon regions.  
476 *Nature* 1984, **312**: 141-143.
- 477 34. Danielsen EF. In situ evidence of rapid, vertical, irreversible transport of lower  
478 tropospheric air into the lower tropical stratosphere by convective cloud turrets and by  
479 larger-scale upwelling in tropical cyclones. *Journal of Geophysical Research:*  
480 *Atmospheres (1984–2012)* 1993, **98**(D5): 8665-8681.
- 481 35. Dai A, Li H, Sun Y, Hong LC, Chou C, Zhou T. The relative roles of upper and lower  
482 tropospheric thermal contrasts and tropical influences in driving Asian summer  
483 monsoons. *Journal of Geophysical Research: Atmospheres* 2013, **118**(13): 7024-7045.
- 484 36. Su H, Neelin JD, Meyerson JE. Sensitivity of tropical tropospheric temperature to sea  
485 surface temperature forcing. *Journal of Climate* 2003, **16**(9): 1283-1301.
- 486 37. Bollasina MA, Ming Y. The general circulation model precipitation bias over the  
487 southwestern equatorial Indian Ocean and its implications for simulating the South  
488 Asian monsoon. *Climate dynamics* 2013, **40**(3-4): 823-838.
- 489 38. Gill AE. Some simple solutions for heat-induced tropical circulation. John Wiley &  
490 Sons, Ltd; 1980. pp. 447-462.
- 491 39. Zhou T, Yu R, Zhang J, Drange H, Cassou C, Deser C, *et al.* Why the western Pacific  
492 subtropical high has extended westward since the late 1970s. *Journal of Climate* 2009,  
493 **22**(8): 2199-2215.
- 494 40. Krishnan R, Ramanathan V. Evidence of surface cooling from absorbing aerosols.  
495 *Geophysical research letters* 2002, **29**(9): 54-51-54-54.
- 496 41. Sanap S, Pandithurai G, Manoj M. On the response of Indian summer monsoon to  
497 aerosol forcing in CMIP5 model simulations. *Climate Dynamics* 2015: 1-13.
- 498 42. Yu R, Wang B, Zhou T. Tropospheric cooling and summer monsoon weakening trend  
499 over East Asia. *Geophysical Research Letters* 2004, **31**(22).

- 500 43. Roxy M, Tanimoto Y, Preethi B, Pascal T, Krishnan R. Intraseasonal SST-precipitation  
501 relationship and its spatial variability over the tropical summer monsoon region.  
502 *Climate Dynamics* 2012, **41**(1): 45-61.
- 503 44. Webster PJ. The coupled monsoon system. *The Asian Monsoon*. Springer, 2006, pp 3-  
504 66.
- 505 45. Bollasina MA, Ming Y, Ramaswamy V. Anthropogenic aerosols and the weakening of  
506 the South Asian summer monsoon. *science* 2011, **334**(6055): 502-505.
- 507 46. Ramanathan V, Chung C, Kim D, Bettge T, Buja L, Kiehl J, *et al.* Atmospheric brown  
508 clouds: Impacts on South Asian climate and hydrological cycle. *Proceedings of the*  
509 *National Academy of Sciences of the United States of America* 2005, **102**(15): 5326-  
510 5333.
- 511 47. Meehl GA, Arblaster JM, Collins WD. Effects of black carbon aerosols on the Indian  
512 monsoon. *Journal of Climate* 2008, **21**(12): 2869-2882.
- 513 48. Menon S, Hansen J, Nazarenko L, Luo Y. Climate effects of black carbon aerosols in  
514 China and India. *Science* 2002, **297**(5590): 2250-2253.
- 515 49. Lau KM, Kim KM. Observational relationships between aerosol and Asian monsoon  
516 rainfall, and circulation. *Geophysical research letters* 2006, **33**(21).
- 517 50. Dong L, Zhou T, Wu B. Indian Ocean warming during 1958–2004 simulated by a  
518 climate system model and its mechanism. *Climate Dynamics* 2014, **42**(1-2): 203-217.
- 519 51. Ning H, Li-Juan L, Bin W. The role of the aerosol indirect effect in the northern Indian  
520 Ocean warming simulated by CMIP5 models. *Atmospheric and Oceanic Science*  
521 *Letters* 2014, **7**(5): 411-416.
- 522 52. Du Y, Xie SP. Role of atmospheric adjustments in the tropical Indian Ocean warming  
523 during the 20th century in climate models. *Geophysical Research Letters* 2008, **35**(8).
- 524 53. Compo GP, Sardeshmukh PD. Removing ENSO-related variations from the climate  
525 record. *Journal of Climate* 2010, **23**(8): 1957-1978.

- 526 54. Du Y, Xie S-P, Huang G, Hu K. Role of Air-Sea Interaction in the Long Persistence of  
527 El Niño-Induced North Indian Ocean Warming. *Journal of Climate* 2009, **22**(8): 2023-  
528 2038.
- 529 55. Lau NC, Nath MJ. Impact of ENSO on the variability of the Asian-Australian monsoons  
530 as simulated in GCM experiments. *Journal of Climate* 2000, **13**(24): 4287-4309.
- 531 56. Chowdary J, Gnanaseelan C. Basin-wide warming of the Indian Ocean during El Niño  
532 and Indian Ocean dipole years. *International Journal of Climatology* 2007, **27**(11):  
533 1421-1438.
- 534 57. Rahul S, Gnanaseelan C. Net Heat Flux Over the Indian Ocean: Trends, Driving  
535 Mechanisms, and Uncertainties. *Geoscience and Remote Sensing Letters, IEEE* 2013,  
536 **10**(4): 776-780.
- 537 58. Ashok K, Sabin T, Swapna P, Murtugudde R. Is a global warming signature emerging  
538 in the tropical Pacific? *Geophysical Research Letters* 2012, **39**(2).
- 539 59. Kumar A, Jha B, L'Heureux M. Are tropical SST trends changing the global  
540 teleconnection during La Niña? *Geophysical Research Letters* 2010, **37**(12).
- 541 60. Goswami BN, Madhusoodanan M, Neema C, Sengupta D. A physical mechanism for  
542 North Atlantic SST influence on the Indian summer monsoon. *Geophysical Research*  
543 *Letters* 2006, **33**(2).
- 544 61. Kucharski F, Bracco A, Yoo J, Molteni F. Atlantic forced component of the Indian  
545 monsoon interannual variability. *Geophysical Research Letters* 2008, **35**(4).
- 546 62. Prodhomme C, Terray P, Masson S, Boschhat G, Izumo T. Oceanic factors controlling  
547 the Indian Summer Monsoon onset in a coupled model. *Climate Dynamics* 2014: 1-26.
- 548 63. Dash SK, Mishra SK, Pattnayak KC, Mangain A, Mariotti L, Coppola E, *et al.*  
549 Projected seasonal mean summer monsoon over India and adjoining regions for the  
550 twenty-first century. *Theoretical and Applied Climatology* 2014: 1-13.
- 551 64. Hoerling M, Kumar A. The perfect ocean for drought. *Science* 2003, **299**(5607): 691-  
552 694.

- 553 65. Ummenhofer CC, D'Arrigo RD, Anchukaitis KJ, Buckley BM, Cook ER. Links  
554 between Indo-Pacific climate variability and drought in the Monsoon Asia Drought  
555 Atlas. *Climate dynamics* 2013, **40**(5-6): 1319-1334.
- 556 66. Luo J-J, Sasaki W, Masumoto Y. Indian Ocean warming modulates Pacific climate  
557 change. *Proceedings of the National Academy of Sciences* 2012, **109**(46): 18701-  
558 18706.
- 559 67. Pai D, Sridhar L, Badwaik M, Rajeevan M. Analysis of the daily rainfall events over  
560 India using a new long period (1901–2010) high resolution (0.25×0.25) gridded rainfall  
561 data set. *Climate Dynamics* 2014: 1-22.
- 562 68. Kendall MG. Rank correlation methods. 1948.
- 563 69. Wang B, Ding Q. Changes in global monsoon precipitation over the past 56 years.  
564 *Geophysical Research Letters* 2006, **33**(6).
- 565 70. Saha S, Moorthi S, Wu X, Wang J, Nadiga S, Tripp P, *et al.* The NCEP climate forecast  
566 system version 2. *Journal of Climate* 2013(2013).

567 **Figure Legends**

568 Figure 1. Observed trend in precipitation ( $\text{mm day}^{-1} 112 \text{ year}^{-1}$ ) in (a) IMD and (b) CRU  
569 datasets, during June-September, for the years 1901-2012. Contours denote regions  
570 significant at the 95% confidence level.

571 Figure 2. Observed trend in mean summer (June–September) SST ( $^{\circ}\text{C} 112 \text{ yr}^{-1}$ ) over the global  
572 tropics during 1901–2012.


573 Figure 3. Correlation between SST over the western Indian Ocean (WIO,  $50\text{-}65^{\circ}\text{E}$ ,  $5^{\circ}\text{S}\text{-}10^{\circ}\text{N}$ )  
574 and precipitation over the South Asian subcontinent, for (a) HadISST and IMD  
575 precipitation, and (b) ERSST and CRU precipitation, for June-September 1901-2012.  
576 Contours denote regions significant at the 95% confidence level. (c) Time series of SST  
577 anomalies ( $^{\circ}\text{C}$ , *red*) over WIO along with CRU (*blue*) precipitation ( $\text{mm day}^{-1}$ ) over  
578 central South Asia ( $60\text{-}100^{\circ}\text{E}$ ,  $20\text{-}30^{\circ}\text{N}$ , inset box in **b**), smoothed with a 10 year moving  
579 average. Note that the correlation coefficient ( $r = -0.34$ ) between HadISST and CRU  
580 precipitation is estimated using non-smoothed time series. Kendall's rank correlation  
581 test for the two variables provided a tau coefficient of -0.3 ( $P < 0.01$ , two tailed). Mann-  
582 Kendall test for the trend in the time series provided a tau coefficient of 0.6 for SST  
583 and -0.2 for precipitation, both significant at 95% confidence level.

584 Figure 4. Observed (a) mean ( $^{\circ}\text{C}$ ) and (b) trend ( $^{\circ}\text{C} 112 \text{ year}^{-1}$ ) of surface temperatures during  
585 June-September, 1901-2012. For the trend, the color shades represent regions  
586 significant at the 95% confidence level. (c) Observed trend ( $^{\circ}\text{C} 65 \text{ year}^{-1}$ ) of upper  
587 tropospheric (200 hPa) temperatures during June-September, 1948-2012. (d) Time  
588 series of the trend in land-sea contrast in the surface ( $T_{surf}$ ) and tropospheric ( $T_{surf}$ , 850-  
589 200hPa average) temperatures ( $^{\circ}\text{C}$ ). The land-sea contrast is estimated as the difference  
590 in the values between the boxes over the South Asian subcontinent ( $70\text{-}85^{\circ}\text{E}$ ,  $10\text{-}30^{\circ}\text{N}$ )  
591 and WIO ( $50\text{-}65^{\circ}\text{E}$ ,  $5^{\circ}\text{S}\text{-}10^{\circ}\text{N}$ ), and the trend is estimated over 31 year sliding periods.


592 The land surface temperature is from HadCRUT, SST from HadISST, and tropospheric  
593 temperature from NCEP reanalysis.

594 Figure 5. **(a)** Trend from NCEP re-analysis ( $\text{Pa s}^{-1} \text{ year}^{-1}$ ) and **(b)** model simulated response  
595 ( $\text{Pa s}^{-1}$ ) to WIO warming in vertical velocity ( $\omega$ ) during northern summer (June-  
596 Sept) along the South Asian domain ( $50\text{-}100^\circ\text{E}$ ). Positive (red) values indicate upward  
597 motion.

598 Figure 6. **(a)** Observed trend in precipitation (CRU data, 1901-2012,  $\text{mm day}^{-1}$ ) and near-  
599 surface winds (NCEP, 1948-2012,  $\text{m s}^{-1}$ ), for June-September. **(b)** Model simulated  
600 mean precipitation ( $\text{mm day}^{-1}$ ) and wind (10m) anomalies ( $\text{m s}^{-1}$ ) in response to  
601 warming over the WIO. The model simulated anomalies are estimated from the  
602 sensitivity run where SST anomalies of the order of  $1.5^\circ\text{C}$  is introduced over the WIO  
603 (CFSv2<sub>WIO</sub>), with respect to a model control run (CFSv2<sub>CTL</sub>).




604  
 605 Figure 1. Observed trend in precipitation (mm day<sup>-1</sup> 112 year<sup>-1</sup>) in (a) IMD and (b) CRU datasets, during  
 606 June-September, for the years 1901-2012. Contours denote regions significant at the 95% confidence  
 607 level.




608

609 Figure 2. Observed trend in mean summer (June–September) SST ( $^{\circ}\text{C } 112 \text{ yr}^{-1}$ ) over the global tropics  
610 during 1901–2012.




611

612 Figure 3. Correlation between SST over the western Indian Ocean (WIO, 50-65°E, 5°S-10°N) and  
 613 precipitation over the South Asian subcontinent, for (a) HadISST and IMD precipitation, and (b)  
 614 ERSST and CRU precipitation, for June-September 1901-2012. Contours denote regions significant at  
 615 the 95% confidence level. (c) Time series of SST anomalies (°C, red) over WIO along with CRU (blue)  
 616 precipitation (mm day<sup>-1</sup>) over central South Asia (60°-100°E, 20°-30°N, inset box in b), smoothed with  
 617 a 10 year moving average. Note that the correlation coefficient ( $r = -0.34$ ) between HadISST and CRU  
 618 precipitation is estimated using non-smoothed time series. Kendall's rank correlation test for the two  
 619 variables provided a tau coefficient of -0.3 ( $P < 0.01$ , two tailed). Mann-Kendall test for the trend in the  
 620 time series provided a tau coefficient of 0.6 for SST and -0.2 for precipitation, both significant at 95%  
 621 confidence level.




622

623 Figure 4. Observed (a) mean (°C) and (b) trend (°C 112 year<sup>-1</sup>) of surface temperatures during June-  
 624 September, 1901-2012. For the trend, the color shades represent regions significant at the 95%  
 625 confidence level. (c) Observed trend (°C 65 year<sup>-1</sup>) of upper tropospheric (200 hPa) temperatures during  
 626 June-September, 1948-2012. (d) Time series of the trend in land-sea contrast in the surface ( $T_{surf}$ ) and  
 627 tropospheric ( $T_{trop}$ , 850-200hPa average) temperatures (°C). The land-sea contrast is estimated as the  
 628 difference in the values between the boxes over the South Asian subcontinent (70-85°E, 10-30°N) and  
 629 WIO (50-65°E, 5°S-10°N), and the trend is estimated over 31 year sliding periods. The land surface  
 630 temperature is from HadCRUT, SST from HadISST, and tropospheric temperature from NCEP  
 631 reanalysis.



632

633 Figure 5. (a) Trend from NCEP re-analysis ( $\text{Pa s}^{-1} \text{ year}^{-1}$ ) and (b) model simulated response ( $\text{Pa s}^{-1}$ ) to  
 634 WIO warming in vertical velocity (omega) during northern summer (June-Sept) along the South Asian  
 635 domain (50-100°E). Positive (red) values indicate upward motion.



636

637 Figure 6. (a) Observed trend in precipitation (CRU data, 1901-2012, mm day<sup>-1</sup>) and near-surface winds  
 638 (NCEP, 1948-2012, m s<sup>-1</sup>), for June-September. (b) Model simulated mean precipitation (mm day<sup>-1</sup>)  
 639 and wind (10m) anomalies (m s<sup>-1</sup>) in response to warming over the WIO. The model simulated  
 640 anomalies are estimated from the sensitivity run where SST anomalies of the order of 1.5°C is  
 641 introduced over the WIO (CFSv2<sub>WIO</sub>), with respect to a model control run (CFSv2<sub>CTL</sub>).