HAL
open science

Reduction Operators and Completion of Rewriting Systems

Cyrille Chenavier

To cite this version:

Cyrille Chenavier. Reduction Operators and Completion of Rewriting Systems. 2016. hal-01325907v1

HAL Id: hal-01325907
https://hal.science/hal-01325907v1

Preprint submitted on 2 Jun 2016 (v1), last revised 20 Feb 2017 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Reduction Operators and Completion of Rewriting Systems

CYRILLE CHENAVIER

Abstract

We propose a functional description of rewriting systems where reduction rules are represented by linear maps called reduction operators. We show that reduction operators admit a lattice structure. Using this structure we define the notions of confluence and of Church-Rosser property. We show that these notions are equivalent. We give an algebraic formulation of completion and show that such a completion exists using the lattice structure. We interpret the confluence for reduction operators in terms of Gröbner bases. Finally, we introduce generalised reduction operators relative to non totally ordered sets.

Contents

1 Introduction **1**

2 Rewriting Properties of Reduction Operators **5**
2.1 Lattice Structure and Confluence 5
2.2 Normal Forms, Church-Rosser Property and Newman's Lemma 10
2.3 Reduction Operators and Abstract Rewriting 13

3 Completion and Presentations by Operator **15**
3.1 Confluence for a Pair of Reduction Operators 15
3.2 Completion 17
3.3 Presentations by Operator 20

4 Generalised Reduction Operators **22**
4.1 Algebraic Structure 22
4.2 Rewriting Properties 24

1 Introduction

Convergent rewriting systems appear in rewriting theory to solve decision problems such as the word problem or the ideal membership problem. Completion algorithms were introduced to compute convergent rewriting systems: the Knuth-Bendix completion algorithm [14] for term rewriting [1] and string rewriting [8] or the Buchberger algorithm for Gröbner bases [9, 20, 7, 6] of commutative algebras [9, 10] or associative algebras [16]. In this paper, we propose an algebraic approach of rewriting: we formulate the completion algebraically and show that it can be obtained with an algebraic construction.

We use the functional point of view considered by Berger [3] for rewriting on non-commutative polynomials. The latter are linear combinations of words. In this introduction, we first explain how the functional approach to string rewriting systems works. In the second part, we introduce *reduction operators* and formulate the confluence and the completion with those. We also make explicit the link between reduction operators and rewriting on non-commutative polynomials, which gives us our algebraic constructions.

A Functional Approach to String Rewriting and Gröbner bases

Confluence for String Rewriting Systems. For string rewriting systems, the method consists in considering an idempotent application modelling the rewrite rules. For that, we say that such a system is *semi-reduced* if

1. the left-hand sides of its rewrite rules are pairwise distinct,
2. no right-hand side of its rules is the left-hand side of another one.

For instance, the string rewriting system with alphabet $\{x, y\}$ and with one rewrite rule $yy \longrightarrow yx$ is semi-reduced.

Given a string rewriting system $\langle X \mid R\rangle$ with alphabet X and set of rewrite rules R, we denote by X^* the set of words over X. Our algebraic constructions require that $\langle X \mid R\rangle$ is equipped with a total termination order $<$. In Theorem 2.1.10, we show, using this order, that $\langle X \mid R\rangle$ can be transformed into a unique semi-reduced string rewriting system, so that we may assume that it has this property. The application modelling its rules is the map $S : X^* \longrightarrow X^*$ defined by

1. $S(l(\alpha)) = r(\alpha)$ for every $\alpha \in R$ with left-hand side $l(\alpha)$ and right-hand side $r(\alpha)$,
2. $S(w) = w$ if no element of R has left-hand side w.

The application associated to our example maps yy to yx and fixes all other words.

The order $<$ guarantees that $\langle X \mid R\rangle$ terminates. Thus, it is sufficient to study whether it is confluent or not to know if it is convergent. In order to obtain the functional formulation of confluence, we consider the *extensions of* S, that is, the applications $S_{p,q}$ defined for every pair of integers (p, q) by

1. $S_{p,q}(w) = w_1 r(\alpha) w_2$, if there exist words $w_1,\ w_2$ of length p and q, respectively, and $\alpha \in R$, such that w is equal to $w_1 l(\alpha) w_2$,
2. $S_{p,q}(w) = w$, otherwise.

In the previous example, $S_{0,1}$ maps yyx to yxx and yyy to yxy, and $S_{1,0}$ maps xyy to xyx and yyy to yyx. These applications enable us to characterise the normal forms for $\langle X \mid R\rangle$: a normal form is a word whose every sub-word is fixed by S, that is, the normal forms are the words fixed by all the extensions of S.

Given a word w, we denote by $[w]$ the class of w for the equivalence relation induced by R. The order $<$ being total and well-founded, $[w]$ admits a smallest element. Let M be the application from X^* to itself mapping a word to this minimum. A word w fixed by all the extensions of S but which is not fixed by M is called an *obstruction* of $\langle X \mid R\rangle$. In other words, an obstruction is a normal form which is not minimal in its equivalence class. Hence, the set of obstructions is empty if and only if each equivalence class contains exactly one normal form, so that we obtain the following functional characterisation of confluence: $\langle X \mid R\rangle$ is confluent if and only if the set of obstructions is empty. Considering our example, we deduce from the diagram

$$\begin{tikzcd}
 & yyy \arrow[dl] \arrow[dr] & \\
yxy & & yyx \arrow[dl] \\
 & yxx &
\end{tikzcd}$$

that yxx and yxy belong to the same equivalence class. Moreover, each sub-word of yxx and yxy is fixed by S. Given a total order on the set of words, yxx is either strictly smaller than yxy or is strictly greater than yxy. In the first case, yxy is an obstruction while in the second case yxx is.

Gröbner Bases and Homogeneous Rewriting Systems. Given a set X, we denote by $\mathbb{K}X^*$ the vector space spanned by X^* over a commutative field $\mathbb{K}$: the non-zero elements of this vector space are the finite linear combinations of words with coefficients in $\mathbb{K}$. Let $<$ be a well-founded total order on X^*. Consider a set of rewrite rules R on $\mathbb{K}X^*$ oriented with respect to $<$: for every $\alpha \in R$, $l(\alpha)$ is a word and is greater than every word occurring in the decomposition of $r(\alpha)$ with respect to the basis X^*. We say that R is a *Gröbner basis* when it induces a convergent rewriting system. The set X^* is naturally embedded into $\mathbb{K}X^*$, so that a string rewriting system $\langle X \mid R\rangle$ induces a unique rewriting system on $\mathbb{K}X^*$. Moreover, $\langle X \mid R\rangle$ is convergent if and only if R, seen as a set of rewrite rules on $\mathbb{K}X^*$, is a Gröbner basis.

The functional characterisation of the confluence for string rewriting systems extends into a functional characterisation of Gröbner bases. The notion of semi-reduced string rewriting system is extended as follows: a rewriting system on $\mathbb{K}X^*$ with set of rewrite rules R is said to be *semi-reduced* if

1. the left-hand sides of the elements of R are pairwise distinct,
2. for every $\alpha,\ \beta \in R$, $l(\alpha)$ does not occur to the decomposition of $r(\beta)$ with respect to the basis X^*.

As for string rewriting systems, Theorem 2.1.10 enables us to conclude that every rewriting system on $\mathbb{K}X^*$ can be transformed into a unique semi-reduced one. The application mapping every left-hand side of a rewrite rule to its right-hand side induces an idempotent linear endomorphism $\overline{S}$ of $\mathbb{K}X^*$. For every pair of integers (p, q), we consider the extension $\overline{S}_{p,q} = \mathrm{Id}_{\mathbb{K}X^{\otimes p}} \otimes \overline{S} \otimes \mathrm{Id}_{\mathbb{K}X^{\otimes q}}$ of $\overline{S}$. Let $\overline{M}$ be the endomorphism of $\mathbb{K}X^*$ mapping every element $f \in \mathbb{K}X^*$ to the smallest element of $[f]$ for the natural multi-set order on $\mathbb{K}X^*$ induced by $<$. When one has a string rewriting system, $\overline{S}$, $\overline{S}_{p,q}$ and $\overline{M}$ are the linear endomorphisms of $\mathbb{K}X^*$ extending S, $S_{p,q}$ and M defined in the previous paragraph, respectively. The reasoning we made also works in this context, so that we obtain: R is a Gröbner basis if and only if the set of obstructions is empty.

Berger considered this functional characterisation of Gröbner bases to study *finitely homogeneous rewriting systems*, that is, the systems such that X is finite and there exists an integer N such that the left-hand side and the right-hand side of every element of R are linear combinations of words of length N. In particular, the endomorphism $\overline{S}$ associated to such a system induces an endomorphism of the vector space $\mathbb{K}X^{\otimes N}$ spanned by the set $X^{(N)}$ of words of length N. More generally, for every integer n, the extensions of $\overline{S}$ induce endomorphisms of the vector spaces spanned by the finite sets $X^{(n)}$. We denote by F_n the set of endomorphisms of $\mathbb{K}X^{\otimes n}$ obtained with the restrictions of the extensions of $\overline{S}$. Moreover, the rewrite rules being homogeneous, for every word w of length n, $\overline{M}(w)$ is a linear combination of words of length n, so that $\overline{M}$ also induces endomorphisms $\overline{M}_n$ of $\mathbb{K}X^{\otimes n}$. Hence, the set of obstructions admits a filtration on the length: an obstruction of length n is a word fixed by every element of F_n but not fixed by $\overline{M}_n$. Using the fact that each set $X^{(n)}$ is finite, Berger proved that $\overline{M}_n$ can be obtained from F_n by an algebraic construction and deduced from this an algebraic formulation of obstructions, and thus, of Gröbner bases for homogeneous rewriting systems.

This formulation enables us to obtain various proofs of *Koszulness* [3, 4, 5, 15, 11]. Koszulness has applications in various topics: representation theory, number theory, algebraic and non-commutative geometry, for instance. We refer the reader to [19, 5] for the definition of Koszul algebras and to [18] for an inventory of references about their applications.

Confluence for Non-Homogeneous Rewriting Systems. Consider a set of rewrite rules on $\mathbb{K}X^*$. When this set is non-homogeneous, $\overline{M}$ does not induce endomorphisms of $\mathbb{K}X^{\otimes n}$, so that we cannot construct it by restrictions on finite-dimensional vector spaces. Our first contribution is to show that it can be constructed globally on $\mathbb{K}X^*$. This construction uses the notion of reduction operator which are generalisations of the endomorphisms associated to a rewriting system on $\mathbb{K}X^*$.

Our Results

Reduction Operators: Lattice Structure and Confluence. Let G be a set and let $<$ be a well-founded total order on G. Typically, when we consider homogeneous rewriting systems, G designates the sets $X^{(n)}$ and when we consider non-homogeneous rewriting systems, G is the set X^*. A reduction operator relative to $(G, <)$ is a linear endomorphism of $\mathbb{K}G$ such that

1. T is idempotent,
2. for every $g \in G$, $T(g)$ is either equal to g or is a linear combination of elements of G strictly smaller than g for $<$.

The set of reduction operators, written $\mathbf{RO}\,(G, <)$, is equipped with a lattice structure. Indeed, the first result of the paper about reduction operators is Proposition 2.1.11 which states that the map $T \longrightarrow \ker(T)$ from $\mathbf{RO}\,(G, <)$ to the set of subspaces of $\mathbb{K}G$ is a bijection. This result extends, with a different method, the one of Berger who obtained it when G is finite. The set of subspaces of $\mathbb{K}G$ admits a lattice structure: the order is the inclusion, the lower bound is the intersection and the upper bound is the sum. Using this structure as well as the bijection induced by the kernel map, we deduce a lattice structure on $\mathbf{RO}\,(G, <)$.

Given a subset F of $\mathbf{RO}\,(G, <)$, let $\wedge F$ be its lower bound. We denote by $\mathrm{Red}\,(T)$ the set of elements of G fixed by a reduction operator T. In Lemma 2.1.15, we show that $\mathrm{Red}\,(\wedge F)$ is included in the intersection of all the $\mathrm{Red}\,(T)$, where T belongs to F. The complement of this intersection is written Obs^F. The set F is said to be *confluent* if Obs^F is empty.

Let X be a set and let R be a set of rewrite rules on $\mathbb{K}X^*$, oriented with respect to a well-founded total order on X^*. The endomorphism $\overline{S}$ associated to R, and more generally all the extensions of $\overline{S}$, are reduction operators relative to $(X^*, <)$. Let F be the set of the extensions of $\overline{S}$. In Section 2, we show that $\wedge F$ maps every element f of $\mathbb{K}X^*$ to the smallest element of $[f]$, that is, $\wedge F$ is the operator $\overline{M}$ and Obs^F is the set of obstructions of $\langle X \mid R\rangle$. We obtain our characterisation of Gröbner bases in terms of reduction operators: R is a Gröbner basis if and only if F is confluent.

Completion. Given a rewriting system on a set of terms, words or non-commutative polynomials with a set of rewrite rules R, the Knuth-Bendix completion algorithm or the Buchberger algorithm provide a new set of rewrite rules R', constructed from R and a termination order, such that

1. R' induces a confluent rewriting system,
2. the equivalence relations induced by R' and R are equal.

Here, what we want to complete is a set F of reduction operators. A *completion of* F is a set F' containing F such that

1. F' is confluent,
2. the two operators $\wedge F'$ and $\wedge F$ are equal.

We show that a completion always exists. For that, we use the lattice structure to define an operator C^F called the *F-complement*. Our main result is Theorem 3.2.6 which states that the set $F \cup \{C^F\}$ is a completion of F. When F is associated to a set of rewrite rules on $\mathbb{K}X^*$, the operator C^F maps every obstruction w to $\wedge F(w)$. In Theorem 3.3.11, we use this operator to construct Gröbner bases with reduction operators.

Reduction Operators without Total Order. The Knuth-Bendix completion algorithm does not require a total order on terms, which implies that it could fail. The same phenomena holds for reduction operators when we do not assume that the order on G is total. In this case, the restriction of the kernel map to reduction operators is not onto. In Section 4, we deduce two important consequences of this

fact. The first one is that it could happen that the lattice structure does not exist. The second one is more subtle: even if a set admits a lower bound, the latter does not necessarily have the "right shape". By right shape, we mean that this lower bound does not necessarily come from the lattice structure on the set of subspaces of $\mathbb{K}G$. As a consequence, the F-complement is not always defined. However, the existence of a lower bound with the right shape is sufficient to guarantee that it exists.

Reduction Operators and Computations. We explain the computational aspects of reduction operators. In order to do computations, we consider reduction operators relative to finite sets. Fix such a set $(G, <)$. Since the lattice structure comes from the bijection between $\mathbf{RO}(G, <)$ and subspaces of $\mathbb{K}G$, we need to have an explicit description of this bijection to compute the various algebraic constructions mentioned above. An online version of this implementation is available [1]. This implementation uses the SageMath software [2], written in Python. In the paper, we illustrate several constructions with examples. The examples for which we do not detail the computations were treated with the online version. Moreover, recall that reduction operators have applications to Koszulness. When an algebra has the Koszulness property, a family of important invariants, called *homology groups* [12], can be expressed in terms of upper bound of reduction operators relative to finite sets. Hence, this is a context where our implementation can be used. The computation of these invariants already exists [2]. Here, we propose an implementation using other techniques.

Organisation.

In Section 2.1, we define the notion of reduction operator relative to a well-ordered set. We also equip the set of these operators with a lattice structure and formulate the notion of confluence. In Section 2.2, we define several notions from abstract rewriting theory in terms of reduction operators: normal forms, Church-Rosser property and local confluence. We show that the notions of confluence, local confluence and Church-Rosser property are equivalent. In Section 2.3, we explain our notion of confluence from the viewpoint of abstract rewriting. Section 3.1 contains results about a pair of reduction operators. These results are necessary in Section 3.2. In the latter, we define the notions of completion, complement and minimal complement. We also show that a minimal complement always exists. In Section 3.3, we formulate the notions of presentation by operator and of confluent presentation by operator. We also link the latter with Gröbner bases. In Section 4.1, we formulate a general definition of reduction operator, which is relative to an ordered set. We show that the set of these operators is an ordered set but does not necessarily admit a lattice structure. In Section 4.2, we define the notion of completable set of reduction operators and study their rewriting properties.

Acknowledgement. This work was supported by the Sorbonne-Paris-Cité IDEX grant Focal and the ANR grant ANR-13-BS02-0005-02 CATHRE.

2 Rewriting Properties of Reduction Operators

2.1 Lattice Structure and Confluence

2.1.1. Notations. We denote by $\mathbb{K}$ a commutative field. We say vector space instead of $\mathbb{K}$-vector space. Let X be a set. We denote by $\mathbb{K}X$ the vector space spanned by X: its non-zero elements are the finite formal linear combinations of elements of X with coefficients in $\mathbb{K}$. An element of X is called a *generator* of $\mathbb{K}X$. For every $v \in \mathbb{K}X \setminus \{0\}$, there exist a unique finite subset S_v of X and a unique family of non zero scalars $(\lambda_x)_{x \in S_v}$ such that v is equal to $\sum_{x \in S_v} \lambda_x x$. The set S_v is the *support* of v.

[1] http://pastebin.com/0YZCfAD4
[2] http://www.sagemath.org

2.1.2. Leading Generator and Leading Coefficient. Let $(G, <)$ be a *well-ordered set*, that is, G is a set and $<$ is a well-founded total order on G. The order on G being total, every finite subset of G admits a greatest element. In particular, for every $v \in \mathbb{K}G \setminus \{0\}$, the support of v admits a greatest element, written $\lg(v)$. We also write $\mathrm{lc}(v) = \lambda_{\lg(v)}$. The elements $\lg(v)$ and $\mathrm{lc}(v)$ are the *leading generator* and the *leading coefficient* of v, respectively. We extend the order $<$ on G into a partial order on $\mathbb{K}G$ in the following way: we have $u < v$ if $u = 0$ and v is different from 0 or if $\lg(u) < \lg(v)$.

Throughout Section 2 we fix a well-ordered set $(G, <)$.

2.1.3. Reduction Operators. A *reduction operator relative to* $(G, <)$ is an idempotent linear endomorphism T of $\mathbb{K}G$ such that for every $g \in G$, we have $T(g) \leq g$. We denote by $\mathbf{RO}(G, <)$ the set of reduction operators relative to $(G, <)$. Given $T \in \mathbf{RO}(G, <)$, a generator g is said to be *T-reduced* if $T(g)$ is equal to g. We denote by $\mathrm{Red}(T)$ the set of T-reduced generators and by $\mathrm{Nred}(T)$ the complement of $\mathrm{Red}(T)$ in G.

2.1.4. Reduction Matrices. In our examples, we sometimes consider the case where $(G, <)$ is a totally ordered finite set. We denote by $G = \{g_1 < \cdots < g_n\}$ such a set. In this case, we use matrix notations to describe linear maps, and thus, to describe reduction operators. For that, given an endomorphism T of $\mathbb{K}G$, the matrix of T with respect to the basis $\{g_1, \cdots, g_n\}$ is called the *canonical matrix of T relative to* $(G, <)$. Moreover, we say that a square matrix M is a *reduction matrix* if the following conditions are fulfilled:

1. M is upper triangular and the elements of its diagonal are equal to 0 or 1,
2. if an element of the diagonal of M is equal to 0, then the other elements of the line to which it belongs are equal to 0,
3. if an element of the diagonal of M is equal to 1, then the other elements of the column to which it belongs are equal to 0.

Using the fact that a reduction matrix is idempotent, we deduce that an endomorphism of $\mathbb{K}G$ is a reduction operator relative to $(G, <)$ if and only if its canonical matrix is a reduction matrix.

2.1.5. Examples.

1. The zero matrix and the identity matrix are reduction matrices. More generally, every diagonal matrix admitting only 0 or 1 on its diagonal is a reduction matrix.
2. The matrix
$$\begin{pmatrix} 1 & 1 & 1 & 1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix},$$
is a reduction matrix.
3. From [13], we consider the presentation of the monoid ${\mathbf{B}_3}^+$ by 3 generators x, y and z submitted to two relations $yz = x$ and $zx = xy$. Let $X = \{x < y < z\}$. Let X^* be the free monoid over X: this is the set of (possibly empty) words written with the alphabet X. This set is totally ordered for the deg-lex order, still denoted by $<$, induced by the order on X. For this order, yz is greater than x and zx is greater than xy. Hence, the endomorphism of $\mathbb{K}X^*$ defined on the basis X^* by $S(yz) = x$, $S(zx) = xy$ and $S(w) = w$ for every $w \in X^* \setminus \{yz, zx\}$, is a reduction operator relative to $(X^*, <)$.

2.1.6. Reduced Basis. Let V be a subspace of $\mathbb{K}G$. A *reduced basis* of V is a basis $\mathscr{B}$ of V satisfying the following conditions:

1. for every $e \in \mathscr{B}$, $\mathrm{lc}\,(e)$ is equal to 1,
2. given two different elements e and e' of $\mathscr{B}$, $\mathrm{lg}\,(e')$ does not belong to the support of e.

2.1.7. Notation. Let V be a subspace of $\mathbb{K}G$ and let $\mathscr{B}$ be a reduced basis of V. Let e and e' be two distinct elements of $\mathscr{B}$. The condition 2 of 2.1.6 implies that $\mathrm{lg}\,(e)$ is different from $\mathrm{lg}\,(e')$. Hence, $\mathscr{B}$ is indexed by the set $\tilde{G} = \{\mathrm{lg}\,(e)\,,\ e \in \mathscr{B}\}$. In the sequel, a reduced basis $\mathscr{B}$ is written $\mathscr{B} = (e_g)_{g\in\tilde{G}}$, where for every $g \in \tilde{G}$, we have $\mathrm{lg}\,(e_g) = g$.

2.1.8. Remark. Let V be a subspace of $\mathbb{K}G$ and let $(e_g)_{g\in\tilde{G}}$ be a reduced basis of V. For every $g \in G$, let V_g be the set of elements of V with leading generator g. The set V_g is non empty if and only if g belongs to $\tilde{G}$. Hence, if $\mathscr{B}_1 = (e_g)_{g\in\tilde{G}_1}$ and $\mathscr{B}_2 = \left(e'_g\right)_{g\in\tilde{G}_2}$ are two reduced bases of V, the two sets $\tilde{G}_1$ ant $\tilde{G}_2$ are equal.

2.1.9. Examples.

1. Let $G = \{g_1 < g_2 < g_3 < g_4\}$ and let V be the subspace of $\mathbb{K}G$ spanned by the elements $v_1 = g_2 - g_1$, $v_2 = g_4 - g_3$ and $v_3 = g_4 - g_2$. The elements $\mathrm{lg}\,(v_2)$ and $\mathrm{lg}\,(v_3)$ are equal to g_4, so that $\{v_1, v_2, v_3\}$ is not a reduced basis of V. Letting ${v_2}' = v_3 - v_2$, that is, ${v_2}' = g_3 - g_2$, the set $\{v_1, {v_2}', v_3\}$ is a basis of V. This is still not a reduced basis because $\mathrm{lg}\,(v_1) = g_2$ appears in the supports of ${v_2}'$ and v_3. Letting ${v_2}'' = {v_2}' + v_1$ and ${v_3}' = v_3 + v_1$, that is, ${v_2}'' = g_3 - g_1$ and ${v_3}' = g_4 - g_1$, the set $\{v_1, {v_2}'', {v_3}'\}$ is a basis of V. This basis is reduced. Using the notation introduced in 2.1.7, it is equal to $\{e_{g_2}, e_{g_3}, e_{g_4}\}$, where e_{g_i} is equal to $g_i - g_1$ for $i \in \{2, 3, 4\}$.
2. A *semi-reduced* string rewriting system is a string rewriting system $\langle X \mid R\rangle$ such that the left-hand sides of the elements of R are pairwise distinct and if no right-hand side of an element of R is the left-hand side of another one. The set $\{w - w',\ w \longrightarrow w' \in R\}$ is a reduced basis of the vector space it spans.

2.1.10. Theorem. *Let $(G, <)$ be a well-ordered set. Every subspace of $\mathbb{K}G$ admits a unique reduced basis.*

Proof. Let V be a subspace of $\mathbb{K}G$. First, we construct by induction on G a reduced basis of V. Let g_0 be the smallest element of G. If V_{g_0} is empty, we let $\mathscr{B}_{g_0} = \emptyset$. In the other case, g_0 belongs to V and we let $\mathscr{B}_{g_0} = \{g_0\}$. Let $g \in G$. Assume by induction that for every $g' < g$ we have built a set $\mathscr{B}_{g'}$ such that the following conditions are fulfilled:

1. For every $g' < g$, the set $\mathscr{B}_{g'}$ contains at most one element.
2. Let
$$I_g = \{g' \in G,\ g' < g \text{ and } \mathscr{B}_{g'} \neq \emptyset\}\,.$$
For every $g' \in I_g$
   (a) the unique element $e_{g'}$ of $\mathscr{B}_{g'}$ belongs to V,
   (b) $\mathrm{lg}\,(e_{g'})$ is equal to g' and $\mathrm{lc}\,(e_{g'})$ is equal to 1,
   (c) for every $\tilde{g} \in I_g$ such that $\tilde{g}$ is different from g', $\tilde{g}$ does not belong to the support of $e_{g'}$,
   (d) the set $V_{g'}$ is included in $\mathbb{K}\{e_{\tilde{g}},\ \tilde{g} \in I_g\}$.

If V_g is empty, we let $\mathscr{B}_g = \emptyset$. If V_g is non empty, let v_g be an element of V_g such that $\mathrm{lc}\,(v_g)$ is equal to 1. In particular, v_g admits a decomposition

$$v_g = \sum_{g' \in J} \mu_{g'} g' + g,$$

where for every $g' \in J$, we have $g' < g$. We let $\mathscr{B}_g = \{e_g\}$, where

$$e_g = v_g - \sum_{g' \in I_g} \mu_{g'} e_{g'}.$$

In particular, $\mathscr{B}_g$ contains at most one element. By construction, e_g belongs to V, $\mathrm{lg}\,(e_g)$ is equal to g and $\mathrm{lc}\,(e_g)$ is equal to 1, so that Point 2a and Point 2b hold. Moreover, for every $g' \in I_g$, g' does not belong to the support of e_g, so that Point 2c holds. It remains to show that V_g is included in the vector space spanned by the elements $e_{g'}$ where g' belongs to $I_g \cup \{g\}$. This vector space is equal to $\mathbb{K}\{e_{\tilde{g}},\ \tilde{g} \in I_g\} \oplus \mathbb{K}e_g$. Let v be an element of V_g. Then, $v - \mathrm{lc}\,(v)\, e_g$ belongs to V and $\mathrm{lg}\,(v - \mathrm{lc}\,(v)\, e_g)$ is strictly smaller than g. By the induction hypothesis, $v - \mathrm{lc}\,(v)\, e_g$ belongs to $\mathbb{K}\{e_{\tilde{g}},\ \tilde{g} \in I_g\}$, so that v belongs to $\mathbb{K}\{e_{\tilde{g}},\ \tilde{g} \in I_g\} \oplus \mathbb{K}e_g$. This inductive construction provides a family of sets $(\mathscr{B}_g)_{g \in G}$ such that $\mathscr{B} = \bigcup_{g \in G} \mathscr{B}_g$ is a generating set of $V = \bigsqcup_{g \in G} V_g$. Moreover, this family is free because the leading generators of its elements are pairwise distinct. Hence, $\mathscr{B}$ is a basis of V. This basis is reduced by construction.

Let us show that such a basis is unique. Let $\mathscr{B}_1 = (e_g)_{g \in \tilde{G}}$ and $\mathscr{B}_2 = \left(e'_g\right)_{\in \tilde{G}}$ be two reduced bases of V. Let $g \in \tilde{G}$. Assume that e_g is different from e'_g. Let

$$e'_g - e_g = \sum_{g' \in I} \lambda_{g'} e'_{g'},$$

be the decomposition of $e'_g - e_g$ with respect to the basis $\mathscr{B}_2$. The leading generator of $e'_g - e_g$ is equal to the greatest element of I, so that it belongs to $\tilde{G}$. Let $\tilde{G}^c$ be the complement of $\tilde{G}$ in G. The condition 2 of the definition of a reduced basis implies that $e_g - g$ and $e'_g - g$ belong to $\mathbb{K}\tilde{G}^c$. Hence, $\mathrm{lg}\left(e'_g - e_g\right)$ belongs to $\tilde{G}^c$, which is contradiction. Thus, for every $g \in \tilde{G}$, the two elements e_g and e'_g are equal, that is, $\mathscr{B}_1$ and $\mathscr{B}_2$ are equal.

□

2.1.11. Proposition. *Let V be a subspace of $\mathbb{K}G$. There exists a unique reduction operator T with kernel V. Moreover,* $\mathrm{Nred}\,(T)$ *is equal to $\tilde{G}$, where $(e_g)_{g \in \tilde{G}}$ is the reduced basis of V*

Proof. Let T be the endomorphism of $\mathbb{K}G$ defined on the basis G in the following way

$$T(g) = \begin{cases} g - e_g, \text{ if } g \in \tilde{G}, \\ g, \text{ ohterwise.} \end{cases}$$

By definition of a reduced basis, T is a reduction operator relative to $(G, <)$. By construction, the kernel of T is equal to V and $\mathrm{Nred}\,(T)$ is equal to $\tilde{G}$.

Let T_1 and T_2 be two reduction operators with kernel V. The two sets $\{g - T_1(g),\ T_1(g) \neq g\}$ and $\{g - T_2(g),\ T_2(g) \neq g\}$ are reduced bases of V. From Theorem 2.1.10, these sets are equal. Hence, $T_1(g)$ is different from g if and only if $T_2(g)$ is different from g and in this case, $T_1(g)$ is equal to $T_2(g)$. It follows that $T_1(g)$ and $T_2(g)$ are equal for every $g \in G$, so that T_1 and T_2 are equal. □

2.1.12. Notation. Proposition 2.1.11 implies that the kernel map induces a bijection between $\mathbf{RO}\,(G, <)$ and the set of subspaces of $\mathbb{K}G$. The inverse of this bijection is written θ.

2.1.13. Lattice Structure. We consider the binary relation on $\mathbf{RO}(G,<)$ defined by

$$T_1 \preceq T_2 \text{ if and only if } \ker(T_2) \subset \ker(T_1).$$

This relation is reflexive and transitive. From Proposition 2.1.11, it is also anti-symmetric. Hence, it is an order relation on $\mathbf{RO}(G,<)$. Moreover, we have the equivalence:

$$T_1 \preceq T_2 \text{ if and only if } T_1 \circ T_2 = T_1. \tag{1}$$

Let us equip $\mathbf{RO}(G,<)$ with a lattice structure. The lower bound $T_1 \wedge T_2$ and the upper bound $T_1 \vee T_2$ of two elements T_1 and T_2 of $\mathbf{RO}(G,<)$ are defined in the following manner:

$$\begin{cases} T_1 \wedge T_2 = \theta\left(\ker(T_1) + \ker(T_2)\right), \\ T_1 \vee T_2 = \theta\left(\ker(T_1) \cap \ker(T_2)\right). \end{cases}$$

2.1.14. Example. Let $G = \{g_1 < g_2 < g_3 < g_4\}$ and $P = (T_1, T_2)$, where

$$T_1 = \begin{pmatrix} 1 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 \end{pmatrix} \text{ and } T_2 = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}.$$

The kernels of T_1 and T_2 are equal to $\mathbb{K}\{g_2 - g_1\} \oplus \mathbb{K}\{g_4 - g_3\}$ and $\mathbb{K}\{g_4 - g_2\}$, respectively. The kernel of $T_1 \wedge T_2$ is the vector space spanned by $v_1 = g_2 - g_1$, $v_2 = g_4 - g_3$ and $v_3 = g_4 - g_2$. This is the vector space of Example 2.1.9, Point 1. Hence, the kernel of $T_1 \wedge T_2$ is the vector space spanned by $\{g_2 - g_1, g_3 - g_1, g_4 - g_1\}$, so that we have

$$T_1 \wedge T_2 = \begin{pmatrix} 1 & 1 & 1 & 1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}.$$

2.1.15. Lemma. *Let T_1 and T_2 be two reduction operators relative to $(G,<)$. Then, we have:*

$$T_1 \preceq T_2 \Longrightarrow \mathrm{Red}(T_1) \subset \mathrm{Red}(T_2).$$

Proof. Let $\mathscr{B}_1 = (e_g)_{g \in \tilde{G}_1}$ and $\mathscr{B}_2 = \left(e'_g\right)_{g \in \tilde{G}_2}$ be the reduced bases of $\ker(T_1)$ and $\ker(T_2)$, respectively. We consider the notations of Remark 2.1.8: given a subspace V of $\mathbb{K}G$, let V_g be the set of elements of V with leading generator g. The sets $\left\{g \in G,\ \ker(T_1)_g \neq \emptyset\right\}$ and $\left\{g \in G,\ \ker(T_2)_g \neq \emptyset\right\}$ are equal to $\tilde{G}_1$ and $\tilde{G}_2$, respectively. Hence, if $\ker(T_2)$ is included in $\ker(T_1)$, then $\tilde{G}_2$ is included in $\tilde{G}_1$. From Proposition 2.1.11, we deduce that $\mathrm{Red}(T_1)$ is included in $\mathrm{Red}(T_2)$.

□

2.1.16. Obstructions. Let F be a subset of $\mathbf{RO}(G,<)$. We let

$$\mathrm{Red}(F) = \bigcap_{T \in F} \mathrm{Red}(T) \text{ and } \wedge F = \theta\left(\sum_{T \in F} \ker(T)\right).$$

For every $T \in F$, we have $\wedge F \preceq T$. Thus, from Lemma 2.1.15, the set $\mathrm{Red}(\wedge F)$ is included in $\mathrm{Red}(T)$ for every $T \in F$, so that it is included in $\mathrm{Red}(F)$. We write

$$\mathrm{Obs}^F = \mathrm{Red}(F) \setminus \mathrm{Red}(\wedge F). \tag{2}$$

2.1.17. Confluence. A subset F of $\mathbf{RO}(G,<)$ is said to be *confluent* if Obs^F is the empty set.

2.1.18. Examples.

1. We consider the pair P of Example 2.1.14. The set $\mathrm{Red}(P)$ is equal to $\{g_1, g_3\}$. Moreover, $\mathrm{Red}(T_1 \wedge T_2)$ is equal to $\{g_1\}$, so that Obs^P is equal to $\{g_3\}$. Hence, P is not confluent.

2. We consider the operator S defined in Point 3 of Example 2.1.5. Let S_1 and S_2 be the restrictions to the vector space spanned by words of length smaller than 3 of $S \otimes \mathrm{Id}_{\mathbb{K}X}$ and $\mathrm{Id}_{\mathbb{K}X} \otimes S$, respectively. Let $P = (S_1, S_2)$. The set of not $\wedge P$-reduced generators is equal to $\{xyz, xzx, yxy, yyz, yzx, yzy, yzz, zxx, zxy, zxz, zyz, zzx\}$. We see that there exists exactly one element of this list which also belongs to $\mathrm{Red}(P)$: yxy. Hence, Obs^P is equal to $\{yxy\}$, so that P is not confluent.

2.2 Normal Forms, Church-Rosser Property and Newman's Lemma

Throughout this section we fix a subset F of $\mathbf{RO}(G,<)$. We denote by $\langle F\rangle$ the submonoid of $(\mathrm{End}(\mathbb{K}G), \circ)$ spanned by F.

2.2.1. Multi-Set Order. Given an element v of $\mathbb{K}G$, let S_v be the support of v. We introduce the order $<_{\mathrm{mul}}$ on $\mathbb{K}G$ defined in the following way: we have $v <_{\mathrm{mul}} v'$ if for every $g \in S_v$ such that g does not belong to $S_{v'}$, there exists an element $g' \in S_{v'}$ not appearing in S_v, such that $g < g'$. The order $<$ being well-founded, this also the case for $<_{\mathrm{mul}}$. Moreover, given $v \in \mathbb{K}G$ and $T \in F$ such that v does not belong to $\mathbb{K}\mathrm{Red}(T)$, $T(v)$ is strictly smaller than v for $<_{\mathrm{mul}}$. Finally, we also remark that 0 is the smallest element of $\mathbb{K}G$ for $<_{\mathrm{mul}}$.

2.2.2. Normal Forms.

1. An *F-normal form* is an element of $\mathbb{K}\mathrm{Red}(F)$.

2. Let v and v' be two elements of $\mathbb{K}G$. We say that v *rewrites into* v' if there exists $R \in \langle F\rangle$ such that v' is equal to $R(v)$.

3. Let v be an element of $\mathbb{K}G$. An *F-normal form of* v is an F-normal form v' such that v rewrites into v'.

2.2.3. Remark. Let $v \in \mathbb{K}G$. If v rewrites into v', then $v - v'$ belongs to $\ker(\wedge F)$.

2.2.4. Church-Rosser Property. We say that F has the *Church-Rosser property* if for every $v \in \mathbb{K}G$, v rewrites into $\wedge F(v)$.

2.2.5. Theorem. *A subset of* $\mathbf{RO}(G,<)$ *is confluent if and only if it has the Church-Rosser property.*

Proof. Let F be a subset of $\mathbf{RO}(G,<)$.

Assume that F has the Church-Rosser property. Let g be an element of $\mathrm{Red}(F)$. For every $T \in F$, $T(g)$ is equal to g. As a consequence, $R(g)$ is equal to g for every $R \in \langle F\rangle$. The set F having the Church-Rosser property, we deduce that g is equal to $\wedge F(g)$, so that g belongs to $\mathrm{Red}(\wedge F)$. That shows that $\mathrm{Red}(F)$ is included in $\mathrm{Red}(\wedge F)$, that is, F is confluent.

Assume that F is confluent. Let us show by induction on $<_{\mathrm{mul}}$ that for every $v \in \mathbb{K}G$, v rewrites into $\wedge F(v)$. If v is equal to 0, this is obvious. Let $v \in \mathbb{K}G$. Assume by induction that for every $v' \in \mathbb{K}G$ such that $v' <_{\mathrm{mul}} v$, v' rewrites into $\wedge F(v')$. If v belongs to $\mathbb{K}\mathrm{Red}(\wedge F)$, then v is equal to $\wedge F(v)$, so that v rewrites into $\wedge F(v)$. Assume that v does not belong to $\mathbb{K}\mathrm{Red}(\wedge F)$. The set F being confluent, v does not belong to $\mathbb{K}\mathrm{Red}(F)$, that is, there exists $T \in F$ such that $T(v)$ is different from v. The element $T(v)$ is strictly smaller than v for $<_{\mathrm{mul}}$. By the induction hypothesis, there exists $R \in \langle F\rangle$

such that $R(T(v))$ is equal to $\wedge F\left(T(v)\right)$. The inequality $\wedge F \preceq T$ implies that $\wedge F \circ T$ is equal to $\wedge F$, so that $\wedge F\left(T(v)\right)$ is equal to $\wedge F(v)$. Hence, $R' = R \circ T$ is an element of $\langle F \rangle$ such that $R'(v)$ is equal to $\wedge F(v)$. Thus, v rewrites into $\wedge F(v)$.

□

2.2.6. Lemma. *Let v be an element of $\mathbb{K}G$ and let $(R_1, \cdots, R_n, \cdots)$ be a sequence of elements of $\langle F \rangle$ such that for every integer n, R_n is a right divisor of R_{n+1} in $\langle F \rangle$. The sequence $\left(R_n(v)\right)_{n\in\mathbb{N}}$ is stationary.*

Proof. We proceed by induction on v. If v is equal to 0, then the sequence $\left(R_n(v)\right)_{n\in\mathbb{N}}$ is constant, equals to 0. Let $v \in \mathbb{K}G$. Assume that Lemma 2.2.6 holds for every $v' \in \mathbb{K}G$ such that v' is strictly smaller than v for $<_{\text{mul}}$. If the sequence $\left(R_n(v)\right)_{n\in\mathbb{N}}$ is constant equals to v, there is nothing to prove. In the other case, there exists n_0 such that $R_{n_0}(v)$ is different from v, so that we have $R_{n_0}(v) <_{\text{mul}} v$. By hypothesis, for every integer n, R_n is a right divisor of R_{n+1} in $\langle F \rangle$, that is, there exists $R'_n \in \langle F \rangle$ such that R_{n+1} is equal to $R'_n \circ R_n$. Let $\left(Q_n\right)_{n\in\mathbb{N}}$ be the sequence of elements of $\langle F \rangle$ defined by

$$Q_1 = R'_{n_0} \text{ and } Q_{n+1} = R'_{n_0+n} \circ Q_n.$$

For every integer n, Q_n is a right divisor of Q_{n+1} in $\langle F \rangle$. By the induction hypothesis, the sequence $\left(Q_n\left(R_{n_0}(v)\right)\right)_{n\in\mathbb{N}}$ is stationary. Moreover, for every integer n, $Q_n \circ R_{n_0}$ is equal to R_{n_0+n}, so that the sequence $\left(R_n(v)\right)_{n\in\mathbb{N}}$ is stationary. Hence, Lemma 2.2.6 holds.

□

2.2.7. Proposition. *Every element of $\mathbb{K}G$ admits an F-normal form.*

Proof. Let v be an element of $\mathbb{K}G$. We have to show that there exists $R \in \langle F \rangle$ such that $R(v)$ belongs to $\mathbb{K}\text{Red}\,(F)$. Assume by way of contradiction that for every $R \in \langle F \rangle$, $R(v)$ does not belong to $\mathbb{K}\text{Red}\,(F)$. The morphism $\text{Id}_{\mathbb{K}G}$ belonging to $\langle F \rangle$, v does not belong to $\mathbb{K}\text{Red}\,(F)$. In particular, there exists $T_1 \in F$ such that v does not belong to $\mathbb{K}\text{Red}\,(T_1)$. Assume that we have constructed elements $T_1, \cdots, T_n$ of F. The morphism $R_n = T_n \circ \cdots \circ T_1$ belongs to $\langle F \rangle$. Hence, the element $R_n(v)$ does not belong to $\mathbb{K}\text{Red}\,(F)$, so that there exists $T_{n+1} \in F$ such that $R_n(v)$ does not belong to $\mathbb{K}\text{Red}\,(T_{n+1})$. This process enables us to obtain a sequence $\left(R_n\right)_{n\in\mathbb{N}}$ of elements of $\langle F \rangle$ such that for every integer n, R_n is a right divisor of R_{n+1} in $\langle F \rangle$ and such that the sequence $\left(R_n(v)\right)_{n\in\mathbb{N}}$ is not stationary. This is a contradiction with Lemma 2.2.6. Hence, Proposition 2.2.7 holds.

□

2.2.8. Notation. For every $v \in \mathbb{K}G$, let $[v]$ be the set of elements $v' \in \mathbb{K}G$ such that $v' - v$ belongs to $\ker\left(\wedge F\right)$.

2.2.9. Lemma. *For every $v \in \mathbb{K}G$, $\wedge F(v)$ is the smallest element of $[v]$ for the order $<_{\text{mul}}$. Moreover, if every element v of $\mathbb{K}G$ admits exactly one F-normal form, this normal form is equal to $\wedge F(v)$.*

Proof. Let us show the first assertion. Let $v \in \mathbb{K}G$ and let $v' \in [v]$. In particular, v' belongs to $[\wedge F(v)]$, that is, there exists $v'' \in \ker\left(\wedge F\right)$ such that

$$v' = \wedge F(v) + v''. \tag{3}$$

The element v'' belonging to $\ker\left(\wedge F\right)$, it admits a decomposition

$$v'' = \sum \lambda_i \left(g_i - \wedge F(g_i)\right), \tag{4}$$

where each g_i is not $\wedge F$-reduced. Let $g \in G$ be an element of the support of $\wedge F(v)$ not appearing to the one of v'. Let us show that there exists an index i such that

1. g_i is greater than g,

2. g_i does not belong to the support of $\wedge F(v)$,

3. g_i belongs to the support of v'.

Relation (3) and the hypothesis on g imply that the latter belongs to the support of v''. Moreover, g belongs to the image of $\wedge F$, that is, it is $\wedge F$-reduced. From Relation (4), we deduce that g belongs to the support of $\wedge F(g_i)$ for some i. The element g_i being not $\wedge F$-reduced, g is smaller than g_i and does not belong to the support of $\wedge F(v)$. Finally, g_i belonging to the support of v'' and not to the one of $\wedge F(v)$, Relation (3) implies that it belongs to the support of v'. Hence, we have $\wedge F(v) <_{\mathrm{mul}} v'$ for every $v' \in [v]$, so that the first part of the lemma holds.

Let us show the second part of the lemma. Assume that every element v of $\mathbb{K}G$ admits a unique F-normal form, written $N(v)$. It is clear that the operator N is idempotent. Moreover, for every $R \in \langle F \rangle$ and for every $g \in G$, we have either $R(g) = g$ or $R(g) < g$. As a consequence, for every $g \in G$, we have either $N(g) = g$ or $N(g) < g$. We conclude that N belongs to $\mathbf{RO}\,(G, <)$. Let us show that N is equal to $\wedge F$. Let T be an element of F and let v be an element of $\ker(T)$. The element v rewrites into 0, so that $N(v)$ is equal to 0. Thus, the kernel of T is included in the kernel of N for every $T \in F$, that is, N is smaller than T for every $T \in F$. Thus, we have the inequality $N \preceq \wedge F$. Moreover, from Relation (1) (see 2.1.13), for every $T \in F$, the operator $\wedge F \circ T$ is equal to $\wedge F$. Hence, for every $R \in \langle F \rangle$, the operator $\wedge F \circ R$ is equal to $\wedge F$. As a consequence, $\wedge F \circ N$ is equal to $\wedge F$. Using again Relation (1), we have $\wedge F \preceq N$. Hence, N is equal to $\wedge F$, so that Lemma 2.2.9 holds.

□

2.2.10. Proposition. *The set F is confluent if and only if every element of $\mathbb{K}G$ admits a unique F-normal form.*

Proof. Assume that F is confluent. Let v be an element of $\mathbb{K}G$ and let v_1 and v_2 be two F-normal forms of v. Let R_1 and R_2 be two elements of $\langle F \rangle$ such that $R_i(v)$ is equal to v_i, for $i = 1$ or 2. The elements $v - v_1$ and $v - v_2$ belong to $\ker(\wedge F)$. Hence, $v_1 - v_2$ belongs to $\ker(\wedge F)$. Moreover, v_1 and v_2 belonging to $\mathbb{K}\mathrm{Red}\,(F)$, $v_1 - v_2$ also belongs to $\mathbb{K}\mathrm{Red}\,(F)$, that is, $\mathbb{K}\mathrm{Red}\,(\wedge F)$ since F is confluent. Thus, $v_1 - v_2$ belongs to the vector space $\mathbb{K}\mathrm{Red}\,(\wedge F) \cap \ker(\wedge F)$. The operator $\wedge F$ being a projector, this vector space is reduced to $\{0\}$. We conclude that v_1 is equal to v_2, so that v admits a unique F-normal form.

Assume that every element of $\mathbb{K}G$ admits a unique F-normal form. Let v be an element of $\mathbb{K}G$. From Lemma 2.2.9, the normal form of v is equal to $\wedge F(v)$. Hence, v rewrites into $\wedge F(v)$. We conclude that F has the Church Rosser-property, that is, F is confluent from Theorem 2.2.5.

□

2.2.11. Local Confluence. We say that F is *locally confluent* if for every $v \in \mathbb{K}G$ and for every $T_1,\ T_2 \in F$, there exists $v' \in \mathbb{K}G$ such that $T_1(v)$ and $T_2(v)$ rewrite into v'.

The last result of this section is the formulation of Newman's Lemma [17] in terms of reduction operators.

2.2.12. Proposition. *The set F is confluent if and only if it is locally confluent.*

Proof. Assume that F is confluent. Let v be an element of $\mathbb{K}G$ and let $T_1,\ T_2 \in F$. Let $i = 1$ or 2. From Theorem 2.2.5, $T_i(v)$ rewrites into $\wedge F\,(T_i(v))$. The latter is equal to $\wedge F(v)$ from Relation (1) (see 2.1.13). Hence, F is locally confluent.

Assume that F is locally confluent. From Proposition 2.2.10, it is sufficient to show that every element v of $\mathbb{K}G$ admits a unique F-normal form. We show this assertion by induction on v. If v is equal to 0, there is nothing to prove. Let v be an element of $\mathbb{K}G$. Assume by induction that for every $v' <_{\mathrm{mul}} v$, v' admits a unique F-normal form. If v belongs to $\mathbb{K}\mathrm{Red}\,(F)$, then v admits a unique F-normal form which is itself. Assume that v does not belong to $\mathbb{K}\mathrm{Red}\,(F)$. Let v_1 and v_2 be two F-normal forms of v. For $i = 1$ or 2, there exists $R_i \in \langle F \rangle$ such that v_i is equal to $R_i(v)$. We write $R_i = R'_i \circ T_i$, where T_i and R'_i belong to F and $\langle F \rangle$, respectively. The operator T_i is chosen in such a way that $T_i(v)$ is different from v. The set F being locally confluent, there exists $u \in \mathbb{K}G$ such that

$T_i(v)$ rewrites into u. From Proposition 2.2.7, u admits an F-normal form $\widehat{u}$. The latter is also an F-normal form of $T_i(v)$. Moreover, v_i is equal to $R'_i(T_i(v))$, so that it is also an F-normal form of $T_i(v)$. The latter is strictly smaller than v for $<_{\mathrm{mul}}$. By the induction hypothesis, it admits a unique F-normal form, so that v_i is equal to $\widehat{u}$ for $i = 1$ or 2. In particular, v_1 is equal to v_2, so that v admits a unique F-normal form.

□

2.3 Reduction Operators and Abstract Rewriting

We fix a subset F of $\mathbf{RO}(G, <)$.

2.3.1. Abstract Rewriting Systems. An *abstract rewriting system* is a pair $\left(A, \longrightarrow\right)$ where A is a set and $\longrightarrow$ is a binary relation on A. We write $a \longrightarrow b$ instead of $(a, b) \in \longrightarrow$. We denote by $\xrightarrow{+}$, $\xrightarrow{*}$ and $\xleftrightarrow{*}$ the transitive closure, the reflexive transitive closure and the reflexive transitive symmetric closure of $\longrightarrow$, respectively. Finally, we recall the notion of normal form: a normal form of $\left(A, \longrightarrow\right)$ is an element a of A such that there does not exist any element of $\longrightarrow$ with the form $a \longrightarrow b$.

2.3.2. Confluence and Church-Rosser Property. Let $\left(A, \longrightarrow\right)$ be an abstract rewriting system. We say that $\longrightarrow$ is *confluent* if for every $a,\ b,\ c \in A$ such that $a \xrightarrow{*} b$ and $a \xrightarrow{*} c$, there exists $d \in A$ such that $b \xrightarrow{*} d$ and $c \xrightarrow{*} d$. We say that $\longrightarrow$ has the *Church-Rosser property* if for every $a,\ b \in A$ such that $a \xleftrightarrow{*} b$, there exists $c \in A$ such that $a \xrightarrow{*} c$ and $b \xrightarrow{*} c$. Recall from [1, Theorem 2.1.5] that $\longrightarrow$ is confluent if and only if it has the Church-Rosser property.

2.3.3. Definition. We consider the abstract rewriting system $\left(\mathbb{K}G, \underset{F}{\longrightarrow}\right)$ defined by $v \underset{F}{\longrightarrow} v'$ if and only if there exists $T \in F$ such that v does not belong to $\mathbb{K}\mathrm{Red}(T)$ and v' is equal to $T(v)$.

2.3.4. Remarks.

1. If we have $v \underset{F}{\longrightarrow} v'$, then we have $v <_{\mathrm{mul}} v'$. The order $<_{\mathrm{mul}}$ being well-founded, the relation $\underset{F}{\longrightarrow}$ is also well-founded.
2. We have $v \xrightarrow[F]{*} v'$ if and only if there exists $R \in \langle F \rangle$ such that v' is equal to $R(v)$, that is, if and only if v rewrites into v'. In particular, $v \xrightarrow[F]{*} v'$ implies that $v - v'$ belongs to $\ker(\wedge F)$.

2.3.5. Lemma. *Let $v_1,\ v_2,\ v_3 \in \mathbb{K}G$ such that $v_1 \xleftrightarrow[F]{*} v_3$. Then, we have $v_1 + v_2 \xleftrightarrow[F]{*} v_2 + v_3$.*

Proof. For every $u_1,\ u_2 \in \mathbb{K}G$ and for every $T \in F$, we have $u_1 + u_2 \xrightarrow[F]{*} T(u_1 + u_2)$ as well as $u_1 + T(u_2) \xrightarrow[F]{*} T(u_1 + u_2)$. Hence, we have

$$u_1 + u_2 \xleftrightarrow[F]{*} u_1 + T(u_2). \tag{5}$$

Let $u_3 \in \mathbb{K}G$ such that $u_2 \xrightarrow[F]{*} u_3$, that is, there exists $R \in \langle F \rangle$ such that u_3 is equal to $R(u_2)$. From Relation (5), we have

$$u_1 + u_2 \xleftrightarrow[F]{*} u_1 + u_3. \tag{6}$$

Let $v_1,\ v_2 \in \mathbb{K}G$ such that $v_1 \overset{*}{\underset{F}{\longleftrightarrow}} v_2$, that is, there exists a zig-zag

$$v_1 = u_1 \overset{*}{\underset{F}{\longrightarrow}} u_2 \overset{*}{\underset{F}{\longleftarrow}} u_3 \overset{*}{\underset{F}{\longrightarrow}} \cdots \overset{*}{\underset{F}{\longrightarrow}} u_{r-1} \overset{*}{\underset{F}{\longleftarrow}} u_r = v_2.$$

Relation (6) implies that for every $v_3 \in \mathbb{K}G$ and for every $i \in \{1, \cdots, r-1\}$, we have $u_i + v_3 \overset{*}{\underset{F}{\longleftrightarrow}} u_{i+1} + v_3$. Thus, Lemma 2.3.5 holds.

□

2.3.6. Proposition. *For every $v_1,\ v_2 \in \mathbb{K}G$, we have*

$$v_1 \overset{*}{\underset{F}{\longleftrightarrow}} v_2 \ \textit{if and only if}\ v_1 - v_2 \in \ker(\wedge F).$$

Proof. Assume that $v_1 \overset{*}{\underset{F}{\longleftrightarrow}} v_2$. Hence, there exists a zig-zag

$$v_1 = u_1 \overset{*}{\underset{F}{\longrightarrow}} u_2 \overset{*}{\underset{F}{\longleftarrow}} u_3 \overset{*}{\underset{F}{\longrightarrow}} \cdots \overset{*}{\underset{F}{\longrightarrow}} u_{r-1} \overset{*}{\underset{F}{\longleftarrow}} u_r = v_2.$$

For every $i \in \{1, \cdots, r-1\}$, $u_i - u_{i+1}$ belongs to $\ker(\wedge F)$. Hence,

$$v_1 - v_2 = (u_1 - u_2) + (u_2 - u_3) + \cdots + (u_{r-2} - u_{r-1}) + (u_{r-1} - u_r),$$

belongs to $\ker(\wedge F)$.

Conversely, assume that $v_1 - v_2$ belongs to the kernel of $\wedge F$. The set

$$\{v - T(v),\ T \in F \text{ and } v \in \mathbb{K}G\},$$

is a generating set of $\ker(\wedge F)$. Thus, there exist $T_1, \cdots, T_n \in F$ and $u_1, \cdots, u_n \in \mathbb{K}G$ such that $v_1 - v_2$ is equal to $\sum_{i=1}^{n} u_i - T_i(u_i)$, so that we have

$$v_1 = \sum_{i=1}^{n} u_i - T_i(u_i) + v_2.$$

For every $i \in \{1, \cdots, n\}$, we have $u_i - T_i(u_i) \overset{*}{\underset{F}{\longleftrightarrow}} 0$. Hence, from Lemma 2.3.5, we have $v_1 \overset{*}{\underset{F}{\longleftrightarrow}} v_2$.

□

2.3.7. Remark. Let v be an element of $\mathbb{K}G$ and let $[v]$ be the set of elements v' such that $v' - v$ belongs to $\ker(\wedge F)$. From Proposition 2.3.6, $[v]$ is the equivalence class of v for the relation $\overset{*}{\underset{F}{\longleftrightarrow}}$. From Lemma 2.2.9, $\wedge F(v)$ is the smallest element of this equivalence class.

2.3.8. Proposition. *The set F has the Church-Rosser property if and only if $\underset{F}{\longrightarrow}$ has the Church-Rosser property.*

Proof. Assume that F has the Church-Rosser property. Let $v,\ v' \in \mathbb{K}G$ such that $v \overset{*}{\underset{F}{\longleftrightarrow}} v'$. From Proposition 2.3.6, $v - v'$ belongs to the kernel of $\wedge F$. We denote by u the common value of $\wedge F(v)$ and $\wedge F(v')$. The set F having the Church-Rosser property, v and v' rewrite into u, that is, we have $v \overset{*}{\underset{F}{\longrightarrow}} u$ and $v' \overset{*}{\underset{F}{\longrightarrow}} u$. Hence, $\underset{F}{\longrightarrow}$ has the Church-Rosser property.

Conversely, assume that $\underset{F}{\longrightarrow}$ has the Church-Rosser property. Let $v \in \mathbb{K}G$. From Proposition 2.3.6, we have $v \overset{*}{\underset{F}{\longleftrightarrow}} \wedge F(v)$. The relation $\underset{F}{\longrightarrow}$ having the Church-Rosser property, there exists $u \in \mathbb{K}G$ such that $v \overset{*}{\underset{F}{\longrightarrow}} u$ and $\wedge F(v) \overset{*}{\underset{F}{\longrightarrow}} u$. Moreover, $\wedge F(v)$ belongs to $\mathbb{K}\mathrm{Red}(F)$, so that it is an F-normal form. As a consequence, u is equal to $\wedge F(v)$. Hence, we have $v \overset{*}{\underset{F}{\longrightarrow}} \wedge F(v)$, that is, v rewrites into $\wedge F(v)$. That shows that F has the Church-Rosser property.

□

2.3.9. Corollary. *The set F is confluent if and only if $\underset{F}{\longrightarrow}$ is confluent.*

Proof. From Theorem 2.2.5, F is confluent if and only if it has the Church-Rosser property. Hence, from Proposition 2.3.8, F is confluent if and only if $\underset{F}{\longrightarrow}$ has the Church-Rosser property, that is, if and only if $\underset{F}{\longrightarrow}$ is confluent.

□

2.3.10. Example. We consider the pair P of Example 2.1.14. We have seen that the pair P is not confluent. The following diagram

$$\begin{tikzcd}
 & g_4 \arrow[dl] \arrow[dr] & \\
T_1(g_4) = g_3 & & T_2(g_4) = g_2 \arrow[dl] \\
 & T_1(g_2) = g_1 &
\end{tikzcd}$$

shows that we have $g_4 \underset{P}{\overset{*}{\longrightarrow}} g_1$ and $g_4 \underset{P}{\overset{*}{\longrightarrow}} g_3$. The two elements g_1 and g_3 are normal forms, so that $\underset{P}{\longrightarrow}$ is not confluent.

3 Completion and Presentations by Operator

The aim of this section is to formulate algebraically the completion using the lattice structure introduced in Section 2.1. We also apply the theory of reduction operators to algebras. Before that, we need to investigate the notion of confluence for a pair of reduction operators.

In Section 3.1 and Section 3.2 we fix a well-ordered set $(G, <)$.

3.1 Confluence for a Pair of Reduction Operators

Throughout this section we fix a pair $P = (T_1, T_2)$ of reduction operators relative to $(G, <)$.

3.1.1. The Braided Products. Given two endomorphisms S and T of $\mathbb{K}G$, we denote by $\langle T, S\rangle^n$ the product $\cdots S \circ T \circ S$ with n factors. Let $g \in G$. From Lemma 2.2.6, there exists an integer n such that $\langle T_2, T_1\rangle^n (g)$ and $\langle T_1, T_2\rangle^n (g)$ are F-normal forms. Let n_g be the smallest integer satisfying the previous condition. Let $\langle T_2, T_1\rangle$ and $\langle T_1, T_2\rangle$ be the two endomorphisms of $\mathbb{K}G$ defined by

$$\langle T_2, T_1\rangle (g) = \langle T_2, T_1\rangle^{n_g} (g) \text{ and } \langle T_1, T_2\rangle (g) = \langle T_1, T_2\rangle^{n_g} (g),$$

for every $g \in G$.

3.1.2. Remark. The vector spaces $\operatorname{im}(\langle T_2, T_1\rangle)$ and $\operatorname{im}(\langle T_1, T_2\rangle)$ are included in $\mathbb{K}\operatorname{Red}(T)$. Hence, every element $v \in \mathbb{K}G$ admits at most two P-normal forms: $\langle T_2, T_1\rangle (v)$ and $\langle T_1, T_2\rangle (v)$.

3.1.3. Lemma. *The pair P is confluent if and only if $\langle T_2, T_1\rangle$ and $\langle T_1, T_2\rangle$ are equal. In this case we have*

$$\begin{aligned} \wedge P &= \langle T_2, T_1\rangle \\ &= \langle T_1, T_2\rangle . \end{aligned}$$

Proof. From Proposition 2.2.10, P is confluent if and only if every element of $\mathbb{K}G$ admits a unique P-normal form. Hence, P is confluent if and only if for every $v \in \mathbb{K}G$, $\langle T_2, T_1\rangle (v)$ and $\langle T_1, T_2\rangle (v)$ are equal. That shows the first part of the proposition. The second part is a consequence of Lemma 2.2.9.

□

3.1.4. Dual Braided Products. Let n be an integer. We show by induction on n that we have

$$\begin{aligned}\langle \mathrm{Id}_{\mathbb{K}G} - T_2, \mathrm{Id}_{\mathbb{K}G} - T_1\rangle^n &= \mathrm{Id}_{\mathbb{K}G} + \sum_{i=1}^{n-1}(-1)^i \left(\langle T_1, T_2\rangle^i + \langle T_2, T_1\rangle^i\right) + (-1)^n \langle T_2, T_1\rangle^n ,\\ \langle \mathrm{Id}_{\mathbb{K}G} - T_1, \mathrm{Id}_{\mathbb{K}G} - T_2\rangle^n &= \mathrm{Id}_{\mathbb{K}G} + \sum_{i=1}^{n-1}(-1)^i \left(\langle T_1, T_2\rangle^i + \langle T_2, T_1\rangle^i\right) + (-1)^n \langle T_1, T_2\rangle^n .\end{aligned} \tag{7}$$

We consider the two operators $\langle \mathrm{Id}_{\mathbb{K}G} - T_1, \mathrm{Id}_{\mathbb{K}G} - T_2\rangle$ and $\langle \mathrm{Id}_{\mathbb{K}G} - T_2, \mathrm{Id}_{\mathbb{K}G} - T_1\rangle$ defined by

$$\begin{aligned}\langle \mathrm{Id}_{\mathbb{K}G} - T_2, \mathrm{Id}_{\mathbb{K}G} - T_1\rangle (g) &= \langle \mathrm{Id}_{\mathbb{K}G} - T_2, \mathrm{Id}_{\mathbb{K}G} - T_1\rangle^{n_g} (g),\\ \langle \mathrm{Id}_{\mathbb{K}G} - T_1, \mathrm{Id}_{\mathbb{K}G} - T_2\rangle (g) &= \langle \mathrm{Id}_{\mathbb{K}G} - T_1, \mathrm{Id}_{\mathbb{K}G} - T_2\rangle^{n_g} (g),\end{aligned}$$

for every $g \in G$.

3.1.5. Remark. We deduce from Lemma 3.1.3 that if the pair P is confluent, then $\langle \mathrm{Id}_{\mathbb{K}G} - T_2, \mathrm{Id}_{\mathbb{K}G} - T_1\rangle$ and $\langle \mathrm{Id}_{\mathbb{K}G} - T_1, \mathrm{Id}_{\mathbb{K}G} - T_2\rangle$ are equal. In the sequel, when P is assumed to be confluent, the common value of $\langle \mathrm{Id}_{\mathbb{K}G} - T_2, \mathrm{Id}_{\mathbb{K}G} - T_1\rangle$ and $\langle \mathrm{Id}_{\mathbb{K}G} - T_1, \mathrm{Id}_{\mathbb{K}G} - T_2\rangle$ is denoted by T.

3.1.6. Lemma. *Assume that P is confluent. Then, $\mathrm{Id}_{\mathbb{K}G} - T$ is a reduction operator relative to $(G, <)$. Moreover, we have*

$$\mathrm{Nred}\,(\mathrm{Id}_{\mathbb{K}G} - T) = \mathrm{Nred}\,(T_1) \cap \mathrm{Nred}\,(T_2)\,.$$

Proof. First, we show that $\mathrm{Id}_{\mathbb{K}G} - T$ is a projector. The operators $\mathrm{Id}_{\mathbb{K}G} - T_1$ and $\mathrm{Id}_{\mathbb{K}G} - T_2$ are projectors. Hence, by definition of T, for every $g \in G$, and for $i = 1$ or 2, we have

$$(\mathrm{Id}_{\mathbb{K}G} - T_i) \circ T(g) = T(g).$$

Hence, T is a projector, so that $\mathrm{Id}_{\mathbb{K}G} - T$ is also a projector.

Let $g \in G$. Let us show that $g - T(g)$ is either equal to g or smaller than g. From Relation (7) of 3.1.4, we have

$$g - T(g) = \sum_{i=1}^{n_g - 1} (-1)^{i+1} \left(\langle T_1, T_2\rangle^i + \langle T_2, T_1\rangle^i\right)(g) + (-1)^{n_g+1} \langle T_2, T_1\rangle^{n_g} (g).$$

Thus, if g belongs to $\mathrm{Nred}\,(T_1) \cap \mathrm{Nred}\,(T_2)$, then $g - T(g)$ is strictly smaller than g. Assume that g does not belong to $\mathrm{Nred}\,(T_1) \cap \mathrm{Nred}\,(T_2)$. Assume that g belongs to $\mathrm{Red}\,(T_1)$ (the case where g belongs to $\mathrm{Red}\,(T_2)$ is analogous). We have

$$\begin{aligned} g - T(g) &= g + T_2(g) + \sum_{i=2}^{n_g - 1} (-1)^{i+1} \left(\langle T_1, T_2\rangle^i + \langle T_1, T_2\rangle^{i-1}\right)(g) + (-1)^{n_g+1} \langle T_1, T_2\rangle^{n_g - 1} (g)\\ &= g.\end{aligned}$$

Hence, $\mathrm{Id}_{\mathbb{K}G} - T$ is a reduction operator relative to $(G, <)$ and the set $\mathrm{Nred}\,(\mathrm{Id}_{\mathbb{K}G} - T)$ is equal to $\mathrm{Nred}\,(T_1) \cap \mathrm{Nred}\,(T_2)$.

□

3.1.7. Lemma. *Assume that P is confluent. Then, $T_1 \vee T_2$ is equal to $\mathrm{Id}_{\mathbb{K}G} - T$.*

Proof. The operator $\mathrm{Id}_{\mathbb{K}G} - T$ being a reduction operator relative to $(G, <)$ and θ being a bijection, it is sufficient to show that the kernel of $\mathrm{Id}_{\mathbb{K}G} - T$ equals the one of $T_1 \vee T_2$. From Relation (7) of 3.1.4, for every $v \in \mathbb{K}G$, we have

$$\begin{aligned} v - T(v) &= \sum_{i=1}^{n-1} (-1)^{i+1} \left(\langle T_1, T_2 \rangle^i + \langle T_2, T_1 \rangle^i \right)(v) + (-1)^{n+1} \langle T_2, T_1 \rangle^n (v) \\ &= \sum_{i=1}^{n-1} (-1)^{i+1} \left(\langle T_1, T_2 \rangle^i + \langle T_2, T_1 \rangle^i \right)(v) + (-1)^{n+1} \langle T_1, T_2 \rangle^n (v), \end{aligned}$$

where n is an integer greater than n_g for every $g \in G$ belonging to the support of v. Hence, $\ker(T_1 \vee T_2) = \ker(T_1) \cap \ker(T_2)$ is included in $\ker(\mathrm{Id}_{\mathbb{K}G} - T)$. Moreover, the operator $\mathrm{Id}_{\mathbb{K}G} - T$ being a projector, its kernel is equal to $\mathrm{im}(T)$, that is, we have

$$\begin{aligned} \ker(\mathrm{Id}_{\mathbb{K}G} - T) &= \mathrm{im}(\langle \mathrm{Id}_{\mathbb{K}G} - T_2, \mathrm{Id}_{\mathbb{K}G} - T_1 \rangle) \\ &= \mathrm{im}(\langle \mathrm{Id}_{\mathbb{K}G} - T_1, \mathrm{Id}_{\mathbb{K}G} - T_2 \rangle). \end{aligned}$$

The latter is included in $\ker(T_1)$ and $\ker(T_2)$, so that it is included in $\ker(T_1) \cap \ker(T_2) = \ker(T_1 \vee T_2)$. □

3.1.8. Lemma. *Assume that P is confluent. Then,* $\mathrm{Nred}(T_1 \vee T_2)$ *is equal to* $\mathrm{Nred}(T_1) \cap \mathrm{Nred}(T_2)$.

Proof. This is a consequence of Lemma 3.1.6 and Lemma 3.1.7. □

3.2 Completion

We fix a subset F of $\mathbf{RO}(G, <)$.

3.2.1. Definitions.

1. A *completion of F* is a subset F' of $\mathbf{RO}(G, <)$ such that
   (a) F' is confluent,
   (b) $F \subset F'$ and $\wedge F' = \wedge F$.
2. A *complement of F* is an element C of $\mathbf{RO}(G, <)$ such that
   (a) $(\wedge F) \wedge C = \wedge F$,
   (b) $\mathrm{Obs}^F \subset \mathrm{Nred}(C)$.

   A complement is said to be *minimal* if the inclusion (2b) is an equality.

3.2.2. Proposition. *Let $C \in \mathbf{RO}(G, <)$ such that $(\wedge F) \wedge C$ is equal to $\wedge F$. The set $F \cup \{C\}$ is a completion of F if and only if C is a complement of F.*

Proof. We let $F' = F \cup \{C\}$. The set F' contains F and is such that $\wedge F' = \wedge F$ by hypothesis. Hence, F' is a completion of F if and only if it is confluent, that is, if and only if $\mathrm{Red}(F')$ is equal to $\mathrm{Red}(\wedge F')$. The set $\mathrm{Red}(F')$ is equal to $\mathrm{Red}(F) \cap \mathrm{Red}(C)$ and $\wedge F'$ is equal to $\wedge F$. Hence, F' is confluent if and only if we have the following relation

$$\mathrm{Red}(F) \cap \mathrm{Red}(C) = \mathrm{Red}(\wedge F).$$

Moreover, $\mathrm{Red}(F)$ is the disjoint union of $\mathrm{Red}(\wedge F)$ and Obs^F. Hence, we have

$$\mathrm{Red}(F) \cap \mathrm{Red}(C) = \Big(\mathrm{Red}(\wedge F) \cap \mathrm{Red}(C) \Big) \bigsqcup \Big(\mathrm{Obs}^F \cap \mathrm{Red}(C) \Big). \tag{8}$$

The hypothesis $(\wedge F) \wedge C$ is equal to $\wedge F$ means that $\wedge F$ is smaller than C. Thus, from Lemma 2.1.15, $\mathrm{Red}\,(\wedge F)$ is included in $\mathrm{Red}\,(C)$. From Relation (8), we have

$$\mathrm{Red}\,(F) \cap \mathrm{Red}\,(C) = \mathrm{Red}\,(\wedge F) \bigsqcup \left(\mathrm{Obs}^F \cap \mathrm{Red}\,(C)\right).$$

Hence, F' is confluent if and only if $\mathrm{Obs}^F \cap \mathrm{Red}\,(C)$ is empty, that is, if and only if C is a complement of F.

□

3.2.3. Examples.

1. The operator $\wedge F$ is a complement of F. In general, this complement is not minimal (see Point 2).
2. We consider the pair P of Example 2.1.14. Let

$$C_1 = \begin{pmatrix} 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \text{ and } C_2 = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}.$$

The sets $\mathrm{Nred}\,(C_1)$ and $\mathrm{Nred}\,(C_2)$ are equal to $\{g_3\}$. The latter is equal to Obs^P (see Point 1 of Example 2.1.18). Moreover, $\ker(C_1)$ and $\ker(C_2)$ are the vector spaces spanned by $g_3 - g_1$ and $g_3 - g_2$, respectively. These two vector spaces are included in $\ker(T_1 \wedge T_2)$. Hence, C_1 and C_2 are greater than $T_1 \wedge T_2$, that is, we have

$$(\wedge P) \wedge C_1 = \wedge P \text{ and } (\wedge P) \wedge C_2 = \wedge P.$$

We conclude that C_1 and C_2 are two minimal complements of P. We also recall that $\mathrm{Nred}\,(T_1 \wedge T_2)$ is equal to $\{g_2, g_3, g_4\}$ (see Example 2.1.14), so that $T_1 \wedge T_2$ is not a minimal complement of P.

3.2.4. The F-Complement. The *F-complement* is the operator

$$C^F = (\wedge F) \vee \left(\vee \overline{F}\right),$$

where $\vee \overline{F}$ is equal to $\theta\,(\mathbb{K}\mathrm{Red}\,(F))$.

3.2.5. Lemma. *The pair $P = \left(\wedge F, \vee \overline{F}\right)$ is confluent.*

Proof. The image of $\wedge F$ is included in $\mathbb{K}\mathrm{Red}\,(F)$ which is equal to the kernel of $\vee \overline{F}$. Thus, $\vee \overline{F} \circ \wedge F$ and $\vee \overline{F} \circ \wedge F \circ \vee \overline{F}$ are equal to the zero operator. Hence, we have the equality

$$\vee \overline{F} \circ \wedge F \circ \vee \overline{F} = \vee \overline{F} \circ \wedge F. \tag{9}$$

Hence, from Lemma 3.1.3, the pair P is confluent.

□

3.2.6. Theorem. *Let F be a subset of $\mathbf{RO}\,(G, <)$. The F-complement is a minimal complement of F.*

Proof. By definition, C^F is greater than $\wedge F$, that is, C^F satisfies (2a) of 3.2.1. Let us show that Obs^F is equal to $\mathrm{Nred}\left(C^F\right)$. From Lemma 3.2.5, the pair $\left(\wedge F, \vee \overline{F}\right)$ is confluent. Hence, from Lemma 3.1.8, we have

$$\begin{aligned} \mathrm{Nred}\left(C^F\right) &= \mathrm{Nred}\left((\wedge F) \vee \left(\vee \overline{F}\right)\right) \\ &= \mathrm{Nred}\,(\wedge F) \cap \mathrm{Nred}\left(\vee \overline{F}\right) \\ &= \mathrm{Nred}\,(\wedge F) \cap \mathrm{Red}\,(F) \\ &= \mathrm{Obs}^F. \end{aligned}$$

□

We end this section with a characterisation of the F-complement. For that, we need the following lemma:

3.2.7. Lemma. *The set* $C^F\left(\mathrm{Obs}^F\right)$ *is included in* $\mathbb{K}\mathrm{Red}\,(\wedge F)$.

Proof. We have seen in Relation (9) (see the proof of Lemma 3.2.5) that we have

$$\vee\overline{F}\circ\wedge F\circ\vee\overline{F}=\vee\overline{F}\circ\wedge F.$$

From Lemma 3.2.5, the pair $P = \left(\wedge F, \vee\overline{F}\right)$ is confluent. Hence, from Lemma 3.1.7, we have

$$C^F=\mathrm{Id}_{\mathbb{K}G}-\left(\mathrm{Id}_{\mathbb{K}G}-\vee\overline{F}\right)\circ\left(\mathrm{Id}_{\mathbb{K}G}-\wedge F\right).$$

The sets Obs^F and $\mathbb{K}\mathrm{Red}\,(\wedge F)$ are included in $\mathbb{K}\mathrm{Red}\,(F)$ which is equal to the kernel of $\vee\overline{F}$. Hence, for every $g\in\mathrm{Obs}^F$, we have

$$\begin{aligned}C^F(g)&=g-\left(\mathrm{Id}_{\mathbb{K}G}-\vee\overline{F}\right)\circ\left(\mathrm{Id}_{\mathbb{K}G}-\wedge F\right)(g)\\&=\wedge F(g)+\vee\overline{F}(g)-\vee\overline{F}\circ\wedge F(g)\\&=\wedge F(g).\end{aligned}$$

That shows that $C^F\left(\mathrm{Obs}^F\right)$ is included in $\mathbb{K}\mathrm{Red}\,(\wedge F)$.

□

3.2.8. Proposition. *The F-complement is the unique minimal complement of F such that* $C^F\left(\mathrm{Obs}^F\right)$ *is included in* $\mathbb{K}\mathrm{Red}\,(F)$.

Proof. Let C be a minimal complement of F such that $C\left(\mathrm{Obs}^F\right)$ is included in $\mathbb{K}\mathrm{Red}\,(F)$. For every $g\in G\setminus\mathrm{Obs}^F$, $C^F(g)$ and $C(g)$ are equal to g. Thus, it is sufficient to show that for every $g\in\mathrm{Obs}^F$, $C(g)$ is equal to $C^F(g)$. The set $C\left(\mathrm{Obs}^F\right)$ being included in $\mathbb{K}\mathrm{Red}\,(F)$ and $\mathrm{Nred}\,(C)$ being equal to Obs^F, $C\left(\mathrm{Obs}^F\right)$ is in fact included in $\mathbb{K}\left(\mathrm{Red}\,(F)\setminus\mathrm{Obs}^F\right)$, that is, it is included in $\mathbb{K}\mathrm{Red}\,(\wedge F)$. From Lemma 3.2.7, $C^F\left(\mathrm{Obs}^F\right)$ is also included in $\mathbb{K}\mathrm{Red}\,(\wedge F)$. Hence, for every $g\in\mathrm{Obs}^F$, we have

$$\wedge F\circ C^F(g)=C^F(g)\text{ and }\wedge F\circ C(g)=C(g). \tag{10}$$

Relation (2a) of 3.2.1 implies that C^F and C are greater than $\wedge F$. Thus, the equivalence (1) (see 2.1.13) implies that $\wedge F\circ C^F$ and $\wedge F\circ C$ are equal to $\wedge F$. Hence, from Relation (10), $C^F(g)$ and $C(g)$ are equal to $\wedge F(g)$ for every $g\in\mathrm{Obs}^F$, so that Proposition 3.2.8 holds.

□

3.2.9. Examples.

1. The operator C_1 of Example 3.2.3, Point 2 is the P-complement.
2. Consider Point 2 of Example 2.1.18. The P-complement maps yxy to xx and fixes all other word.

3.3 Presentations by Operator

3.3.1. Algebras. An associative unitary $\mathbb{K}$-algebra is a $\mathbb{K}$-vector space $\mathbf{A}$ equipped with a $\mathbb{K}$-linear map, called *multiplication*, $\mu : \mathbf{A} \otimes \mathbf{A} \longrightarrow \mathbf{A}$ which is associative and for which there exists a unit $1_{\mathbf{A}}$. We say algebra instead of unitary associative $\mathbb{K}$-algebra. Given a set X, let X^* be the set of words written with X. This set admits a monoid structure, where the multiplication is given by concatenation of words and the unit is the empty word. Moreover, the free algebra over X is the vector space $\mathbb{K}X^*$ spanned by X^* equipped with the multiplication induced by the one of the monoid X^*.

From now on, we fix an algebra $\mathbf{A}$.

3.3.2. Monomial Orders. Let X be a set. A *monomial order on* X^* is a well-founded total order $<$ on X^* such that the following conditions are fulfilled:

1. $1 < w$ for every word w different from 1,
2. for every $w_1,\ w_2,\ w,\ w' \in X^*$ such that $w < w'$, we have $w_1 w w_2 < w_1 w' w_2$.

In particular, $(X^*, <)$ is a well-ordered set. In the sequel, given an element $f \in \mathbb{K}X^*$, we write $\mathrm{lm}\,(f)$ (for leading monomial) instead of $\mathrm{lg}\,(f)$.

3.3.3. Gröbner Bases. Let X be a set and let $<$ be a monomial order on X^*. Given a subset E of $\mathbb{K}X^*$, we let $\mathrm{lm}\,(E) = \{\mathrm{lm}\,(f)\,,\ f \in E\}$. Let I be a two-sided ideal of $\mathbb{K}X^*$. A subset R of I is called a *Gröbner basis of* I if the semi-group ideal spanned by $\mathrm{lm}\,(R)$ is equal to $\mathrm{lm}\,(I)$. In other words, R is a Gröbner basis of I if and only if for every $w \in \mathrm{lm}\,(I)$, there exist $w' \in \mathrm{lm}\,(R)$ and $w_1,\ w_2 \in X^*$ such that w is equal to $w_1 w' w_2$.

3.3.4. Definition. A *presentation by operator* of $\mathbf{A}$ is a triple $\langle X, <|\ S\rangle$, where

1. X is a set and $<$ is a monomial order on X^*,
2. S is a reduction operator relative to $(X^*, <)$,
3. we have an isomorphism of algebras

$$\mathbf{A} \simeq \frac{\mathbb{K}X^*}{I\,(\ker\,(S))},$$

where $I\,(\ker\,(S))$ is the two-sided ideal of $\mathbb{K}X^*$ spanned by $\ker\,(S)$.

3.3.5. Reduction Family of a Presentation. Let X be a set and let n be an integer. We denote by $X^{(n)}$ and $X^{(\leq n)}$ the set of words of length n and of length smaller than n, respectively. Let $\langle X, <|\ S\rangle$ be a presentation by operator of $\mathbf{A}$. For every integers n and m such that (n, m) is different from $(0, 0)$, we let

$$T_{n,m} = \mathrm{Id}_{\mathbb{K}X^{(\leq n+m-1)}} \ \oplus\ \mathrm{Id}_{\mathbb{K}X^{(n)}} \otimes S \otimes \mathrm{Id}_{\mathbb{K}X^{(m)}}.$$

Explicitly, given $w \in X^*$, $T_{n,m}(w)$ is equal to w if the length of w is strictly smaller than $n + m$. If the length of w is greater than $n + m$, we let $w = w_1 w_2 w_3$, where w_1 and w_3 have length n and m, respectively. Then, $T_{n,m}(w)$ is equal to $w_1 S(w_2) w_3$. We also let $T_{0,0} = S$. The *reduction family* of $\langle X, <|\ S\rangle$ is the set $\{T_{n,m},\ 0 \leq n,\ m\}$.

3.3.6. Lemma. *Let $\langle X, <| S\rangle$ be a presentation by operator of* $\mathbf{A}$. *Let n and m be two integers. Then, $T_{n,m}$ is a reduction operator relative to $(X^*, <)$ and its kernel is equal to $\mathbb{K}X^{(n)} \otimes \ker(S) \otimes \mathbb{K}X^{(m)}$.*

Proof. First, we show that $T_{n,m}$ is a reduction operator relative to $(X^*, <)$. The operator S being a projector, $T_{n,m}$ is also a projector. Let $w \in X^*$. If the length of w is strictly smaller than $n+m$, then $T_{n,m}(w)$ is equal to w. If the length of w is greater than $n+m$, we write $w = w_1w_2w_3$, where w_1 and w_3 have length n and m, respectively. If w_2 belongs to $\mathrm{Red}(S)$, then $T_{n,m}(w)$ is equal to w. In the other case, let

$$S(w_2) = \sum \lambda_i w_i,$$

be the decomposition of $S(w_2)$ with respect to the basis X^*. We have

$$T_{n,m}(w) = \sum \lambda_i w_1 w_i w_3.$$

For every $i \in I$, w_i is strictly smaller than w_2. The order $<$ being a monomial order, $w_1w_iw_3$ is strictly smaller than w. Hence, $T_{n,m}$ is a reduction operator relative to $(X^*, <)$.

Let us show the second part of the lemma. Given an integer k, we denote by $X^{(\geq k)}$ the set of words of length greater than k. For every $f \in \mathbb{K}X^*$, we write $f = f_1 + f_2$, where f_1 and f_2 are the images of f by the natural projections of $\mathbb{K}X^*$ on $\mathbb{K}X^{(\leq n+m-1)}$ and $\mathbb{K}X^{(\geq n+m)}$, respectively. These two vector spaces are stabilised by $T_{n,m}$ and $T_{n,m}(f_1)$ is equal to f_1. Thus, f belongs to $\ker(T_{n,m})$ if and only if f_1 is equal to 0 and f_2 belongs to $\ker(T_{n,m})$. Moreover, f_2 admits a unique decomposition with shape

$$f_2 = \sum_{i \in I} w_i f_i w'_i,$$

where for every $i \in I$

1. w_i and w'_i are words of length n and m, respectively,
2. for every $j \in I$ such that j is different from i, the pair (w_i, w'_i) is different from (w_j, w'_j),
3. f_i is a non zero element of $\mathbb{K}X^*$.

We have

$$T_{n,m}(f_2) = \sum_{i \in I} w_i S(f_i) w'_i.$$

Thus, f_2 belongs to $\ker(T_{n,m})$ if and only if for every $i \in I$, f_i belongs to $\ker(S)$. Hence, the kernel of $T_{n,m}$ is equal to $\mathbb{K}X^{(n)} \otimes \ker(S) \otimes \mathbb{K}X^{(m)}$.

□

3.3.7. Remark. Let $\langle X, <| S\rangle$ be a presentation by operator of $\mathbf{A}$. From Lemma 3.3.6, its reduction family is a subset of $\mathbf{RO}(X^*, <)$. Moreover, the kernel of $\wedge F$ is the sum of the vector spaces $\mathbb{K}X^{(n)} \otimes \ker(S) \otimes \mathbb{K}X^{(m)}$, that is, $\ker(\wedge F)$ is the two-sided ideal spanned by $\ker(S)$.

3.3.8. Confluent Presentation. A *confluent presentation by operator* of $\mathbf{A}$ is a presentation by operator of $\mathbf{A}$ such that its reduction family is confluent.

3.3.9. Lemma. *Let $\langle X, <| S\rangle$ be a presentation by operator of* $\mathbf{A}$ *and let F be its reduction family. Let R be the reduced basis of* $\ker(S)$. *A word belongs to* $\mathrm{Red}(F)$ *if and only if it does not belong to the semi-group ideal spanned by* $\mathrm{lm}(R)$.

Proof. The set $\mathrm{Red}(F)$ is the set of words w such that every sub-word of w belongs to $\mathrm{Red}(S)$. From Proposition 2.1.11, a word belongs to $\mathrm{Red}(S)$ if and only if it does not belong to $\mathrm{lm}(R)$, so that Lemma 3.3.9 holds.

□

3.3.10. Proposition. *Let $\langle X, <| S\rangle$ be a presentation by operator of* $\mathbf{A}$. *Let R be the reduced basis of* $\ker(S)$. *The presentation $\langle X, <| S\rangle$ is confluent if and only if R is a Gröbner basis of $I(R)$.*

Proof. Let F be the reduction family of $\langle X, <| S\rangle$.

Assume that $\langle X, <| S\rangle$ is not confluent. Let $w \in \mathrm{Obs}^F$. The element $w - \wedge F(w)$ belongs to $I(\ker(S))$. The latter is equal to $I(R)$. Moreover, the leading monomial of $w - \wedge F(w)$ is equal to w, so that it belongs to $\mathrm{Red}(F)$. Hence, from Lemma 3.3.9, w does not belong to the semi-group ideal spanned by $\mathrm{lm}(R)$. Thus, R is not a Gröbner basis of $I(R)$.

Assume that $\langle X, <| S\rangle$ is confluent. Let $f \in I(R)$. We assume that $\mathrm{lc}(f)$ is equal to 1. The kernel of $\wedge F$ being equal to $I(R)$, $\wedge F(f)$ is equal to 0. Hence, $\wedge F(\mathrm{lm}(f))$ is equal to $\wedge F(\mathrm{lm}(f) - f)$. The leading monomial of $\mathrm{lm}(f) - f$ is strictly smaller than $\mathrm{lm}(f)$. In particular, $\mathrm{lm}(f)$ is not $\wedge F$-reduced. The set F being confluent, that implies that $\mathrm{lm}(f)$ does not belong to $\mathrm{Red}(F)$. From Lemma 3.3.9, $\mathrm{lm}(f)$ belongs to the semi-group ideal spanned by $\mathrm{lm}(R)$. Thus, R is a Gröbner basis of $I(R)$.

□

3.3.11. Theorem. *Let $\langle X, <| S\rangle$ be a presentation by operator of* $\mathbf{A}$ *and let C be a complement of its reduction family. The triple $\langle X, <| S \wedge C\rangle$ is a confluent presentation of* $\mathbf{A}$.

Proof. We denote by F the reduction family of $\langle X, <| S\rangle$.

First, we show that $\langle X, <| S \wedge C\rangle$ is a presentation of $\mathbf{A}$. For that, we need to show that $I(\ker(S \wedge C))$ is equal to $I(\ker(S))$. The vector space $\ker(S)$ being included in $\ker(S \wedge C)$, $I(\ker(S))$ is included in $I(\ker(S \wedge C))$. Moreover, by definition of a complement, $\wedge F$ is smaller than C, that is, the kernel of C is included in the one of $\wedge F$. The latter is equal to $I(\ker(S))$. Thus, $\ker(S \wedge C)$ is also included in $I(\ker(S))$, so that $I(\ker(S \wedge C))$ is included in $I(\ker(S))$.

Let us show that this presentation is confluent. Writing $\tilde{S} = S \wedge C$, we let $\tilde{T}_{0,0} = \tilde{S}$, and for every integers n and m such that $n + m$ is greater than 1, we let

$$\tilde{T}_{n,m} = \mathrm{Id}_{\mathbb{K}X^{(\leq n+m-1)}} \ \oplus \ \mathrm{Id}_{\mathbb{K}X^{(n)}} \otimes \tilde{S} \otimes \mathrm{Id}_{\mathbb{K}X^{(m)}}.$$

Thus, $\tilde{F} = \left\{\tilde{T}_{n,m},\ 0 \leq n,\ m\right\}$ is the reduction family of $\langle X, <| S \wedge C\rangle$. The kernel of $\wedge\tilde{F}$ is equal to $I(\ker(S \wedge C))$. We have seen that the latter is equal to $I(\ker(S))$ which is equal to $\ker(\wedge F)$. Hence, $\wedge\tilde{F}$ is equal to $\wedge F$. As a consequence, we have to show that $\mathrm{Red}\left(\tilde{F}\right)$ is included in $\mathrm{Red}(\wedge F)$. For every integers n and m, $T_{n,m}$ is greater than $\tilde{T}_{n,m}$. Hence, from Lemma 2.1.15, $\mathrm{Red}\left(\tilde{T}_{n,m}\right)$ is included in $\mathrm{Red}(T_{n,m})$, so that $\mathrm{Red}\left(\tilde{F}\right)$ is included in $\mathrm{Red}(F)$. Moreover, $\mathrm{Red}\left(\tilde{F}\right)$ is also included in $\mathrm{Red}(S \wedge C)$ which is itself included in $\mathrm{Red}(C)$. Hence, $\mathrm{Red}\left(\tilde{F}\right)$ is included in $\mathrm{Red}(F) \cap \mathrm{Red}(C)$. From Proposition 3.2.2, the set $F \cup \{C\}$ is confluent, so that $\mathrm{Red}(F) \cap \mathrm{Red}(C)$ is equal to $\mathrm{Red}(\wedge F)$. That shows that the presentation $\langle X, <| S \wedge C\rangle$ is confluent.

□

4 Generalised Reduction Operators

So far, we have been studying reduction operators relative to a well-ordered set. In this section, we investigate the more general case where we do not consider a total order. We fix an ordered set $(G, <)$.

4.1 Algebraic Structure

The general definition of reduction operator is stated as follows:

4.1.1. Definition. A *reduction operator relative to* $(G, <)$ is and idempotent endomorphism T of $\mathbb{K}G$ such that for every $g \in G$, we have either $T(g) = g$, or for every g' occurring in the support of $T(g)$, we have $g' < g$. As in the case of well-ordered sets, the set of reduction operators relative to $(G, <)$ is denoted by $\mathbf{RO}\,(G, <)$. Given a reduction operator T, the set of T-reduced generators is also denoted by $\mathrm{Red}\,(T)$ and its complement in G is denoted by $\mathrm{Nred}\,(T)$.

4.1.2. From Projectors to Reduction Operators. As an application, we want to consider a set G together with a subset F of the set of linear idempotent endomorphisms of $\mathbb{K}G$. We want to equip G with an order $<$ making F a subset of $\mathbf{RO}\,(G, <)$. For that, consider the binary relation $\lesssim_F$ on G defined by $g' \lesssim_F g$ if there exists $T \in F$ such that $T(g)$ is different from g and such that g' belongs to the support of $T(g)$. The reflexive transitive closure of $\lesssim_F$ is denoted by $<_F$. This relation is not necessarily anti-symmetric. Indeed, let $G = \{g_1, g_2\}$ and consider $F = (T_1, T_2)$, where T_1 and T_2 are defined by $T_1(g_2) = g_1$, $T_1(g_1) = g_1$, $T_2(g_1) = g_2$ and $T_2(g_2) = g_2$, respectively. Then, we have $g_1 <_F g_2$ and $g_2 <_F g_1$. However, if $<_F$ is well-founded, then it is an order relation, and F is a subset of $\mathbf{RO}\,(G, <_F)$.

4.1.3. Absence of Reduced Basis. We would like to equip the set $\mathbf{RO}\,(G, <)$ with a lattice structure. We cannot use the argument of Section 2.1 because a subspace of $\mathbb{K}G$ does not necessarily admits a reduced basis. Indeed, consider $G = \{g_1, g_2, g_3\}$ ordered such: $g_1 < g_3$ and $g_2 < g_3$. The subspace of $\mathbb{K}G$ spanned by $g_3 - g_1$ and $g_3 - g_2$ does not admit any reduced basis.

In order to equip $\mathbf{RO}\,(G, <)$ with an order relation, we need the following lemma:

4.1.4. Lemma. *Let T_1 and T_2 be two reduction operators relative to $(G, <)$ such that $\ker(T_1)$ is included in $\ker(T_2)$. Then, $\mathrm{Red}\,(T_2)$ is included in $\mathrm{Red}\,(T_1)$.*

Proof. Assume by way of contradiction that there exists $g \in \mathrm{Red}\,(T_2)$ not belonging to $\mathrm{Red}\,(T_1)$. The element $g - T_1(g)$ belongs to the kernel of T_1, so that it belongs to the one of T_2. Hence, $T_2(g)$ is equal to $T_2\,(T_1(g))$. The generator g belongs to $\mathrm{Red}\,(T_2)$, so that $T_2(g)$ is equal to g. Moreover, g being not T_1-reduced, every generator appearing in the support of $T_1(g)$ is strictly smaller than g, so that every generator belonging to the support of $T_2\,(T_1(g))$ is also strictly smaller than g. Thus, we reach a contradiction.

□

4.1.5. Order Relation. The binary relation defined by $T_1 \preceq T_2$ if $\ker(T_2) \subset \ker(T_1)$ is clearly reflexive and transitive. Moreover, from Lemma 4.1.4, if two reduction operators have the same kernel, then they have the same image, so that they are equal. Hence, $\preceq$ is anti-symmetric, so that it is an order relation on $\mathbf{RO}\,(G, <)$.

4.1.6. Absence of a Lattice Structure. The order introduced in 4.1.5 does not induce a lattice structure. Consider $G = \{g_1, g_2, g_3, g_4, g_5\}$ ordered such: $g_1 < g_3$, $g_1 < g_4$, $g_2 < g_3$, $g_2 < g_4$, $g_3 < g_5$ and $g_4 < g_5$. Let T_1, T_2 be the two reduction operators defined by $\mathrm{Red}\,(T_i) = \{g_1, g_2, g_3, g_4\}$ for $i = 1$ or 2, $T_1(g_5) = g_3$ and $T_2(g_5) = g_4$. Consider the two reductions operators U_1 and U_2 defined by $\mathrm{Red}\,(U_i) = \{g_1, g_2\}$ for $i = 1$ or 2, $U_1(g_j) = g_2$ and $U_2(g_j) = g_1$ for $j \in \{3, 4, 5\}$. The vector space $\ker(T_1) + \ker(T_2) = \mathbb{K}\{g_5 - g_4\} \oplus \mathbb{K}\{g_5 - g_3\}$ is included in $\ker(U_i)$ for $i = 1$ or 2, that is, U_i is smaller than T_1 and T_2. Moreover, there does not exist a reduction operator with kernel $\mathbb{K}\{g_5 - g_4\} \oplus \mathbb{K}\{g_5 - g_3\}$, so that U_1 and U_2 are two maximal elements smaller than T_1 and T_2. Hence, T_1 and T_2 do not admit a lower bound. Moreover, even when the lower bound exists, its kernel is not necessarily the sum of the kernels. Consider the example from 4.1.3: $G = \{g_1, g_2, g_3\}$ with $g_1 < g_3$ and $g_2 < g_3$, and let $P = (T_1, T_2)$ where, for $i = 1$ or 2, $\mathrm{Nred}\,(T_i) = \{g_3\}$ and $T_i(g_3) = g_i$. Then, we check that T_1 and T_2 admit a lower bound which is the zero operator, that is, the kernel of this lower bound is equal to $\mathbb{K}G$.

4.2 Rewriting Properties

In this section, we investigate the rewriting properties associated to generalised reduction operators. We have seen in the previous section that, given a subset F or $\mathbf{RO}(G,<)$, there does not not necessarily exist a reduction operator with kernel $\sum_{T\in F}\ker(T)$. Hence, in order to define the notion of confluence as it was done in 2.1.17, we have to consider subsets of $\mathbf{RO}(G,<)$ for which such a reduction operator exists. For that, we introduce the following definition:

4.2.1. Completable Sets. A subset F of $\mathbf{RO}(G,<)$ for which there exists a reduction operator with kernel $\sum_{T\in F}\ker(T)$ is said to be *completable.*

4.2.2. Confluence and Church-Rosser Property. Let F be a completable set. The reduction operator whose kernel is equal to $\sum_{T\in F}\ker(T)$ is denoted by $\wedge F$. From Lemma 4.1.4, $\mathrm{Red}(\wedge F)$ is included in $\mathrm{Red}(T)$, so that the set Obs^F is well-defined. The set F is said to be *confluent* if Obs^F is the empty set. We say that v *rewrites into* v' as it was done in Section 2.2 and that F has the *Church-Rosser property* if for every $v\in\mathbb{K}G$, v rewrites into $\wedge F(v)$. Finally, the binary relation $\underset{F}{\longrightarrow}$ on $\mathbb{K}G$ is defined as it was done in Section 2.3.

4.2.3. Normalising Relations. In Theorem 4.2.4, we use the notion of *normalising relation.* Let us recall it. Given an abstract rewriting system $\left(A,\longrightarrow\right)$, the relation $\longrightarrow$ is said to be normalising if every element of A admits at least one normal form.

4.2.4. Theorem. *Let F be a completable subset of $\mathbf{RO}(G,<)$. The following assertions are equivalent:*

1. *F is confluent and $\underset{F}{\longrightarrow}$ is normalising,*
2. *F has the Church-Rosser property,*
3. *$\underset{F}{\longrightarrow}$ is confluent.*

Proof. The proof of the equivalence between 2 and 3 is the same as in Proposition 2.3.8 (indeed, we check that in Section 2.3 we only require the existence of the operator $\wedge F$). As in Theorem 2.2.5, we show that 2 implies that F is confluent. Moreover, if 2 holds, every element v of $\mathbb{K}G$ rewrites into $\wedge F(v)$, that is, we have $v\overset{*}{\underset{F}{\longrightarrow}}\wedge F(v)$. The latter belongs to $\mathbb{K}\mathrm{Red}(F)$, so that it is a normal form for $\underset{F}{\longrightarrow}$. Hence, $\underset{F}{\longrightarrow}$ is normalising. Thus, 2 implies 1. Assume 1 and let us show 3. Let $v_1,\ v_2,\ v_3\in\mathbb{K}G$ such that $v_1\overset{*}{\underset{F}{\longrightarrow}}v_2$ and $v_1\overset{*}{\underset{F}{\longrightarrow}}v_3$. The relation $\underset{F}{\longrightarrow}$ being normalising, v_2 and v_3 admit normal forms. Let $\widehat{v_2}$ and $\widehat{v_3}$ be normal forms of v_2 and v_3, respectively. We have $\widehat{v_2}\overset{*}{\underset{F}{\longleftrightarrow}}\widehat{v_3}$, so that $\widehat{v_2}-\widehat{v_3}$ belongs to the kernel of $\wedge F$. Moreover, $\widehat{v_2}$ and $\widehat{v_3}$ being normal forms, they belong to $\mathbb{K}\mathrm{Red}(F)$, that is, $\mathbb{K}\mathrm{Red}(\wedge F)$ since F is confluent. Hence, $\widehat{v_2}-\widehat{v_3}$ also belongs to $\mathbb{K}\mathrm{Red}(\wedge F)$, that is, it belongs to the image of $\wedge F$. Thus, $\widehat{v_2}-\widehat{v_3}$ belongs to $\ker(\wedge F)\cap\mathrm{im}(\wedge F)$ which is reduced to $\{0\}$ since $\wedge F$ is a projector. We conclude that $\widehat{v_2}$ is equal to $\widehat{v_3}$, so that $\underset{F}{\longrightarrow}$ is confluent.

□

4.2.5. Completion. Given a completable set F, the notion of *complement* is stated as follows: a complement of F is a reduction operator C satisfying

1. $\wedge F\preceq C$,
2. $\mathrm{Obs}^F\ \subset\ \mathrm{Nred}(C)$.

4.2.6. Completion and Rewriting. Let C be a reduction operator, greater than $\wedge F$. We write $F' = F \cup \{C\}$. The vector space $\sum_{T \in F'} \ker(T)$ is equal to $\sum_{T \in F} \ker(T)$. Hence, the set F' is also completable and $\wedge F'$ is equal to $\wedge F$. In particular, the two equivalence relations $\overset{*}{\underset{F}{\longleftrightarrow}}$ and $\overset{*}{\underset{F'}{\longleftrightarrow}}$ are equal. Using the arguments of the proof of Proposition 3.2.2, we get

4.2.7. Proposition. *Let $C \in \mathbf{RO}(G, <)$ such that C is greater than $\wedge F$. The set $F \cup \{C\}$ is confluent if and only if C is a complement of F.*

4.2.8. Remark. We have seen that the absence of a total order on G implies that the reduction operator $\wedge F$ does not necessarily exist. In particular, the notion of complement is not necessarily defined, like the Knuth-Bendix completion algorithm in term rewriting does not necessarily succeed. Consider the example from 4.1.3: $G = \{g_1, g_2, g_3\}$ with $g_1 < g_3$ and $g_2 < g_3$, and let $P = (T_1, T_2)$ where, for $i = 1$ or 2, $\mathrm{Nred}(T_i) = \{g_3\}$ and $T_i(g_3) = g_i$. We have seen in 4.1.3 that $T_1 \wedge T_2$ does not exist. Moreover, the Knuth-Bendix completion algorithm does not work because as shown on the following diagram

$$\begin{tikzcd} & g_3 \arrow[dl] \arrow[dr] & \\ T_1(g_3) = g_1 & & T_2(g_3) = g_2 \end{tikzcd}$$

g_3 admits two distinct normal forms g_1 and g_2, and these two normal forms cannot be compared.

References

[1] Franz Baader and Tobias Nipkow. *Term rewriting and all that.* Cambridge University Press, Cambridge, 1998.

[2] Jörgen Backelin, Svetlana Cojocaru, and Victor Ufnarovski. The computer algebra package Bergman: current state. In *Commutative algebra, singularities and computer algebra (Sinaia, 2002)*, volume 115 of *NATO Sci. Ser. II Math. Phys. Chem.*, pages 75–100. Kluwer Acad. Publ., Dordrecht, 2003.

[3] Roland Berger. Confluence and Koszulity. *J. Algebra*, 201(1):243–283, 1998.

[4] Roland Berger. Weakly confluent quadratic algebras. *Algebr. Represent. Theory*, 1(3):189–213, 1998.

[5] Roland Berger. Koszulity for nonquadratic algebras. *J. Algebra*, 239(2):705–734, 2001.

[6] George M. Bergman. The diamond lemma for ring theory. *Adv. in Math.*, 29(2):178–218, 1978.

[7] Leonid A. Bokut′. Imbeddings into simple associative algebras. *Algebra i Logika*, 15(2):117–142, 245, 1976.

[8] Ronald V. Book and Friedrich Otto. *String-rewriting systems.* Texts and Monographs in Computer Science. Springer-Verlag, New York, 1993.

[9] Bruno Buchberger. Ein Algorithmus zum Auffinden der Basiselemente des Restklassenrings nach einem nulldimensionalen Polynomideal. *Universitat Innsbruck, Austria, Ph. D. Thesis*, 1965.

[10] Bruno Buchberger. History and basic features of the critical-pair/completion procedure. *J. Symbolic Comput.*, 3(1-2):3–38, 1987. Rewriting techniques and applications (Dijon, 1985).

[11] Cyrille Chenavier. Confluence algebras and acyclicity of the Koszul complex. *Algebr. Represent. Theory*, 2016.

[12] Peter J. Hilton and Urs Stammbach. *A course in homological algebra*, volume 4 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, second edition, 1997.

[13] Deepak Kapur and Paliath Narendran. A finite Thue system with decidable word problem and without equivalent finite canonical system. *Theoret. Comput. Sci.*, 35(2-3):337–344, 1985.

[14] Donald E. Knuth and Peter B. Bendix. Simple word problems in universal algebras. In *Computational Problems in Abstract Algebra (Proc. Conf., Oxford, 1967)*, pages 263–297. Pergamon, Oxford, 1970.

[15] Benoit Kriegk and Michel Van den Bergh. Representations of non-commutative quantum groups. *Proc. Lond. Math. Soc. (3)*, 110(1):57–82, 2015.

[16] Teo Mora. An introduction to commutative and noncommutative Gröbner bases. *Theoret. Comput. Sci.*, 134(1):131–173, 1994. Second International Colloquium on Words, Languages and Combinatorics (Kyoto, 1992).

[17] Maxwell H. A. Newman. On theories with a combinatorial definition of "equivalence.". *Ann. of Math. (2)*, 43:223–243, 1942.

[18] Alexander Polishchuk and Leonid Positselski. *Quadratic algebras*, volume 37 of *University Lecture Series*. American Mathematical Society, Providence, RI, 2005.

[19] Stewart B. Priddy. Koszul resolutions. *Trans. Amer. Math. Soc.*, 152:39–60, 1970.

[20] Anatoly I. Shirshov. Some algorithmic problems for lie algebras. In *Selected Works of AI Shirshov*, pages 125–130. Springer, 2009.

Cyrille Chenavier
INRIA, équipe πr^2
Laboratoire IRIF, CNRS UMR 8243
Université Paris-Diderot
Case 7014
75205 PARIS Cedex 13
chenavier@pps.univ-paris-diderot.fr