



HAL
open science

Sporulation boundaries and spore formation kinetics of *Bacillus* spp. as a function of temperature, pH and aw.

Eugénie Baril, Louis Coroller, Olivier Couvert, Mohammed El Jabri, Ivan Leguerinel, Florence Postollec, Christophe Boulais, Frédéric Carlin, Pierre Mafart

► To cite this version:

Eugénie Baril, Louis Coroller, Olivier Couvert, Mohammed El Jabri, Ivan Leguerinel, et al.. Sporulation boundaries and spore formation kinetics of *Bacillus* spp. as a function of temperature, pH and aw.. Food Microbiology, 2012, 32 (1), pp.79-86. 10.1016/j.fm.2012.04.011 . hal-01330230

HAL Id: hal-01330230

<https://hal.science/hal-01330230v1>

Submitted on 29 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License

Accepted Manuscript



Sporulation boundaries and spore formation kinetics of *Bacillus spp.* as a function of temperature, pH and a_w

E. Baril, L. Coroller, O. Couvert, M. El Jabri, I. Leguerinel, F. Postollec, C. Boulais, F. Carlin, P. Mafart

PII: S0740-0020(12)00093-7

DOI: [10.1016/j.fm.2012.04.011](https://doi.org/10.1016/j.fm.2012.04.011)

Reference: YFMIC 1784

To appear in: *Food Microbiology*

Received Date: 8 August 2011

Revised Date: 18 April 2012

Accepted Date: 19 April 2012

Please cite this article as: Baril, E., Coroller, L., Couvert, O., El Jabri, M., Leguerinel, I., Postollec, F., Boulais, C., Carlin, F., Mafart, P., Sporulation boundaries and spore formation kinetics of *Bacillus spp.* as a function of temperature, pH and a_w , *Food Microbiology* (2012), doi: 10.1016/j.fm.2012.04.011.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

1 **Sporulation boundaries and spore formation kinetics of *Bacillus spp.* as a function of**
2 **temperature, pH and a_w**

3 E. Baril^{1,2,3*}, L. Coroller^{1,2}, O. Couvert^{1,2}, M. El Jabri³, I. Leguerinel^{1,2}, F. Postollec³, C.
4 Boulais⁴, F. Carlin^{5,6}, P. Mafart^{1,2}

5 ¹ : Université Européenne de Bretagne, France

6 ²: Université de Brest, EA3882, Laboratoire Universitaire de Biodiversité et Ecologie
7 Microbienne, UMT 08.3 PHYSI'Opt, IFR148 ScInBioS, 6 rue de l'Université, F-29334
8 Quimper, France

9 ³: ADRIA Développement, UMT 08.3 PHYSI'Opt, Z.A. Creac'h Gwen, F-29196
10 Quimper Cedex, France

11 ⁴: Danone Research, Avenue de la Vauve, RD128, F-91767 Palaiseau Cedex, France

12 ⁵: INRA, UMR408, Sécurité et Qualité des Produits d'Origine Végétale, F-84000
13 Avignon, France

14 ⁶: Université d'Avignon et des Pays de Vaucluse, UMR408, Sécurité et Qualité des
15 Produits d'Origine Végétale, F-84000 Avignon, France

16 * Corresponding author

17 Address: Université Européenne de Bretagne, France - Université de Brest, EA3882,
18 Laboratoire Universitaire de Biodiversité et Ecologie Microbienne, IFR148 ScInBioS
19 6, rue de l'université,
20 29 334 Quimper, France

21 Phone: 33 (0)2 98 64 19 30

22 Fax: 33 (0)2 98 64 19 69

23 E-mail: Eugenie.baril@univ-brest.fr

24 Abstract

25 Sporulation niches in the food chain are considered as a source of hazard and are not
26 clearly identified. Determining the sporulation environmental boundaries could contribute
27 to identify potential sporulation niches. Spore formation was determined in a Sporulation
28 Mineral Buffer. The effect of incubation temperature, pH and water activity on time to
29 one spore per mL, maximum sporulation rate and final spore concentration was
30 investigated for a *Bacillus weihenstephanensis* and a *B. licheniformis* strain.

31 Sporulation boundaries of *B. weihenstephanensis* and of *B. licheniformis* were similar to,
32 or included within, the range of temperatures, pH and water activities supporting growth.
33 For instance, sporulation boundaries of *B. weihenstephanensis* were evaluated at 5°C,
34 35°C, pH 5.2 and a_w 0.960 while growth boundaries were observed at 5°C, 37°C, pH 4.9
35 and a_w 0.950. Optimum spore formation was determined at 30°C pH 7.2 for *B.*
36 *weihenstephanensis* and at 45°C pH 7.2 for *B. licheniformis*. Lower temperatures and pH
37 delayed the sporulation process. For instance, the time to one spore per mL was tenfold
38 longer when sporulation occurred at 10°C and 20°C, for each strain respectively, than at
39 optimum sporulation temperature. The relative effect of temperature and pH on
40 sporulation rates and on growth rates is similar. This work suggests that the influence of
41 environmental factors on the quantitative changes in sporulation boundaries and rates was
42 similar to their influence on changes in growth rate.

43

44 Key words: *Bacillus weihenstephanensis*, *Bacillus licheniformis*, sporulation boundaries,
45 growth, temperature, pH, water activity, modelling

46

47 **1 Introduction**

48 Contamination and persistence of spores of *Bacillus* and related genera in foods is a real
49 concern for the food industry, despite a constant implementation of hygienic practices.
50 Bacterial spores are common contaminants of food raw materials and ingredients (dairy
51 products, flour, cocoa, bakery products, spices, meat products, vegetables or fruits,)
52 (Anonymous, 2005; Carlin, 2011). In favourable conditions spores germinate, grow and
53 may cause spoilage and quality loss. Pathogenic species may cause foodborne diseases.
54 *B. cereus* and *B. licheniformis* for instance are well known agents of spoilage in dairy
55 products such as sweet curdling and bitty cream (De Jonghe *et al.*, 2010; Meer *et al.*,
56 1991). *B. cereus* emetic or diarrheal toxins are regularly at the origin of foodborne
57 poisonings in which a large diversity of foods are implicated (Anonymous, 2005;
58 Stenfors Arnesen *et al.*, 2008). The spore formation occurs in diverse natural or linked to
59 agricultural and industrial activities niches (Carlin, 2011). As sources of spore
60 contamination in the food chain, sporulation niches are potential food safety hazards. The
61 determination of environmental conditions allowing sporulation and of sporulation
62 boundaries may contribute to identify the main contamination sources throughout the
63 food chain.

64 Sporulation is the latest cellular response of bacteria to nutrient depletion (Burkholder
65 and Grossman, 2000). It generally occurs at high population density and involves cellular
66 communication by accumulation of various extracellular oligopeptides (Auchtung and
67 Grossman, 2008; Grossman, 1995; Sonenshein, 1999). Spore formation may be observed
68 in laboratory with specific broth and agar media as well as in natural environments.

69 Indeed a complete life cycle of germination-growth-sporulation was observed in the
70 mouse intestinal tract (Tam *et al.*, 2006) and thermophilic spore formation was observed
71 in heat exchangers and evaporators (Scott *et al.*, 2007).

72 Spore formation is delayed and spore yield is generally lower at sub-optimum growth
73 temperatures (Baweja *et al.*, 2008; Garcia *et al.*, 2010; Lechowich and Ordal, 1962;
74 Nguyen Thi Minh *et al.*, 2011). For instance, time to achieve more than 80% of free
75 spores of three *Bacillus cereus* strains was two to three fold longer at 20°C than at 30°C
76 to 45°C (Gonzalez *et al.*, 1999). Similarly, acidic pH delays sporulation and reduces
77 sporulation yield (Baweja *et al.*, 2008; Craven, 1990; Nguyen Thi Minh *et al.*, 2011;
78 Yazdany and Lashkari, 1975). For three *Bacillus cereus* strains, time to reach 5 to 10% of
79 free spores was at pH 6.0 two to four fold longer than time needed to achieve more than
80 80% of free spores at pH 6.5 and 7.0 (Mazas *et al.*, 1997). These variations in sporulation
81 time and in spore concentration were also dependent on water activity (a_w) (Jakobsen and
82 Murrell, 1977). At a_w 0.993, 4 days were needed to obtain 9.10^9 spore/mL of *B. subtilis*,
83 while 17 days were needed to form only 3.10^7 spore/mL at a_w 0.950 (Nguyen Thi Minh *et*
84 *al.*, 2008).

85 The aim of this work was to evaluate the influence of temperature, pH and a_w on
86 sporulation boundaries and sporulation kinetics as follows: (i) identify the environmental
87 conditions where sporulation could be observed, the changes in maximum spore
88 concentration of *B. weihenstephanensis* and *B. licheniformis* were followed in two
89 sporulation media to determine sporulation boundaries; ii) quantify the influence of
90 environmental conditions on sporulation kinetic parameters by fitting a mathematical

91 model; and iii) compare the impact of temperature, pH and aw on sporulation boundaries
92 and sporulation rates to those on growth boundaries and growth rates.

93

94 **2 Materials and Methods**

95 **2.1 Bacterial strains and culture conditions**

96 The *Bacillus weihenstephanensis* strain KBAB4 isolated from a forest soil was kindly
97 provided by the Institut National de la Recherche Agronomique (INRA, Avignon,
98 France) (Vilas-Boas *et al.*, 2002) . This strain belongs to the psychrotrophic group VI of
99 *B. cereus sensu lato* (Guinebretiere *et al.*, 2008). *Bacillus licheniformis* strain AD978 was
100 isolated from raw dairy ingredients and was kindly provided by ADRIA Développement
101 (Quimper, France).

102 Both strains were frozen (-20°C) as 1 mL aliquots in nutrient broth diluted in 50%
103 glycerol (v/v). A preculture was performed by diluting 1 mL aliquot into 100 mL nutrient
104 broth (NB) (Biokar Diagnostics, Beauvais, France) in 250 mL flasks for 8 h, followed by
105 a 1/100 preculture incubated for 16 h. Incubation was performed under shaking (100 rpm)
106 at 30°C (45°C) for *B. weihenstephanensis* KBAB4 (for *B. licheniformis* AD978). Finally
107 a 0.1 mL culture of *B. weihenstephanensis* KBAB4 or a 0.01 mL culture of *B.*
108 *licheniformis* AD978 was diluted in 100 mL NB and incubated for 6 h to reach
109 concentrations of approximately 7.7 log₁₀(CFU/mL) for both strains with less than 2.6
110 log₁₀(CFU/mL) of spores, which was the limit of spore quantification (20 colonies per
111 plates for a 50 µL inoculation of undiluted sample).

112

113 **2.2 Determination of cardinal growth values of vegetative cells of *Bacillus sp.* by**
114 **turbidimetry measurements**

115 Growth boundaries were considered as the minimum and maximum temperatures,
116 minimum pH and minimum a_w values at which growth was observed (Ross *et al.*, 2011).
117 They were observed during the estimation of cardinal growth values by using the
118 turbidimetry method (Membré *et al.*, 2002) to compare the influence of temperature, pH
119 and a_w on maximum growth rates. At least ten levels of temperature, pH or a_w were
120 studied from 5°C to 55°C, from pH 4.6 to 7.5 and from a_w 0.920 to 0.996, for both
121 strains. Throughout the study, experimental data sets were collected from univariate
122 experiments. To test the effect of one environmental factor, other factors were fixed at
123 constant values: temperature was fixed at 30°C for *B. weihenstephanensis* KBAB4 and at
124 45°C for *B. licheniformis* AD978, pH was fixed at 7.2 and a_w was fixed at 0.996.

125 Growth kinetics were performed in brain heart infusion supplemented with yeast extract
126 (0.3%) and D-glucose (0.2%) (BHIYG) (all from Biokar diagnostics, Beauvais, France).
127 The pH value was adjusted with HCl (1 or 5 N) and the a_w was adjusted with glycerol
128 (Achard *et al.*, 1992). Broth sterilization was performed by 0.22 μm filtration (Steritop
129 system; Millipore Corporation, Billerica, MA). The wells of the 10x10 honeycomb plate
130 (Oy growth curves AB Ltd; Helsinki, Finland) were filled with 200 μL of pH- a_w -
131 adjusted BHIYG, with one well column (10 wells) per pH or a_w condition. The vegetative
132 cell suspension (see section 2.1) was diluted in 10 mL BHIYG to obtain a suspension of
133 approximately 5.0 \log_{10} (CFU/mL). 200 μL of this suspension was inoculated in the first
134 wells of each well column of the honeycomb plate. For each well column, twofold
135 dilution series were made up from the first to the eighth well (Biesta-Peters *et al.*, 2010;

136 Membré *et al.*, 2002). The ninth and the tenth wells were used as sterility control. The
 137 honeycomb plate was then incubated in Bioscreen C (Labsystems, Helsinki, Finland) at
 138 the tested temperature with continuous moderate shaking. The absorbance was
 139 automatically measured at 660 nm every 10 min. Growth at 5°C was observed by
 140 turbidimetry measurement in 10 mL BHIYG, inoculated at approximately 5.0
 141 log₁₀(CFU/mL). When no increase of absorbance was observed after 30 days of
 142 incubation, no-growth was concluded.

143 The statistical analysis of Bioscreen curves was done according to previous studies
 144 (Cuppers and Smelt, 1993; Membré *et al.*, 2002). For each well, the detection time was
 145 determined as the time to reach the absorbance value of 0.4, meaning the middle of the
 146 linear phase of the turbidimetry curve (Membré *et al.*, 2005). The slope of the linear
 147 relationship between detection times and logarithm of initial concentrations corresponded
 148 to the maximum growth rates (μ_{max}) (Biesta-Peters *et al.*, 2010; Le Marc *et al.*, 2002;
 149 Membré *et al.*, 2002) (regress function, Statistical Toolbox, MATLAB 7.9.0, The Math-
 150 works, Natick, USA). To estimate cardinal growth values, the maximum growth rates
 151 were then fitted to the growth cardinal model according to the equation 1 (Augustin and
 152 Carlier, 2000; Rosso *et al.*, 1995) (fminsearch function, Optimization Toolbox,
 153 MATLAB 7.9.0, The Math-works, Natick, USA).

$$154 \quad \mu_{max} = \mu_{opt} \cdot CM_2(T) \cdot CM_1(pH) \cdot CM_1(a_w) \quad (1)$$

155 with

$$156 \quad CM_n(X) = \begin{cases} 0 & X \leq X_{min} \\ \frac{(X - X_{max})(X - X_{min})^n}{(X_{opt} - X_{min})^{(n-1)} [(X_{opt} - X_{min})(X - X_{opt}) - (X_{opt} - X_{max})((n-1)X_{opt} + X_{min} - nX)]} & X_{min} < X < X_{max} \\ 0 & X \geq X_{max} \end{cases} \quad (2)$$

158 Where μ_{max} (h^{-1}) is the maximum growth rate, μ_{opt} (h^{-1}) is the maximum growth rate at
159 optimum conditions (optimum temperature, pH and a_w), T ($^{\circ}\text{C}$) is the temperature, X_{min} ,
160 X_{opt} and X_{max} are respectively minimum, optimum and maximum growth values.

161 As described by Efron and Tibshirani (1993) and Gauchi *et al.* (2010), the 95%
162 confidence intervals were computed by the bootstrap resampling method of 1000 samples
163 (bootstrap function, Statistical Toolbox, MATLAB 7.9.0, The Math-works, Natick, USA).

164

165 **2.3 Spore production**

166 To evaluate the dependence of the boundaries on the composition of the medium, spores
167 were produced in Sporulation Mineral Buffer (SMB) and in Modified Nutrient Agar
168 (MNA). Both media were prepared at various initial pH and a_w levels and incubated at
169 various temperatures. In particular SMB is a buffered medium, which allows the control
170 of pH during the sporulation process (Baril *et al.*, 2012). SMB was composed of
171 phosphate buffer ($\text{K}_2\text{HPO}_4/\text{KH}_2\text{PO}_4$, 39 mM) (Sigma-Aldrich, Saint Quentin Fallavier,
172 France) supplemented with $\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$ (8.0 mg/L) and $\text{MnSO}_4 \cdot \text{H}_2\text{O}$ (1.5 mg/L)
173 (Sigma-Aldrich, Saint Quentin Fallavier, France) (Baril *et al.*, 2011). pH was adjusted by
174 modifying the acid-base balance of the buffer and a_w was adjusted with glycerol (Achard
175 *et al.*, 1992). SMB was then sterilized by filtration on a 0.22 μm filter. The vegetative cell
176 cultures of *B. weihenstephanensis* KBAB4 and *B. licheniformis* AD978 (see section 2.1)
177 were centrifuged at 6000 x g for 10 min at 12 $^{\circ}\text{C}$. The pellets were suspended in 100 mL
178 SMB. The final concentration was approximately 7.7 $\log_{10}\text{CFU/mL}$, with less than 2.6
179 $\log_{10}\text{CFU/mL}$ of spores. The SMB flasks were then incubated at the tested temperatures
180 and shaken at 100 rpm.

181 The MNA was composed of nutrient agar (Biokar diagnostics, Beauvais, France)
182 modified by the supplementation with $\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$ (100.0 mg/L) and $\text{MnSO}_4 \cdot \text{H}_2\text{O}$ (50.0
183 mg/L) (Gaillard *et al.*, 1998). The pH was adjusted with HCl (1 or 5 N) and a_w with
184 glycerol (Achard *et al.*, 1992). The MNA was then sterilized by autoclaving at 120°C for
185 15 min. MNA plates were inoculated with 0.5 mL of the vegetative cell cultures (see
186 section 2.1) and were incubated at the tested temperatures.

187 Spores of both strains were produced from univariate experiments at temperatures
188 ranging from 5°C to 40°C for *B. weihenstephanensis* KBAB4 (5°C to 50°C for *B.*
189 *licheniformis* AD978), at pH ranging from 5.2 to 8.5 and at a_w ranging from 0.935 to
190 0.996. The pH and a_w of sporulation media were checked immediately after the
191 inoculation respectively by a pH-meter (PHM210, MeterLab; Radiometer Copenhagen)
192 and an a_w -meter (FA-st1 GBX; France Scientific Instrument, Romans, France).

193

194 **2.4 Determination of sporulation boundaries and kinetics of *Bacillus* sp.**

195 Total counts corresponded in this work to the total viable microbial population, which is
196 vegetative cells plus spores. Total counts were performed on unheated suspension. Spores
197 were defined as resisting cells to a heat treatment at 70°C for 5 min (Baril *et al.*, 2011).

198 To enumerate spores, a 1 mL suspension was heated at 70°C for 5 min in a glass tube by
199 immersion into a water bath. Diluted suspensions were then spread on nutrient agar plates
200 using a spiral plater (WASP1; Don Whitley, Shipley, West Yorkshire, United Kingdom).

201 Plates were incubated for 24 h at 30°C for *B. weihenstephanensis* and at 45°C for *B.*
202 *licheniformis*. As mentioned by the manufacturer, the limit of quantification corresponds

203 to 20 colonies per plates for a 50 μ L inoculation of the least diluted sample, i.e. 2.6
204 \log_{10} (CFU/mL), for both total counts and spore counts.

205 Sporulation boundaries were considered as the minimum and maximum temperatures,
206 minimum pH and minimum a_w values at which spores were detected. To determine the
207 sporulation boundaries, final spore concentrations were evaluated at various
208 temperatures, pH and a_w in both sporulation media. The study was performed as
209 univariate experiments for both microorganisms at temperatures ranging from 5°C to
210 40°C for *B. weihenstephanensis* KBAB4 (5°C to 50°C for *B. licheniformis* AD978) with
211 a 5°C interval, at pH ranging from 5.2 to 8.5 with a 0.3 pH unit interval and at a_w ranging
212 from 0.935 to 0.996 with a 0.005 a_w unit interval. When at least 95% of free spores were
213 observed in the sporulation medium under phase contrast microscope (Olympus BX50,
214 Olympus Optical Co., Ltd, Hamburg, Germany), total counts and spore counts were
215 enumerated in 2 mL aliquots of SMB, or after scrapping MNA plates and suspending
216 spores in 5 mL of distilled water. When only a few cells were observed under phase
217 contrast microscope, meaning concentration lower than 6 \log_{10} CFU/mL, total counts and
218 spore counts were performed after 30 days of incubation. pH and a_w values of the
219 sporulation media were checked after microbial counting.

220 To quantify the influence of temperature and pH on spore formation, sporulation kinetics
221 were performed in SMB at two temperature and pH levels. Sporulation kinetics were
222 performed, as univariate experiments, at optimum conditions of temperature and pH
223 (30°C and pH 7.2 for *B. weihenstephanensis*, 45°C and pH 7.2 for *B. licheniformis*) and
224 at suboptimum conditions (10°C or pH 5.9 for *B. weihenstephanensis*, 20°C or pH 6.3 for
225 *B. licheniformis*). Total counts and spore counts were performed at appropriate time

226 intervals. Three independent (different dates of experiment) sporulation kinetics were
 227 done for each tested temperature and pH condition.

228 The following sporulation kinetic model, inspired from a logistic model using a
 229 hyperbolic tangent function, was proposed to fit sporulation curves:

$$230 \quad \log_{10}(N_s) = \log_{10}(N_{sf}) \frac{1 - \exp(-\mu_s(t - t_{1s}))}{1 + \exp(-\mu_s(t - t_{1s}))}, t > t_{1s} \quad (3)$$

231 where N_s is the spore concentration (CFU/mL) at the time t (h), N_{sf} is the final spore
 232 concentration (CFU/mL), μ_s is the maximum sporulation rate (h^{-1}) and t_{1s} is the
 233 incubation time (h) at which N_s is theoretically equal to one spore per mL.

234 Model parameters were estimated by minimizing the sum of squared errors (SSE)
 235 between the logarithm of the observed and the estimated spore concentration ($\log_{10}(N_s)$).

236 These estimations were computed by non linear regressions (lsqcurvefit function,
 237 Optimization Toolbox, MATLAB 7.9.0, The Math-works, Natick, USA), as well as the

238 estimation of the 95% confidence intervals (nlparci, Statistical Toolbox, MATLAB 7.9.0,
 239 The Math-works, Natick, USA). Since the estimated values of the three replicates were

240 comparable for all environmental conditions, sporulation kinetics were fitted once for
 241 each triplicate experimental data set to estimate only one set of parameters per

242 environmental condition (i.e. $\log_{10}(N_{sf})$, μ_s and t_{1s}). The goodness of the model fit was
 243 checked by the root mean square error (RMSE) (Equation 4).

$$244 \quad RMSE = \sqrt{\frac{SSE}{n - p}} \quad (4)$$

245 where SSE is the sum of squared errors, n is the number of experimental data set and p is
 246 the number of estimated parameters.

247 The Kruskal-Wallis test was computed to test the influence of the sporulation
248 temperature, pH and a_w on final spore concentrations and on estimated parameters
249 (kruskalwallis and ranksum functions, Statistics Toolbox, MATLAB 7.9.0; The Math-
250 works). For each statistical analysis, a significant difference between compared data was
251 stated at the risk α of 5%.

252 **3 Results**

253 **3.1 Growth abilities**

254 Growth of *B. weihenstephanensis* KBAB4 was observed from 5°C to 37°C, however at
255 37°C growth was only observed for one out of two experiments (Figure 1A). Growth was
256 observed from pH 4.9 to 7.5, and not at pH 4.6 and 4.8 (Figure 1C). Growth was also
257 observed from a_w 0.950 to 0.996, and not at a_w 0.920, 0.930 and 0.940 (Figure 1E). Thus
258 growth boundaries of *B. weihenstephanensis* KBAB4 were at 5°C, 37°C, pH 4.9 and a_w
259 0.950. Growth cardinal values of *B. weihenstephanensis* KBAB4 were estimated at
260 0.9°C, 33.1°C and 38.6°C, respectively for minimum, optimum and maximum
261 temperatures, at pH 4.6 and a_w 0.946 respectively for minimum growth pH and a_w . The
262 maximum growth rate (μ_{max}) was estimated at 1.68 h⁻¹ (Table 1).

263 Growth of *B. licheniformis* AD978 was observed from 20°C to 55°C. No growth at 7°C,
264 10°C and 15°C was observed (Figure 1B). Growth was also observed from pH 4.9 to 7.5
265 (Figure 1D) and from a_w 0.920 to 0.996 (Figure 1F). Growth boundaries of *B.*
266 *licheniformis* AD978 were at 20°C, 55°C, pH 4.9 and a_w 0.920. Growth cardinal values
267 estimated from the equation 1 were 8.6°C, 50.9°C, 56.1°C for minimum, optimum,
268 maximum temperatures, at pH 5.3 and a_w 0.915 respectively for minimum pH and a_w .
269 The maximum growth rate was estimated at 3.12 h⁻¹ (Table 1).

270 **3.2 Sporulation abilities**

271 **Effect of temperature on sporulation abilities**

272 For *B. weihenstephanensis*, the average final spore concentration in SMB was slightly but
273 significantly higher at 30°C (7.3 log₁₀CFU/mL) than at 10°C and 35°C (6.3 log₁₀CFU/mL
274 and 6.4 log₁₀CFU/mL, respectively) ($p < 0.05$) (Figure 2A). Out of this range, the final
275 spore concentration largely decreased. At 5°C, spores were formed in only one out of
276 three replicates indicating that the sporulation temperature boundary was reached.
277 Similarly, at 40°C, no spore formation was detected. For *B. licheniformis*, the final spore
278 concentration in SMB was not significantly different between 20°C and 45°C ($p > 0.05$)
279 and was comprised between 6.9 and 7.5 log₁₀CFU/mL (Figure 2B). Out of this range, a
280 sharp decrease in the final spore concentration was observed. At 5°C, 10°C and 15°C, no
281 spore was detected. At 50°C, the final spore concentration of 4.6 log₁₀CFU/mL was
282 significantly lower than at 45°C ($p < 0.05$).

283 The same trend was observed in MNA for both strains. Sporulation temperature
284 boundaries ranged from 5°C to 35°C for *B. weihenstephanensis* KBAB4 and from 20°C
285 to 50°C for *B. licheniformis* AD978. Therefore, sporulation boundaries are close to
286 temperature growth boundaries, which were observed at 5°C and 37°C for *B.*
287 *weihenstephanensis* KBAB4 and from 20°C to 50°C for *B. licheniformis* AD978.

288

289 **Effect of pH on sporulation abilities**

290 Variations of the pH along the sporulation periods were less pronounced in SMB than in
291 MNA. In SMB, the pH increased of 0.4 units for an initial pH of 5.2 and decreased of 0.2
292 units for an initial pH of 8.5. Lower variations were observed at initial pH between 5.6

293 and 8.0. In MNA, whatever the initial sporulation pH was, ranging from 5.6 to 7.2, the
294 final sporulation pH was measured at 8.7 (± 0.1).

295 For *B. weihenstephanensis* KBAB4, the final spore concentrations in SMB were not
296 different from pH 8.5 to 7.2 and were close to 7.0 log₁₀CFU/mL (Figure 2C). The final
297 spore concentration decreased progressively as pH decreased. For example, no significant
298 differences were noticed between the final spore concentration at pH 7.2 and 5.9. and
299 between pH 5.9 and 5.6 ($p > 0.05$), however significant differences were observed
300 between the final spore concentration at pH 7.2 and 5.6 ($p < 0.05$). The final spore
301 concentration decreased sharply from 6.4 log₁₀CFU/mL at pH 5.6 to 4.0 log₁₀CFU/mL at
302 pH 5.2. For *B. licheniformis* AD978, the final spore concentration in SMB remained
303 unchanged from pH 8.5 to 7.0 and reached 7.5 log₁₀CFU/mL (Figure 2D). At pH 6.3 the
304 final spore concentration decreased significantly to 5.0 log₁₀CFU/mL ($p < 0.05$). No
305 spore was detected at pH 5.2, 5.5 and 5.8. In MNA, final spore concentrations were not
306 significantly different from pH 5.6 to 7.2 for *B. weihenstephanensis* KBAB4 and from pH
307 6.3 to 7.2 for *B. licheniformis* AD978 ($p > 0.05$). Sporulation pH boundaries in SMB
308 were evaluated at pH 5.2 for *B. weihenstephanensis* KBAB4 and at pH 5.8 for *B.*
309 *licheniformis* AD978. These sporulation boundaries are higher than the growth
310 boundaries for pH (at pH 4.9 for both strains).

311

312 **Effect of a_w on sporulation abilities**

313 In SMB, no significant difference of a_w values was observed from the beginning to the
314 end of sporulation period. In MNA, a_w values increased during the incubation. The
315 maximum increase (0.034 a_w unit) was observed at an initial a_w of 0.955.

316 For *B. weihenstephanensis* KBAB4, the final spore concentration in SMB increased
317 significantly from 3.2 log₁₀CFU/mL to 7.3 log₁₀CFU/mL as a_w increased from 0.965 and
318 0.996 ($p < 0.05$) (Figure 2E). At a_w 0.960, spores were formed in one out of two
319 replicates. For *B. licheniformis* AD978, the final spore concentration in SMB increased
320 largely from 3.1 to 7.5 log₁₀CFU/mL between a_w 0.985 and 0.996 ($p < 0.05$) (Figure 2F).
321 At a_w 0.975 and a_w 0.980, spores were detected in one out of two or three replicates. The
322 same trend was observed in MNA, but on a wider range of a_w for both strains comprised
323 between a_w 0.935 and 0.996. The sporulation a_w boundaries were in SMB at a_w 0.960 for
324 *B. weihenstephanensis* KBAB4 and at 0.975 for *B. licheniformis* AD978. These
325 sporulation boundaries are higher than the growth boundaries being respectively 0.950
326 and 0.920 for each strain.

327

328 **3.3 Sporulation kinetics**

329 Because the a_w range allowing spore formation was too narrow (between 0.960 and 0.996
330 for *B. weihenstephanensis* and between 0.975 and 0.996 for *B. licheniformis*) and final
331 spore concentration was close to the limit of quantification, sporulation kinetics were not
332 performed at different a_w levels.

333 As shown on Figure 3, equation 3 gave a good description of the observed spore
334 concentration and allowed the quantification of the time to achieve one spore per mL
335 (t_{1S}), the maximum sporulation rate (μ_s) and the final spore concentration (N_{sf}).

336 As previously observed, the final spore concentrations of *B. weihenstephanensis* KBAB4
337 were significantly lower at 10°C than at 30°C while no significant difference was
338 observed between pH 5.9 and 7.2 ($p > 0.05$) (Table 2). The time to observe one spore per

339 mL (t_{IS}) of *B. weihenstephanensis* KBAB4 was estimated at 148.0 h, 14.5 h and 6.7 h
340 respectively at 10°C pH 7.2, at 30°C pH 5.9 and at 30°C pH 7.2 (Table 2). This time (t_{IS})
341 was significantly prolonged when temperature and pH decreased from 30°C to 10°C and
342 from pH 7.2 to 5.9 ($p < 0.05$). The maximum sporulation rate (μ_S) was significantly lower
343 at 10°C (0.05 h^{-1}) than at 30°C (0.60 h^{-1}) but was not significantly different between pH
344 5.9 (0.50 h^{-1}) and pH 7.2 (0.60 h^{-1}) ($p > 0.05$).

345 For *B. licheniformis* AD978, the final spore concentration was no significantly different
346 between 20°C and 45°C but was significantly lower at pH 6.3 than at pH 7.2 (Table 2) (p
347 < 0.05). t_{IS} was estimated at 45.9 h, 4.2 h and 1.6 h for sporulation at 20°C pH 7.2, at
348 45°C pH 6.3 and at 45°C pH 7.2, respectively (Table 2). t_{IS} increased when temperature
349 or pH decreased ($p < 0.05$). The maximum sporulation rate (μ_S) was significantly lower at
350 20°C than at 45°C (0.02 h^{-1} and 0.37 h^{-1} respectively), but significantly higher at pH 6.3
351 than at pH 7.2 (0.55 h^{-1} and 0.37 h^{-1} respectively) ($p < 0.05$) (Table 2).

352

353 4 Discussion

354 This study shows that the range of temperature, pH and a_w allowing sporulation of *B.*
355 *weihenstephanensis* KBAB4 and of *B. licheniformis* AD978 is similar to, or lying within,
356 the range of temperature, pH and a_w allowing growth. These results are in agreement with
357 previous works. For instance, *B. sphaericus* grew from 10°C to 40°C while sporulation
358 was only observed between 10°C and 30°C (De Pieri and Ludlow, 1992). Similarly,
359 growth of three *B. cereus* strains was lying between pH 5.0 and pH 9.5 when their
360 sporulation was only possible between pH 6.5 and 8.0 (Mazas *et al.*, 1997).
361 Consequently, as a first approximation, it seems that temperature, pH and a_w growth

362 boundaries could be as well used as an estimation of sporulation boundaries even if it
363 would be a conservative estimation. For tested conditions, the final spore concentration
364 markedly decreased as temperature, pH and a_w were close to growth boundaries (Figure 1
365 and 2). It must be noticed that care should be taken when inferring sporulation boundaries
366 from growth boundaries of *Bacillus* strains because differences due to strain type,
367 physiological state of cells and composition of the media used might occur.

368 A primary model was proposed to quantify the sporulation kinetics. Spore formation was
369 delayed as sporulation temperature and pH decreased (Figure 3). For instance, a tenfold
370 decrease of the time to one spore per mL (t_{1S}) was observed at 10°C as compared to 30°C
371 for *B. weihenstephanensis* KBAB4 and at 20°C as compared to 45°C for *B. licheniformis*
372 AD978. Although the maximum sporulation rate decreased with the sporulation
373 temperature, such trend was not observed with the decrease of pH. The ratios between
374 maximum sporulation rates at sub-optimum temperature and reference temperature were
375 calculated to assess the relative effect of environmental factors on sporulation rates (Ross
376 and Dalgaard, 2004) ($\mu_{S10^\circ C}/\mu_{S30^\circ C}$ for *B. weihenstephanensis* for instance) (Table 3).
377 These ratios were compared to the ratios of maximum growth rates, estimated at the same
378 temperatures from the growth cardinal model (Equation 1) and from cardinal growth
379 values (Table 1) ($\mu_{G10^\circ C}/\mu_{G30^\circ C}$). For *B. weihenstephanensis* KBAB4 cells, the ratio of
380 maximum growth rates was evaluated at 0.12 for temperature, and that of sporulation
381 rates was evaluated at 0.08. The ratios of maximum growth rates and sporulation rates
382 were also calculated for the pH. For *B. weihenstephanensis* KBAB4, these ratios were
383 respectively evaluated at 0.82 ($\mu_{GpH5.9}/\mu_{GpH7.2}$), and at 0.83 ($\mu_{SpH5.9}/\mu_{SpH7.2}$). For *B.*
384 *licheniformis* AD978, same trends were observed with a lower growth ratio

385 ($\mu_{GpH6.3}/\mu_{GpH7.2}=0.92$) than sporulation ratio ($(\mu_{SpH6.3}/\mu_{SpH7.2}=1.49)$). The slope of the
386 linear regression between the ratio on μ_S values and the ratio on μ_G values was not
387 significantly different of 1 (t-test, $p > 0.1$). Even though a model validation at
388 intermediate growth and sporulation conditions is certainly necessary, these results
389 suggest that in a general way the quantitative influence of temperature and pH on growth
390 and sporulation are highly similar. As a consequence, as a first approximation, the
391 knowledge of factors determining growth of spore forming bacteria may also be used for
392 the determination of environmental conditions favorable to spore formation and for the
393 determination of sporulation rates.

394 Food processes may induce prolonged residence times of food material and therefore a
395 higher exposure probability of spore-forming bacteria to environments potentially
396 favourable to cells multiplication or spore formation in-line (Oomes *et al.*, 2007). To
397 prevent spore formation, it is recommended to limit the niches where growth of *Bacillus*
398 *sp.* could occur. Determining sporulation kinetics, either by specific experiments or from
399 growth parameters (cardinal growth temperatures, pH or a_w for instance) may contribute
400 to the identification of critical points in processing lines, and finally to a better control of
401 spore-forming bacteria in foods.

402

403 **Acknowledgments**

404 This work was supported by the Agence Nationale de la Recherche (ANR) (France) as
405 part of an ANR-07-PNRA-027-07 MEMOSPORE contract, by the industrial association
406 BBA (Bretagne Biotechnologies Alimentaires) and by the French National Association of
407 the Technical Research (ANRT).

408

409 **References**

- 410 Achard, C., Gros, J. B., Dussap, C. G. (1992). Prédiction de l'activité de l'eau, des températures
411 d'ébullition et de congélation de solutions aqueuses de sucres par un modèle UNIFAC. *Industries*
412 *Alimentaires et Agricoles*, 109, 93-101.
- 413
414 Anonymous (2005). Opinion of the scientific panel on biological hazards on *Bacillus cereus* and
415 other *Bacillus spp.* in foodstuffs. *The EFSA Journal*, 175, 1-48.
- 416
417 Auchtung, J. M., Grossman, A. D. (2008). Extracellular peptide signaling and quorum responses
418 in development, self-recognition and horizontal transfer in *B. subtilis*. In *Chemical*
419 *communication among bacteria*, pp. 13-30. Edited by S. C. Winans and B. L. Bassler.
420 Washington: ASM Press.
- 421
422 Augustin, J.-C., Carlier, V. (2000). Modelling the growth rate of *Listeria monocytogenes* with a
423 multiplicative type model including interactions between environmental factors. *International*
424 *Journal of Food Microbiology*, 56, 53-70.
- 425
426 Baril, E., Coroller, L., Couvert, O., Leguerinel, I., Postollec, F., Boulais, C., Carlin, F., Mafart, P.
427 (2012). Modeling heat resistance of *Bacillus weihenstephanensis* and *Bacillus licheniformis*
428 spores as function of sporulation temperature and pH. *Food Microbiology*, 30, 29-36.
- 429
430 Baril, E., Coroller, L., Postollec, F., Leguerinel, I., Boulais, C., Carlin, F., Mafart, P. (2011). The
431 wet-heat resistance of *Bacillus weihenstephanensis* KBAB4 spores produced in a two-step
432 sporulation process depends on sporulation temperature but not on previous cell history.
433 *International Journal of Food Microbiology*, 146, 57-62.
- 434
435 Baweja, R., Zaman, M., Mattoo, A., Sharma, K., Tripathi, V., Aggarwal, A., Dubey, G., Kurupati,
436 R., Ganguli, M., Chaudhury, N., Sen, S., Das, T., Gade, W., Singh, Y. (2008). Properties of
437 *Bacillus anthracis* spores prepared under various environmental conditions. *Archives of*
438 *Microbiology*, 189, 71-79.
- 439
440 Biesta-Peters, E. G., Reij, M. W., Joosten, H., Gorris, L. G. M., Zwietering, M. H. (2010).
441 Comparison of two optical-density-based methods and a plate count method for estimation of
442 growth parameters of *Bacillus cereus*. *Applied and Environmental Microbiology*, 76, 1399-1405.
- 443
444 Burkholder, W. F., Grossman, A. D. (2000). Regulation of the initiation of endospore formation
445 in *Bacillus subtilis*. In *Prokaryotic development*, pp. 151-166. Edited by Y. V. Brun and L. J.
446 Shimkets. ASM Press: Washington, D.C.
- 447
448 Carlin, F. (2011). Origin of bacterial spores contaminating foods. *Food Microbiology*, 28, 177-
449 182.
- 450
451 Craven, S. E. (1990). The effect of the pH of the sporulation environment on the heat resistance
452 of *Clostridium perfringens* spores. *Current Microbiology*, 20, 233-237.
- 453
454 Cuppers, H., Smelt, J. (1993). Time to turbidity measurement as a tool for modeling spoilage by
455 *Lactobacillus*. *Journal of Industrial Microbiology*, 12, 168-171.

- 456
457 De Jonghe, V., Coorevits, A., De Block, J., Van Coillie, E., Grijspeerdt, K., Herman, L., De Vos,
458 P., Heyndrickx, M. (2010). Toxinogenic and spoilage potential of aerobic spore-formers isolated
459 from raw milk. *International Journal of Food Microbiology*, 136, 318-325.
- 460
461 De Pieri, L. A., Ludlow, I. K. (1992). Relationship between *Bacillus sphaericus* spore heat
462 resistance and sporulation temperature. *Letters in Applied Microbiology*, 14, 121-124.
- 463
464 Efron, B., Tibshirani, R. J. (1993). An introduction to the bootstrap. London: Chapman & Hall.
- 465
466 Gaillard, S., Leguerinel, I., Mafart, P. (1998). Model for combined effects of temperature, pH and
467 water activity on thermal inactivation of *Bacillus cereus* spores. *Journal of Food Science*, 63,
468 887-889.
- 469
470 Garcia, D., Van der Voort, M., Abee, T. (2010). Comparative analysis of *Bacillus*
471 *weihenstephanensis* KBAB4 spores obtained at different temperatures. *International Journal of*
472 *Food Microbiology*, 140, 146-153.
- 473
474 Gauchi, J. P., Vila, J. P., Coroller, L. (2010). New prediction interval and band in the nonlinear
475 regression model : Application to predictive modeling in Foods. *Communication in Statistics -*
476 *Simulation and Computation*, 39, 322-334.
- 477
478 Gonzalez, I., Lopez, M., Martinez, S., Bernardo, A., Gonzalez, J. (1999). Thermal inactivation of
479 *Bacillus cereus* spores formed at different temperatures. *International Journal of Food*
480 *Microbiology*, 51, 81-84.
- 481
482 Grossman, A. D. (1995). Genetic networks controlling the initiation of sporulation and the
483 development of genetic competence in *Bacillus subtilis*. *Annual Review of Genetics*, 29, 477-508.
- 484
485 Guinebretiere, M.-H., Thompson, F. L., Sorokin, A., Normand, P., Dawyndt, P., Ehling-Schulz,
486 M., Svensson, B., Sanchis, V., Nguyen-The, C., Heyndrickx, M., De Vos, P. (2008). Ecological
487 diversification in the *Bacillus cereus* Group. *Environmental Microbiology*, 10, 851-865.
- 488
489 Jakobsen, T., M., Murrell, W. G. (1977). The effect of water activity and a_w -controlling solute on
490 sporulation of *Bacillus cereus*. *Journal of Applied Microbiology*, 43, 239-245.
- 491
492 Le Marc, Y., Huchet, V., Bourgeois, C. M., Guyonnet, J. P., Mafart, P., Thuault, D. (2002).
493 Modelling the growth kinetics of *Listeria* as a function of temperature, pH and organic acid
494 concentration. *International Journal of Food Microbiology*, 73, 219-237.
- 495
496 Lechowich, R. V., Ordal, Z. J. (1962). The influence of the sporulation temperature on the heat
497 resistance and chemical composition of bacterial spores. *Canadian Journal of Microbiology* 8,
498 287-295.
- 499
500 Mazas, M., Lopez, M., Gonzalez, I., Bernardo, A., Martin, R. (1997). Effects of sporulation pH
501 on the heat resistance and the sporulation of *Bacillus cereus*. *Letters in Applied Microbiology*, 25,
502 331-334.
- 503
504 Meer, R. R., Baker, J., Bodyfelt, F. W., Griffiths, M. W. (1991). Psychotrophic *Bacillus spp.* in
505 fluid milk products: a review. *Journal of Food Protection*, 54, 969-979.

- 506
507 Membré, J.-M., Leporq, B., Vialette, M., Mettler, E., Perrier, L., Thuault, D., Zwietering, M.
508 (2005). Temperature effect on bacterial growth rate: quantitative microbiology approach
509 including cardinal values and variability estimates to perform growth simulations on/in food.
510 *International Journal of Food Microbiology*, 100, 179-186.
511
- 512 Membré, J. M., Leporc, B., Vialette, M., Mettler, E., Perrier, L., Zwietering, M. H. (2002).
513 Experimental protocols and strain variability of cardinal values (pH and aw) of bacteria using
514 Bioscreen C: microbial and statistical aspects. In *Microbial adaptation to changing environments*,
515 pp. 143-146. Edited by L. Alexon, E. S. Tronrud and K. J. Merok. Matforsk Norwegian Food
516 Research Institute, Lillehammer, Norway.
517
- 518 Nguyen Thi Minh, H., Perrier-Cornet, J.-M., Gervais, P. (2008). Effect of the osmotic conditions
519 during sporulation on the subsequent resistance of bacterial spores. *Applied Microbiology and*
520 *Biotechnology*, 80, 107.
521
- 522 Nguyen Thi Minh, H., Durand, A., Loison, P., Perrier-Cornet, J.-M., Gervais, P. (2011). Effect of
523 sporulation conditions on the resistance of *Bacillus subtilis* spores to heat and high pressure.
524 *Applied Microbiology and Biotechnology*, 90, 1409-1417.
525
- 526 Oomes, S. J. C. M., van Zuijlen, A., Hehenkamp, J. O., Witsenboer, H., Van der Vossen, J., Brul,
527 S. (2007). The characterisation of *Bacillus* spores occurring in the manufacturing of (low acid)
528 canned products. *International Journal of Food Microbiology*, 120, 85-94.
529
- 530 Ross, R., Dalgaard, P. (2004). Secondary models. In *Modeling microbial responses in food*, pp.
531 63-150. Edited by R. C. Mc Kellar and X. Lu. Washington D. C.: CRC Press.
532
- 533 Ross, T., Olley, J., McMeekin, T. A., Ratkowsky, D. A. (2011). Letter to the Editor. *International*
534 *Journal of Food Microbiology*, 147, 78-80.
535
- 536 Rosso, L., Lobry, J. R., Bajard, S., Flandrois, J. P. (1995). Convenient model to describe the
537 combined effects of temperature and pH on microbial growth. *Applied and Environmental*
538 *Microbiology*, 61, 610-616.
539
- 540 Scott, S. A., Brooks, J. D., Rakonjac, J., Walker, K. M. R., Flint, S. H. (2007). The formation of
541 thermophilic spores during the manufacture of whole milk powder. *International Journal of*
542 *Dairy Technology*, 60, 109-117.
543
- 544 Sonenshein, A. L. (1999). Endospore-forming bacteria: an overview. In *Prokaryotic*
545 *development*, pp. 133-150. Edited by Y. V. Brun and L. J. Shimkets. ASM Press: Washington,
546 D.C.
547
- 548 Stenfors Arnesen, L. P., Fagerlund, A., Granum, P. E. (2008). From soil to gut: *Bacillus cereus*
549 and its food poisoning toxins. *FEMS Microbiology Reviews*, 32, 579-606.
550
- 551 Tam, N. K. M., Uyen, N. Q., Hong, H. A., Duc, L. H., Hoa, T. T., Serra, C. R., Henriques, A. O.,
552 Cutting, S. M. (2006). The intestinal life cycle of *Bacillus subtilis* and close relatives. *Journal of*
553 *Bacteriology*, 188, 2692-2700.
554

- 555 Vilas-Boas, G., Sanchis, V., Lereclus, D., Lemos, M. V. F., Bourguet, D. (2002). Genetic
556 differentiation between sympatric populations of *Bacillus cereus* and *Bacillus thuringiensis*.
557 *Applied and Environmental Microbiology*, 68, 1414-1424.
558
- 559 Yazdany, S., Lashkari, K. B. (1975). Effect of pH on sporulation of *Bacillus stearothermophilus*.
560 *Applied microbiology*, 30, 1-3.

ACCEPTED MANUSCRIPT

561

562 **Figures**

563 Figure 1. Effect of temperature, pH and a_w on the maximum growth rate of *B.*
564 *weihenstephanensis* KBAB4 (A, C, E) and on *B. licheniformis* AD978 (B, D, F). Filled
565 symbols correspond to growth observation. Empty symbols correspond to no growth. The
566 lines correspond to the estimation of maximum growth rate by the equation 1, which was
567 fitted only on growth observation.

568

569 Figure 2. Effect of sporulation temperature, pH and a_w on the final spore concentration of
570 *B. weihenstephanensis* KBAB4 strain (A, C, E) and on *B. licheniformis* AD978 strain (B,
571 D, F). Filled symbols correspond to sporulation in SMB, empty symbols correspond to
572 sporulation in MNA. The dashed lines correspond to the limit of quantification (2.6
573 \log_{10} CFU/mL). Symbols under these lines correspond to undetectable spore
574 concentrations.

575

576 Figure 3. Influence of the sporulation temperature and pH on sporulation kinetics of *B.*
577 *weihenstephanensis* (A, C,) and *B. licheniformis* (B, D) in SMB. (A) Sporulation of *B.*
578 *weihenstephanensis* at 10°C (filled symbols) and 30°C (empty symbols). (C) Sporulation
579 of *B. weihenstephanensis* at pH 5.9 (filled symbols) and pH 7.2 (empty symbols). (B)
580 Sporulation of *B. licheniformis* at 20°C (filled symbols) and 45°C (empty symbols). (D)
581 Sporulation of *B. licheniformis* at pH 6.3 (filled symbols) and pH 7.2 (empty symbols).
582 Bars represent standard deviation of the three independent triplicates. Lines correspond to
583 the estimated spore concentrations from the sporulation kinetic model (Equation 3).
584 Dashed lines correspond to the limit of quantification.

- 1 Table 1. Estimated cardinal growth parameters of *B. weihenstephanensis* KBAB4 and *B.*
 2 *licheniformis* AD978 and quality of fit criterions.

	<i>B. weihenstephanensis</i> KBAB4	<i>B. licheniformis</i> AD978
μ_{max}	1.68 [1.52 ; 2.19]*	3.12 [2.77 ; 3.90]
T_{min}	0.9 [-2.5 ; 5.9]	8.6 [2.7 ; 19.4]
T_{opt}	33.1 [31.9 ; 33.9]	50.9 [49.2 ; 52.6]
T_{max}	38.6 [37.8 ; 43.0]	56.1 [55.3 ; 68.9]
pH_{min}	4.6 [4.5 ; 4.8]	5.3 [5.2 ; 5.3]
pH_{opt}	6.9 [6.7 ; 7.1]	6.9 [6.7 ; 7.1]
a_{wmin}	0.946 [0.943 ; 0.950]	0.915 [0.908 ; 0.927]
a_{wopt}	1.000 [0.995 ; 1.000]	1.000 [0.994 ; 1.000]
Number of data**	45	48
RMSE	0.073	0.190

- 3 * Estimated value [confidence interval of the estimated parameter value at 95%]
 4 ** Number of data fitted to the model. Data corresponding to no-growth were excluded to
 5 the fit.

- 1 Table 2. Estimated sporulation kinetic parameters of *B. weihenstephanensis* KBAB4 and of *B. licheniformis* AD978 and quality of fit
 2 criteria (Equation 3).

	<i>B. weihenstephanensis</i> KBAB4			<i>B. licheniformis</i> AD978		
Temperature	10°C	30°C*	30°C	20°C	45°C*	45°C
pH	7.2	7.2*	5.9	7.2	7.2*	6.3
t_{1S}	148.0 [140.7 ; 155.3]** ^a	6.7 [6.3 ; 7.0] ^b	14.5 [13.6 ; 15.3] ^c	45.9 [34.2 ; 57.5] ^d	1.6 [1.2 ; 2.1] ^e	4.2 [3.7 ; 4.7] ^f
μ_S	0.05 [0.04 ; 0.07] ^a	0.60 [0.50 ; 0.70] ^b	0.50 [0.35 ; 0.64] ^b	0.02 [0.02 ; 0.03] ^d	0.37 [0.32 ; 0.42] ^e	0.55 [0.45 ; 0.66] ^f
$\text{Log}_{10}(N_{SF})$	6.5 [6.3 ; 6.7] ^a	7.5 [7.3 ; 7.6] ^b	7.7 [7.4 ; 8.0] ^b	6.8 [6.6 ; 7.1] ^d	7.1 [6.9 ; 7.3] ^d	5.5 [5.3 ; 5.6] ^e
Number of data	22	51	26	54	89	84
RMSE	0.380	0.371	0.373	0.226	0.549	0.328

3 * Reference sporulation temperature and pH


4 ** Estimated value [confidence interval of the estimated parameter value at 95%]


5 ^{a-f} For each line and each strain, the same superscript letter indicates that the parameter values are not significantly different ($p > 0.05$)

- 1 Table 3. Ratio of maximum growth rates and sporulation rates of *B. weihenstephanensis*
- 2 KBAB4 and *B. licheniformis* AD978 at different temperatures and pH.





	<i>B. weihenstephanensis</i> KBAB4		<i>B. licheniformis</i> AD978	
	$\mu_{10^{\circ}\text{C}} / \mu_{30^{\circ}\text{C}}$	$\mu_{\text{pH } 5.9} / \mu_{\text{pH } 7.2}$	$\mu_{20^{\circ}\text{C}} / \mu_{45^{\circ}\text{C}}$	$\mu_{\text{pH } 6.3} / \mu_{\text{pH } 7.2}$
Growth	0.12 [0.05; 0.17]*	0.82 [0.78; 0.87]	0.11 [0.00; 0.17]	0.92 [0.82; 1.02]
Sporulation	0.08	0.83	0.05	1.49

- 3 * Estimated ratio [confidence interval of the estimated ratio value at 95%]





A



ACCEPTED

1 **Research highlights**

2 - Temperature, pH and a_w growth limits are reasonable predictions of sporulation

3 boundaries

4 - Temperature and pH determined time to first spore

5 - Temperature and pH determined final spore concentration

6 - Sporulation rates and growth rates are similarly affected by environmental factors

7

ACCEPTED MANUSCRIPT