



HAL
open science

NEW FORMULAS FOR THE SPECTRAL RADIUS VIA ALUTHGE TRANSFORM

Fadil Chabbabi, Mostafa Mbekhta

► **To cite this version:**

Fadil Chabbabi, Mostafa Mbekhta. NEW FORMULAS FOR THE SPECTRAL RADIUS VIA ALUTHGE TRANSFORM. 2016. hal-01334044

HAL Id: hal-01334044

<https://hal.science/hal-01334044>

Preprint submitted on 20 Jun 2016

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

NEW FORMULAS FOR THE SPECTRAL RADIUS VIA ALUTHGE TRANSFORM

FADIL CHABBABI AND MOSTAFA MBEKHTA

ABSTRACT. In this paper we give several expressions of spectral radius of a bounded operator on a Hilbert space, in terms of iterates of Aluthge transformation, numerical radius and the asymptotic behavior of the powers of this operator.

Also we obtain several characterizations of normaloid operators.

1. INTRODUCTION

Let H be complex Hilbert spaces and $\mathcal{B}(H)$ be the Banach space of all bounded linear operators from H into it self.

For $T \in \mathcal{B}(H)$, the spectrum of T is denoted by $\sigma(T)$ and $r(T)$ its spectral radius. We denote also by $W(T)$ and $w(T)$ the numerical range and the numerical radius of T .

As usually, for $T \in \mathcal{B}(H)$ we denote the module of T by $|T| = (T^*T)^{1/2}$ and we shall always write, without further mention, $T = U|T|$ to be the unique polar decomposition of T , where U is the appropriate partial isometry satisfying $\mathcal{N}(U) = \mathcal{N}(T)$. The Aluthge transform introduced in [1] as

$$\Delta(T) = |T|^{\frac{1}{2}}U|T|^{\frac{1}{2}}, \quad T \in \mathcal{B}(H),$$

to extend some properties of hyponormal operators. Later, in [8], Okubo introduced a more general notion called λ -Aluthge transform which has also been studied in detail.

For $\lambda \in [0, 1]$, the λ -Aluthge transform is defined by,

$$\Delta_\lambda(T) = |T|^\lambda U |T|^{1-\lambda}, \quad T \in \mathcal{B}(H).$$

Notice that $\Delta_0(T) = U|T| = T$, and $\Delta_1(T) = |T|U$ which is known as Duggal's transform. It has since been studied in many different contexts and considered by a number of authors (see for instance, [1, 2, 3, 7, 6, 5] and some of the references there). The interest of the Aluthge transform lies in the fact that it respects many properties of the original operator. For example, (see [5, Theorems 1.3, 1.5])

$$(1.1) \quad \sigma(\Delta_\lambda(T)) = \sigma(T), \text{ for every } T \in \mathcal{B}(H),$$

Another important property is that $\text{Lat}(T)$, the lattice of T -invariant subspaces of H , is nontrivial if and only if $\text{Lat}(\Delta(T))$ is nontrivial (see [5, Theorem 1.15]).

2000 *Mathematics Subject Classification.* 47A13, 47A30, 47B37.

Key words and phrases. spectral radius, polar decomposition, λ -Aluthge transform, normaloid operator

This work was supported in part by the Labex CEMPI (ANR-11-LABX-0007-01).

Moreover, Yamazaki ([12]) (see also, [11, 10]), established the following interesting formula for the spectral radius

$$(1.2) \quad \lim_{n \rightarrow \infty} \|\Delta_\lambda^n(T)\| = r(T)$$

where Δ_λ^n the n -th iterate of Δ_λ , i.e $\Delta_\lambda^{n+1}(T) = \Delta_\lambda(\Delta_\lambda^n(T))$, $\Delta_\lambda^0(T) = T$.

In this paper we give several expressions of the spectral radius of an operator. Firstly in terms of the Aluthge transformation (section 2), and secondly, in section 3, we give several expressions of spectral radius, based on numerical radius and Aluthge transformation. Also, We infer several characterizations of normaloid operators (i.e. $r(T) = \|T\|$).

2. FORMULAS OF SPECTRAL RADIUS VIA ALUTHGE TRANSFORM

In this section, we use the of Rota's Theorem, order to obtain new formulas of spectral radius via Aluthge transformation

Theorem 2.1. For every operator $T \in \mathcal{B}(H)$, we have

$$\begin{aligned} r(T) &= \inf\{\|\Delta_\lambda(XTX^{-1})\|, X \in \mathcal{B}(H) \text{ invertible}\} \\ &= \inf\{\|\Delta_\lambda(e^A T e^{-A})\|, A \in \mathcal{B}(H) \text{ self adjoint}\}. \end{aligned}$$

Proof. For every invertible operator $X \in \mathcal{B}(H)$, by (1.1), we have

$$\sigma(\Delta_\lambda(XTX^{-1})) = \sigma(XTX^{-1}) = \sigma(T).$$

It follows that

$$r(T) = r(\Delta_\lambda(XTX^{-1})) \leq \|\Delta_\lambda(XTX^{-1})\| \text{ for every invertible operator } X \in \mathcal{B}(H).$$

Hence

$$\begin{aligned} r(T) &\leq \inf\{\|\Delta_\lambda(XTX^{-1})\|; X \in \mathcal{B}(H) \text{ invertible}\} \\ &\leq \inf\{\|\Delta_\lambda(\exp(A)T \exp(-A))\|; A \in \mathcal{B}(H) \text{ self adjoint}\}, \end{aligned}$$

In the other hand, for $\varepsilon > 0$, we have

$$r\left(\frac{T}{r(T) + \varepsilon}\right) = \frac{r(T)}{r(T) + \varepsilon} < 1.$$

From Rota's Theorem [9, Theorem 2], $\frac{T}{r(T) + \varepsilon}$ is similar to a contraction. Thus there exists an invertible operator $X_\varepsilon \in \mathcal{B}(H)$ such that

$$(2.1) \quad \|\Delta_\lambda(X_\varepsilon T X_\varepsilon^{-1})\| \leq \|X_\varepsilon T X_\varepsilon^{-1}\| \leq r(T) + \varepsilon.$$

Now, let $X_\varepsilon = U_\varepsilon |X_\varepsilon|$ be the polar decomposition of X_ε . Clearly U_ε is a unitary operator, and $|X_\varepsilon|$ is invertible. Therefore there exist $\alpha > 0$ such that $\sigma(|X_\varepsilon|) \subseteq [\alpha, +\infty[$. Consequently $A_\varepsilon = \ln(|X_\varepsilon|)$ exists and it is self adjoint, we also have

$$|X_\varepsilon| = e^{A_\varepsilon} \text{ and } |X_\varepsilon|^{-1} = e^{-A_\varepsilon}.$$

Thus

$$\begin{aligned}\|\Delta_\lambda(X_\varepsilon T X_\varepsilon^{-1})\| &= \|\Delta_\lambda(U e^{A_\varepsilon} T e^{-A_\varepsilon} U^*)\| \\ &= \|U \Delta_\lambda(e^{A_\varepsilon} T e^{-A_\varepsilon}) U^*\| \\ &= \|\Delta_\lambda(e^{A_\varepsilon} T e^{-A_\varepsilon})\|.\end{aligned}$$

Hence

$$\|\Delta_\lambda(e^{A_\varepsilon} T e^{-A_\varepsilon})\| \leq \|X_\varepsilon T X_\varepsilon^{-1}\| \leq r(T) + \varepsilon.$$

It follows that for all $\varepsilon > 0$,

$$\begin{aligned}r(T) &\leq \inf\{\|\Delta_\lambda(X T X^{-1})\|, X \in \mathcal{B}(H) \text{ invertible}\} \\ &\leq \inf\{\|\Delta_\lambda(e^A T e^{-A})\|, A \in \mathcal{B}(H) \text{ self adjoint}\} \\ &\leq \|\Delta_\lambda(e^{A_\varepsilon} T e^{-A_\varepsilon})\| \leq \|X_\varepsilon T X_\varepsilon^{-1}\| \\ &\leq r(T) + \varepsilon.\end{aligned}$$

Finally, since $\varepsilon > 0$ is arbitrary, we obtain

$$\begin{aligned}r(T) &= \inf\{\|\Delta_\lambda(X T X^{-1})\|, X \in \mathcal{B}(H) \text{ invertible}\} \\ &= \inf\{\|\Delta_\lambda(e^A T e^{-A})\|, A \in \mathcal{B}(H) \text{ self adjoint}\}.\end{aligned}$$

Therefore the proof of Theorem is complete. \square

As immediate consequence of the Theorem 2.1, we obtain the following corollary which gives a formula of the spectral radius based on n -th iterate of Δ_λ .

Corollary 2.1. If $T \in \mathcal{B}(H)$, then for every $n \geq 0$,

$$\begin{aligned}r(T) &= \inf\{\|\Delta_\lambda^n(X T X^{-1})\|, X \in \mathcal{B}(H) \text{ invertible}\} \\ &= \inf\{\|\Delta_\lambda^n(e^A T e^{-A})\|, A \in \mathcal{B}(H) \text{ self adjoint}\}.\end{aligned}$$

Proof. First, note that $\|\Delta_\lambda(T)\| \leq \|T\|$, consequently we have

$$(2.2) \quad \|\Delta_\lambda^n(T)\| \leq \|\Delta_\lambda^{n-1}(T)\| \leq \dots \leq \|\Delta_\lambda(T)\| \leq \|T\|, \forall n \in \mathbb{N}^*.$$

Now, clearly $\sigma(\Delta_\lambda^n(T)) = \sigma(T)$, for all $n \in \mathbb{N}$. It follows that, for every invertible operator $X \in \mathcal{B}(H)$ we have

$$\begin{aligned}r(T) &= r(\Delta_\lambda^n(X T X^{-1})) \\ &\leq \|\Delta_\lambda^n(X T X^{-1})\| \\ &\leq \|\Delta_\lambda(X T X^{-1})\|.\end{aligned}$$

Therefore

$$\begin{aligned}r(T) &\leq \inf\{\|\Delta_\lambda^n(X T X^{-1})\|, X \in \mathcal{B}(H) \text{ invertible}\} \\ &\leq \inf\{\|\Delta_\lambda^n(e^A T e^{-A})\|, A \in \mathcal{B}(H) \text{ self adjoint}\} \\ &\leq \inf\{\|\Delta_\lambda(e^A T e^{-A})\|, A \in \mathcal{B}(H) \text{ self adjoint}\} \\ &= r(T).\end{aligned}$$

\square

An operator T is said to be normaloid if $r(T) = \|T\|$.

As immediate consequence of the Corollary 2.1, we obtain the following corollary which is a characterization of normaloid operators via λ -Aluthge transformation :

Corollary 2.2. If $T \in \mathcal{B}(H)$, then the following assertions are equivalent

- (i) T is normaloid;
- (ii) $\|T\| \leq \|\Delta_\lambda(XTX^{-1})\|$, for all invertible $X \in \mathcal{B}(H)$;
- (iii) $\|T\| \leq \|\Delta_\lambda^n(XTX^{-1})\|$, for all invertible $X \in \mathcal{B}(H)$ and for all natural number n .

As immediate consequence of the Corollary 2.2, we obtain a new characterization of normaloid operators

Corollary 2.3. If $T \in \mathcal{B}(H)$, then the following assertions are equivalent

- (i) T is normaloid;
- (ii) $\|T\| \leq \|XTX^{-1}\|$, for all invertible $X \in \mathcal{B}(H)$;

Theorem 2.2. Let $T \in \mathcal{B}(H)$. Then for each natural number n , we have

$$\begin{aligned} r(T) &= \lim_k \|\Delta_\lambda^n(T^k)\|^{1/k} \\ &= \lim_k \|\Delta_\lambda(T^k)\|^{1/k}. \end{aligned}$$

Proof. Note that,

$$(2.3) \quad r(T) = r(\Delta_\lambda^n(T)) \leq \|\Delta_\lambda^n(T)\| \leq \|\Delta_\lambda(T)\| \leq \|T\| \quad \forall n \in \mathbb{N}^*.$$

Let $k \in \mathbb{N}$ be arbitrary, we have

$$r(T)^k = r(T^k) = r(\Delta_\lambda^n(T^k)) \leq \|\Delta_\lambda^n(T^k)\| \leq \|\Delta_\lambda(T^k)\| \leq \|T^k\| \quad \forall n \in \mathbb{N}.$$

Hence

$$r(T) \leq \|\Delta_\lambda^n(T^k)\|^{1/k} \leq \|\Delta_\lambda(T^k)\|^{1/k} \leq \|T^k\|^{1/k}.$$

Therefore

$$r(T) \leq \lim_k \|\Delta_\lambda^n(T^k)\|^{1/k} \leq \lim_k \|\Delta_\lambda(T^k)\|^{1/k} \leq \lim_k \|T^k\|^{1/k} = r(T).$$

Which completes the proof. □

As immediate consequence of Theorem 2.2, we obtain the following corollary which is a new characterization of normaloid operators

Corollary 2.4. If $T \in \mathcal{B}(H)$, then the following assertions are equivalent

- (i) T is normaloid;
- (ii) $\|T\|^k = \|\Delta_\lambda(T^k)\|$, for all natural number k ;
- (iii) $\|T\|^k = \|\Delta_\lambda^n(T^k)\|$, for every natural number k, n

3. SPECTRAL RADIUS VIA NUMERICAL RADIUS AND ALUTHGE TRANSFORM

For $T \in \mathcal{B}(H)$, we denote the numerical range and numerical radius of T by $W(T)$ and $w(T)$, respectively.

$$W(T) = \{ \langle Tx, x \rangle; \|x\| = 1 \} \quad \text{and} \quad w(T) = \sup\{|\lambda|; \lambda \in W(T)\}.$$

In the following theorem, we obtain a new expression of the spectral radius by means of the numerical radius and Aluthge transform.

Theorem 3.1. For every operator $T \in \mathcal{B}(H)$ and for each natural number n , we have

$$\begin{aligned} r(T) &= \inf\{w(\Delta_\lambda^n(XTX^{-1})), X \in \mathcal{B}(H) \text{ invertible} \} \\ &= \inf\{w(\Delta_\lambda^n(e^A T e^{-A})), A \in \mathcal{B}(H) \text{ self adjoint} \}. \end{aligned}$$

Proof. It is well known that $r(T) \leq w(T) \leq \|T\|$. Thus, for all $X \in \mathcal{B}(H)$, invertible and for each natural number n , we have

$$r(T) = r(\Delta_\lambda^n(XTX^{-1})) \leq w(\Delta_\lambda^n(XTX^{-1})) \leq \|\Delta_\lambda^n(XTX^{-1})\|$$

It follows that

$$\begin{aligned} r(T) &\leq \inf\{w(\Delta_\lambda^n(XTX^{-1})); X \in \mathcal{B}(H) \text{ invertible} \} \\ &\leq \inf\{w(\Delta_\lambda^n(\exp(A)T \exp(-A))); A \in \mathcal{B}(H) \text{ self adjoint} \}, \\ &\leq \inf\{\|\Delta_\lambda^n(\exp(A)T \exp(-A))\|; A \in \mathcal{B}(H) \text{ self adjoint} \} \\ &= r(T) \quad (\text{by Corollary 2.1}). \end{aligned}$$

Hence we obtain the desired equalities. \square

For a bounded linear operator S , we will write $\mathcal{R}e(S) = \frac{1}{2}(S + S^*)$, the real part of S . And we denote by $\overline{W}(S)$ the closure of the numerical range of S . Then we have the following result

Theorem 3.2. For every operator $T \in \mathcal{B}(H)$, there exists $\theta \in \mathbb{R}$ such that for all natural number n ,

$$\begin{aligned} r(T) &= \inf\{w(\mathcal{R}e(\Delta_\lambda^n(\exp(i\theta)XTX^{-1}))), X \in \mathcal{B}(H) \text{ invertible} \} \\ &= \inf\{\|\mathcal{R}e(\Delta_\lambda^n(\exp(i\theta)XTX^{-1}))\|, X \in \mathcal{B}(H) \text{ invertible} \}. \end{aligned}$$

Proof. First assume that $r(T) \in \sigma(T)$. Then for all invertible $X \in \mathcal{B}(H)$, we have

$$r(T) \in \mathcal{R}e(\sigma(T)) = \mathcal{R}e(\sigma(\Delta_\lambda^n(XTX^{-1}))).$$

Thus

$$r(T) \in \mathcal{R}e(\sigma(\Delta_\lambda^n(XTX^{-1}))) \subseteq \mathcal{R}e(\overline{W}(\Delta_\lambda^n(XTX^{-1}))) = \overline{W}(\mathcal{R}e(\Delta_\lambda^n(XTX^{-1})))$$

which implies

$$\begin{aligned} r(T) &\leq w(\mathcal{R}e(\Delta_\lambda^n(XTX^{-1}))) \\ &\leq \|\mathcal{R}e(\Delta_\lambda^n(XTX^{-1}))\| \\ &\leq \|\Delta_\lambda^n(XTX^{-1})\|. \end{aligned}$$

Since the last inequalities are satisfied for all $X \in \mathcal{B}(H)$ invertible, we obtain

$$\begin{aligned} r(T) &\leq \inf\{w(\mathcal{R}e(\Delta_\lambda^n(XTX^{-1}))) \mid X \in \mathcal{B}(H) \text{ invertible}\} \\ &\leq \inf\{\|\mathcal{R}e(\Delta_\lambda^n(XTX^{-1}))\| \mid X \in \mathcal{B}(H) \text{ invertible}\} \\ &\leq \inf\{\|\Delta_\lambda^n(XTX^{-1})\| \mid X \in \mathcal{B}(H) \text{ invertible}\} \\ &= r(T) \quad (\text{by Corollary 2.1}). \end{aligned}$$

We have shown that if $r(T) \in \sigma(T)$ then

$$\begin{aligned} r(T) &= \inf\{w(\mathcal{R}e(\Delta_\lambda^n(XTX^{-1}))) \mid X \in \mathcal{B}(H) \text{ invertible}\} \\ &= \inf\{\|\mathcal{R}e(\Delta_\lambda^n(XTX^{-1}))\| \mid X \in \mathcal{B}(H) \text{ invertible}\} \end{aligned}$$

Now, if T is an arbitrary operator, then there exists $z \in \sigma(T)$ such that, $|z| = r(T)$. Put $\theta = -\arg(z)$. Then $r(T) = z \exp(i\theta) \in \sigma(\exp(i\theta)T)$. Hence by the first part of de proof, we conclude that

$$\begin{aligned} r(T) = r(\exp(i\theta)T) &= \inf\{w(\mathcal{R}e(\Delta_\lambda^n(\exp(i\theta)XTX^{-1}))), X \in \mathcal{B}(H) \text{ invertible}\} \\ &= \inf\{\|\mathcal{R}e(\Delta_\lambda^n(\exp(i\theta)XTX^{-1}))\|, X \in \mathcal{B}(H) \text{ invertible}\}. \end{aligned}$$

This completes the proof of the theorem. \square

As immediate consequence of Theorem 3.2, we obtain the following corollary which is a characterization of normaloid operators

Corollary 3.1. If $T \in \mathcal{B}(H)$, and a natural number n , then the following assertions are equivalent

- (i) T is normaloid;
- (ii) there exists $\theta \in \mathbb{R}$ such that, for all $X \in \mathcal{B}(H)$ invertible

$$\|T\| \leq w(\mathcal{R}e(\Delta_\lambda^n(\exp(i\theta)XTX^{-1})));$$

- (iii) there exists $\theta \in \mathbb{R}$ such that, for all $X \in \mathcal{B}(H)$ invertible

$$\|T\| \leq \|\mathcal{R}e(\Delta_\lambda^n(\exp(i\theta)XTX^{-1}))\|.$$

We end this paper by the following theorem which gives a new formula of the spectral radius of T , in terms of the asymptotic behavior of powers and the numerical radius of T

Theorem 3.3. For every operator $T \in \mathcal{B}(H)$ and for each natural number n , we have

$$r(T) = \lim_k w(\Delta_\lambda^n(T^k))^{1/k}.$$

Proof. For each natural number n and k , we have

$$r(T)^k = r(T^k) = r(\Delta_\lambda^n(T^k)) \leq w(\Delta_\lambda^n(T^k)) \leq \|\Delta_\lambda^n(T^k)\|$$

Hence

$$r(T) \leq w(\Delta_\lambda^n(T^k))^{1/k} \leq \|\Delta_\lambda^n(T^k)\|^{1/k}.$$

By Theorem 2.2, we deduce that

$$r(T) = \lim_k w(\Delta_\lambda^n(T^k))^{1/k}.$$

Which completes the proof. \square

REFERENCES

- [1] A. ALUTHGE, *On p -hyponormal operators for $0 < p < 1$* , Integral Equations Operator Theory 13 (1990), 307-315.
- [2] T. Ando and T. Yamazaki, *The iterated Aluthge transforms of a 2-by-2 matrix converge*, Linear Algebra Appl. 375 (2003), 299-309
- [3] A.J. ANTEZANA, P. MASSEY AND D. STOJANOFF, *λ -Aluthge transforms and Schatten ideals*, Linear Algebra Appl, 405 (2005), 177-199.
- [4] T. Furuta, *Invitation to linear operators*, Taylor Francis, London 2001.
- [5] I. JUNG, E. KO, AND C. PEARCY , *Aluthge transform of operators*, Integral Equations Operator Theory 37 (2000), 437-448.
- [6] I. JUNG, E. KO, C. PEARCY , *Spectral pictures of Aluthge transforms of operators*, Integral Equations Operator Theory 40 (2001), 52-60.
- [7] I. JUNG, E. KO, C. PEARCY , *The iterated Aluthge transform of an operator*, Integral Equations Operator Theory 45 (2003), 375-387.
- [8] K. OKUBO, *On weakly unitarily invariant norm and the Aluthge transformation*, Linear Algebra Appl. 371 (2003), 369–375.
- [9] G. ROTA, *On models for linear operators*, comm. Pure Appl. Math. 13 (1960), 496–472.
- [10] T. TAM, *λ - Aluthge iteration and spectral radius*, Integral Equations Operator Theory 60 (2008), 591-596.
- [11] D.WANG, *Heinz and McIntosh inequalities, Aluthge transformation and the spectral radius*, Math. Inequal. Appl. 6 (2003), 121-124.
- [12] T. YAMAZAKI, *An expression of the spectral radius via Aluthge transformation*, Proc. Amer. Math. Soc. 130 (2002), 1131-1137.

UNIVERSITÉ LILLE1, UFR DE MATHÉMATIQUES, LABORATOIRE CNRS-UMR 8524 P. PAINLEVÉ, 59655 VILLENEUVE D'ASCQ CEDEX, FRANCE

E-mail address: Fadil.Chabbabi@Math.univ-lille1.fr

E-mail address: Mostafa.Mbekhta@math.univ-lille1.fr