



HAL
open science

Exceedance of critical loads and of critical limits impacts tree nutrition across Europe

Peter Waldner, Anne Thimonier, Elisabeth Graf Pannatier, Sophia Etzold, Maria Schmitt, Aldo Marchetto, Pasi Rautio, Kirsti Derome, Tiina Maileena Nieminen, Seppo Nevalainen, et al.

► To cite this version:

Peter Waldner, Anne Thimonier, Elisabeth Graf Pannatier, Sophia Etzold, Maria Schmitt, et al.. Exceedance of critical loads and of critical limits impacts tree nutrition across Europe. *Annals of Forest Science*, 2015, 72 (7), pp.929-939. 10.1007/s13595-015-0489-2. hal-01337297

HAL Id: hal-01337297

<https://hal.science/hal-01337297>

Submitted on 24 Jun 2016

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Exceedance of critical loads and of critical limits impacts tree nutrition across Europe

Peter Waldner¹ · Anne Thimonier¹ · Elisabeth Graf Pannatier¹ · Sophia Etzold¹ · Maria Schmitt¹ · Aldo Marchetto² · Pasi Rautio³ · Kirsti Derome⁴ · Tiina Maileena Nieminen⁵ · Seppo Nevalainen⁶ · Antti-Jussi Lindroos⁵ · Päivi Merilä⁴ · Georg Kindermann⁷ · Markus Neumann⁷ · Nathalie Cools⁸ · Bruno de Vos⁸ · Peter Roskams⁸ · Arne Verstraeten⁸ · Karin Hansen⁹ · Gunilla Pihl Karlsson¹⁰ · Hans-Peter Dietrich¹¹ · Stephan Raspe¹¹ · Richard Fischer¹² · Martin Lorenz¹² · Susanne Iost¹² · Oliver Granke¹³ · Tanja G. M. Sanders¹⁴ · Alexa Michel¹⁴ · Hans-Dieter Nagel¹⁵ · Thomas Scheuschner¹⁵ · Primož Simončič¹⁶ · Klaus von Wilpert¹⁷ · Henning Meesenburg¹⁸ · Stefan Fleck¹⁸ · Sue Benham¹⁹ · Elena Vanguelova¹⁹ · Nicholas Clarke²⁰ · Morten Ingerslev²¹ · Lars Vesterdal²¹ · Per Gundersen²¹ · Inge Stupak²¹ · Mathieu Jonard²² · Nenad Potočić²³ · Mayte Minaya²⁴

Received: 30 June 2014 / Accepted: 25 May 2015 / Published online: 25 June 2015
© The Author(s) 2015. This article is published with open access at Springerlink.com

Abstract

• **Key message** Exceedance of critical limits in soil solution samples was more frequent in intensively monitored forest plots across Europe with critical loads for acidity and eutrophication exceeded compared to other plots from the same network. Elevated inorganic nitrogen concentrations in soil solution tended to be related to less favourable nutritional status.

Handling Editor: Marco Ferretti

Contribution of the co-authors M. Lorenz: Coordination of Futmon and ICP Forests; P. Waldner, A. Thimonier, E. Graf Pannatier and A. Marchetto: concept and data evaluations; K. Hansen and R. Fischer: idea; P. Rautio, A. Fürst: checking foliar data; M. Neumann, G. Kindermann, S. Etzold: checking growth data; P. Roskams: checking crown condition data; N. Cools: checking soil data; all authors: contributions to surveys, data checking and writing of the manuscript.

• **Context** Forests have been exposed to elevated atmospheric deposition of acidifying and eutrophying sulphur and nitrogen compounds for decades. Critical loads have been identified, below which damage due to acidification and eutrophication are not expected to occur.

• **Aims** We explored the relationship between the exceedance of critical loads and inorganic nitrogen concentration, the base cation to aluminium ratio in soil solutions, as well as the nutritional status of trees.

• **Methods** We used recent data describing deposition, elemental concentrations in soil solution and foliage, as well as the level of damage to foliage recorded at forest plots of the ICP Forests intensive monitoring network across Europe.

• **Results** Critical loads for inorganic nitrogen deposition were exceeded on about a third to half of the forest plots. Elevated

✉ Peter Waldner
peter.waldner@wsl.ch

Anne Thimonier
anne.thimonier@wsl.ch

Elisabeth Graf Pannatier
elisabeth.pannatier@wsl.ch

Sophia Etzold
sophia.etzold@wsl.ch

Maria Schmitt
maria.schmitt@wsl.ch

Aldo Marchetto
a.marchetto@ise.cnr.it

Pasi Rautio
pasi.rautio@luke.fi

Kirsti Derome
kirsti.derome@luke.fi

Tiina Maileena Nieminen
tiina.nieminen@luke.fi

Seppo Nevalainen
seppo.nevalainen@luke.fi

inorganic nitrogen concentrations in soil solution occurred more frequently among these plots. Indications of nutrient imbalances, such as low magnesium concentration in foliage or discolouration of needles and leaves, were seldom but appeared more frequently on plots where the critical limits for soil solution were exceeded.

• **Conclusion** The findings support the hypothesis that elevated nitrogen and sulphur deposition can lead to imbalances in tree nutrition.

Keywords Inorganic nitrogen concentration in soil solution · Base cation to aluminium ratio · Tree nutrition · Foliage · ICP Forests

1 Introduction

Forest ecosystems have been exposed to elevated atmospheric deposition of sulphur (S) and nitrogen (N), mainly as sulphate (SO_4^{2-}) and inorganic N, for more than five decades. The main reason being a large increase in the anthropogenic

emissions of N and S compounds in the second half of the last century. The elevated deposition of N and S affects forest ecosystems through several processes.

Inorganic N and SO_4^{2-} deposition may accelerate acidification of forest soils through leaching of strong acid anions, mainly nitrate (NO_3^-) and SO_4^{2-} , accompanied by base cations (BC) such as calcium (Ca^{2+}), magnesium (Mg^{2+}), and potassium (K^+) (Ulrich et al. 1980). Soil acidification may result in (i) depletion of soil BC and (ii) mobilisation of aluminium (Al^{3+}) into soil solution, possibly with adverse effects on fine roots and associated mycorrhizal fungi (e.g. de Wit et al. 2010). Toxic effects of dissolved Al are reduced by the presence of dissolved base cations. The molar ratio Bc/Al (molc molc⁻¹) in soil solution, where Bc is the sum of the molar concentrations of the base cations Ca^{2+} , Mg^{2+} and K^+ , and Al that of dissolved aluminium species in soil solution, has been suggested as a criterion to assess Al toxicity (Cronan and Grigal 1995).

Enhanced N supply may stimulate tree growth in N limited stands. In excess, however, N may induce (i) nutrient imbalances; (ii) increased sensitivity to frost, insects, and fungi; and (iii) elevated NO_3^- and ammonium (NH_4^+) leaching from the root zone (Aber et al. 1989). The concentration of inorganic N in soil

Antti-Jussi Lindroos
antti.lindroos@luke.fi

Päivi Merilä
paivi.merila@luke.fi

Georg Kindermann
georg.kindermann@bfiw.gv.at

Markus Neumann
markus.neumann@bfiw.gv.at

Nathalie Cools
nathalie.cools@inbo.be

Bruno de Vos
bruno.devos@inbo.be

Peter Roskams
peter.roskams@inbo.be

Arne Verstraeten
arne.verstraeten@inbo.be

Karin Hansen
karin.hansen@ivl.se

Gunilla Pihl Karlsson
gunilla.pihl.karlsson@ivl.se

Hans-Peter Dietrich
hans-peter.dietrich@lwf.bayern.de

Stephan Raspe
stephan.raspe@lwf.bayern.de

Richard Fischer
richard.fischer@ti.bund.de

Martin Lorenz
martin.lorenz@ti.bund.de

Susanne Iost
susanne.iost@ti.bund.de

Oliver Granke
granke@digisland.de

Tanja G. M. Sanders
tanja.sanders@ti.bund.de

Alexa Michel
alexa.michel@ti.bund.de

Hans-Dieter Nagel
hans.dieter.nagel@oekodata.com

Thomas Scheuschner
thomas.scheuschner@oekodata.com

Primož Simončič
primoz.simoncic@gozdis.si

Klaus von Wilpert
klaus.wilpert@forst.bwl.de

Henning Meesenburg
henning.meesenburg@nw-fva.de

Stefan Fleck
stefan.fleck@nw-fva.de

Sue Benham
sue.benham@forestry.gsi.gov.uk

Elena Vanguelova
elena.vanguelova@forestry.gsi.gov.uk

Nicholas Clarke
Nicholas.clarke@skogoglandskap.no

Morten Ingerslev
moi@ign.ku.dk

Lars Vesterdal
lv@ign.ku.dk

Per Gundersen
pgu@ign.ku.dk

solution ($N_{min} = NO_3^- + NH_4^+$, mg L⁻¹), NO₃⁻ leaching, the nutritional status of trees, as well as the organic carbon (C) to N ratio in the forest floor (C/N , kg kg⁻¹) have all been suggested as indicators of the N saturation status of forests (Gundersen et al. 2006; Dise et al. 2009).

In the frame of the UNECE Convention on Long-range Transboundary Air Pollution (CLRTAP), long-term effects of atmospheric deposition on ecosystems are generally assessed based on the concepts of critical loads and critical limits. Such critical loads and critical limits are defined as quantitative estimates of an exposure to deposition loads or levels below which significant harmful effects on specified sensitive elements of the environment do not occur according to current knowledge (Nilsson and Grennfelt 1988).

Two approaches are widely used for estimating critical loads for atmospheric N deposition. The first approach is to compile empirical observations to provide a range of typical critical loads for N deposition for each ecosystem type, e.g. 5 to 15 kg N ha⁻¹ year⁻¹ for coniferous woodland and 10 to 20 kg N ha⁻¹ year⁻¹ for broadleaved deciduous woodland (empirical critical loads for N deposition) (Bobbink and Hettelingh 2011). The second approach derives critical loads for N deposition from a criterion applied to nutrient fluxes or

levels in an ecosystem model and is generally implemented with a steady-state mass balance (SSMB) of input sources and output sinks (Sverdrup and de Vries 1994). Pools are excluded and assumed to be irrelevant under long-term considerations. A common criterion for calculating critical loads for N deposition using the SSMB is that the leaching flux of N below the root zone should not exceed an acceptable level (Spranger et al. 2004). This acceptable flux itself is often based on critical limits for N_{min} . Such N_{min} thresholds have been defined as a criterion for nutrient imbalances, elevated nitrate leaching or enhanced sensitivity to frost and fungal diseases (Table 1, Spranger et al. 2004; Iost et al. 2012).

A general threshold of $Bc/Al=1$ is a widely used criterion to derive SSMB critical loads for acidity, assuming that higher Bc/Al values will not damage tree roots (Spranger et al. 2004). In addition, species-specific threshold values (Bc/Al , Table 1) have been defined by Sverdrup and Warfvinge (1993). When the Bc/Al ratio falls below the thresholds, we denominate this case as exceedance of the critical limit for Al toxicity in the subsequent text.

These approaches enable the determination and mapping of critical loads for N deposition and critical loads for acidity and their exceedances for Europe (Reis et al. 2012). However, there may be a time lag between the start of exceedance of

Inge Stupak
ism@ign.ku.dk

Mathieu Jonard
mathieu.jonard@uclouvain.be

Nenad Potočić
nenadp@sumins.hr

Mayte Minaya
minaya@inia.es

¹ WSL, Swiss Federal Institute for Forest, Snow and Landscape Research, Zürcherstrasse 111, CH-8903 Birmensdorf, Switzerland

² CNR-Istituto per lo Studio degli Ecosistemi, Largo Tonolli 50, I-28922 Verbania Pallanza, VB, Italy

³ Luke, Natural Resources Institute Finland, P.O.Box 16, FI-96301 Rovaniemi, Finland

⁴ Luke, Natural Resources Institute Finland, P.O. Box 413, FI-90014 Oulu, Finland

⁵ Luke, Natural Resources Institute Finland, P.O.Box 18, FI-01301 Vantaa, Finland

⁶ Luke, Natural Resources Institute Finland, P.O.Box 68, FI-80101 Joensuu, Finland

⁷ BFW, Federal Research Centre for Forests, Seckendorff-Gudent-Weg 8, A-1131 Vienna, Austria

⁸ INBO, Research Institute for Nature and Forests, Kliniekstraat 25, BE-1070 Brussels, Belgium

⁹ IVL Swedish Environmental Research Institute, SE-100 31 Stockholm, Sweden

¹⁰ IVL Swedish Environmental Research Institute, SE-400 14 Göteborg, Sweden

¹¹ LWF, Bavarian State Institute of Forestry, Hans-Carl-von-Carlowitz-Platz 1, D-85354 Freising, Germany

¹² TI, Thünen-Institute, Leuschnerstrasse 91, D-21031 Hamburg, Germany

¹³ DigSyLand, Institute for digital System Analyses and Landscape Diagnosis, Zum Dorfteich 6, D-24975 Husby, Germany

¹⁴ TI, Thünen Institute of Forest Ecosystems, Alfred-Möller-Strasse 1, D-16225 Eberswalde, Germany

¹⁵ OEKO-DATA, Hegermühlenstraße 58, D-15344 Strausberg, Germany

¹⁶ GIS, Slovenian Forestry Institute, Vecna pot 2, SI-1000 Ljubljana, Slovenia

¹⁷ FVA-BW, Forest Research Institute of Baden-Wuerttemberg, Wonnhaldestraße 4, D-79100 Freiburg, Germany

¹⁸ NW-FVA, Northwest German Forest Research Station, Grätzelstr. 2, D-37079 Göttingen, Germany

¹⁹ Forest Research, Alice Holt Lodge, Wrecclesham, Farnham, Surrey GU10 4LH, UK

²⁰ Norwegian Forest and Landscape Institute, P.O. Box 115, N-1431 Ås, Norway

²¹ University of Copenhagen, Rolighedsvej 23, DK-1958 Frederiksberg C, Denmark

²² UCL-ELI, Université Catholique de Louvain, Croix du Sud 2, BE-1348 Louvain-la-Neuve, Belgium

²³ Croatian Forest Research Institute, Cvjetno naselje 41, HR-10450 Jastrebarsko, Croatia

²⁴ CIFOR-INIA, Center for Forest Investigations, Carretera Coruna km 7.5, ES-28040 Madrid, Spain

Table 1 Ranges of the ‘adequate to optimal’ (A/O) nutritional class for tree foliar concentrations (mg g^{-1}) of nitrogen (*N*), potassium (*K*) and magnesium (*Mg*) as compiled by Stefan et al. (1997), ratios of *N/Mg* and *N/K* according to Mellert and Göttlein (2012); species-specific critical limits of the molar ratio of base cations to total aluminium (*Bc/Al*)

and of the inorganic N concentration in the mineral top soil solution (*Nmin*, mg L^{-1}) regarding nutrient imbalances, and the critical limit for the mineral N concentration in the soil solution below the rooting zone (*Nmin*) regarding nitrogen saturation

Species group ^a	Foliage					Soil solution		
	<i>N</i> ^a	<i>K</i> ^a	<i>Mg</i> ^a	<i>N/Mg</i> ^b	<i>N/K</i> ^b	<i>Bc/Al</i> ^c	<i>Nmin</i> ^d	<i>Nmin</i> ^d
Spruce	12–17	3.5–9	0.6–1.5	10.7–21	1.7–3.3	1.2	0.2	1
Pine	12–17	3.5–10	0.6–1.5	10.8–22.9	2–4	1.2	0.2	1
Silver fir	12–17	3.5–9	0.6–1.5	10.7–22.9*	1.7–4*	1.2	0.2	1
Douglas fir	12–17	3.5–10	0.6–1.5	10.7–22.9*	1.7–4*	0.3	0.2	1
Other conifers	12–17	3.5–10	0.6–1.5	10.7–22.9*	1.7–4*	1.2	0.2	1
Beech	18–25	5–10	1–1.5	8.2–21.8	1.9–3.8	0.6	0.4	1
Birch	18–25	5–10	1–1.5	8.1–21.8*	1.7–3.8*	0.8	0.4	1
Oak	15–25	5–10	1–2.5	8.1–21.8	1.7–3.7	0.6	0.4	1
Other broadleaves	15–25	5–10	1–2.5	8.1–21.8*	1.7–3.8*	0.6	0.4	1

^a Forest Foliar Coordinating Centre of the Expert Panel on Foliage and Litterfall of ICP Forests (Stefan et al. 1997)

^b Mellert and Göttlein (2012)

^c Sverdrup and Warfvinge (1993) and Lorenz et al. (2008)

^d Iost et al. (2012)

*Adapted

critical loads and the start of the exceedance of the underlying critical limits, as well as between exceedance of critical limits and effects, e.g. on tree nutritional status.

Foliar analyses have been used to assess the nutritional status of trees based on the concentration ranges for nutrition classes compiled from various experiments and expert knowledge by the Expert Panel on Foliage and Litterfall of the International Cooperative Programme on Assessment and Monitoring of Air Pollution Effects on Forests (ICP Forests) (Stefan et al. 1997) or more recently by Mellert and Göttlein (2012). Severe nutrient deficiencies may cause visual symptoms differing by species and nutrient-specific discolouration patterns (ICP Forests 2010).

The aim of this study was to carry out an exploratory investigation of the effects of high S and N deposition on tree nutrition, using recent data from intensively monitored forest plots of the ICP Forests Level II plot network (Ferretti and Fischer 2013) and currently used critical loads concepts (Spranger et al. 2004).

- Our first hypothesis was that the exceedance of critical loads for acidity and nitrogen had lasted long enough to result in an exceedance of critical limits for Al toxicity and the inorganic N concentration in soil solution.
- Our second hypothesis was that the exceedance of critical limits affects the nutritional status of trees as assessed from nutrient contents in leaves and needles.

2 Material and methods

The study was based on measurements of atmospheric deposition, soil solution chemistry, foliar nutrition, soil chemistry and tree growth as well as on the assessment of crown condition, carried out in the period from 2006 to 2009 on 201 forest plots of the ICP Forests intensive monitoring (Level II) network (ICP Forests 2010) as well as 43 forest plots of the similar Swedish Throughfall Monitoring Network (SWETHRO: Pihl Karlsson et al. 2011).

Bulk deposition (*BD*) and throughfall deposition (*TF*) were continuously collected in an open field and below tree canopy, respectively, at weekly, bi-weekly or in a few cases monthly sampling intervals. The bulk deposition volume was used to derive the precipitation quantity (*P*). Annual *BD*, *TF* and *P* values were calculated for each plot and year as described in Waldner et al. (2014).

Soil solution was generally sampled with suction lysimeters, typically at the same time intervals as deposition. The *Nmin* concentration and the *Bc/Al* ratio were calculated for each sample using the NO_3^- , NH_4^+ , *Bc* and total Al concentrations. Annual mean concentrations as well as the proportion of samples (f_{ss}) exceeding critical limits (Table 1) were calculated for each plot, depth and year and aggregated for the top 40-cm layer of soil as specified by Iost et al. (2012). The same parameters were calculated for the depth of the deepest lysimeters per plot (from 10- to 250-cm depth).

Soil was sampled at fixed depths: 0 to 10, 10 to 20 and 20 to 40 cm, respectively, by taking about 24 samples, which were pooled to at least three composite samples per depth and analysed (dataset version AFSCDB.LII.2.1: Cools and de Vos 2010). We calculated fine earth content weighted averages for organic C/N ratio (C/N) and base saturation BS (%) between 0- and 40-cm depth as well as averages of the C/N ratios of the organic layer.

Tree density (*trees*, ha^{-1}) was calculated based on the growth survey typically carried out every 5 years (ICP Forests 2010).

Foliage from the upper third of the sun exposed tree canopy was sampled from at least five trees of the main tree species of each plot. In the case of deciduous species, fully developed leaves were sampled during the second half of the growing season and before the beginning of the autumn senescence. Evergreen foliage was sampled during the dormancy period. The nutritional status of the trees was assessed by comparing the mean foliage concentrations of N, magnesium (Mg) and potassium (K) to the adjacent ranges of three classes (Stefan et al. 1997) referred to as 'low/deficient' (L/D), 'adequate to optimal' (A-O) and 'high to surplus' (H/S) (c.f. Table 1). The foliar concentration ratios N/Mg and N/K were compared to classes of ratios established by Mellert and Göttlein (2012). The same abbreviations as for concentrations were used to refer to ratios within the adequate to optimal range (A-O), below the lower end (L/D) and above the upper end of the adequate range (H/S) (Table 1).

Crown condition and damage cause assessments of trees on the plot (typically adjacent to the foliage sample trees) include the description of the damage symptoms and the determination of the possible cause. The presence of observed symptoms was reported to various levels of detail, e.g. only symptom and cause class or Latin name of causing agent (insect, fungus etc.). When a specific symptom was reported for at least one tree in a given country for a given year, then we assumed that it would have been reported for any other tree that showed this symptom in that country and year. Consequently, trees with no mention of this specific symptom in that country and year were treated as trees showing no symptom. We focussed on the symptom 'light green to yellow discolouration' and calculated the proportion of trees per plot and year (f_y) that showed this symptom.

We used values of exceedance of the SSMB of critical loads for N deposition and for acidity that were calculated by Nagel et al. (2011), Waldner et al. (2007) and Marchetto et al. (2010) based on measurements on Level II plots.

As a simplified estimate for the exceedance of empirical critical loads for N deposition, we used the criterion that throughfall deposition of inorganic N (TFN , $\text{kg ha}^{-1} \text{ year}^{-1}$) exceeds $15 \text{ kg ha}^{-1} \text{ year}^{-1}$. The critical loads apply to total deposition, which is difficult to determine because of N uptake

in the canopy. Total deposition is thus typically a factor of 1 to 2 times higher than TFN .

2.1 Evaluations and statistical analyses

Not all variables were available for all plots and all years for the period 2006 to 2009. The available annual values for this period for throughfall, bulk deposition, P and f_{ss} were aggregated per plot. The values for f_{ss} , N , Mg , and K in foliage, and f_y were aggregated per plot and tree species group. We used the tree species groups defined by Stefan et al. (1997) and assigned *Fagus sylvatica* (L.) including *Fagus moesiaca* (MALÝ) to the group referred to hereafter as 'beech', *Abies alba* (MILL.) and *Abies borisii-regis* (MATTF.) to 'fir', *Quercus robur* (L.), *Quercus petraea* (LIEBL.) and *Quercus cerris* (L.) to 'oak', *Betula sp.* and *Fraxinus sp.* to 'other broad-leaves', *Pinus sp.* to 'pine', *Picea abies* (KARST.) to 'spruce' and *Pseudotsuga menziesii* (MIRB.) to 'other conifers'.

The relationship between deposition and soil solution was investigated based on these aggregates by comparing percentages of plots in three classes of the frequency of the exceedance of critical limits ($Bc/Al < 1$ and $N_{min} > 1 \text{ mg L}^{-1}$) in soil solution samples ($f_{SS} = 0$ %, $f_{SS} > 0$ % to 50 % and $f_{SS} > 50$ %) for plots grouped according to exceedance of critical loads. The relationship between soil solution quality and tree nutritional status was investigated by comparing percentages of plots in tree nutritional classes as well as the frequencies of symptoms for plots grouped according to their frequency class of exceedances of species-specific critical limits in soil solution (Table 1).

Linear regression models were applied on means from 2006 to 2009 (function 'lm' in R Development Core Team 2009). Foliar N and Mg concentrations were used as response variables, while inorganic N and Mg concentrations in soil solution, topsoil base saturation and C/N ratio, tree density, precipitation, longitude, latitude and altitude were used as predictors. Terms that were (clearly) not relevant at a significance level of 90 % ($p > 0.10$) were excluded from the full model. We compared models with and without inorganic N concentration in soil solution and N throughfall deposition. S deposition was not used.

The relationship between deposition and soil solution was investigated on 234 plots in total, SSMB critical loads for acidity and Bc/Al were available for 62 plots, SSMB critical loads for N deposition and N_{min} in soil solution for 71 plots, and throughfall N deposition and N_{min} in soil solution for 231 plots of which 109 had available values for C/N .

3 Results

Regarding acidification, the SSMB critical loads were exceeded at 11 out of 62 plots (17 %). Exceedance of the general critical limit for aluminium toxicity ($Bc/Al < 1$) in at least one soil solution sample ($f_{ss} > 0$ %) was reported from 5 of the 11 plots (45 %) where the SSMB critical load for acidity was exceeded and similarly from 27 of the 51 plots (53 %) where it was not exceeded (Fig. 1). However, the percentage of plots with exceedance of this critical limit in the majority of the soil solution samples ($f_{ss} > 50$ %) was higher among the plots where the SSMB critical load for acidity was exceeded (36 %) than among the plots where this critical load was not exceeded (14 %, Fig. 1).

Regarding eutrophication, the SSMB critical load for N deposition was exceeded at 37 out of the 71 plots for which values were available for both, SSMB critical load for N deposition and $Nmin$. The percentage of plots with exceedance of the threshold $Nmin = 1 \text{ mg L}^{-1}$ in at least one soil solution sample from the deepest lysimeters was higher (65 %) among the 37 plots where the SSMB critical load for N deposition was exceeded than in the plots where it was not (38 %, Fig. 1). A similar result was found for the much larger sample of 231 plots, when throughfall N deposition $> 15 \text{ kg N ha}^{-1} \text{ year}^{-1}$ was used as a proxy for the exceedance of empirical critical load for N deposition (Fig. 1).

Values of $C/N < 25$ in the organic layer of the soil were more common among the plots with throughfall N deposition $> 15 \text{ kg ha}^{-1} \text{ year}^{-1}$ (46 %), than among the plots with throughfall N deposition $< 15 \text{ kg ha}^{-1} \text{ year}^{-1}$ (36 %) (Fig. 1). Samples from the deepest lysimeters per plot with soil

solution $Nmin > 1 \text{ mg L}^{-1}$ were reported for 88 % of the 16 plots with throughfall N deposition $> 15 \text{ kg ha}^{-1} \text{ year}^{-1}$ and organic soil layer $C/N < 25$; for 65 % of the 46 plots with either throughfall N deposition $> 15 \text{ kg N ha}^{-1} \text{ year}^{-1}$ or organic layer $C/N < 25$; and 43 % of the 47 plots with neither throughfall N deposition $> 15 \text{ kg N ha}^{-1} \text{ year}^{-1}$ nor $C/N < 25$ (Fig. 1).

Conifers had foliage N concentration in the class L/D more often than broadleaved species. Higher N in foliage for spruce, pine, fir and oak as well as lower Mg in foliage for pine were observed on plots where the critical limits for $Nmin$ in soil solution were exceeded compared to plots without exceedance of these critical limits (Table 2). In conifers, Mg concentrations in the L/D class were recorded only on plots where critical limits for $Nmin$ in soil solution were exceeded. In beech, the percentage of plots with Mg in the L/D class was higher at plots with these critical limits exceeded compared to other plots. Similarly, the ratios N/Mg and N/K were more frequently in the H/S class in spruce and pine (Table 2, N/K not shown).

Distribution in foliar nutrition classes showed little difference between groups of plots built according to the exceedance of the species-specific critical limit for Al toxicity ($Bc/Al < \text{threshold}$) (Table 2). However, in spruce and fir, Mg concentrations in foliage were only in the L/D class on plots with exceedance of the species-specific critical limit for Al toxicity ($Bc/Al < 1.2$, c.f. Table 1). Potassium was in the L/D class of spruce at four plots which had critical limits exceeded, but this was seen for only two other plots including all tree species (not shown). The ratios N/Mg and N/K were more often in the less favourable H/S class for spruce, pine, fir (only N/Mg) and

Fig. 1 Number of plots with exceedance of the critical limits for Al toxicity ($Bc/Al < 1$) and N saturation ($Nmin > 1 \text{ mg L}^{-1}$) in none ($f_{ss} = 0$ %), a minority ($f_{ss} > 0$ to 50 %) and a majority ($f_{ss} > 50$ %) of the soil solution samples from lysimeters in 0- to 40-cm depth and the deepest lysimeters, respectively. Number of plots (displayed as bars) and percentages of plots (displayed as arc length of the pie graphs disks) were compiled for groups of plots with and without exceedance of SSMB critical loads for acidity, exceedance of SSMB critical loads for N deposition and throughfall N deposition exceeding $TFN = 15 \text{ kg ha}^{-1} \text{ year}^{-1}$ ($TFN > 15$) based on a mean of the available annual values of the period 2006 to 2009




Table 2 Percentages of plots in nutrition classes for groups of plots with species-specific critical limits for inorganic N (*Nmin*) and Al (*Bc/Al*<threshold) exceeded in none (0%), in a minority (>0%–50%) and in the majority (>50%) of the soil solution samples from the inorganic topsoil (0- to 40-cm depth)

Species Group ^a	<i>Nmin</i> exceedance	<i>N in foliage</i>			<i>Mg in foliage</i>			<i>N/Mg in foliage</i>			Plots n
		L/D	A-O	H/S	L/D	A-O	H/S	L/D	A-O	H/S	
Spruce	0 %	45	55	0	0	100	0	36	64	0	11
	>0 %–50 %	12	88	0	0	85	15	32	68	0	26
	>50 %	3	97	0	3	90	7	20	63	17	30
Pine	0 %	11	78	11	0	100	0	11	89	0	9
	>0 %–50 %	0	95	5	5	95	0	0	75	25	20
	>50 %	7	57	36	7	86	7	14	57	29	14
Fir	0 %	100	0	0	0	100	0	100	0	0	1
	>0 %–50 %	14	86	0	14	43	43	86	0	14	7
	>50 %	0	100	0	0	67	33	100	0	0	3
Beech	0 %	0	33	67	25	50	25	0	33	67	4
	>0 %–50 %	0	78	22	26	35	39	0	61	39	23
	>50 %	0	65	35	36	27	36	0	50	50	22
Oak	0 %	0	100	0	0	100	0	0	100	0	5
	>0 %–50 %	0	17	83	0	100	0	0	100	0	6
	>50 %	0	46	54	0	100	0	0	85	15	13
Species group ^a	<i>Bc/Al</i> < threshold	<i>N in foliage</i>			<i>Mg in foliage</i>			<i>N/Mg in foliage</i>			Plots n
		L/D	A-O	H/S	L/D	A-O	H/S	L/D	A-O	H/S	
Spruce	0 %	14	86	0	0	86	14	33	67	0	21
	>0 %–50 %	23	77	0	0	87	13	32	64	5	23
	>50 %	5	95	0	5	95	0	18	64	18	22
Pine	0 %	0	78	22	11	89	0	0	78	22	9
	>0 %–50 %	0	94	6	0	100	0	0	88	12	17
	>50 %	8	62	31	8	92	0	8	54	38	13
Fir	0 %	29	71	0	0	71	29	100	0	0	7
	>0 %–50 %	0	100	0	0	0	100	100	0	0	1
	>50 %	0	100	0	50	0	50	50	0	50	2
Beech	0 %	0	66	34	32	35	32	0	53	47	34
	>0 %–50 %	0	75	25	38	25	38	0	50	50	8
	>50 %	0	100	0	100	0	0	0	0	100	1
Oak	0 %	0	45	55	0	100	0	0	91	9	22
	>0 %–50 %	0	100	0	0	100	0	0	100	0	1
	>50 %	0	0	0	0	0	0	0	0	0	0

Legend: *Nmin*=exceedance of species-specific critical limits for *Nmin* in soil solution; *Bc/Al*<threshold=species-specific critical limit for Al toxicity for soil solution: 0 %=exceeded in any sample, >0–50 %=exceeded in a minority of samples, >50 %=exceeded in a majority of samples. Foliar concentration ranges: L/D=low/deficient, A-O=adequate to optimum, H/S=high/surplus; n=number of plots

^aSpecies grouped as suggested by Stefan et al. (1997)

beech on the plots where the species-specific critical limits for Al toxicity (*Bc/Al* below threshold) were exceeded compared with other plots (Table 2, N/K not shown).

Linear regression modelling based on the plot-wise aggregated dataset indicated that *Nmin* and *TFN* were important predictors of foliar N and Mg concentration (Table 3). For foliar N in spruce and foliar Mg in pine and fir, the explained variance increased when inorganic N concentration in soil solution (*Nmin*) and throughfall N deposition were included into the models (compare adjusted *R*² of *m*=1 with those of *m*=2 and 3).

Discolouration with light green to yellow foliage was more frequently reported for plots with the critical limit for *Nmin* in soil solution exceeded than for other plots, in particular for spruce (Fig. 2).

4 Discussion

On a European scale, this study explored and showed relations between the exceedance of critical loads and the exceedance of critical limits in soil solution, as well as tree nutritional

Table 3 Linear regression models m to explain foliar N and Mg concentrations without ($m=1$) and with soil solution inorganic N in inorganic topsoil $Nmin$ ($m=2$) and throughfall N deposition TFN ($m=3$) for each tree species group

Species group ^a	m	Estimates for intercept and regression coefficients for significant variables (at $p<0.1$)	sd	$sd\ resid$	$Adj\ R^2$	n
N concentration in foliage						
Spruce	1	13.4***+0.0002 z	1.56	1.56	-0.02	39
	2	14.0***+0.614 $Nmin$ ***+0.0001 z		1.24	0.34	
	3	13.5***+0.520 $Nmin$ **+0.040 TFN +0.00006 z		1.2	0.36	
Pine	1	11.9*-0.371 x ***+0.165 y -0.008 z **+0.119 BS *	2.41	1.31	0.65	27
	2	12.3*+0.447 $Nmin$ -0.299 x **+0.150 y -0.006 z +0.074 BS		1.23	0.67	
	3	6.2+0.110 $Nmin$ +0.203 TFN ** -0.152 x +0.167 y -0.006 z **+0.077 BS		1.04	0.76	
Fir	1	11.5-0.0106 x +0.0635 y +0.0002 z -0.0097 BS	0.64	0.27	0.45	7
	2	11.5-0.0040 $Nmin$ -0.0105 x +0.0648 y +0.0002 z -0.0099 BS		0.27	-0.09	
Beech	1	23.1***-0.137 x *+0.002 z *+0.003 $trees$	1.99	1.63	0.26	30
	2	23.1***-0.022 $Nmin$ -0.136 x *+0.002 z *+0.003 $trees$		1.63	0.23	
	3	22.9***-0.021 $Nmin$ +0.008 TFN -0.136 x +0.002 z *+0.003 $trees$		1.63	0.19	
Oak	1	30.3***-0.00003 z -0.03148 BS -0.00729 $trees$	3.49	2.54	0.29	13
	2	31.0***-0.3419 $Nmin$ +0.0008 z -0.0388 BS -0.0087 $trees$		2.5	0.23	
	3	28.6***-0.31952 $Nmin$ +0.14423 TFN +0.00003 z -0.02825 BS -0.00806 $trees$		2.42	0.18	
Mg concentration in foliage						
Spruce	1	0.965***-0.00007 P +0.01697 CN	0.23	0.21	0.09	39
	2	0.968***+0.00283 $Nmin$ -0.00007 P +0.01705 CN		0.21	0.06	
	3	0.967***-0.0017 $Nmin$ +0.00209 TFN -9 10^{-5} P +0.01642 CN		0.21	0.04	
Pines	1	2.06***-0.001 P ***-0.002 CN	0.31	0.23	0.4	27
	2	2.18***+0.0671 $Nmin$ -0.0017 P ***+0.0005 CN		0.21	0.48	
	3	2.08***+0.095 $Nmin$ * -0.012 TFN -0.0013 P ** -0.0002 CN		0.2	0.51	
Fir	1	5.02*-0.0007 P -0.1768 CN	0.57	0.35	0.41	7
	2	6.10*+0.166 $Nmin$ -0.001 P -0.195 CN *		0.24	0.63	
	3	5.68*+0.132 $Nmin$ +0.046 TFN -0.002 P -0.150 CN		0.16	0.77	
Beech	1	3.11*+0.1828 Mg_{ss} ***+0.0361 x ** -0.0476 y -0.0003 z +0.0034 BS	0.51	0.3	0.6	30
	2	1.21***+0.0673 $Nmin$ +0.1229 Mg_{ss} *+0.0050 BS *-0.0002 P		0.34	0.48	
	3	1.41***+0.0620 $Nmin$ -0.0193 TFN +0.1368 Mg_{ss} *+0.0047 BS *-0.0002 P		0.34	0.48	
Oak	1	1.08***+0.194 Mg_{ss} *+0.004 BS	0.33	0.23	0.41	13
	2	1.07***-0.008 $Nmin$ +0.200 Mg_{ss} *+0.004 BS		0.23	0.34	
	3	1.23** -0.005 $Nmin$ -0.010 TFN +0.194 Mg_{ss} +0.004 BS		0.23	0.29	

m model number, sd standard deviation of data ($mg\ g^{-1}$), $sd\ resid$ standard deviation of residuals ($mg\ g^{-1}$), $adj\ R^2$ adjusted R^2 , n number of plots, * $p<0.05$, ** $p<0.01$, *** $p<0.001$, x longitude, y latitude, z lower boundary of altitude class (m), P precipitation (mm), CN C/N ratio of the inorganic topsoil from 0- to 40-cm depth, BS base saturation (%) in the inorganic topsoil from 0- to 40-cm depth, Mg_{ss} Mg concentration in the soil solution from the inorganic topsoil from 0- to 40-cm depth, $trees$ tree density ($trees\ ha^{-1}$), $Nmin$ inorganic N concentration in soil solution ($mg\ L^{-1}$), TFN throughfall deposition of N ($kg\ ha^{-1}\ year^{-1}$)

^a Species grouped as suggested by Stefan et al. (1997)

status based on recent measurement data after several decades of high deposition loads.

In line with our first hypothesis, we observed a relationship between the exceedance of critical loads for N deposition and the frequency of elevated $Nmin$ concentrations in soil solution. About half of the plots with exceedance of SSMB critical loads for N deposition showed signs of N saturation in the period 2006 to 2009. Similar results were found when we used a $15\ kg\ ha^{-1}\ year^{-1}$ threshold applied to throughfall N deposition as a proxy for the empirical critical load for total N

deposition on a larger number of plots. Note that total deposition is generally higher than throughfall deposition because a fraction of the deposited N is directly taken up by the canopy. Furthermore, the empirical critical load for N deposition depends on forest type and ranges from 5 to $20\ kg\ ha^{-1}\ year^{-1}$. Hence, it is likely that the exceedance of the empirical critical load for N deposition is similar or even more frequent than the number of plots with throughfall N deposition $>15\ kg\ ha^{-1}\ year^{-1}$ may suggest. Assuming that the deepest lysimeters represent the depth of the rooting zone, it could be




Fig. 2 Percentage of trees per plot ($f_y > 50\%$) with reported foliage discolouration (light green to yellow) for the species groups spruce, pine, beech and oak for plots with and without exceedance of the critical limits for inorganic N in soil solution ($N_{min} < 1$) in the majority of samples ($f_{ss} > 50\%$) from the depth of the deepest lysimeters per plot

suggested that one third of plots with exceedance of N deposition critical loads show temporary indications and one third permanent indications of N saturation.

We found percentages of plots with insufficient or imbalanced nutrition status in the period 2006 to 2009 that were comparable to those obtained by Stefan et al. (1997) in a survey carried out in the mid 1990s. The generally lower N nutrition status of conifers compared to broadleaves may be due to the fact that conifers are more abundant at higher altitudes and latitudes, in regions with generally lower N deposition (Thimonier et al. 2010), higher rainfall and on poorer, more acidic soils.

A correlation between N deposition and N in foliage had already been found by de Vries et al. (2003) based on measurements of the period from 1994 to 1999. They reported that about 44–63, 33–71 and 26–38 % of the spatial variations in foliar N, Mg and K concentrations, respectively, were explained in a regression model that included stand age, soil type, altitude, precipitation, soil chemistry and deposition. In addition, we observed a tendency towards less favourable N/Mg and N/K ratios in foliage in high N deposition areas

in the more recent data. This tendency seemed to be more strongly related to N concentration in soil solution and N deposition than to the other factors considered. The tendency towards less favourable foliage nutrition at plots with high N concentration in soil solution is in line with our second hypothesis that the exceedance of critical limits affects the nutritional status of trees. It remains to be investigated whether the temporal trends of mineral nutrition in foliage determined by Jonard et al. (2015) could be explained by changes in N deposition or N concentrations in soil solution.

The higher proportion of soil solution samples with the general critical limit for Al toxicity exceeded ($Bc/Al < 1$) on plots with critical loads for acidity exceeded (Fig. 1) is in line with our first hypothesis. However, this result is based on relatively few plots with critical loads for acidity exceeded. Frequent ratios of $Bc/Al < 1$ suggest acidified soils according to Sverdrup and de Vries (1994) and were found on approximately one third of these plots. However, we did not differentiate between different Al species. Determination of the Al speciation carried out for some of the plots (Graf Pannatier et al. 2011) showed that the most important toxic form, Al^{3+} , was typically about 30 to 100 % of total dissolved Al, whereas Hansen et al. (2007) found Al^{3+} to be up to 82 to 95 % of total dissolved Al. In some cases, the Al^{3+} concentration might thus be lower than total Al and less harmful than the Bc/Al suggests (e.g. Lange et al. 2006).

The relationship between the Bc/Al ratio in soil solution and foliar nutritional status was not clear, and thus, it is difficult to draw conclusions for our second hypothesis regarding acidification based on the data. The Mg and K concentrations in conifer foliage tended to be lower on plots with species-specific critical limit for Al toxicity exceeded, indicating a possible depletion of base cations due to soil acidification. No such tendency was observed for broadleaves. Values of $Bc/Al < 1$ have been related to damage to fine roots but rarely related to mature forest nutritional problems, due to tree roots' ability to chelate, detoxify and prevent some of the Al from being taken up (Richter et al. 2013). Augustin et al. (2005) found stronger relationships between foliar nutrition, soil pH and base saturation than between foliar nutrition and exceedances of critical loads in an investigation of ICP Forests data from Germany, and explained this with the indirect and delayed nature of the effects. In line with this, the results of the regression analyses reported here suggested that the relationships between foliar nutrition and inorganic N concentration in soil solution and base saturation were stronger than those between foliar nutrition and throughfall N deposition.

5 Conclusions

This study showed that there were differences in the frequency of exceedance of critical limits for soil solution between

groups of plots built according to current exceedance of critical loads, with these exceedances probably having persisted for several decades. Similar differentiation was found for tree nutritional status for groups built according to exceedances of critical limits for soil solution. The findings support the hypothesis that eutrophying or acidifying effects of inorganic N and S deposition may lead to imbalances in tree nutrition.

Further insight might be gained with supplementary analyses, e.g. by calculating SSMB critical loads for all plots, by comparing temporal trends in the variables, by using non-linear models, and by further investigation of reported symptoms.

Acknowledgments We gratefully acknowledge the Programme Coordinating Centre of ICP Forests and all observers, technicians and scientists who performed sampling in the fields, analyses and data handling. The evaluation was mainly based on data that are part of the UNECE ICP Forests PCC Collaborative Database (see <http://icp-forests.net>). In particular, data from France (14 plots), Belgium (7 plots), the Netherlands (3 plots), Germany (77 plots), Italy (10 plots), the United Kingdom (8 plots), Ireland (2 plots), Denmark (6 plots), Greece (3 plots), Spain (3 plots), Austria (2 plots), Finland (17 plots), Switzerland (7 plots), Hungary (1 plot), Romania (4 plots), Poland (1 plot), the Slovak Republic (5 plots), Norway (8 plots), Lithuania (2 plots), the Czech Republic (11 plots), Estonia (5 plots), Slovenia (2 plots), Latvia (1 plot), Cyprus (2 plots) and Sweden (43 plots) were part of the analyses. In addition, throughfall data from the SWETHRO network were used for the 43 plots in Sweden. For soil, we used and acknowledge the aggregated forest soil condition database (AFSCDB.LII.2.1) compiled by the ICP Forests Forest Soil Coordinating Centre. The long-term collection of forest monitoring data was to a large extent funded by national research institutions and ministries, with support from governmental bodies, services and land owners. It was partially funded by the European Union under the Regulation (EC) No. 2152/2003 concerning monitoring of forests and environmental interactions in the Community (Forest Focus) and the project LIFE 07 ENV/D/000218 “Further Development and Implementation of an EU-level Forest Monitoring System (FutMon)”. We gratefully acknowledge the contribution of Matthias Dobbertin who passed away within the duration of this study.

Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

References

- Aber JD, Nadelhoffer KJ, Steudler P, Melillo JM (1989) Nitrogen saturation in northern forest ecosystems. *Bioscience* 39:378–386
- Augustin S, Bolte A, Holzhausen M, Wolff B (2005) Exceedance of critical loads of nitrogen and sulphur and its relation to forest conditions. *Eur J For Res* 124:289–300
- Bobbink R, Hettelingh J-P (2011) Review and revision of empirical critical loads and dose–response relationships, Coordination Centre for Effects, National Institute for Public Health and the Environment (RIVM), www.rivm.nl/cce
- Cools N, de Vos B (2010) Availability and evaluation of European forest soil monitoring data in the study on the effects of air pollution on forests. *iForest* 4:205–211
- Cronan CS, Grigal DF (1995) Use of calcium aluminum ratios as indicators of stress in forest ecosystems. *J Environ Qual* 24:209–266
- de Vries W, Vel E, Reinds GJ, Deelstra H, Klap JM, Leeters EEJM, Hendriks CMA, Kerkvoorden M, Landmann G, Herkendell J, Haussmann T, Erisman JW (2003) Intensive monitoring of forest ecosystems in Europe: 1. Objectives, set-up and evaluation strategy. *For Ecol Manag* 174:77–95
- de Wit HA, Eldhuset TD, Mulder J (2010) Dissolved Al reduces Mg uptake in Norway spruce forest: results from a long-term field manipulation experiment in Norway. *For Ecol Manag* 259:2072–2082
- Dise NB, Rothwell JJ, Gauci V, van der Salm C, de Vries W (2009) Predicting dissolved inorganic nitrogen leaching in European forests using two independent databases. *Sci Total Environ* 407:1798–1808
- Ferretti M, Fischer R (2013) Forest monitoring: Methods for terrestrial investigations in Europe with an overview of North America and Asia, vol 12, Developments in Environmental Science. Elsevier, Oxford, 507 p
- Graf Pannatier E, Thimonier A, Schmitt M, Walthert L, Waldner P (2011) A decade of monitoring at Swiss Long-term Forest Ecosystem Research (LWF) sites: can we observe trends in atmospheric acid deposition and in soil solution acidity? *Environ Monit Assess* 174: 3–30
- Gundersen P, Schmidt IK, Raulund-Rasmussen K (2006) Leaching of nitrate from temperate forests - effects of air pollution and forest management. *Environ Rev* 14:1–57
- Hansen K, Vesterdal L, Bastrup-Birk A, Bille-Hansen J (2007) Are indicators for critical load exceedance related to forest condition. *Water Air Soil Pollut* 183:293–308
- ICP Forests (2010) Manual on methods and criteria for harmonized sampling, assessment, monitoring and analysis of the effects of air pollution on forests. International Co-operative Programme on Assessment and Monitoring of Air Pollution Effects on Forests (ICP Forests). Convention on Long-Range Transboundary Air Pollution (LRTAP). UNECE, Hamburg, 578 p
- Iost S, Rautio P, Lindroos A-J (2012) Spatio-temporal trends in soil Solution Bc/Al and N in relation to critical limits in European forest soils. *Water Air Soil Pollut* 223:1467–1479
- Jonard N, Fürst A, Verstraeten A, Thimonier A, Timmermann V, Potočić N, Waldner P, Benham S, Hansen K, Merilä P, Ponette Q, de la Cruz AC, Roskams P, Nicolas M, Croisé L, Ingerslev M, Matteucci G, Decinti B, Bascietto M, Rautio P (2015) Tree mineral nutrition is deteriorating in Europe. *Glob Chang Biol* 21:418–430
- Lange H, Solberg S, Clarke N (2006) Aluminium dynamics in forest soil waters in Norway. *Sci Total Environ* 367:942–957
- Lorenz M, Nagel H-D, Granke O, Kraft P (2008) Critical loads and their exceedances at intensive forest monitoring sites in Europe. *Environ Pollut* 155:426–435
- Marchetto A, Arisci S, Tartari G, Matteucci G, de Cinti B, Fabbio G (2010) Nitrogen deposition and critical loads in the FutMon plots in Italy. CNR Institute for Ecosystem Studies, Verbania Pallanza, 25 p
- Mellert K, Göttlein A (2012) Comparison of new foliar nutrient thresholds derived from van den Burg’s literature compilation with established central European references. *Eur J For Res* 131:1461–1472
- Nagel H-D, Scheuschner T, Schlutow A, Granke O, Clarke N, Fischer R (2011) Exceedance of critical loads for acidity and nitrogen and scenarios for the future development of soil solution chemistry. In: R Fischer, M Lorenz (eds.), Forest Condition in Europe, 2011 Technical Report of ICP Forests and FutMon. Work Report of the Institute for World Forestry 2011/1, ICP Forests, Hamburg, p. 97–113

- Nilsson J, Grennfelt P (1988) Critical loads for sulphur and nitrogen. Workshop in Skokloster, Sweden, 19–24 March 1988, Nordic Council of Ministers, Copenhagen, 418 p
- Pihl Karlsson G, Akselsson C, Hellsten S, Karlsson PE (2011) Reduced European emissions of S and N – effects on air concentrations, deposition and soil water chemistry in Swedish forests. *Environ Pollut* 159:3571–3582
- R Development Core Team (2009) R: a Language and Environment for Statistical Computing. Reference Index Version 2.10.1. R Foundation for Statistical Computing. Vienna, Austria. ISBN 3-900051-07-0
- Reis S, Grennfelt P, Klimont Z, Amann M, ApSimon H, Hettelingh J-P, Holland M, LeGall A-C, Maas R, Posch M, Spranger T, Sutton MA, Williams M (2012) From acid rain to climate change. *Science* 338: 1153–1154
- Richter AK, Hajdas I, Frossard E, Brunner I (2013) Soil acidity affects fine root turnover of European beech. *Plant Biosyst* 147:50–59
- Spranger T, Smith R, Fowler D, Mills G, Posch M, Hall J, Schütze G, Hettelingh J-P, Slootweg J (2004) Modelling and mapping critical loads and levels and air pollution effects, risks and trends. International Co-operative Programme for Modelling and Mapping. Convention on Long-range Transboundary Air Pollution (LRTAP). UNECE, www.icpmapping.org, 236 p
- Stefan K, Fürst A, Hacker R, Bartels U (1997) Forest foliar condition in Europe - Results of large-scale foliar chemistry surveys (survey 1995 and data from previous years). EC - UNECE, Austrian Federal Forest Research Centre, Brussels, 207 p
- Sverdrup H, de Vries W (1994) Calculating critical loads for acidity with the simple mass balance method. *Water Air Soil Pollut* 72:143–162
- Sverdrup H, Warfvinge P (1993) The effect of soil acidification on the growth of trees, grasses and herbs as expressed by the (Ca+Mg+K) / Al ratio. Reports in Ecology and Environmental Engineering 1993: 2, Department of Chemical Engineering II, Lund University, Sweden
- Thimonier A, Graf Pannatier E, Schmitt M, Waldner P, Walthert L, Schleppe P, Dobberrin M, Kräuchi N (2010) Does exceeding the critical loads for nitrogen alter nitrate leaching, the nutrient status of trees and their crown condition at Swiss Long-term Forest Ecosystem Research (LWF) sites? *Eur J For Res* 129:443–461
- Ulrich B, Mayer R, Khanna PK (1980) Chemical changes due to acid precipitation in a loess-derived soil in central Europe. *Soil Sci* 130: 193–199
- Waldner P, Schaub M, Graf Pannatier E, Schmitt M, Thimonier A, Walthert L (2007) Atmospheric deposition and ozone levels in Swiss forests: are critical values exceeded? *Environ Monit Assess* 128:5–17
- Waldner P, Marchetto A, Thimonier A, Schmitt M, Rogora M, Granke O, Mues V, Hansen K, Pihl Karlsson G, Žilindra D, Clarke N, Verstraeten A, Lazdins A, Schimming C, Iacoban C, Lindroos A-J, Vanguelova E, Benham S, Meesenburg H, Nicolas M, Kowalska A, Apuhtin V, Napa U, Lachmanová Z, Kristoefel F, Bleeker A, Ingerslev M, Vesterdal L, Molina J, Fischer U, Seidling W, Jonard M, O’Dea P, Johnson J, Fischer R, Lorenz M (2014) Detection of temporal trends in atmospheric deposition of inorganic nitrogen and sulphate to forests in Europe. *Atmos Environ* 95:363–374