



HAL
open science

Jumplets

Roman Andreev

► **To cite this version:**

| Roman Andreev. Jumplets. 2016. hal-01338101

HAL Id: hal-01338101

<https://hal.science/hal-01338101>

Preprint submitted on 27 Jun 2016

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

JUMPLETS

ROMAN ANDREEV[†]

ABSTRACT. In this note we describe a locally supported Riesz basis consisting of “jumplets” for the orthogonal complement of continuous splines on an interval in the space of discontinuous ones.

1. INTRODUCTION

Let $T > 0$ and write $J := [0, T]$. For $N \geq 1$ let a finite partition of J be given by

$$(1) \quad \mathcal{T} = \{0 =: t_0 < t_1 < \dots < t_N < t_{N+1} := T\}.$$

Let \mathbb{P}_d denote the space of real-valued polynomials on \mathbb{R} of degree at most d . We write

$$(2) \quad F := \{f \in L_2(J) : f|_{(t_{n-1}, t_n)} \in \mathbb{P}_{p_n}\} \quad \text{and} \quad E := F \cap H^1(J)$$

for the space F of piecewise polynomials on the interval $[0, T]$ of given polynomial degrees $p_n \geq 0$ and for its subset $E \subset F$ of the continuous ones. Here and throughout, L_2 and H^1 refer to the usual Lebesgue and Sobolev spaces of real-valued functions on J .

In the next Section 2 we construct a locally supported Riesz basis for the L_2 orthogonal complement of E in F . Its dimension is obviously N , which is the number of interior nodes of \mathcal{T} . We refer to E as the continuous part of F , and to its L_2 -orthogonal complement in F as the discontinuous part of F . An application to adaptive approximation in L_2 is discussed in Section 3.

2. CONSTRUCTION OF THE BASIS

If $I \subset \mathbb{R}$ is an interval, we write $(\cdot, \cdot)_I$ for the $L_2(I)$ scalar product, and $\|\cdot\|_I$ for the norm. The euclidean norm is denoted by $|\cdot|$. For integer $d \geq 0$ let $\ell_d \in \mathbb{P}_d$ be the Legendre polynomial on the unit interval $[0, 1]$ characterized by $\ell_d(1) = \sqrt{2d+1}$ and $\int_0^1 \ell_d(t)q(t)dt = 0$ for any $q \in \mathbb{P}_{d-1}$. Then $\{\ell_k : 0 \leq k \leq d\}$ is an orthonormal basis for $\mathbb{P}_d \cap L_2([0, 1])$.

For integer $d \geq 0$ define the polynomial $R_d := \sum_{k=0}^d \ell_k(1)\ell_k$. Its key property is that $(R_d, q)_{[0,1]} = q(1)$ for any polynomial $q \in \mathbb{P}_d$, as can be verified by expanding q into the Legendre basis. Further, $\|R_d\|_{[0,1]} = 1 + d$. Together, these observations imply $R_d(1) = (R_d, R_d)_{[0,1]} = (1 + d)^2$. Using $\ell_k(0) = (-1)^k \ell_k(1)$ in the definition of R_d we also obtain $R_d(0) = (-1)^d(1 + d)$.

For each $n = 1, \dots, N + 1$, let $J_n := (t_{n-1}, t_n)$ be the n -th subinterval in the partition \mathcal{T} from (1). Write $\alpha_n := |t_n - t_{n-1}|^{-1}$ for the inverse of its length, and define the constants $\beta_n := \alpha_n(1 + p_n)^2$, where p_n is the polynomial degree from (2). Define the functions ψ_n^- and ψ_{n-1}^+ in F having the values

$$(3) \quad \psi_n^-(t) := \alpha_n R_{p_n}(\alpha_n(t - t_{n-1})) \quad \text{and} \quad \psi_{n-1}^+(t) := \alpha_n R_{p_n}(\alpha_n(t_n - t))$$

if $t \in J_n$, and zero else. Note that ψ_{n-1}^+ is the reflection of ψ_n^- about the vertical at the middle of the subinterval J_n . Since β_n is associated with both, ψ_{n-1}^+ and ψ_n^- , we also write $\beta_{n-1}^+ := \beta_n^- := \beta_n$. These functions are designed towards the following observation.

Lemma 2.1. $(\psi_n^\pm, f)_J = \lim_{\epsilon \rightarrow 0} f(t_n \pm |\epsilon|)$ for all $f \in F$ and $n = 1, \dots, N$.

Proof. If $f \in F$ then $q : t \mapsto f(t_{n-1} + s/\alpha_n)$ is a polynomial of degree p_n on the unit interval $[0, 1]$. Since ψ_n^- is supported on J_n , we employ the definition of ψ_n^- and the properties of R_d to obtain $(\psi_n^-, f)_J = (R_{p_n}, q)_{[0,1]} = q(1) = \lim_{\epsilon \rightarrow 0} f(t_n - |\epsilon|)$. The proof for ψ_n^+ is analogous. \square




FIGURE 1. The basis function ψ_2 from (4) for polynomial degrees $p = 1$ (left) / $p = 2$ (right).

The announced basis for the discontinuous part of F is now defined by

$$(4) \quad \psi_n := \frac{\psi_n^- - \psi_n^+}{\sqrt{\beta_n^- + \beta_n^+}}, \quad n = 1, \dots, N.$$

Proposition 2.2. *Assume that $p_n \geq p_{\min} \geq 1$ in (2). Then, $\Psi := \{\psi_n : n = 1, \dots, N\}$ is a Riesz basis for the L_2 -orthogonal complement of E in F . More precisely, for all vectors $c \in \mathbb{R}^N$,*

$$(5) \quad C_- |c|^2 \leq \|\Psi^T c\|_J^2 \leq C_+ |c|^2 \quad \text{with} \quad C_{\pm} = 1 \pm \frac{1}{1 + p_{\min}}.$$

Proof. Set $\psi_n^\circ := \psi_n^- - \psi_n^+$. Then $(\psi_n^\circ, e)_J = \lim_{\epsilon \rightarrow 0} e(t_n - |\epsilon|) - \lim_{\epsilon \rightarrow 0} e(t_n + |\epsilon|) = 0$ for any $e \in E$ by continuity of e . Hence, each ψ_n is L_2 -orthogonal to E . By definition, $\psi_n \in F$. Therefore, once (5) has been established, the fact that the L_2 -orthogonal complement of E in F is N -dimensional shows that Ψ is a basis for it. Now we compute

$$(6) \quad \gamma_{n,n-1}^\circ := (\psi_n^\circ, \psi_{n-1}^\circ)_J = -(\psi_n^-, \psi_{n-1}^+)_J = -\lim_{\epsilon \rightarrow 0} \psi_n^-(t_{n-1} + |\epsilon|) = \alpha_n (-1)^{1+p_n} (1 + p_n),$$

and similarly

$$(7) \quad \beta_n^\circ := (\psi_n^\circ, \psi_n^\circ)_J = \|\psi_n^-\|_J^2 + \|\psi_n^+\|_J^2 = \beta_n^- + \beta_n^+.$$

Set $\delta_n := \gamma_{n,n-1}^\circ / \sqrt{\beta_n^\circ \beta_{n-1}^\circ}$. Then the Gramian of Ψ is the tridiagonal symmetric $N \times N$ matrix

$$(8) \quad M_\Psi := \begin{pmatrix} 1 & \delta_2 & & & \\ \delta_2 & 1 & \delta_3 & & \\ & \delta_3 & 1 & \ddots & \\ & & \ddots & \ddots & \\ & & & & 1 \end{pmatrix}.$$

Note $c^T M_\Psi c = |c|^2 + \sum_{n=2}^N 2\delta_n c_n c_{n-1}$ for any $c \in \mathbb{R}^N$, so we wish to compare the magnitude of the second term to $|c|^2$. Observe

$$(9) \quad |2\delta_n c_n c_{n-1}| = |\gamma_{n,n-1}^\circ| \times 2 \frac{|c_n|}{\sqrt{\beta_n^\circ}} \frac{|c_{n-1}|}{\sqrt{\beta_{n-1}^\circ}} \leq \frac{\beta_n}{1 + p_{\min}} \times \left(\frac{c_n^2}{\beta_n^\circ} + \frac{c_{n-1}^2}{\beta_{n-1}^\circ} \right),$$

so that $\sum_{n=2}^N |2\delta_n c_n c_{n-1}| \leq \frac{1}{1+p_{\min}} |c|^2$. Therefore, $C_- |c|^2 \leq c^T M_\Psi c \leq C_+ |c|^2$, with $C_{\pm} = 1 \pm \frac{1}{1+p_{\min}}$. \square

On a uniform mesh \mathcal{T} and with uniform polynomial degree p one obtains $\delta_n = -\frac{1}{2} \frac{(-1)^p}{1+p}$ for each n in the Gramian (8). It is a tridiagonal symmetric Toeplitz matrix with eigenvalues $\{1 + \frac{1}{1+p} \cos \frac{k\pi}{N+1} : k = 1, \dots, N\}$. Letting $N \rightarrow \infty$ shows that the constants in (5) cannot be improved in general.

The fact that the Gramian (8) is well-conditioned (5), together with Lemma 2.1, allows for fast computation of the discontinuous part of a function $f \in F$. Let $\Delta \in \mathbb{R}^N$ denote the vector of jumps of f , whose components are $\Delta_n = \lim_{\epsilon \rightarrow 0} \{f(t_n - |\epsilon|) - f(t_n + |\epsilon|)\}$ for $n = 1, \dots, N$. Then Lemma




FIGURE 2. Left: L_2 error achieved by the adaptive approximation algorithm for $g : t \mapsto t^{-1/3}$ on $J = (0, 1)$ by piecewise constant functions, and the error of the L_2 best approximation of g on a uniform mesh, as a function of the mesh size. Right: g and its adaptive approximation g_4 for $\theta = 3/4$.

2.1 and definition (4) of ψ_n imply $b_n := (\psi_n, f)_J = \Delta_n / \sqrt{\beta_n^- + \beta_n^+}$. The coefficients $c \in \mathbb{R}^N$ of the discontinuous part of f with respect to the basis (4) satisfy the linear system $M_\Psi c = b$, which can be quickly solved approximatively by the conjugate gradient method. This may be cheaper than projecting onto E and then taking the difference because the dimension of E is significantly larger than N when the polynomial degrees p_n in (2) are large.

3. APPLICATION: ADAPTIVE APPROXIMATION

As an application of the above construction we describe an algorithm for the adaptive approximation of a given function $g \in L_2(J)$. Suppose \mathcal{T}_i , $i = 0, 1, 2, \dots$, is a sequence of meshes as in the introduction, which are nested, $\mathcal{T}_i \subset \mathcal{T}_{i+1}$. Let $F_i \subset Y$ be the space of piecewise affine splines on with respect to \mathcal{T}_i , and set $E_i := F_i \cap H^1(J)$, which is then the space of continuous piecewise affine splines. Note that $F_i = E_i' + E_i$, where the prime denotes the derivative. Let E_i^\perp denote the L_2 -orthogonal complement of E_i in F_i , and let $Q_i^\perp : L_2(J) \rightarrow E_i^\perp$ be the surjective L_2 -orthogonal projection.

Suppose $g_i \in E_i'$ is an approximation of g . Then we consider $g_i^\perp := Q_i^\perp g_i \in E_i^\perp$, and use the coefficients of g_i^\perp with respect to the Riesz basis (4) for E_i^\perp as error indicators for the marking of subintervals to be adaptively refined. The adaptive algorithm is as follows. Let \mathcal{T}_0 be given. Fix a threshold parameter $\theta \in (0, 1]$. For each $i = 0, 1, 2, \dots$, do: **1)** Compute the L_2 -orthogonal projection g_i of g onto E_i' , or an approximation thereof. **2)** Compute the projection $g_i^\perp := Q_i^\perp g_i$. Set $N_i = \#(\mathcal{T}_i \cap (0, T))$ for the number of interior nodes in \mathcal{T}_i , and $[N_i] := \{1, \dots, N_i\}$. Let $c \in \mathbb{R}^{N_i}$ be the vector of the coefficients of $g_i^\perp \in E_i^\perp$ with respect to the Riesz basis (4) for E_i^\perp . Select $M_i \subset [N_i]$ of minimal size such that $\sum_{n \in M_i} c_n^2 \geq \theta^2 |c|^2$. **3)** If $\mathcal{T}_i = \{0 = t_0 < t_1 < \dots < t_{N_i} < t_{N_i+1} = T\}$, let the new mesh \mathcal{T}_{i+1} contain \mathcal{T}_i , and the new nodes $\frac{1}{2}(t_{n-1} + t_n)$ and $\frac{1}{2}(t_n + t_{n+1})$ for all $n \in M_i$.

The Riesz basis property guarantees that the part of the indicator g_i^\perp corresponding to the marked subset M_i carries a fraction of its total $L_2(J)$ norm that is comparable to θ .

As an example we take $g : t \mapsto t^{-1/3}$ on $J := [0, T]$ with $T := 1$. We set $\mathcal{T}_0 := \{0, \frac{1}{4}, \frac{1}{2}, \frac{3}{4}, 1\}$ for the initial mesh. Instead of the L_2 -orthogonal projection we use the 4-point Gauss–Legendre quadrature rule on each subinterval to obtain g_i . The error $\|g - g_i\|_J$ for the above adaptive is shown in Figure 2. Due to the lack of smoothness of the given function g , it is approximated by piecewise constant functions on a uniform mesh with a rate of $\approx 1/6$ with respect to the mesh size $\#\mathcal{T}_i$, while adaptivity recovers the asymptotic rate of one.

This note was mainly written while at RICAM, Linz (AT), 2014.